

Irradiated asymmetric Friedmann branes

László A. Gergely and Zoltan Keresztes

Departments of Theoretical and Experimental Physics,
University of Szeged, 6720 Szeged, Dom ter 9, Hungary

(Dated: March 26, 2024)

Abstract

We consider a Friedmann brane moving in a bulk impregnated by radiation. The setup is strongly asymmetric, with only one black hole in the bulk. The radiation emitted by this bulk black hole can be reflected, absorbed or transmitted through the brane. Radiation pressure accelerates the brane, behaving as dark energy. Absorption however generates a competing effect: the brane becomes heavier and gravitational attraction increases. We analyse the model numerically, assuming a total absorption on the brane for $k = 1$. We conclude that due to the two competing effects, in this asymmetric scenario the Hawking radiation from the bulk black hole is not able to change the recollapsing fate of this brane-world universe. We show that for light branes and early times the radiation pressure is the dominant effect. In contrast, for heavy branes the self-gravity of the absorbed radiation is a much stronger effect. We find the critical value of the initial energy density for which these two effects roughly cancel each other.

Electronic address: gergely@physx.u-szeged.hu \ zkeresztes@titan.physx.u-szeged.hu

I. INTRODUCTION

According to brane-world models, our universe is a hypersurface (the brane) embedded in a 5-dimensional spacetime (the bulk), in which Einstein's gravity holds. The attempts to increase the number of dimensions in physics have a long history, originating with the pioneering works of Kaluza and Klein. In the more recent models presented in [1], the (at) extra dimensions are still compactified. The novelty of the brane-world models generalizing the original proposal [2] of Randall and Sundrum (RS) consist in allowing for one additional noncompact dimension. Instead of compactifying, the bulk is warped.

Gravitational dynamics on the brane is given by a modified Einstein equation, which arises from the projection of the 5-dimensional dynamics and the junction conditions across the brane. A major difficulty in this theory is that due to a new source term, arising from the projection of the bulk Weyl tensor to the brane, the system of differential equations describing gravitational evolution is not closed on the brane. This new gravitational theory allows general relativity as the low energy density limit (as compared to the brane tension).

Brane-world models were initially Z_2 -symmetric, with identical copies of the bulk on the two sides of the brane. Equivalently, the brane could be imagined as a domain boundary. This assumption, although it considerably simplifies the model, seems unnecessarily restrictive in the context of brane-world models with curved branes (generalized RS type 2 models). In this context, our observable universe may be imagined as a Friedmann brane moving either in a Schwarzschild-anti de Sitter bulk (if the bulk is vacuum), or in a 5-dimensional Vaidya-anti de Sitter (VAdS5) spacetime (if the bulk contains radiation). The modified Einstein equation in the Z_2 -symmetric case was given covariantly in [3], and generalized for asymmetric embedding in [4]. The effects of asymmetric embedding were also analyzed in a more generic class of models, containing induced gravity contributions [5]. There it was shown that the late time universe behaves differently under the introduction of a small asymmetry of the embedding: on the two branches of such theories. While on the RS branch asymmetry produces late-time acceleration, on the DGP branch the late-time acceleration is diminished by asymmetry.

Whenever there is radiation in the bulk, it can arise from both brane and bulk sources. The situation when the brane radiates into the bulk was already considered both in a

symmetric [6] and in an asymmetric scenario [7], [8] in the framework of generalized RS theories. In both scenarios a radial emission of radiation into the bulk was considered. A more realistic set-up, allowing for the emitted gravitons to follow geodesic paths, was advanced recently in [9].

In this paper we discuss the other possibility, when the radiation is emitted by sources in the bulk. For this, we embed the Friedmann brane asymmetrically into $VA dS_5$ spacetime. The symmetry of the embedding is severely broken by allowing for a radiating black hole on one side and no black hole on the other side of the brane. Due to this asymmetry, there is a radiation pressure acting on one side of the brane. This radiation pressure continuously accelerates the motion of the brane, leading to accelerated cosmic expansion. In principle, this mechanism can be a dark energy candidate.

In brane-world models, standard matter is confined to the brane. Therefore the radiation coming from the bulk should be of non-standard nature. We discuss here the case when this is the Hawking radiation of the single bulk black hole. The expression of the Hawking radiation in a 5-dimensional Schwarzschild-anti de Sitter ($SA dS_5$) spacetime was derived for the curvature index $k = 1$ in [10]–[12]. This energy density was employed, but for $k = 0$ in [13] for the study of asymmetrically embedded Friedmann branes into $VA dS_5$.

Some part of the radiation reaching the brane will be absorbed, some part will be reflected and the remaining part will go through. We will study here only the models without reflection in order to have the $VA dS_5$ spacetime in both bulk regions (Fig 1). We disregard the possibility of reflection, because there is no known exact solution with cosmological constant describing a cross flow of radiation streams. Such a spacetime with both incoming and outgoing null dust streams, but without a cosmological constant is known in 4-dimensions [14], however no generalization including a cosmological constant has been found. Thus we discuss the case when the reflected component can be neglected.

We keep however the absorption, as an essential element of our model, in contrast with the one presented in [13], where there is a black hole on each side of the brane, but their radiation is completely transmitted through the brane.

By retaining the absorption on the brane, we obtain novel features. The absorbed radiation appears on the brane as continuously emerging energy. Thus the energy density on the brane increases, strengthening the gravitational selfattraction of our universe. As consequence of absorption, cosmic expansion slows down.

Therefore there are two competing effects in our model: radiation pressure accelerates, while the absorbed radiation decelerates cosmic expansion. We would like to study how their balance affects the Friedmann brane-world. We derive the relevant equations of the model in Section II and we comment on the physical interpretation in Section III.

As our model emphasizes the effects of the absorption (as opposed to [13]), we finally choose a total absorption on the brane in Section IV. We also employ $k = 1$, as the formula for the energy density of the Hawking radiation was derived for this case [10]–[12]. Then we introduce new dimensionless variables, adapted to our choice of $k = 1$.

We provide quantitative results by numerical analysis in Section V. First we discuss cosmological evolution in the case of a non-radiating bulk black hole. Then, by switching on the radiation, we see how the two competing effects modulate the cosmological evolution. We show that a critical behaviour can appear, when these two competing effects roughly cancel each other. We discuss the relevance of these results in a broader context in Section VI.

Throughout the paper a tilde distinguishes the quantities defined on the 5-dimensional spacetime and a hat denotes dimensionless quantities and units with $c = 1 = \hbar$ are chosen.

II. FRIEDMANN BRANE IN VADYA-ANTIDESITTER BULK

The Friedmann brane representing our observable universe is embedded in the VAdS5 spacetime:

$$ds^2 = f(v; r; k) dv^2 - 2dvdr + r^2 d\Omega^2 + H^2(\tilde{r}; k) d\tilde{t}^2 + \sin^2 \tilde{t} d\Omega^2; \quad (1)$$

where

$$H(\tilde{r}; k) = \begin{cases} \tilde{r}^2 & ; \quad k = 1 \\ \tilde{r} & ; \quad k = 0 \\ \sinh \tilde{r} & ; \quad k = -1 \end{cases} \quad (2)$$

and

$$f(v; r; k) = k \frac{2m(v)}{r^2} - \frac{e^2 e}{6} r^2; \quad (3)$$

Here $e^2 = 8 G_{(5)}$ is the gravitational coupling constant in the bulk ($G_{(5)}$ representing the 5-dimensional gravitational constant) and it has the dimension of $(\text{length}^4 \text{mass}^{-1} \text{time}^2)$. The null coordinate v is ingoing (the $v = \text{const.}$ lines are outgoing). As a simplifying assumption and because we would like to focus on the radiation effects, we choose the same bulk cosmological constant ϵ on the two sides of the brane. The mass function $m(v)$ is freely specifiable, however different on the left and right sides of the brane: $m_L(v) \neq m_R(v)$. We note that, as defined here, the mass function $m(v)$ has the dimension of length^2 . This is, because it denotes $G_{(5)} M(v) = c^2$, with $M(v)$ the true mass. Since we have chosen $\hbar = 1 = c$, dimensionally speaking, length, time and mass¹ are the same. Thus technically the mass function $m(v)$ represents the cumulated mass of the bulk black hole and radiation multiplied by $G_{(5)}$ (of dimension mass³).

The source of the metric (1) is pure radiation considered in the geometrical optics limit (null dust), with energy-momentum tensor

$$T_{ab}^{ND} = (v;r) l_a l_b : \quad (4)$$

Here $(v;r)$ is the energy density and l is a (future-oriented) null 1-form :

$$l = \frac{dv}{\underline{v}} = n + u : \quad (5)$$

Here n is the dual 1-form of the unit normal to the brane and u the dual 1-form of the timelike (negative) unit vector. We define these vectors in what follows. We give the brane by the embedding relations $v = v(\)$ and $r = a(\)$, with ϵ the cosmological time. Then we choose the timelike unit vector u such that ϵ is adapted to u :

$$u = \frac{\partial}{\partial \epsilon} = \underline{v} \frac{\partial}{\partial v} + \underline{r} \frac{\partial}{\partial r} : \quad (6)$$

(A dot denotes derivatives with respect to ϵ). As consequence of $u^\alpha u_\alpha = -1$, we have on the brane

$$f \underline{v} = \underline{a} + \underline{a}^2 + f^{1=2} : \quad (7)$$

Here we have chosen the positive root in order to have $\underline{v}$ positive. Now we can write the 1-form ϵ as

$$u = \underline{r}^2 + f^{1=2} dv - \underline{v} dr = d\epsilon : \quad (8)$$




FIG .1: The radiation from the left bulk black hole is partially absorbed on the brane and partially transmitted through into the other bulk region containing no black hole.

The unit normal 1-form to the brane can be expressed as

$$n = \underline{r}dv + \underline{v}dr : \quad (9)$$

The bulk Einstein equation for the metric (1) and the source term

$$e_{\alpha\beta} = e_{g_{ab}} + T_{ab}^{N D} ; \quad (10)$$

establishes a connection between the energy density and the time derivative of the mass function $\underline{m} = (dm/dv) \underline{v}$ as

$$= \frac{3\underline{m} \underline{v}}{e^2 a^3} ; \quad (11)$$

where $\underline{v}$ is given by Eq. (7).

The square root from $\underline{v}$ can be eliminated by using the relations derived in [4]:

$$\begin{aligned} \underline{a}^2 + f_{R,L}^{1=2} &= \overline{B} + \frac{B}{2} ; \\ \overline{B} &= \frac{e^2 a}{6} (+) ; \\ B &= \frac{6 m}{e^2 a^3 (+)} : \end{aligned} \quad (12)$$

(In the above and the forthcoming formulae the first subscript refers to the upper sign and always is understood as the difference taken between the R and L regions. An overbar

denotes the average of a physical quantity, taken on the two sides of the brane.) Therefore Eqs. (7), (11) and (12) imply

$$\rho_{R,L} = \frac{3\bar{m}_{R,L}}{e^2 a^3 f_{R,L}} = \frac{a}{6} \left(\frac{e^2 a}{6} \left(\frac{1}{a} + p \right) + \frac{3\bar{m}}{e^2 a^3 \left(\frac{1}{a} + p \right)} \right) ; \quad (13)$$

Why do we need such a detailed expression for $\rho_{R,L}$? The answer is that radiation energy density appears in two of the key equations governing cosmological evolution. (These equations emerge from the modified Einstein equation.) The energy densities $\rho_{R,L}$ are present both in the energy-balance equation

$$-\frac{\dot{a}}{a} + 3\frac{\dot{a}}{a} \left(\frac{1}{a} + p \right) = -\ddot{a} ; \quad (14)$$

and in the Raychaudhuri equation:

$$\begin{aligned} \frac{\dot{a}}{a} = \frac{\dot{a}_0}{3} - \frac{\ddot{a}}{6} \left(1 + \frac{2}{a} \right) + 3p \left(1 + \frac{2}{a} \right) - \\ - \frac{2\bar{m}}{a^4} + \frac{27(p - \frac{1}{a}) \left(\frac{1}{a} \right)^2}{e^4 a^8 \left(\frac{1}{a} + p \right)^3} \\ - \frac{e^2 \bar{m}}{3} - \frac{3\bar{m}}{e^2 a^4 \left(\frac{1}{a} + p \right)^2} ; \end{aligned} \quad (15)$$

However, the third essential equation, the Friedmann equation is independent of $\ddot{a}$:

$$\begin{aligned} \frac{\dot{a}^2 + k}{a^2} = \frac{\dot{a}_0^2}{3} + \frac{\ddot{a}}{3} \left(1 + \frac{2}{a} \right) \\ + \frac{2\bar{m}}{a^4} + \frac{9 \left(\frac{1}{a} \right)^2}{e^4 a^8 \left(\frac{1}{a} + p \right)^2} ; \end{aligned} \quad (16)$$

(The above equations are particular cases of the equations derived in [4] in a more generic set-up.) In the above two equations $\dot{a}_0$ denotes the 4-dimensional cosmological constant given in terms of the brane tension σ and bulk cosmological constant Λ as:

$$2\dot{a}_0^2 = -\Lambda + e^2 \sigma ; \quad (17)$$

The system of equations (13)–(16) governs the motion of the brane, seen from the brane point of view as cosmological evolution.

III. THE FREE FUNCTIONS OF THE MODEL

There are a number of free functions in our model. Among them, by straightforward algebra on Eq. (13) the quantities $\rho_{R,L}$ and $\bar{m}$ can be expressed in terms of $\bar{m}$, m , $\bar{m} = \bar{m}$

and $(m) = \underline{m}$ as

$$= \frac{3}{2a^3} \underline{a} - \frac{3m}{e^2 a^3 (+)} \underline{2m} \underline{F} + \underline{m} \underline{F} + \frac{1}{4a^2} (+) \underline{2m} \underline{F} + \underline{m} \underline{F} ; \quad (18)$$

$$= \frac{3}{4e^2 a^3} \underline{a} - \frac{3m}{e^2 a^3 (+)} \underline{2m} \underline{F} + \underline{m} \underline{F} + \frac{1}{8a^2} (+) \underline{2m} \underline{F} + \underline{m} \underline{F} ; \quad (19)$$

where we have introduced the shorthand notations

$$\underline{F} = \frac{1}{f_R} - \frac{1}{f_L} = \frac{2a^2 m}{a^2 \bar{f}^2 (m)^2} ; \quad (20)$$

$$\underline{F}_+ = \frac{1}{f_R} + \frac{1}{f_L} = \frac{2a^4 \bar{f}}{a^2 \bar{f}^2 (m)^2} ; \quad (21)$$

$$a^2 \bar{f} = \frac{2}{6} - \frac{0}{3} a^4 + ka^2 \underline{2m} : \quad (22)$$

Then in Eqs. (14)–(16) the role of the free functions $\bar{m}$ and $\bar{m}_+$ is taken by $\underline{m}$ and $\underline{m}_+$.

In what follows, we discuss the physical interpretation of these free functions. First we show that m is determined by three quantities. These are the energy density of the brane, the scale factor and Hubble parameter. Second, that $\underline{m}_+$ is given by the energy emission of the bulk black-hole, evaluated at the left side of the brane. Finally we interpret $\underline{m}$ as the energy absorption on the brane.

A . The energy content of the brane

By rearranging the Friedmann equation, we find m as function of the cosmological perfect fluid energy density ρ , scale factor a and Hubble parameter $H = \dot{a}/a$:

$$(m)^2 = \frac{2}{3} a^8 (+) H^2 + \frac{\bar{f}}{a^2} - \frac{2}{6} (+)^2 : \quad (23)$$

In the derivation we have employed the relation among the brane tension and the gravitational coupling constants in 5 and 4-dimensions:

$$\frac{2}{3} = \frac{e^4}{6} : \quad (24)$$

It is easy to argue, that $m_R > m_L$ as m_R contains in addition to m_L the contribution from the brane energy momentum. Therefore $m > 0$.

We stress here that m is not strictly the mass of the brane. Rather, it represents the difference of the mass functions of the VAdS5 metric, evaluated on the two sides of the brane. Although, dimensionally speaking, m is mass^2 , it is directly proportional to the difference in the cumulated masses $M(v)$ of the bulk black hole and radiation on the two sides of the brane. The reason why $m_R \neq m_L$ is that the brane contributes toward the mass function both through its energy momentum and its curvature. In a less strict sense then m can be regarded as a measure of the energy content of the brane.

B. Hawking radiation in the bulk

The rate of decrease in mass of an evaporating black hole in an SAdS5 spacetime with $k = 1$ was computed in [10]–[12]. In [13] the assumption was advanced that this agrees with the rate of decrease of the mass function $m_L(v)$ in the VAdS5 spacetime, even for arbitrary k . Numerical results were given in [13] for $k = 0$. We adopt here the same identification, but only for $k = 1$, for which the proof of [10]–[12] holds. Then the energy density of the radiation escaping from the black hole, evaluated at the left side of the brane is:

$$\rho_L = \frac{h^{5(5)}}{24^9 a^3 m_L} \frac{1}{\underline{a} + (\underline{a}^2 + f_L)^{1/2}} ; \quad (25)$$

where ζ is the Riemann-zeta function. Here m_L contains both the mass of the black hole and the energy of the Hawking radiation. This radiation spreads away with the velocity of light and it overtakes the brane, which has a subluminal motion, therefore $\dot{m}_{R,L} < 0$. Moreover, as part of the energy radiated away is captured on the brane, there is less radiation escaping in the R region than there would be in the absence of the brane. Therefore $\dot{m}_L < \dot{m}_R < 0$ and in consequence $\dot{m} > 0$. For the same reason $\dot{m} < 0$ holds.

C. Energy absorption on the brane

The decomposition of the pure radiation energy-momentum tensor (4) with respect to the brane can be done by employing Eq. (5). We obtain:

$$T_{R\mu L}^{ND} = \rho_{R\mu L} u_a u_b + 2u_{(a} n_{b)} + n_a n_b ; \quad (26)$$

On the L region the radiation escaping the bulk black hole is characterized by ρ_L . Some of this radiation will pass unaffected through the brane into the R region (ρ_R). Then we have

to interpret the difference of the transmitted and incident energy-momentum tensors:

$$\begin{aligned} T_{ab}^{ND} &= u_a u_b + 2u_{(a} n_{b)} + n_a n_b \\ &= {}^{\text{rad}}u_a u_b + 2q_a^{\text{rad}} u_b + p^{\text{rad}} n_a n_b \quad (y) : \end{aligned} \quad (27)$$

Therefore the difference in the energy density of the radiation arriving from left and escaping into the right regions appears on the brane as a dust with energy density ${}^{\text{rad}}\rho > 0$. The energy flux arriving from the bulk into the brane is $q_a^{\text{rad}} n_a$. Finally, the radiation pressure $p^{\text{rad}} > 0$ drives the brane into an accelerated motion. This in turn accelerates the cosmic expansion.

With $\epsilon < 0$, the acceleration caused by the radiation pressure is manifest from the last term of the Raychaudhuri equation (15).

IV. TOTAL ABSORPTION ON THE BRANE

In this section we consider a simple model, in which $k = 1$ and $\epsilon_0 = 0$. We allow the brane to absorb all of the radiation escaping from the bulk black hole. With this choice the equations simplify considerably and we can study the effects of the absorption, which were not taken into account in previous treatments.

A. Brane dynamics

For total absorption $\epsilon_R = 0$ and m_R becomes a constant. Two free functions are then left, which can be chosen as $m = m$ and $\epsilon = \epsilon_L$. The energy-balance and Raychaudhuri equations (14) and (15) become

$$0 = -\frac{a}{3} + 3\frac{a}{a} (\epsilon + p) ; \quad (28)$$

$$\begin{aligned} \frac{a}{a} &= \frac{2}{6} \left(1 + \frac{2}{\epsilon} + 3p \right) + \frac{2m_R}{a^4} + \frac{m}{a^4} + \frac{27(p - \epsilon)(m)^2}{e^4 a^8 (\epsilon + p)^3} \\ &\quad + \frac{e^2}{6} + \frac{3m}{e^2 a^4 (\epsilon + p)^2} : \end{aligned} \quad (29)$$

We remark that with the assumption $\epsilon_R = 0$ both ϵ and m become determined. The Friedmann equation (16) emerges now as a consequence of the energy-balance equation

(28) and the Raychaudhuri equation (29). Stated otherwise, only two of the Friedmann, Raychaudhuri and energy-balance equations are now independent. (The interdependence of these three equations is a generic feature of the standard cosmological models, but not of brane-world universes. Indeed, prior to assuming the total absorption, the three equations were independent.)

From Eqs. (7), (11) and (25) the variation in time of the brane mass function is found:

$$\dot{m} = \frac{5}{72} \frac{(5) e^2}{(m_R - m)} : \quad (30)$$

This equation is an ordinary differential equation which can be integrated for $t > t_0$. The result is:

$$m = m_R - \frac{r}{(m_R - m_0)^2} \frac{5}{36} \frac{(5) e^2}{9} (t - t_0) ; \quad (31)$$

where by m_0 we have denoted the brane mass function before the Hawking radiation reaches the brane (this happens at $t = t_0$): The brane mass function m is monotonically increasing in time until $t_c = t_0 + 36 \frac{9}{5} (m_R - m_0)^2 = 5 \frac{(5) e^2}{9}$, when it reaches its maximal value $m = m_R$: When this happens, the bulk black hole has already completely evaporated and the resulting Hawking radiation is entirely absorbed by the brane.

The square root in Eq. (25) can be eliminated by employing Eq. (12). We obtain

$$= \frac{h}{24} \frac{5}{9 a^3 (m_R - m)} \frac{(5) e^2}{a + \frac{e^2 a}{6} \left(\frac{1}{a} + \frac{3m}{e^2 a^3 \left(\frac{1}{a} + \frac{3m}{e^2 a^3} \right)} \right)} : \quad (32)$$

With this both free functions m and a are given explicitly as functions of time. This has been achieved by specifying both the rate of absorption and the energy density of the black hole radiation Eq. (25).

B. Dimensionless variables

We introduce dimensionless variables in a slightly different way than in [6] and [7]. The reason for this is that the chosen comoving radial coordinate has to be dimensionless in the $k = 1$ cases. Therefore the scale factor has to carry the dimension, unless in the case $k = 0$, discussed in [6] and [7]. The dimensionless variables introduced below are well suited for $k = 1$ but they also apply for $k = 0$ if in this latter case the scale factor, rather than

the x -coordinate is regarded dimensional. These variables are:

$$\begin{aligned} t &= Ct; a = Ca; H = \frac{H}{C}; \\ b &= -; p = \frac{p}{C}; \rho = \frac{\rho}{C}; \\ \mu &= C^2 m; \mu_R = C^2 m_R; \end{aligned} \quad (33)$$

Here $C = \frac{p}{m}$ and its inverse represents a distance scale. (The dimensions of p and m are length/mass and mass/length³, respectively.) We note here that the dimension of ρ given by Eqs. (11) or (25) is consistent only by choosing $\rho = 1 = c$, implying that the dimension of mass is length³. Then C can be thought as a mass scale as well. Alternatively, by the choice $c = 1$ (therefore length and time having the same dimension), C becomes a basic energy scale:

Written in the variables (33), the system of equations describing cosmological evolution contains the following dimensionless parameter: $e^{8/3} = 36^{-4} = 2304^{-2} = M_p^4$, where M_p denotes the 4-dimensional Planck mass. By denoting the brane tension as $\mu = M_T^4$, the free parameter can be chosen as $M_T = M_p$. However for later notational convenience we introduce the dimensionless parameter $\mu = 40^{-5} (M_T = M_p)^4 = (3^{-7})$. The energy-balance and Raychaudhuri equations (14) and (15), written in terms of the dimensionless variables (33) and parameter μ read

$$0 = \dot{b}^0 + 3H(\dot{b} + p)\dot{b}; \quad (34)$$

$$\begin{aligned} \dot{H}^0 &= H^2 - \frac{1}{6}b(1+2b) - \frac{1}{2}p(1+b) \\ &\quad - \frac{2\mu_R}{b^4} + \frac{\mu}{b^4} + \frac{9(p-1)\mu^2}{2b^8(b+1)^3} \\ &\quad - \frac{b}{6} + \frac{3\mu b}{6b^4(b+1)^2}; \end{aligned} \quad (35)$$

with

$$b = \frac{h}{b^4(\mu_R - \mu)H + \frac{b+1}{6} + \frac{3\mu}{6b^4(b+1)}} \quad (36)$$

and

$$\dot{\mu}^0 = \frac{2}{6(\mu_R - \mu)}; \quad (37)$$

Here a prime denotes derivatives with respect to the dimensionless time $\mathfrak{t}$. Eq. (37) integrated gives the temporal evolution of the dimensionless mass function of the brane

$$\mathfrak{m} = \mathfrak{m}_R + \frac{(\mathfrak{m}_R - \mathfrak{m}_0)^2}{\mathfrak{p} - \frac{4}{6}} \mathfrak{t} - \mathfrak{t}_0 : \quad (38)$$

The generalized Friedmann equation (16) in dimensionless variables readily follows:

$$\mathfrak{H}^2 = \frac{1}{\mathfrak{b}^2} + \frac{\mathfrak{b}}{3} \left(1 + \frac{\mathfrak{b}}{2} + \frac{2\mathfrak{m}_R}{\mathfrak{b}^4} \right) + \frac{\mathfrak{m}}{\mathfrak{b}^4} + \frac{3(\mathfrak{m})^2}{2\mathfrak{b}^8 (\mathfrak{b} + 1)^2} : \quad (39)$$

V. NUMERICAL RESULTS

In this section we give numerical results on the evolution of the brane-world universe in the presence of one radiating bulk black hole. We assume the movement of the brane starts on the apparent horizon

$$\mathfrak{b}_{AH} = \frac{\mathfrak{q}}{3 + \frac{\mathfrak{p}}{9 + 12\mathfrak{m}_L}} \quad (40)$$

of the bulk black hole (therefore $\mathfrak{t}_0 = 0$). This horizon is dynamical, as $\mathfrak{m}_L$ decreases due to the escaping Hawking radiation. The brane leaves the horizon at $\mathfrak{m}_{L0} = \mathfrak{m}_R - \mathfrak{m}_0$. In order to have clear graphical solutions, we have found convenient to chose the dimensionless mass values of $\mathfrak{m}_R = 0.3$ and $\mathfrak{m}_0 = 0.29$. Then the initial position of the apparent horizon is at 0.1411870178. Away from the horizon, the brane absorbs the Hawking radiation. We investigate a radiation dominated universe, so $\mathfrak{p} = \mathfrak{b} = 3$. We choose the free parameter as $M_T = M_P = 1 = 10$. In what follows, we compare the evolutions of the brane in the absence and in the presence of the radiation coming from the bulk black hole.

A. The evolution of the brane-world without bulk black hole radiation

In the simplest case when we switch off the radiation from the bulk black hole, beyond the standard density and curvature source terms in the Friedmann equation (39), the model includes dark radiation and a strong asymmetry due to the chosen setup of a single left black hole in the bulk. We would like to see the evolution of such a brane universe, for later




FIG . 2: The evolution of the scalefactor in the absence of the Hawking radiation shows the characteristic pattern of a closed universe. The plot is for the initial brane density $b_0 = 520$.




FIG . 3: Evolution in the absence of the Hawking radiation of the Hubble parameter (dotted line) and brane energy density (solid line). The initial brane density is $b_0 = 520$.

comparison with the radiating case. With the Hawking radiation from the bulk black hole switched on, $\rho_b = \text{const.}$

Our numerical analysis shows that none of the brane effects (quadratic energy density, dark radiation and asymmetry source terms) will change significantly the usual behaviour of a closed ($k = 1$) universe, as can be seen from the behaviour of the scale-factor, represented in Fig 2. The Hubble parameter and the brane energy density evolve accordingly. The Hubble parameter continuously decreases, while the energy density tends to infinity both at




FIG . 4: Evolution of the dark radiation (DRST, dotted line) and asymmetry (AST, solid line) source terms of the Friedmann equation (for the initial brane density $\rho_0 = 520$).

the beginning and at the end of this brane-world universe. These behaviours are shown in Fig 3.

The evolutions of the dark radiation source term

$$\frac{2\dot{b}_R}{b^4} = \frac{\dot{b}}{b^4} ; \quad (41)$$

and of the asymmetry source term

$$\frac{3(\dot{b})^2}{2b^8(b+1)^2} \quad (42)$$

are represented in Fig 4. Both contributions are positive, so they act in a similar way to ordinary matter sources in the Friedmann equation. The dark radiation dominates over the asymmetry during the whole evolution, the difference being less when the universe reaches its maximum size. As the initial value of the brane energy density is lowered, these two source terms become comparable (Fig 5). For even lower initial energy densities the asymmetry source term dominates at both early and late times (Fig 6).

B . The energy density of the Hawking radiation

The energy density ρ^b (evaluated near the brane, on the left region) of the Hawking radiation from the bulk black hole evolves; cf Fig 7. In the expanding era ρ^b shows a sharp decrease. This is due to the fact that the distance between the brane and the horizon




FIG . 5: Similar evolutions of the dark radiation (dotted line) and asymmetry (solid line) source terms in the Friedmann equation at $b_0 = 250$.




FIG . 6: For light branes ($b_0 = 100$) the asymmetry (solid line) source term dominates over the dark radiation (dotted line) source term in the Friedmann equation for both early and late universes.

increases. Interestingly b starts to increase before the maximal size of the closed universe is reached. Later on b keeps increasing as the universe contracts. The higher values of b towards the end of contraction mathematically can be explained by the presence of the forever decreasing factor H in the denominator of b (see Eq. (36)). Stated otherwise, from Eq. (11) $\sim \sqrt{t}$ so that from a physical point of view, the increase in b at late times, exceeding its value at early times can be explained as a Doppler effect. The contracting brane absorbs more energy in unit time than the expanding brane at the same scale factor value.




FIG . 7: The evolution of the energy density of Hawking radiation for $b_0 = 520$.

We have checked, that with the Hawking radiation switched on, the basic behaviour of the closed universe will be very similar to that in the case without this radiation, discussed in the previous subsection. Even in the presence of the radiation the cosmological evolution shows the same patterns. Although there are modifications, which will be discussed in detail in the following subsection, they behave as small perturbations during the whole evolution of this asymmetric brane-world universe.

C . The evolution of the brane-world with the bulk black hole radiation

The evolution of the Hubble parameter, the scalefactor, the brane energy density and different source terms of the Friedmann equation, in the presence of the radiation from the bulk black hole is very similar to their evolutions in the absence of the radiation. This is due to a delicate balance between two effects. First, the brane is pushed away by the radiation pressure, which tends to accelerate it. Second, as all of the radiation is absorbed on the brane, the self-gravity of the brane increases, which tends to decelerate it. Still, there are interesting, through small effects due to the radiation. We illustrate them by representing the differences in the scalefactor, the brane energy density and different source terms of the Friedmann equation in the radiating and non-radiating cases. More specifically, we denote by A the quantity A in the radiating case and we plot the evolution of $A = A - A$.

In Fig 8 we show the evolution of the differences in the scalefactors when the radiation is switched on and off, for three different initial brane energy densities. Surprisingly, we have




FIG. 8: The difference between the scale factors in the radiating and non-radiating case versus time. For $b_0 = 520$ a critical-like behavior is observed (solid line), when the increase of the self-gravity of the brane due to absorption is roughly compensated by the radiation pressure. For initial densities much smaller (e.g. $b_0 = 100$, upper dotted line), the effect of the radiation pressure is dominant. For higher initial densities than the critical one ($b_0 = 2000$, lower dotted line) the increase of self-gravity due to absorption dominates over the radiation pressure.

found a critical-like behaviour for $b_0 = 520$. The difference in the scale factors is extremely close to zero, but it shows a sinusoidal-like pattern. For higher initial densities the scale factor is smaller in the radiating case throughout the cosmological evolution. This means that the self-gravity of the absorbed radiation dominates over the radiation pressure. For initial pressures higher than the critical one, the scale factor is higher in the radiating case during the whole cosmological evolution. This means that the radiation pressure dominates over the self-gravity of the absorbed radiation.

The behaviour of the difference in the energy densities is illustrated in Fig 9. Among the previous graphs, Figs 2, 3, 5 and 9 were plotted for the critical initial density. For other density values these graphs would be merely scaled, and would show similar features.

This is not true for the source terms of the Friedmann equation. We have seen various behaviours even in the non-radiating case (Figs 4-6), when the initial brane energy density is varied. In Figs 10-12 we have plotted the evolution of the difference, in the radiating and non-radiating cases, of the asymmetry and dark radiation source terms, for the critical initial energy density (Fig 10), for a much lighter brane (Fig 11) and for a much heavier




FIG. 9: The difference between the brane energy densities in the radiating and non-radiating case for $b_0 = 520$.

brane (Fig 12). With the exceptions of the very early and very late stages of the evolution of the universe, the two contributions roughly cancel each other for the critical initial energy density. For light branes the increase in the asymmetry term at early times is greater than the decrease in the dark radiation term. For heavy branes, by contrast, the decrease in the dark radiation term due to the Hawking radiation from the bulk exceeds the increase of the asymmetry term.

V I. D I S C U S S I O N

Table-top experiments [15] on possible deviations from Newton's law currently probe gravity at sub-millimeter scales and as a result they constrain the characteristic curvature scale of the bulk to 1.0 ± 0.1 mm. Expressed in units of inverse energy (when $c = 1 = \hbar$) this curvature scale is

$$l_{\text{max}} = 506.77 \text{ eV}^{-1} = 0.50677 \cdot 10^{12} \text{ GeV}^{-1} : \quad (43)$$

This is usually known as $1 \cdot 10^{12} \text{ GeV}^{-1}$ [16].

The 4-dimensional coupling constant e^2 and 4-dimensional gravitational constant G are related to the 4-dimensional Planck mass M_P as $e^2 = 8\pi G = 8\pi M_P^{-2}$, with $M_P = 10^{19} \text{ GeV}$. The 5-dimensional Planck mass is defined as $e^2 = 8\pi M_{(5)}^3$. As $M_{(5)}$ depends on both M_P




FIG . 10: The difference between source terms in the Friedmann equation computed when the radiation is switches on and off. The asymmetry source term (AST, solid line) is increased, while the dark radiation source term (DRST, dotted line) is decreased by radiation. The plot is for the "critical" initial density $b_0 = 520$.




FIG . 11: The same as in Fig 10, but for $b_0 = 100$. The main difference with respect to Fig 10 is that at early times the increase in the asymmetry term is faster than the decrease in the dark radiation term.

and on the characteristic curvature scale l as $M_{(5)}^3 = M_P^2 = 1$ [2], we get

$$M_{(5) \min} = 6.65 \cdot 10^8 \text{ GeV} : \quad (44)$$

This result is referred usually as $M_P > M_{(5)} > 10^8 \text{ GeV}$ [16]. Due to the presence of the extra dimension accessible via gravity, brane-world theories allow for the effective 4-dimensional




FIG. 12: The same as in Fig 10, but for $b_0 = 2000$. The decrease in the dark radiation term dominates over the increase of the asymmetry term. This effect is more accentuated than at the critical initial density, shown on Fig 10.

Planck scale on the brane to be much higher than the true, 5-dimensional Planck scale.

The knowledge of both l and M_P gives a characteristic lower limit for the brane tension in the two-brane model [17]. In units $c = 1 = \hbar$ this is

$$m_{\text{in}} = \frac{3M_P^2}{4l^2} = 138.59 \text{ TeV}^4; \quad (45)$$

This result is frequently quoted as $> 1 \text{ TeV}^4$ [16]. On the other hand Big Bang Nucleosynthesis (BBN) constraints result in a much milder lower limit, $> 1 \text{ MeV}^4$ [18]. There is also an astrophysical limit for ρ , this being sensitive to the equation of state of a neutron star on the brane [19]. For a typical neutron star (with uniform density) this is $> 5 \cdot 10^8 \text{ MeV}^4$, in between the two previous lower limits.

Our choice for $M_T = M_P = 10^{19} \text{ GeV}$ employed in the numerical analysis of the previous Section and the numerical value of the Planck mass $M_P = 10^{19} \text{ GeV}$ implies $M_T = 10^{15} \text{ TeV}$. In consequence $\rho = (M_T)^4 = 10^{60} \text{ TeV}^4$, well above all previously enlisted lower bounds. This choice is in agreement with the range of ρ represented in Fig. (1) of Ref. [18]. The reason we have chosen such a high value of the brane tension is the following. Decreasing ρ would mean to decrease M_T accordingly. As ρ is proportional to $(M_T = M_P)^4$ (cf Eq. (36)), by choosing a significantly smaller value of ρ the effect of the radiation energy density becomes very small, in the range of the numerical errors.

In this paper our intention was to focus on the effects of the asymmetric setup and bulk

radiation absorbed on the brane. Therefore we have simplified the model by setting $\alpha = 0$. By means of our choices of ρ_R and ρ_0 the dark radiation term was chosen of comparable size to the matter energy density source term, slightly above the limits established in [20] and [21] from BBN for the dark radiation term. We note however that those limits were derived in a symmetric setup, and in the presence of a cosmological constant, which is not the case here.

The question arises of whether the dark radiation term can take the role of a cosmological constant producing cosmic acceleration. Contrarily to optimistic expectations, this is not exactly happening in brane-world theories. Although the dark radiation term $2\bar{m}a^4$ is positive (for $\bar{m} > 0$) in the Friedmann equation (16), it appears with negative sign in the Raychaudhuri equation (15). Therefore dark radiation increases both the cosmic expansion rate and the cosmic deceleration as compared to standard cosmology. A slower expansion rate and additional acceleration is gained for negative values of $\bar{m}$, but such a possibility is not allowed by our present model.

Would the bulk radiation pushing away the brane and therefore causing cosmic expansion be able to produce inflation on the brane for any set of numerical data? For the range of numerical parameters employed in the preceding Section such an effect does not occur. Without radiation, the scale factor shows the typical evolution of a closed universe; cf Fig 2: Inflation would mean the occurrence of an exponential expansion, but this is not happening by switching on the radiation. Indeed, a glance on the scales of Figs 2 and 8, respectively (in the latter we have represented the differences in the scale factor with and without radiation), shows that radiation is not changing significantly the evolution of the scale factor. The deviations of physical quantities evaluated in the radiating case as compared to the nonradiating case are typically of the order of 10^{-4} . We have also checked that the radiation is not changing significantly the evolution of the Hubble-parameter depicted in Fig 3, therefore inflation due to this mechanism is excluded.

Stated more generically and independently of the specific numerical choices in our analysis, we can pose the question, whether the absorbed radiation could behave like a scalar field in a slow-roll regime. Eq. (27) shows that the absorbed radiation is better interpreted as dark dust matter, rather than dark radiation. Indeed, it is a pressureless fluid. Therefore, the attempt to describe it in terms of a scalar field would yield $\rho = 2V$ and the slow-roll condition could not be satisfied.

As we are considering closed cosmological models with $k = 1$, it is obvious that we should ask whether the radiation pressure would be able to stop the recollapse of the brane-world universe, producing a bounce? Our numerical investigations suggest a negative answer. For the whole range of initial brane energy densities that we have checked, the recollapse will inevitably occur. The explanation is again the double role of the radiation: it gives rise to both a radiation pressure, which is accelerating the motion of brane in the bulk, and an increase of brane self-gravity, which is decelerating it. As consequence, the Hubble parameter is zero only once during the whole cosmological evolution, when the expansion ceases and the recollapse begins.

Although radiation pressure can in principle accelerate the cosmological expansion, it will always encounter the competing effect of the brane becoming heavier as it absorbs more and more radiation. In our model radiation pressure acts as dark energy during the whole cosmological evolution; however, dark matter (some form of dust) is produced as well – cf Eq. (27). This is counterbalancing the accelerating effect.

Our numerical investigations show, that the increase in the energy density ρ^{rad} , which is due to the absorbed radiation is not significant as compared to the energy density of conventional matter. This can be seen from our Fig 9, representing the variation in ρ due to the bulk black hole radiation. The variation is of order 10^{-4} as compared to the energy density of the brane in the non-radiating case, shown in Fig 3. It is important to mention this point, as in our model we have studied a radiation dominated universe. By contrast, all dark matter born from the Hawking radiation absorbed on the brane is dust, cf. Eq. (27). As the absorbed energy density merely represents a small fraction of the original brane energy density, our assumption of a radiation-dominated universe holds true.

As we have seen, even the tiny variations of order 10^{-4} in the physical quantities can be achieved only at the price of having a huge brane tension $\sigma = 10^{60} \text{ TeV}^4$. With the Hawking radiation energy density so small, we do not expect drastical changes to occur in the mass of the bulk black hole, although theoretically a complete evaporation may occur at t_c (see the remarks following Eq. (31)). Indeed, the coefficient of $(t - t_0)$ in Eq. (31) is very small, therefore the mass of the brane will not increase significantly due to the absorption of this radiation, even seen during the whole cosmological evolution. Accordingly, the bulk black hole will not completely evaporate, rather its mass function will decrease only slightly during the lifetime of the closed universe. How small the rate of evaporation of the bulk black hole

is can be better seen from Eq. (38), containing dimensionless variables. There, with our choice $M_T = M_P = 1=10$, the coefficient of $\mathfrak{e} = \mathfrak{e}_0$ can be evaluated to give

$$\frac{4}{\mathfrak{e}} = \frac{160}{3} \frac{(5)}{\mathfrak{e}^7} 10^{-4} = 7.475 \cdot 10^{-7} : \quad (46)$$

This means that starting with the initial (dimensionless) mass function of the bulk black hole of $\mathfrak{m}_L(0) = 0.01$, towards the end of the recollapse the mass function decreases to the values presented in Table I. We note that the value of the initial brane energy density has a certain influence on the final value of the dimensionless mass function, the evaporation being maximal for the critical brane energy density.

TABLE I: Due to Hawking radiation, the dimensionless mass function of the bulk black hole slightly decreases during cosmological evolution. Towards the end of the recollapse (at $\mathfrak{e} = 1.115$) the dimensionless bulk black hole mass function is minimal for the critical value of the initial brane energy density.

$\mathfrak{m}_L$	$b_0 = 100$	$b_0 = 520$	$b_0 = 2000$
$\mathfrak{e} = 0$	0.01	0.01	0.01
$\mathfrak{e} = 1.115$	0.0099582386	0.0099582385	0.0099582387

VII. CONCLUDING REMARKS

In this paper we have considered a scenario which is able to deal with the possibility of a Hawking radiation escaping a bulk black hole and we have investigated the consequences of such a radiation on the cosmological evolution of a closed $k = 1$ universe.

For this purpose first we have developed a formalism suitable for describing the asymptotic brane-world containing a single radiating bulk black hole on the left bulk region. In the most generic scenario the radiation can be partially absorbed, partially transmitted and partially reflected on the brane. We have considered here the case of negligible reflection, because we wanted to dispose of an exact solution of the Einstein equations in the bulk. By suppressing the reflection, and considering the geometrical optics limit of the radiation, the bulk can be described by the VAdS5 metric.

The part of radiation absorbed on the brane generates quite interesting effects, which were not studied before. These can be related to both dark energy and dark matter. In order to concentrate on these effects, the absorption was chosen as maximal. By suppressing transmission, the equations could be partially integrated, cf. Eq. (31). Then we have introduced dimensionless quantities in order to perform a complex numerical analysis.

We have carefully compared the cosmological evolutions of the models both without and with radiation and found that the Hawking radiation of the bulk black hole represents merely a perturbation of the non-radiating model. This holds true during the entire cosmological evolution.

We have studied these perturbations in detail and shown that in the presence of the radiation, there are two competing effects. The absorption on the brane contributes towards the self-gravity of the brane, which then tends to recollapse faster. This phenomenon appears as continuously emerging dark matter on the brane. By contrast, radiation pressure tends to accelerate away the brane from the bulk black hole, producing an equivalent of dark energy. For the numerical data employed, we have found the critical value of the initial brane energy density for which these two competing effects roughly annihilate each other. We have also analysed cosmological evolution for both lighter and heavier branes and found that (at early times) the radiation pressure is dominant for light branes, while for heavy branes the self-gravity dominates during the whole cosmological evolution.

As a main result, we have proved that the asymmetry introduced in the model is not able to change the recollapse of the universe in the $k = 1$ case, regardless of whether the radiation is switched off or on. Our numerical analysis has shown that the bulk Hawking radiation cannot change the final fate of the recollapsing universe either.

VIII. ACKNOWLEDGMENTS

We thank Roy Maartens for useful comments and for stimulating interactions, and Ibolya Kepiro for assistance in producing the graphs. This work was supported by OTKA grants no. T 046939 and T S044665. LAG wishes to thank the support of the Janos Bolyai Scholarship

of the Hungarian Academy of Sciences.

- [1] Arkani-Hamed N, Dimopoulos S and Dvali G, The Hierarchy Problem and New Dimensions at a Millimeter, 1998 Phys. Lett. B 429, 263
Arkani-Hamed N, Dimopoulos S and Dvali G, Phenomenology, Astrophysics and Cosmology of Theories with Sub-Millimeter Dimensions and TeV Scale Quantum Gravity, 1999 Phys. Rev. D 59, 086004
- [2] Randall L and Sundrum R, An Alternative to Compactification, 1999 Phys. Rev. Lett. 83, 4690
- [3] Shiromizu T, Maeda K and Sasaki M, The Einstein Equations on the 3-Brane World, 2000 Phys. Rev. D 62, 024012
- [4] Gergely L A, Generalized Friedmann branes, 2003 Phys. Rev. D 68, 124011
- [5] Gergely L A and Maartens R, Asymmetric brane-worlds with induced gravity, 2005 Phys. Rev. D 71, 024032
- [6] Langlois D, Sorbo L and Rodriguez-Martinez M, Cosmology of a brane radiating gravitons into the extra dimension, 2002 Phys. Rev. Lett. 89, 171301
- [7] Gergely L A, Leeper E and Maartens R, Asymmetric radiating brane-world, 2004 Phys. Rev. D 70, 104025
- [8] Jennings D, Vernon I R, Graviton emission into non- Z_2 symmetric brane world spacetimes, 2005 JCAP 0507 011
- [9] Langlois D, Is our Universe brany?, 2005 [hep-th/0509231]
- [10] Emparan R, Horowitz G T and Myers R C, Black Holes Radiate Mainly on the Brane, 2000 Phys. Rev. Lett. 85, 499
- [11] Hemming S and Keski-Vakkuri E, Hawking radiation from AdS black holes, 2001 Phys. Rev. D 64, 044006
- [12] Guedens R, Clancy D and Liddle A R, Primordial black holes in braneworld cosmologies: Formation, cosmological evolution and evaporation, 2002 Phys. Rev. D 66, 043513
- [13] Jennings D, Vernon I R, Davis A C and van de Bruck C, Bulk black holes radiating in non- Z_2 brane-world spacetimes, 2005 JCAP 0504 013
- [14] Gergely L A, Spherically symmetric static solution for colliding null dust, 1998 Phys. Rev. D

58, 084030

- [15] Long J C et al., New experimental limits on macroscopic forces below 100 microns, 2003 Nature 421, 922
- [16] M aartens R , Brane-world gravity, 2004 Living Rev. Rel. 7, 1
- [17] Randall L and Sundrum R , Large mass hierarchy from a small extra dimension, 1999 Phys. Rev. Lett. 83, 3370
- [18] M aartens R , Wands D , Bassett B A , and Heard I P C , Chaotic Inflation on the brane, 2000, Phys. Rev. D 62, 041301 (R)
- [19] Germani C and M aartens R , Stars in the braneworld, 2001 Phys. Rev. D 64, 124010
- [20] Binetruy P , Dey C , Ellwanger U and Langlois D , Brane cosmological evolution in a bulk with cosmological constant, 2000 Phys.Lett. B 477, 285
- [21] Ichiki K , Yahiro M , Kajino T , Orito M and Mathews G J , Observational Constraints on Dark Radiation in Brane Cosmology, 2002, Phys.Rev. D 66, 043521