Holographic Derivation of Entanglement Entropy from AdS/CFT

Shinsei Ryu and Tadashi Takayanagi

K avli Institute for Theoretical Physics, University of California, Santa Barbara, CA 93106, USA (Dated: October 18, 2021)

A holographic derivation of the entanglement entropy in quantum (conformal) eld theories is proposed from AdS/CFT correspondence. We argue that the entanglement entropy in d+1 dimensional conformal eld theories can be obtained from the area of d dimensional minimal surfaces in AdS_{d+2} , analogous to the Bekenstein-Hawking from the area of d dimensional minimal surfaces in proposal perfectly reproduces the correct entanglement entropy in 2D CFT when applied to AdS_3 . We also compare the entropy computed in AdS_5 S⁵ with that of the free N = 4 super Yang-Mills.

I. IN TRODUCTION

One of the most remarkable successes in gravitational aspects of string theory is the microscopic derivation of the Bekenstein-Hawking entropy $\rm S_{B\,H}$

$$S_{BH} = \frac{A \text{ rea of horizon}}{4G_{N}};$$
 (1.1)

for BPS black holes [1]. This idea relates the gravitational entropy with the degeneracy of quantum—eld theory as its microscopic description. Taking near horizon limit, we can regard this as a special example of AdS/CFT correspondence [2, 3, 4]. It claims that the d+1 dimensional conformal—eld theories (CFT $_{d+1}$) are equivalent to the (super)gravity on d+2 dimensionalantideSitter space AdS $_{d+2}$. We expect that each CFT is sitting at the boundary of AdS space.

On the other hand, there is a dierent kind of entropy called entanglement entropy (von-Neumann entropy) in quantum mechanical systems. The entanglement entropy

$$S_A = t_A \log_A ; \quad A = t_B j \text{ ih } j; \quad (1.2)$$

provides us with a convenient way to measure how closely entangled (or how \quantum ") a given wave function j i is. Here, the total system is divided into two subsystems A and B and A is the reduced density matrix for the subsystem A obtained by taking a partial trace over the subsystem B of the total density matrix = j ih j. Intuitively, we can think S_A as the entropy for an observer who is only accessible to the subsystem A and cannot receive any signals from B. In this sense, the subsystem B is analogous to the inside of a black hole horizon for an observer sitting in A , i.e., outside of the horizon. Indeed, an original motivation of the entanglement entropy was its similarity to the Bekenstein-Hawking entropy [5, 6].

The entanglement entropy is of growing importance in many elds of physics in our exploration for better understanding of quantum systems. For example, in a modern trend of condensed matter physics it has been becoming clear that quantum phases of matter need to be characterized by their pattern of entanglement encoded in many-body wave functions of ground states, rather than conventional order parameters [7, 8, 9]. Recently, the entanglement entropy has been extensively studied in low-dimensional quantum many-body systems as a new

tool to investigate the nature of quantum criticality (refer to [10] and references therein for example).

For one-dimensional (1D) quantum many-body systems at criticality (i.e. 2D CFT), it is known that the entanglement entropy is given by [10, 11]

$$S_A = \frac{c}{3} \quad \log \frac{L}{a} \sin \frac{1}{L} \quad ; \tag{1.3}$$

where l and L are the length of the subsystem A and the total system A [B (both ends of A [B are periodically identi ed), respectively; a is a ultra violet (UV) cuto (lattice spacing); c is the central charge of the CFT. When we are away from criticality, Eq. (1.3) is replaced by [7,10]

$$S_A = \frac{C}{6} \quad A \quad \log g \tag{1.4}$$

where is the correlation length and A is the number of boundary points of A (e.g. A = 2 in the setup of (1.3)).

In spite of these recent developments, and its similarity to the black hole entropy, a comprehensive gravitational interpretation of the entanglement entropy has been lacking so far. Here, we present a simple proposal on this issue in the light of AdS/CFT duality. Earlier discussions from dierent view points can be found in e.g. papers [12, 13]. De ne the entanglement entropy \S_{A} in a CFT on $\mathbb{R}^{1/d}$ (or \mathbb{R}^{-3}) for a subsystem A that has an arbitrary d 1 dimensional boundary @A 2 \mathbb{R}^d (or \mathbb{S}^d). In this setup we propose the following area law'

$$S_A = \frac{A \text{ rea of }_A}{4G_N^{(d+2)}};$$
 (1.5)

where $_{\rm A}$ is the d dimensional static minimal surface in AdS_{d+2} whose boundary is given by @A, and ${\sf G}_{\rm N}^{(d+2)}$ is the d+2 dimensional Newton constant. Intuitively, this suggests that the minimal surface $_{\rm A}$ plays the role of a holographic screen for an observer who is only accessible to the subsystem A. We show explicitly the relation (1.5) in the lowest dimensional case d=1, where $_{\rm A}$ is given by a geodesic line in AdS $_{\rm 3}$. We also compute S $_{\rm A}$ from the gravity side for general d and compare it with eld theory results, which is successful at least qualitatively. From (1.5), it is readily seen that the basic properties of the entanglement entropy (i) $S_{\rm A}=S_{\rm B}$ (B is the complement




FIG. 1: (a) $A\,dS_3$ space and $C\,F\,T_2$ living on its boundary and (b) a geodesics $_A$ as a holographic screen.

of A) and (ii) $S_{A_1}+S_{A_2}$ $S_{A_1\,[\,A_2\,]}$ (subadditivity) are satis ed.

We can also de ne the entanglement entropy at nite temperature $T=\frac{1}{1}$. E.g. in a 2D CFT on a in nitely long line, it is given by replacing L in Eq. (1.3) with i [10]. We argue that Eq. (1.5) still holds in T>0 cases. Note that $S_A=S_B$ is no longer true if T>0 since is in a mixed state generically. At high temperature, we will see that this occurs due to the presence of black hole horizon in the dual gravity description.

II. ENTANGLEMENT ENTROPY IN AdS3=CFT2

Let us start with the entanglement entropy in 2D CFTs. According to AdS/CFT correspondence, gravitational theories on AdS_3 space of radius R are dual to (1+1)D CFTs with the central charge [14]

$$C = \frac{3R}{2G_N^{(3)}} : (2.1)$$

The metric of AdS_3 in the global coordinate (t;;) is

$$ds^2 = R^2$$
 $\cosh^2 dt^2 + d^2 + \sinh^2 d^2$: (2.2)

At the boundary = 1 of AdS $_3$ the metric is divergent. To regulate physical quantities we put a cuto $_0$ and restrict the space to the bounded region $_0$. This procedure corresponds to the UV cuto in the dual CFTs [15]. If L is the total length of the system with both ends identied, and a is the lattice spacing (or UV cuto) in the CFTs, we have the relation (up to a numerical factor)

$$e^{0}$$
 L=a: (2.3)

The (1+1)D spacetime for the CFT $_2$ is identiced with the cylinder (t;) at the boundary = $_0$. The subsystem A is the region 0 = $_0$. Then $_A$ in Eq. (1.5) is identiced with the static geodesic that connects the boundary points = 0 and 2 l=L with t exed, traveling inside AdS $_3$ (Fig. 1 (a)). With the cuto $_0$ introduced above, the geodesic distance L $_A$ is given by

$$\cosh \frac{L_{A}}{R} = 1 + 2 \sinh^{2}_{0} \sin^{2} \frac{1}{L};$$
(2.4)

Assuming the large UV cuto e° 1, the entropy (1.5) is expressed as follows, using Eq. (2.1)

$$S_{A}' \frac{R}{4G_{A}^{(3)}} \log e^{2} \sin^{2} \frac{1}{L} = \frac{c}{3} \log e^{0} \sin \frac{1}{L} : (2.5)$$

This entropy precisely coincides with the known CFT result (1.3) after we remember the relation Eq. (2.3).

This proposed relation (1.5) suggests that the geodesic (orm inimal surface in the higher dimensional case) $_{\rm A}$ is analogous to an event horizon as if B were a black hole, though the division into A and B is just articial. In other words, the observer, who is not accessible to B, will probe $_{\rm A}$ as a holographic screen [16], instead of B itself (Fig. 1 (b)). The minimal surface provides the severest entropy bound when we x its boundary condition. In our case it saturates the bound.

M ore generally, we can consider a subsystem A which consists of multiple disjoint intervals as follows

$$A = fx_{3}x_{2} [r_{1}; s_{1}] [r_{2}; s_{2}] [N_{1}; s_{1}x_{2}]$$
 (2.6)

where 0 $\,$ $r_{\rm l} < s_1 < r_2 < s_2 < \,$ $_{\rm N} < {\rm rs_N}$ $\,$ L. In the dual A dS $_3$ description, the region (2.6) corresponds to $\,$ 2 [$_{\rm i=1}^{\rm N}$ [$_{\rm L}^{\rm 2}$; $_{\rm L}^{\rm 2}$] at the boundary. In this case it is not straightforward to determ ine m inim al (geodesic) lines responsible for the entropy. However, we cannot the answer from the entanglement entropy computed in the CFT side. The general prescription of calculating the entropy for such systems is given in [10] using conformal mapping. For our system (2.6), we not, when rewritten in the AdS $_3$ language, the following expression of S $_A$

$$S_{A} = \frac{P}{\underset{i,j}{L_{r_{j};s_{i}}}} \frac{P}{\underset{i < j}{L_{r_{j};r_{i}}}} \frac{P}{\underset{i < j}{L_{s_{j};s_{i}}}}; (2.7)$$

where $L_{a,b}$ is the geodesic distance between two boundary points a and b. We can think that the correct denition of minimal surface is given by the numerator in Eq. (2.7).

Next we turn to the entanglement entropy at nite temperature. We assume the spacial length of the total system L is in nitely long st. =L 1. At high temperature, the gravity dual of the CFT is the Euclidean BTZ black hole [17] with the metric given by

$$ds^{2} = (r^{2} + r^{2})d^{2} + \frac{R^{2}}{r^{2} + r^{2}}dr^{2} + r^{2}dr^{2}; \qquad (2.8)$$

The Euclidean time is compactified as $+\frac{2~R}{r_+}$ to obtain a smooth geometry in addition to the periodicity '+2. Looking at its boundary, we not the relation $\frac{1}{L}=\frac{R}{r_+}$ 1 between the CFT and the BTZ black hole.

The subsystem A is defined by 0 $^{\prime}$ 2 l=L at the boundary. Then we expect that the entropy can be computed from the geodesic distance between the boundary points $^{\prime}$ = 0;2 l=L at a xed time. To nd the geodesic line, it is useful to remember the familiar fact that the Euclidean BTZ black hole at temperature T_{BTZ} is equivalent to the therm alAdS $_3$ at temperature $1=T_{BTZ}$. This

equivalence can be interpreted as a modular transformation in the CFT side [18]. If we de ne the new coordinates $r=r_+\cosh$; $r_+=R$; $r_+'=Rt$; then the metric (2.8) indeed becomes the Euclidean version of $A\,dS_3$ metric (2.2). Thus the geodesic distance can be found in the same way as in Eq. (2.4): $\cosh(L_A=R)=1+2\cosh^2_0 \sinh^2_-\frac{1}{2}$, where the UV cuto is interpreted as e 0 =a. Then the area law (1.5) reproduces the known CFT result [10]

$$S_A() = \frac{c}{3} \quad \log \frac{1}{a} \sinh \frac{1}{a} : \qquad (2.9)$$

We can extend these arguments to the multi interval cases and nd the same formula (2.7) as before.

It is instructive to repeat the same analysis in the Poincare metric $ds^2=R^2z^2$ (dz^2 $dx_0^2+dr^2$): We dene the subsystem A by the region l=2 r l=2 at the boundary z=0. The geodesic line A is given by

$$(r;z) = \frac{1}{2} (\cos s; \sin s);$$
 (s); (2.10)

where the in nitesimal is the UV cuto 2a=1 (or equally z_{UV} a). We obtain the entropy $S_{\!A}$ as follows

$$S_A = \frac{L_A}{4G_N^{(3)}} = \frac{R}{2G_N^{(3)}} = \frac{Z_{=2}}{\frac{ds}{\sin s}} = \frac{c}{3} \log \frac{1}{a};$$
 (2.11)

This reproduces the small limit of Eq. (1.3) [11].

When we perturb a CFT by a relevant perturbation, the RG ow generically drives the theory to a trivial R xed point. We denote the correlation length in the latter theory. In the AdS dual, this massive deformation corresponds to capping othe IR region, restricting the allowed values of z to z. In the large l limit, we not

$$S_A = \frac{1}{4G_N^{(3)}}^{Z_{2}=1} \frac{ds}{\sin s} = \frac{c}{6} \log \frac{1}{a}$$
: (2.12)

This agrees with the CFT result with A = 1 (1.4) [7, 10].


III. HIGHER DIM ENSIONAL CASES

M otivated by the success in our gravitational derivation of the entanglem ent entropy for d=1, it is interesting to extend the idea to higher dimensional cases (d 2). A natural thing to do is to replace geodesic lines with minimal surfaces. The computations are analogous to the evaluation of Wilson loops [19], though the dimension of relevant minimal surfaces is different.

W e will work in the Poincare m etric for A dS $_{\mbox{\scriptsize d+}\mbox{\scriptsize 2}}$

$$ds^{2} = R^{2}z^{2} (dz^{2} dz^{2} dz^{2} + dx_{0}^{2} + dx_{1}^{2});$$
 (3.1)

We consider two examples for the shape of A. The stone is a straight belt $A_S = fx_i jx_1 2 \ [1=2; 1=2]; x_2; 3;$






FIG. 2: M in im alsurfaces in AdS $_{d+\,2}\colon$ (a) A $_{S}$ and (b) A $_{D}$.

[1 ;1]g at the boundary z = 0 (Fig. 2 (a)). In this case the minimal surface is de ned by dz=dx = $\frac{p}{z^{2d}}$ z^{2d} = z^d ; where z is determined by l=2 = $\frac{z}{z}$ dzz^d (z^{2d} z^{2d}) z^{2d} z^{2d}

A rea_{A_S} =
$$\frac{2R^{d}}{d} \frac{L}{d} \frac{L}{a} \frac{d^{1}}{d} \frac{2^{d-d=2}R^{d}}{d} \frac{\frac{d+1}{2d}}{\frac{1}{2d}} \frac{d^{1}}{d} \frac{L}{1} ;$$
(3.2)

where L is the length of A $_{S}$ in the $x_{2;3}$; $_{i\bar{c}}$ direction. The second example is the disk A $_{D}$ de nedpby A $_{D}$ = fx_{1} jr fx_{1} lg (Fig. 2 (b)) in the polar coordinate fx_{1} dx fx_{1} = fx_{2} dr fx_{2} dr fx_{1} . The minimal surface is a drdimensional ball B fx_{2} de ned by fx_{2} .10). Its area is

$$A rea_{A_{D}} = C \frac{Z_{1}}{a=1} dy \frac{(1 - y^{2})^{(d-2)=2}}{y^{d}}$$

$$= p_{1} (l=a)^{d-1} + p_{3} (l=a)^{d-3} +$$

$$p_{d-1} (l=a) + p_{d} + 0 (a=1); d: even;$$

$$p_{d-2} (l=a)^{2} + q log (l=a) + 0 (1); d: odd,$$

$$(3.3)$$

where $C=2^{d=2}R^{d}=p$ (d=2) and $p=C=(d-1)^{1}$ etc. For deven, $p_{d}=C=(2^{D-1})^{-1}$ $\frac{d}{2}$ $\frac{1}{2}^{d}$ and for d odd, $q=C=(-1)^{(d-1)=2}$ (d=2)!!=(d=1)!!.

From these results, the entanglement entropy can be calculated by Eq. (1.5). Each of (3.2) and (3.3) has a UV divergent term a^{d+1} that is proportional to the area of the boundary @A. This agrees with the known area law' of the entanglement entropy in quantum eld theories [5,6]. Note that this area law' is related to ours Eq. (1.5) via the basic property of holography.

W e m ay prefer a physical quantity that is independent of the cuto (i.e. universal). The second term in Eq. § 2) has this property. In general, when A is a nite size, there is a universal and conform al invariant constant contribution to S_A if d is even (see [20] for properties of m in in al surfaces in AdS). In (2+1)D topological eld theories the constant contribution to S_A encodes the quantum dimension and is called the topological entanglement entropy [8, 9]. If d is odd, the coeclient of the logarithm ic term

log(l=a) is universal as in Eq. (13). Let us apply the previous results to a spe

Let us apply the previous results to a speci c string theory setup. Type IIB string on $A\,dS_5$ S^5 is dual to

4D N = 4 SU (N) super Yang-M ills theory [2]. The radius of A dS $_5$ and S 5 are given by the same value R = (4 g_s 12 N) $^{1\over 4}$. The 5D New ton constant is related to the 10D one via $G_N^{(10)}$ = 3 R 5 G $_N^{(5)}$. Then we obtain from Eqs. (3.2) and (3.3)

$$S_{A_s} = \frac{N^2 L^2}{2 a^2} \quad \stackrel{p}{2} - \frac{\frac{2}{3}}{\frac{1}{6}} \cdot \frac{N^2 L^2}{1^2}; \quad (3.4)$$

$$S_{A_D} = N^2 \frac{1^2}{a^2} \log \frac{1}{a} + 0 (1) : (3.5)$$

It is interesting to compare the nite universal term in Eq. (3.4) with the eld theory one. For free real scalars and ferm ions in general dimensions, one way to compute S_{A_S} is presented in [21] (see also [22]). Indeed, this leads to the same behavior in a and las in Eq. (3.4). Following this approach, we can estimate nite contributions from 6 scalars and 4 M a prana ferm ions in the N = 4 Yang-Mills multiplet. In the end, we obtain Sfinite (0.068 + g) ${}^{2}NL^{2}=1^{2}$, where g is the contribution from the gauge eld (g = 0.010) if we treat the gauge eld as 2 scalars). On the other hand, our AdS result (3.4) leads to SAdS finite 0.051 ${}^{2}NL^{2}=1^{2}.Wemay$ think this is a good agreem ent if we rem em ber that the gravity description corresponds to the strongly coupled gauge theory instead of the free theory as in [23].

W e can also compute the entanglement entropy for the nearhorizon limit $A dS_4 = S^7 (A dS_7 = S^4)$ of N M 2-branes (M 5-branes)

$$S_{A_s}^{M_2} = \frac{P_{\overline{2}}}{3} N^{3=2} \frac{L}{a} \frac{4^3}{(1=4)^4} \frac{L}{1} ;$$
 (3.6)

$$S_{A_D}^{M_2} = \frac{P_{\overline{2}}}{3} N^{3=2} \frac{1}{a} 1;$$
 (3.7)

$$S_{A_s}^{M_5} = \frac{2}{3^2} N^3 \frac{L^4}{a^4} 16^{5=2} \frac{(3=5)^5}{(1=10)^5} \frac{L^4}{1^4}; (3.8)$$

$$S_{A_D}^{M_5} = \frac{32}{9} N^3 \frac{1}{4} \frac{1^4}{a^4} \frac{3}{4} \frac{1^2}{a^2} + \frac{3}{8} \log \frac{1}{a}$$
: (3.9)

Note that the constant term s in Eqs. (3.6), (3.7) and (3.8) do not depend on the choice of the cuto a.

IV. YANG-M ILLS AT FINITE TEMPERATURE

As the nalexample, we discuss the N = 4 super Y ang-Mills theory on \mathbb{R}^3 at a nite temperature T, which is dual to the AdS black hole de ned by the metric [24]

$$ds^{2} = R^{2} \frac{du^{2}}{hu^{2}} + u^{2} \quad hdt^{2} + dx_{1}^{2} + dx_{2}^{2} + dx_{3}^{2} + d \frac{2}{5} ;$$
(4.1)

where h=1 $u_0^4=u^4$; $u_0=T$. For the straight belt A $_S$, the area is given by (putting the cut o $u z^1$ a)

A rea_{A_S} =
$$2R^3L^2$$
 $\frac{Z_{a^{-1}}}{u}$ $\frac{du \ u^6}{(u^4 \ u^4)(u^6 \ u^6)}$; (4.2)




FIG. 3: (a) M in im al surfaces $_{\rm A}$ for various sizes of A. (b) $_{\rm A}$ and $_{\rm B}$ w rap the di event parts of the horizon.

where u satis es $l=2=\frac{R_1}{u}$ du $[(u^4 \quad u_0^4) (u^6=u^6 \quad 1)]^{1=2}$: Eq. (4.2) contains the UV divergence a 2 as before. As in the analogous computation of Wilson loops [25], we also expect a term which is proportional to the area of A. Indeed, when lis large $(u \quad u_0)$ we not the constant term $^3R^3L^2lT^3$. This leads to the nite part of S_A

$$S_{finite}' \frac{2}{2} N^2 T^3 L^2 l = \frac{2N^2 T^3}{2}$$
 (A rea of A_S):

We can regard this entropy as a part of the Bekenstein-Hawking entropy of black 5-branes [23], which is proportional to the area of horizon situated at $u = u_0$. Thus we can interpret the part (4.3) as a therm alentropy contribution to the total entanglement entropy at nite temperature. In our gravitational description, this part arises because the minimal surface wraps a part of the black hole horizon (Fig. 3 (a)). If we expand the size of A until it coincides with the total system (in the global coordinate), A wraps the horizon completely and A becomes equal to the Bekenstein-Hawking entropy as expected. In a sense, the overall normalization in Eq. (1.5) is xed from Eq. (1.1) once we consider the entanglem ententropy at nite tem perature. Note that at nite tem perature, $S_A = S_B$ does not hold generically. In the present situation, this occurs because the surfaces $_{\mathrm{A}}$ and $_{\mathrm{B}}$ wrap two di erent parts of the horizon (Fig. 3 (b)).

As argued in [13,26], the AdS black hole can be dual to an entanglem ent of two dierent CFTs at the two boundaries. It is interesting to look at the minimal surfaces that connect them. As a specienchimit, we may think the black hole entropy is the same as the entanglement entropy of the CFTs as the minimal surface wrap the horizon.

V. CONCLUSION

In this paper we proposed a holographic description of the entanglement entropy in quantum (conformal) eld theories via $A\,dS_{d+\,2}/C\,F\,T_{d+\,1}$ correspondence. This is sum marized as the area law relation (1.5). Based on our proposal we computed the entanglement entropy for various systems, e.g. 2D CFTs and the 4D large-N N = 4 super Yang-M ills theory. We checked that in the lowest dimensional case (d = 1), our formula exactly coincides with the entropy computed directly from CFT.

In higher dimensions (d 2), a quantitative comparison is not easy because the gravity description corresponds to the strongly coupled gauge theory whose entanglement entropy is not known at present. Nevertheless, we found that our computation for $A \, dS_5$ S^5 reproduces the same functional form of entanglement entropy as the one in the 4D free N=4 super Yang-Mills. Their numerical coecients only dier by a factor $\frac{3}{2}$ in a rough estimation (assuming that the gauge eld contribution is the same as that of two real scalar elds). This is similar to the well-known result of the thermal entropy [23].

Finally we computed the entanglement entropy at nite temperature employing AdS black hole geometry. In this case them inimal surface wraps a fraction of the black hole horizon and this part is responsible for the thermal contribution.

We would like to thank M. Einhom, T. Harmark, G. Horowitz, Y. Hyakutake, D. Marolf, T. Okuda, S.-J. Rey, M. Shigemori and Y. Sugawara for useful discussions. This research was supported in part by the National Science Foundation under Grant No. PHY 99-07949.

- [L] A. Strom inger and C. Vafa, \M icroscopic Origin of the Bekenstein-Hawking Entropy," Phys. Lett. B 379, 99 (1996) [arX iv hep-th/9601029].
- [2] J. M. Maldacena, \The large N limit of superconformal eld theories and supergravity," Adv. Theor. Math. Phys. 2, 231 (1998) [Int. J. Theor. Phys. 38, 1113 (1999)] [arX iv hep-th/9711200].
- [3] S.S.Gubser, I.R.K lebanov and A.M. Polyakov, \Gauge theory correlators from non-critical string theory, "Phys. Lett. B 428, 105 (1998) [arX iv hep-th/9802109].
- [4] E. Witten, \Anti-de Sitter space and holography," Adv. Theor. Math. Phys. 2, 253 (1998) [arXiv:hep-th/9802150].
- [5] L. Bombelli, R. K. Koul, J. H. Lee and R. D. Sorkin, \A Quantum Source Of Entropy For Black Holes," Phys. Rev. D 34, 373 (1986).
- [6] M. Srednicki, \Entropy and area," Phys. Rev. Lett. 71, 666 (1993) [arX iv hep-th/9303048].
- [7] G. Vidal, J. I. Latorre, E. Rico and A. Kitaev, \Entanglement in quantum critical phenomena, Phys. Rev. Lett. 90, 227902 (2003) [arXiv quant-ph/0211074].
- [8] A.K itaev and J.P reskill, \Topological entanglem ent entropy," arX iv hep-th/0510092.
- [9] M. Levin and X.G. Wen, \D etecting topological order in a ground state wave function," arX iv cond-m at/0510613.
- [10] P. Calabrese and J. Cardy, \Entanglement entropy and quantum eld theory," J. Stat. Mech. 0406, P002 (2004) [arX iv hep-th/0405152].
- [11] C. Holzhey, F. Larsen and F. Wilczek, \Geometric and renormalized entropy in conformal eld theory," Nucl. Phys. B 424, 443 (1994) [arXiv:hep-th/9403108].
- [12] S. Hawking, J. M. Maldacena and A. Strom inger, \D e-Sitter entropy, quantum entanglement and AdS/CFT," JHEP 0105,001 (2001) [arX iv hep-th/0002145].
- [13] J.M. Maldacena, \Etemalblack holes in Anti-de-Sitter," JHEP 0304,021 (2003) [arX iv hep-th/0106112].
- [14] J. D. Brown and M. Henneaux, \Central Charges In The Canonical Realization Of A sym ptotic Sym metries ,"Commun.Math.Phys.104,207 (1986).
- [15] L. Susskind and E. W itten, \The holographic bound in anti-de Sitter space," arX iv hep-th/9805114.
- [16] G. 't Hooff, \D im ensional reduction in quantum gravity," arX iv gr-qc/9310026; L. Susskind, \The W orld as a hologram," J. M ath. Phys. 36, 6377 (1995) [arX iv hep-th/9409089]; R. Bousso, \A Covariant Entropy Conjecture," JHEP 9907 (1999) 004

- [arX iv:hep-th/9905177].
- [17] M. Banados, C. Teitelboim and J. Zanelli, \The Black hole in three-dimensional space-time," Phys. Rev. Lett. 69, 1849 (1992) [arX iv hep-th/9204099].
- [18] J.M. Maldacena and A. Strom inger, \AdS (3) black holes and a stringy exclusion principle," JHEP 9812, 005 (1998) [arX iv hep-th/9804085].
- [19] S. J. Rey and J. T. Yee, \Macroscopic strings as heavy quarks in large N gauge theory and anti-de Sitter supergravity," Eur. Phys. J. C 22, 379 (2001) [arX iv hep-th/9803001]; J.M. Maldacena, \W ilson loops in large N eld theories," Phys. Rev. Lett. 80, 4859 (1998) [arX iv hep-th/9803002]; D. Berenstein, R. Corrado, W. Fischler and J. M. Maldacena, \The operator product expansion for W ilson loops and surfaces in the large N limit," Phys. Rev. D 59, 105023 (1999) [arX iv hep-th/9809188].
- [20] C.R.Graham and E.W itten, \Conformal anomaly of submanifold observables in AdS/CFT correspondence," Nucl. Phys. B 546, 52 (1999) [arX iv hep-th/9901021].
- [21] H.Casini and M.Huerta, \Entanglement and alpha entropies for a massive scalar eld in two dimensions," arX iv cond-mat/0511014.
- [22] D.V. Fursaev, \Entanglement entropy in critical phenomena and analogue models of quantum gravity," arX iv hep-th/0602134.
- [23] S. S. Gubser, I. R. Klebanov and A. W. Peet, \Entropy and Temperature of Black 3-Branes," Phys. Rev. D 54, 3915 (1996) [arX iv:hep-th/9602135].
- 24] E.W itten, \Anti-de Sitter space, them alphase transition, and con nement in gauge theories," Adv. Theor. Math. Phys. 2, 505 (1998) [arX iv hep-th/9803131].
- [25] S. J. Rey, S. Theisen and J. T. Yee, \W ilson-Polyakov loop at nite temperature in large N gauge theory and anti-de Sitter supergravity," Nucl. Phys. B 527, 171 (1998) [arX iv:hep-th/9803135]; A. Brandhuber, N. Itzhaki, J. Sonnenschein and S. Yankielowicz, \W ilson loops in the large N limit at nite temperature," Phys. Lett. B 434, 36 (1998) [arX iv:hep-th/9803137]; \W ilson loops, con nement, and phase transitions in large N gauge theories from supergravity," JHEP 9806, 001 (1998) [arX iv:hep-th/9803263].
- [26] R. Brustein, M. B. Einhom and A. Yarom, \Entanglement interpretation of black hole entropy in string theory," JHEP 0601,098 (2006) [arX iv hep-th/0508217].