Sym m etry A spects in N on relativistic M ulti-Scalar Field M odels and Application to a C oupled Two-Species D ilute Bose G asz

RudneiO Ram os¹ and M arcus B P into²

¹ Departamento de F sica Teorica, Universidade do Estado do Rio de Janeiro, 20550-013 Rio de Janeiro, RJ, Brazil

 2 D epartam ento de F $\,$ sica, U n iversidade Federal de Santa C atarina, 88040–900 F lorianopolis, SC , B razil

E-m ail: rudnei@uerj.br, marcus@fsc.ufsc.br

A bstract. We discuss unusual aspects of sym metry that can happen due to entropic e ects in the context of multi-scalar eld theories at nite temperature. We present their consequences, in special, for the case of nonrelativistic models of hard core spheres. We show that for nonrelativistic models phenom ena like inverse sym metry breaking and sym metry non-restoration cannot take place, but a reentrant phase at high temperatures is shown to be possible for some region of parameters. We then develop a model of interest in studies of Bose-E instein condensation in dilute atom ic gases and discuss about its phase transition patterns. In this application to a Bose-E instein condensation model, how ever, no reentrant phases are found.

PACS num bers: 11.10 W x,03.75 F i,05.30 Jp Published: J. Phys. A:M ath. Gen. 39, 6687 (2006)

1. Introduction

One of the most interesting aspects concerning the studies of multi-eld models at nite temperature is the possibility of emergence of a much richer phase diagram than one would usually nd in one eld type of models. The possibility that unusual symmetry patterns could emerge in those models, for some speci c region of parameters, has attracted considerable attention in the literature (see, for instance, [1, 2] and references therein and [3] for a short review).

Form ost of the standard physical system s we know in nature, we have a good sense of how symmetries seem to change as the temperature is changed. Typically, the larger is the temperature the larger is the symmetry exhibited by the system and vice-versa. Examples of this behaviour are expected to happen in the particle physics models, like in the electroweak phase transition and possibly also in G rand-U ni ed models. The same is

z Based on a talk given by R.O.Ram os at the QFEXT05 workshop, Barcelona, Spain, Septem ber 5-9, 2005

expected in the much lower energy systems, like those of condensed matter. For example, the phase transition in ferrom agnets, superconductors, Bose-Einstein condensation of atom ic gases, etc., just to name a few systems. In all these examples we always go from an ordered (less symmetric) phase below some critical temperature of phase transition to a disordered (more symmetrical) phase above the critical temperature, or the opposite, if the temperature is decreased from a high temperature (or symmetry restored) state [4].

However, the above symmetry aspects seem not to be the rule. In fact we are also becoming increasingly aware that entropic e ects in multi-eld models may show other patterns of symmetry breaking and restoration that are less usual. For example, m any condensed m atter system s, like spin glasses, com pounds know n as the m anganites, liquid crystals and many others, commonly show phenomena like reentrant phases of low er sym m etries at higher tem peratures and, therefore, they can exhibit unusual phase diagram s that we would otherwise not expect. M any of these system s have recently been reviewed in [5]. As concerned to quantum eld theory models, the possibility of other phase transition patterns was also shown to be possible in the context of multiscalar eld theories at nite tem peratures [1]. These models show the possibility of a sym m etry that is not broken at low tem peratures, getting broken at high tem peratures (what is called an inverse symmetry breaking). Other case that seems possible is a symmetry that is broken at lower temperatures, not getting restored at all as we go to higher tem peratures (what is called a symmetry non-restoration). The problem of how a symmetry broken or restored phase may emerge as a reentrant phase in the system, like it is seen in many low energy condensed matter systems, was recently analyzed in [2] in the context of a coupled nonrelativistic model of two scalar elds.

The plan of this paper is as follows. In the next section we brie y review the results obtained in [2] for the case of a nonrelativistic model of two scalar elds with overall symmetry U (1) U (1) and show that it adm its reentrant phases for some region of parameters. One possible physical realization of this kind of model is for example in the description of a coupled two species dilute atom ic gas system. This is a kind of system in atom ic physics that has been of great interest recently concerning studies (both theoretical and experimental) of Bose-E instein condensation. In Sec. 3 we o er a quantum eld theory description for this problem. We then analyze the possibility of emergence of reentrant phases in the quantum eld formulation for these systems. This would be a novel symmetry behaviour that could be of great interest, given its possible in plementation in the laboratory. Our results allow us to conclude, at the level of our approximations, on the non appearance of such reentrant phases in these kind of coupled dilute atom ic gases system s.

2. Reentrant Phases in Nonrelativistic Multi-Scalar Field Models

W e start our discussion by considering the following nonrelativistic Lagrangian density model, of two (complex) scalar elds and , with global symmetry U (1) U (1),

L(;;;) =
$$i\theta_{t} + \frac{1}{2m}r^{2}$$
 $\frac{g}{3!}()^{2}$
+ $i\theta_{t} + \frac{1}{2m}r^{2}$ $\frac{g}{3!}()^{2}$
g()(): (1)

Them odel (1) can be thought as coming from the nonrelativistic limit of a corresponding relativistic counterpart, as shown explicitly in [2]. In (1) the interaction parameters q and q describe two-body self-interaction terms, as commonly considered for dilute and cold (low energy) systems of particles [6], in which case only binary type interactions, i.e., hard core type of interactions of the form shown in (1), are relevant. g is the cross-coupling between the two elds, which we consider here as a quadratic type of interaction. The mass parameters m and m are the masses for the elds (or particles). The one-body type of interactions, of magnitude and , can either represent the e ect of external potentials (for example a magnetic eld) on the system, internal energy term s (like the internal molecular energy relative to free atom s in which case the elds in the Lagrangian would be related to molecular dimers), an explicit gap of energy in the system (like in superconductors), or just chem ical potentials added to the action in the grand-canonical formulation to enforce nite density (or xed number of particles) for both and . The latter will be the case for our application of (1) to the coupled atom ic gas problem in Sec. 3. Here we will just consider the i parameters as constant one-body parameters added to our model such that the possible symmetry breaking patterns, depending on the sign of i (i = ;), can easily be determined from the potential term in (1). This is just what is done in spontaneous symmetry breaking studies performed on the relativistic analogous models. Therefore, for $_{i} > 0$, we have an initially symmetry restored phase in both and directions, while $_{i} < 0$ corresponds (at zero tem perature) to symmetry broken phases for both and .

We require the model (1) to be overall bounded from below, which then gives the constraint condition on the two-body interaction terms, g > 0, g > 0 and $g g > 9g^2$. This is the same condition in posed on the analogous relativistic problem [1, 2]. This boundness constraint will be observed in all our results below. Non-trivial phase transitions can emerge for negative values of the cross-coupling g, which is allowed by the above boundness constraint. This was shown in diverse instances to be the case in the relativistic analogous models [1, 3].

In the analysis below, we will also restrict, for simplicity, to an initial symmetry restored phase (at zero temperature) for both elds, $_{i} > 0$ and leave the symmetry broken case for the Bose-E instein condensation problem studied in Sec. 3. The phase structure of the model is then determined by the sign of the temperature dependent one-body terms, $_{i}(T) = _{i} + _{i}$, where $_{i}$ is the eld temperature dependent self-energy. We look for an intermediate (reentrant) phase at some interval of temperature and parameters in which the symmetry in one of the eld directions is broken. This analysis can in principle be carried out within perturbation theory, as described in

a companion paper $[\beta]$, which indeed shows the possibility of appearance of reentrant phases for a system of hard-core particles described by (1). However, the results in [2, 3] also shows that as the (perturbative) tem perature corrections are considered for the twobody term s, g (T); g (T), these e ective couplings run to negative values above som e tem perature T_{neg}. This then violates the initial condition of boundness for the potential when considering them odel in equilibrium in a therm albath with tem perature $T > T_{neg}$. At the leading order perturbative calculation and assuming = = and m == m, T_{neq} / min 12 $=(2m^{3})g = (5g^{2} + 9g^{2}); 12$ $=(2m^{3})q = (5q^{2} + 9q^{2})$. This is rem in iscent of the breakdown of perturbation theory in quantum eld theory at nite tem perature, which is well known in relativistic models (see e.g. [7] and references in there). Nonperturbative methods are then called for a proper interpretation of the results and to con m that the appearance of reentrant phases in our m odel is not just an artifact of perturbation theory.

The problem of the self-couplings running to negative values at high tem peratures can be solved, e.g., by resumming all leading order bubble corrections to the couplings. This is naturally done in the context of the renormalization group, by solving the ow equations for all couplings and parameters of the model. A simpler and equivalent approach was also shown in [2], where this resummation is also accomplished by solving a set of self-consistent hom ogeneous linear equations for all elective couplings, g (T);g (T) and g(T), and the result of these equations feeded back in the equations for the elective one-body terms, $_{i}(T)$. We refer the interested reader to [2] for the details and we give here only the main results of this approach.

We consider, for illustrative purposes, the parameters (at T = 0): g = 210 ¹⁵eV ²; g = 10 ¹⁶eV ², m ' m = 1GeV and = = 1 neV. These values of couplings and masses could for instance be representative of some dilute Bose gas atom ormolecule (see next section). The temperature and the tree-level value of g are then changed and we look for regions of sym metry broken phase in the eld direction (for the values of parameters considered, it easy to show that the sym metry remains always restored in the direction [3]). For all values of parameters and temperature considered we check the boundness condition extended for the elective couplings (temperature dependent), as obtained by the ow equations described previously. That is, g (T)g (T) > 9g²(T), or that the ratio R^{NR}(T) = g (T)g (T)=[9g²(T)] > 1. The resulting phase diagram, as a function of g at T = 0 and the (log of the) temperature, is shown in Fig. 1.

Fig. 1 shows clearly the possibility of reentrant phases in the system, through an inverse symmetry breaking, in direction. For instance, for g (T = 0) = 10¹⁶eV², we nd a reentrant symmetry broken phase starting at the temperature $T_{c;}^{(ISB)}$, '3:4 10⁴eV (or 4 K) and ending at $T_{c;}^{(SR)}$, '1:4 10²eV (or 161 K), through symmetry restoration. In this region of temperature and parameters, $R^{NR}(T) > 1$ and the (e ective) potential is still bounded from below. In the direction there is no symmetry breaking or reentrant phases at any temperature for the parameters considered.




Figure 1. The phase diagram of the system in term s of g and tem perature, for the parameters considered in the text. The dark gray region denotes a reentrant phase with symmetry breaking (SB) in the direction, (T) < 0. The region below it, in light gray, is the unstable region, R^{NR} (T) < 1 and above it is a symmetry restored (SR) phase, (T) > 1. Tem perature is given in units of eV.

3. A pplication to a C oupled T wo-Species D ilute and H om ogeneous A tom ic Bose G as M odel

Let us now consider the case of model (1) as describing two coupled Bose gases of xed densities and , respectively. The model could then be describing a system composed by a mixture of coupled atom ic gases, like the ones recently produced [8], with same chemical element in two dierent hyper ne states, or even two dierent m ono-atom ic B ose gases in the hom ogeneous case [9]. Here and are then explicitly chem ical potentials added in the grand-canonical form alism to ensure the xed densities for each Bose atom gas. Here we start describing the system in the broken phase in both and directions. Therefore, ! and ! in (1) and these chemical potentials are taken as positive quantities and with their values determ ined by the usual therm odynam ic relation, in term s of the pressure P (T; ;),

$$= \frac{(P (T; ;))}{(P (T; ;))}; = \frac{(P (T; ;))}{(P (T; ;))};$$
(2)

where the pressure is de ned as the negative of the e ective potential com puted at its m inim a (which is the therm odynam ic free energy of the system),

P P (T; ;) =
$$V_{e}$$
 (T; ₀; ₀) _{0 = m; 0 = m}; (3)

where m and m are the values of 0 and 0 that extrem izes (corresponding to a minimum of) the elective potential,

Symmetry A spects in Nonrelativistic Multi-Scalar Field Models and Application

$$\frac{@V_{e}(T; 0; 0)}{@0} = m; 0 = m = 0; \quad \frac{@V_{e}(T; 0; 0)}{@0} = m; 0 = m = 0: (4)$$

The elds around the vacuum expectation values, $h = {}_{0} = 2$ and $h = {}_{0} = 2$, and i = eds around the vacuum expectation values, $h = {}_{0} = 2$ and $h = {}_{0} = 2$, and i = eds around i = eds around the one-bop approximation in a standard computation of quantum eld theory at nite temperature (for the one-eld case, see for instance [10]). At the tree-level, m and m are given in terms of the minima of the potential in (1),

$${}^{2}_{m} = 6 \frac{g}{g} \frac{3g}{g} \frac{3g}{9g^{2}}; {}^{2}_{m} = 6 \frac{g}{g} \frac{3g}{g} \frac{3g}{9g^{2}};$$
 (5)

At nite temperature, the equations for $_{m}$ and $_{m}$ are given by analogous expressions to (5), but in term softhe e ective chemical potentials instead, and , that are de ned by the solution of the self-consistent equations, = ; and = ;, given in term softhe and eld self-energies .

The explicit expression for the pressure at nite temperature, obtained from the e ective potential as described above and that follows from some length but straightforward calculation, is given by [11]

$$P(T; ;) = \frac{3}{2 (g g 9g^2)} g^2 2; + g^2 2;$$

+ 6g(;; ;)] $\frac{Z}{(2)^3} A_+ + A_{0} A_+$

where

$$A^{2} = \frac{H G + H G}{2} \qquad \frac{1}{2} H G \qquad H G^{2} + 4g^{2} g^{2} g^{2} g^{2} G G \qquad (7)$$

and

$$H = \frac{q^2}{2m} + \frac{g}{3} \frac{2}{m}; \quad G = \frac{q^2}{2m}; \quad H = \frac{q^2}{2m} + \frac{g}{3} \frac{2}{m}; \quad G = \frac{q^2}{2m}; \quad (8)$$

H $_{\rm i}$ and G $_{\rm i}$ denote the H iggs and G oldstone m odes, respectively, for each $\,$ eld in the broken phase.

From (2) and after some algebra to eliminate the dependence of these expressions on the chemical potentials, we ind the expressions relating the total densities and with (the condensate densities) m and m, as given by [11]

$$= \frac{\frac{2}{m}}{2} - \frac{1}{2} \frac{1}{2} \frac{d^{3}q}{(2)^{3}} \frac{d^{3}q}{d^{2}} + \frac{d^{3}q}{d^{2}$$

and

$$= \frac{\frac{2}{m}}{2} - \frac{1}{2} \frac{1}{2} \frac{d^{3}q}{(2)^{3}} \frac{d^{4}}{(2)^{3}} + \frac{1}{2} + 2n_{A_{+}} + \frac{dA_{-}}{(2)^{3}} + 2n_{A_{-}} +$$

where $n_A = 1 = [exp(A) \ 1]$ and the partial derivatives of A with respect to and are dened by

$$\frac{@A_{+}}{@} = \frac{H + G}{4A_{+}} \begin{cases} 2 & H & G & H & G & 2g^{2} \frac{2}{m} \frac{2}{m} G & 7\\ H & G & H & G & 2g^{2} \frac{2}{m} \frac{2}{m} G & 6\\ H & G & H & G & 2g^{2} \frac{2}{m} \frac{2}{m} G & 6\\ H & G & H & G & 2g^{2} \frac{2}{m} \frac{2}{m} G & 6\\ H & G & H & G & 2g^{2} \frac{2}{m} \frac{2}{m} G & 7\\ H & G & H & 2g^{2} \frac{2}{m} \frac{2}{m} G & 7\\ H & G & H & 2g^{2} \frac{2}{m} \frac{2}{m} G & 7\\ H & G & H & 2g^{2} \frac{2}{m} \frac{2}{m} G & 7\\ H & G & H & 2g^{2} \frac{2}{m} \frac{2}{m} G & 7\\ H & G & H & 2g^{2} \frac{2}{m} \frac{2}{m} G & 7\\ H & G & H & 2g^{2} \frac{2}{m} \frac{2}{m} G & 7\\ H & G & H & 2g^{2} \frac{2}{m} \frac{2}{m} G & 7\\ H & G & H$$

The coupled equations (9) and (10) give completely the phase diagram for the condensates $_{\rm m}$ and $_{\rm m}$ as a function of the temperature and the densities. A qualitative analysis of the phase structure is also possible to be deduced already at this level from equations (9), (10) and (11). Note that for g = 0, from (7) we obtain that A_{+}^{2} (g = 0) = H G , A^{2} (g = 0) = H G , and we obtain the Bogoliubov spectrum [6] for each eld in the uncoupled case. Also, (9) and (10) decouples and we obtain as a result, for example for ,

$$= \frac{\frac{2}{m}}{2} + \frac{1}{2} \frac{d^{3}q}{(2)^{3}} + \frac{\frac{q^{2}}{2m} + \frac{g}{m}}{\frac{q^{2}}{2m} + \frac{g}{m}} + \frac{g}{m} + \frac{g}{m}}{\frac{q^{2}}{2m} + \frac{g}{m}} + \frac{g}{m} + \frac{g}{m}} + \frac{g}{m} + \frac{g}{m} + \frac{g}{m} + \frac{g}{m}} + \frac{g}{m} + \frac{g}{m} + \frac{g}{m} + \frac{g}{m}}{\frac{q^{2}}{2m} + \frac{g}{m} + \frac{g}{m}} + \frac{g}{m}} + \frac{g}{m} + \frac{g}{m} + \frac{g}{m} + \frac{g}{m}} + \frac{g}{m} + \frac$$

with analogous equation for . Taking (12) at the critical point, $T = T_{c;}$, we have that $_{m}$ ($T = T_{c;}$) = 0, since the condensate density at T_{c} vanishes and (12) gives = $^{R} d^{3}q = (2)^{3} [\exp (q^{2} = (2m T_{c;})) 1]^{1}$, or inverting it, $T_{c;} = 2 = m [= (3=2)^{2=3}$, where (3=2)' 2:612. This is the standard result for the critical temperature of an hom ogeneous ideal B ose gas. This result emerges because of the level of approximation we are considering. It is only modied by corrections due to the self-interactions through nonperturbative methods and it requires at least second order corrections in the selfenergy (see, for instance, [12] and references therein).

Note also that, at the level of approximation we are considering, from the equations (7), (9), (10) and (11), if any of the elds go above the transition point (either $_{m} = 0$

or $_{m} = 0$) the two equations (9) and (10) also decouple, becoming independent of each other, since the cross-coupling term in (7) and (11) always appears multiplying both $_{m}$ and $_{m}$. As a result, no reentrant phase at high temperatures seem s to be possible here. A computation performed in the restored phase case (similar to the one done in Sec. 2) also seem s to con m this result. This comes about as a consequence of the strong temperature dependence introduced by the chemical potentials through the relation (2) in both the broken and symmetric cases. In the broken (BEC) phase, it is also seen that the system exhibits a small dependence on the cross-coupling term (at one-bop order it is even insensitive to the sign of g). A throughout analysis of the phase structure coming from the coupled set of equations (9) and (10), including higher order term s, will be presented elsewhere [11].

A cknow ledgm ents

The authors were partially supported by CNPq-Brazil. ROR was also partially supported by FAPERJ.W e also would like the organizers of the QFEXT05 workshop in Barcelona, Spain, for the enjoyable conference atm osphere where this work has been presented.

References

- [1] Pinto M B and Ram os R O 2000 Phys. Rev. D 61 125016 (Preprint hep-ph/9912273)
- [2] Pinto M B, Ramos R O and Parreira J E 2005 Phys. Rev. D 71 123519 (Preprint hep-th/0506131)
- [3] Pinto M B and Ram os R O 2006 J. Phys. A: Math. Gen. 39 6649 (Preprint cond-mat/0605508)
- [4] Goldenfeld N Lectures on Phase Transitions and The Renorm alization G roup, Frontiers in Physics, V. 85 (Addison-Wesley, NY, 1992)
- [5] Schupper N and Shnerb N M 2005 Phys. Rev. E 72 046107
- [6] Courteille Ph W, Bagnato V S and Yukalov V I 2001 Laser Phys. 11 659; D alfovo F, G iorgini S, Pitaevskii L P and Stringari S 1999 Rev. M od. Phys. 71 463
- [7] Gleiser M and Ram os R O 1993 Phys. Lett. B 300 271 (Preprint hep-ph/9211219)
- [8] M yatt C J et al. 1997 Phys. Rev. Lett. 78 586; M atthews M R et al. 1998 Phys. Rev. Lett. 81 243
- [9] Pu H and Bigelow N P 1998 Phys. Rev. Lett. 80 1130; A o P and Chuis T 1998 Phys. Rev. A 58 4836
- [10] Haugset T, Haugerud H and Ravndal F 1998 Ann. Phys. (N.Y.) 266 27
- [11] Pinto M B, Ramos R O and Souza Cruz F F, Preprint cond-m at/0606041
- [12] de Souza Cruz F F, Pinto M B, Ram os R O and Sena P 2002 Phys. Rev. A 65 053613 (Preprint cond-m at/0112306); Kneur J-L, Pinto M B and Ram os R O 2002 Phys. Rev. Lett. 89 210403 (Preprint cond-m at/0207089); 2003 Phys. Rev. A 68 043615 (Preprint cond-m at/0207295)