

SIT-LP-06/11
hep-th/0611072
November, 2006

On Origin of Mass and Supersymmetry

KAZUNARI SHIMA ^{*} and MOTOMU TSUDA [†]

^a*Laboratory of Physics, Saitama Institute of Technology
Fukaya, Saitama 369-0293, Japan*

Abstract

We show that the nonlinear supersymmetric general relativity gives new insights into the origin of mass and elucidates the mysterious relations between the cosmology and the (low energy) particle physics.

^{*}e-mail: shima@sit.ac.jp

[†]e-mail: tsuda@sit.ac.jp

In the previous papers we have shown that Nonlinear Supersymmetric General Relativity (NLSUSY GR) [1] may give a new insight into the unsolved problems of the cosmology, e.g. the dark matter, the dark energy and the mysterious relations between the cosmology and the (low energy) particle physics, etc. [2]. In this article we study further these problems in self-contained and explicit arguments.

NLSUSY GR is obtained by extending the geometrical arguments of Einstein general relativity (EGR) to new (called *SGM* from composite viewpoints [1]) spacetime just inspired by NLSUSY, where *tangent spacetime* is specified not only by the ordinary SO(3,1) Minkowski coordinates x^a but also by the SL(2,C) Grassmann coordinates ψ turning subsequently to the dynamical Nambu-Goldstone (NG) fermion (called *superon* hereafter) of NLSUSY Volkov-Akulov (VA) model [3]. NLSUSY GR ($N = 1$ SGM) action [1] is written as the following geometrical form of (empty) SGM spacetime (of everything):

$$L(w) = \frac{c^4}{16\pi G} |w| (\Omega(w) - \Lambda), \quad (1)$$

where

$$|w| = \det w^a{}_\mu = \det(e^a{}_\mu + t^a{}_\mu(\psi)), \quad t^a{}_\mu(\psi) = \frac{\kappa^2}{2i} (\bar{\psi} \gamma^a \partial_\mu \psi - \partial_\mu \bar{\psi} \gamma^a \psi). \quad (2)$$

$w^a{}_\mu(x)$, $\Omega(w)$, $s_{\mu\nu} \equiv w^a{}_\mu \eta_{ab} w^b{}_\nu$ and $s^{\mu\nu}(x) \equiv w^\mu{}_a(x) w^{\nu a}(x)$ are the fundamental unified quantities defining SGM spacetime [4, 5], i.e. the (composite) vierbein, the scalar curvature and the unified metric tensors of SGM spacetime. $e^a{}_\mu$ for the local SO(3,1) and $t^a{}_\mu(\psi)$ for the local SL(2,C) are the ordinary vierbein of EGR and the mimic vierbein of the stress-energy-momentum tensor of superon $\psi(x)$, whose dynamical parallel alignments for the spacetime orientations of SGM spacetime are encoded in $w^a{}_\mu(x)$. G is the Newton gravitational constant, Λ is a (*small*) cosmological term. Remarkably κ , which is the arbitrary constant of NLSUSY VA model with the dimension $(length)^2$ and subsequently becomes the strength of the superon-vacuum coupling constant in the (low energy theorem) particle physics viewpoint, is now related to G and Λ by $\kappa^2 = (\frac{c^4 \Lambda}{8\pi G})^{-1}$ in the NLSUSY GR (SGM) scenario. NLSUSY GR action (1) is invariant at least under [new NLSUSY] \otimes [local GL(4, R)] \otimes [local Lorentz] \otimes [local spinor translation] for spacetime symmetries and [global SO(N)] \otimes [local U(1) N] for the internal symmetries in N (-superons ψ^i ($i = 1, 2, \dots, N$))-extended NLSUSY GR [4].

NLSUSY GR is unstable due to the NLSUSY structure of tangent space and decays spontaneously (called *Big Decay* of spacetime) to ordinary Riemann spacetime described by the Einstein-Hilbert action coupled with massless superon (matter) ψ of NLSUSY VA model, which is written formally as follows,

$$L(e, \psi) = \frac{c^4}{16\pi G} |e| \{R(e) - \Lambda + T(e, \psi)\}, \quad (3)$$

where $T(e, \psi)$ is the kinetic term and the gravitational interaction of superon. The second and the third terms produce (N -extended) NLSUSY VA action in asymptotic Riemann-flat ($e^a{}_\mu(x) \rightarrow \delta^a{}_\mu$) spacetime with $\kappa^2 = (\frac{c^4 \Lambda}{8\pi G})^{-1}$, which is different from SGM-flat ($w^a{}_\mu(x) \rightarrow \delta^a{}_\mu$) spacetime for (1).

As some NL theories can be recasted into equivalent linear (L) renormalizable theories, NLSUSY is also the case and the equivalence of $N = 1$ VA model to the free action of $N = 1$ scalar multiplet or of $N = 1$ axial vector multiplet are studied in detail [6]. The linearization of the highly nonlinear SGM action (3) may be inevitable to extract the physical contents of SGM, where the broken LSUSY is realized on the supermultiplet containing mass gaps. To investigate the mass (gap) generation mechanism in SGM scenario it is useful to go to asymptotic flat spacetime. We have shown [7] that (the more realistic) $N = 2$ NLSUSY VA model in terms of two superons (Majorana NG fermion fields) ψ^i ($i = 1, 2$), which corresponds to the SGM action (3) for $N = 2$ in asymptotic Riemann-flat spacetime, is (algebraically) equivalent to the spontaneously broken $N = 2$ LSUSY massless supermultiplet with component fields denoted by $(v_a, \lambda^i, \phi, F^I)$ ($I = 1, 2, 3$) for a U(1) gauge field, two Majorana spinor fields of $\underline{2}$ of SU(2), a complex scalar field and three auxiliary real scalar fields of $\underline{3}$ of SU(2); indeed, the $N = 2$ NLSUSY VA action,

$$L_{\text{VA}} = -\frac{1}{2\kappa^2}|w| = -\frac{1}{2\kappa^2} \left\{ 1 + t^a{}_a + \frac{1}{2}(t^a{}_a t^b{}_b - t^a{}_b t^b{}_a) + \cdots \right\} \quad (4)$$

with $|w| = \det w^a{}_b = \det(\delta^a{}_b + t^a{}_b)$ and $t^a{}_b = -i\kappa^2 \bar{\psi}^i \gamma^a \partial_b \psi^i$, is recasted into the equivalent spontaneously broken $N = 2$ LSUSY action

$$L_0 = \partial_a \phi \partial^a \phi^* - \frac{1}{4} F_{ab}^2 + \frac{i}{2} \bar{\lambda}^i \not{\partial} \lambda^i + \frac{1}{2} (F^I)^2 - \frac{1}{\kappa} \xi^I F^I, \quad (5)$$

where $F_{ab} = \partial_a v_b - \partial_b v_a$, the ξ^I are arbitrary real parameters for the induced global SO(3) (SU(2)) rotation satisfying $(\xi^I)^2 = 1$ and L_0 means the (almost) free action. As usual in the linearization of NLSUSY the Fayet-Iliopoulos (FI) F terms in (5) indicating spontaneous SUSY breaking with the vacuum expectation value $\langle F^I \rangle = \frac{\xi^I}{\kappa}$ are generated *automatically*. And all component fields of the LSUSY supermultiplet can be constructed as the composites of superons ψ^i (called SUSY invariant relations [6]), in such a way as the familiar LSUSY transformations [8] defined on the elementary supermultiplet are reproduced on the abovementioned composite supermultiplet in terms of the NLSUSY transformations, $\delta_\zeta \psi^i = \frac{1}{\kappa} \zeta^i - i\kappa \bar{\zeta}^j \gamma^a \psi^j \partial_a \psi^i$, on the constituent superons ψ^i [7]. For $\xi^1 = \xi^2 = 0$ the above supermultiplet is the ordinary $N = 1$ (axial) vector multiplet containing the U(1) gauge field.

We have tried to extract the cosmological implications of SGM scenario described by (1) and the subsequent Big Decay to Riemann spacetime with massless superon (3) igniting the Big Bang [5]. The variation of (3) with respect to $e^a{}_\mu$ gives the

equation of motion for $e^a{}_\mu$ depicted formally as follows

$$R_{\mu\nu}(e) - \frac{1}{2}g_{\mu\nu}R(e) = \frac{8\pi G}{c^4} \left\{ T_{\mu\nu}(e, \psi) - g_{\mu\nu} \frac{c^4 \Lambda}{8\pi G} \right\}, \quad (6)$$

where $T_{\mu\nu}(e, \psi)$ represents the stress-energy-momentum of superon including the gravitational interactions. Note that $-\frac{c^4 \Lambda}{8\pi G}$ can be interpreted as *the constant negative energy density of empty spacetime*, i.e. *the dark energy density* $-\rho_D$ responsible for the observed present acceleration of the universe. While, on tangent spacetime, the low energy theorem of the particle physics gives the following superon-vacuum coupling $\langle \psi^j{}_\alpha(q) | J^{k\mu}{}_\beta | 0 \rangle = i\sqrt{\frac{c^4 \Lambda}{8\pi G}} (\gamma^\mu)_{\alpha\beta} \delta^{jk} e^{iqx} + \dots$, where $J^{k\mu} = i\sqrt{\frac{c^4 \Lambda}{8\pi G}} \gamma^\mu \psi^k + \dots$ is the conserved supercurrent of NLSUSY VA action [9] and $\sqrt{\frac{c^4 \Lambda}{8\pi G}}$ is *the coupling constant g_{sv} of superon with the vacuum*. Further as shown before the right hand side of (6) for $N = 2$ flat case is essentially $N = 2$ NLSUSY VA action. And it is equivalent to the broken LSUSY action (5) which contains the SUSY breaking mass (gap) M_{SUSY} for the (composite) fields of the (massless) LSUSY supermultiplet

$$M_{SUSY}^2 \sim \langle F^I \rangle \sim \sqrt{\frac{c^4 \Lambda}{8\pi G}} \xi^I \sim g_{sv} \xi^I. \quad (7)$$

Suppose that among the LSUSY supermultiplet the stable and the lightest particle retains the mass of the order of the spontaneous SUSY breaking M_{SUSY} . And if the neutrino is such a particle $\lambda(x)$, i.e. $m_\nu^2 \sim \sqrt{\frac{c^4 \Lambda}{8\pi G}}$, then SGM predicts the observed value of the (dark) energy density of the universe and naturally explains the mysterious coincidence between m_ν and $\rho^{obs} \sim \rho_D$,

$$\rho^{obs} \sim (10^{-12} \text{GeV})^4 \sim m_\nu^4. \quad (8)$$

The tiny values of neutrino mass and of the energy density of the universe are the direct evidence of SUSY breaking in the spontaneous phase transition of SGM spacetime (Big Dickey) preceding the Big Bang.

In the above discussions the realistic mass generation mechanism in the LSUSY supermultiplet is speculative, for unfortunately the ordinary Yukawa interaction term is absent in the equivalent LSUSY theory (5). Recently we have found that the ordinary LSUSY invariant mass terms as well as the Yukawa interaction terms vanish identically under the SUSY invariant relation in the linearization of NLSUSY. Therefore we can add these terms to the linearized free action (5) *without violating the equivalence* and the invariance [10]. Now we can investigate explicitly the realistic mass generation mechanism in SGM scenario.

For simplicity and without loss of generality we consider $N = 2$, $D = 2$ SO(2) SUSY and $D = 2$ SGM. The NLSUSY VA model has the same form as (4) except

L_{VA} terminates at $\mathcal{O}(t^2)$. The equivalent $D = 2$ LSUSY theory [10] consists of the supermultiplet with component fields $(v^a, \lambda^i, A, \phi, F)$ and the following Lagrangian

$$L_0 = \frac{1}{2}(\partial_a A)^2 - \frac{1}{4}(F_{ab})^2 + \frac{i}{2}\bar{\lambda}^i \not{\partial} \lambda^i + \frac{1}{2}(\partial_a \phi)^2 + \frac{1}{2}F^2 - \frac{1}{\kappa}\xi F. \quad (9)$$

We constructed the action (9) based on the arguments in $N = 1, D = 2$ super-Yang-Mills theory. The $D = 2$ SUSY invariant relations for those component fields are constructed as composites of ψ^i in all orders as follows [10],

$$v^a = -\frac{i}{2}\xi\kappa\epsilon^{ij}\bar{\psi}^i\gamma^a\psi^j|w|, \quad (10)$$

$$\lambda^i = \xi \left[\psi^i|w| - \frac{i}{2}\kappa^2\partial_a\{\gamma^a\psi^i\bar{\psi}^j\psi^j(1 - i\kappa^2\bar{\psi}^k\not{\partial}\psi^k)\} \right], \quad (11)$$

$$A = \frac{1}{2}\xi\kappa\bar{\psi}^i\psi^i|w|, \quad (12)$$

$$\phi = -\frac{1}{2}\xi\kappa\epsilon^{ij}\bar{\psi}^i\gamma_5\psi^j|w|, \quad (13)$$

$$F = \frac{\xi}{\kappa}|w| - \frac{1}{8}\xi\kappa^3\Box(\bar{\psi}^i\psi^i\bar{\psi}^j\psi^j). \quad (14)$$

The most general LSUSY invariant (Yukawa) interaction and mass terms are given by

$$L_f = f(A\bar{\lambda}^i\lambda^i - \epsilon^{ij}\phi\bar{\lambda}^i\gamma_5\lambda^j + A^2F - \phi^2F - \epsilon^{ab}A\phi F_{ab}) \quad (15)$$

and

$$L_m = -\frac{1}{2}m(\bar{\lambda}^i\lambda^i - 2AF + \epsilon^{ab}\phi F_{ab}), \quad (16)$$

where f is an arbitrary constant with the dimension $(mass)^1$. Substituting the SUSY invariant relations from (10) to (14) into (15) and (16) we see they vanish identically, i.e. $L_f(\psi) = L_m(\psi) \equiv 0$ in $D = 2$ (and $D = 4$ as well). Therefore the most general form of the renormalizable LSUSY invariant theory equivalent to $N = 2, D = 2$ NLSUSY VA action is written as follows,

$$L_{\text{VA}} = L_0 + L_f + L_m, \quad (17)$$

which is equipped with the self-contained spontaneous SUSY breaking.

Now at the tree level we can easily see the mediation of mass generation in the spontaneous SUSY breaking from the spacetime origin for NLSUSY to the (Yukawa-)Higgs origin for LSUSY as follows. For simplicity and corresponding to the chiral

LSUSY supermultiplet we put $L_m = 0$. The configuration of the scalar fields for the vacuum determines the contents of the LSUSY theory. The potential is given by

$$V(A, \phi, F) = -\frac{1}{2}F^2 + \frac{\xi}{\kappa}F - f(A^2F - \phi^2F). \quad (18)$$

Substituting the field equation for the auxiliary field, $F = \frac{\xi}{\kappa} - f(A^2 - \phi^2)$, into (18) we obtain

$$V(A, \phi) = \frac{1}{2} \left\{ f(A^2 - \phi^2) - \frac{\xi}{\kappa} \right\}^2 \geq 0. \quad (19)$$

In the (A, ϕ) plane, the massless mode and the massive mode appear parallel and perpendicular to the hyperbolic curve, $A^2 - \phi^2 = \frac{\xi}{f\kappa}$, respectively and some staffs of the LSUSY supermultiplet become massive. To see this we express the scalar fields A and ϕ as follows,

$$A(x) = -\sqrt{\frac{\xi}{f\kappa}} \{1 + \varphi(x)\} \cosh \theta(x) \quad (20)$$

$$\phi(x) = \sqrt{\frac{\xi}{f\kappa}} \{1 + \varphi(x)\} \sinh \theta(x) \quad (21)$$

and substitute these expressions into $L_0 + L_f$. We obtain after rescaling appropriately

$$\begin{aligned} L_0 + L_f = & \frac{1}{2}(\partial_a \varphi)^2 + \frac{1}{2}(\partial_a \theta)^2 - \frac{1}{4}F_{ab}^2 + \frac{i}{2}\bar{\lambda}^i \not{\partial} \lambda^i - \sqrt{\frac{f\xi}{\kappa}} \bar{\lambda}^i \lambda^i - \frac{2f\xi}{\kappa} \varphi^2 \\ & + \dots (\text{interaction terms}), \end{aligned} \quad (22)$$

where we assume $\varphi, \theta \ll 1$ for simplicity. We see from Eq.(22) that the fermions λ^i and the radial scalar mode φ get the following masses in $D = 2$ SGM scenario with $\kappa^2 = (\frac{c^4 \Lambda}{8\pi G})^{-1}$,

$$m_{\lambda^i}^2 = m_{\varphi}^2 = 4 \sqrt{\frac{c^4 \Lambda}{8\pi G}} f \xi, \quad (23)$$

while θ and v_a remain massless. These arguments show that the relation (7) and (8) hold in $D = 2$ and $D = 4$ as well, provided $f\xi \sim \mathcal{O}(1)$. (The same order contribution of the see-saw mechanism are anticipated [11].) These results show explicitly the mediation of mass generation in the spontaneous SUSY breaking in SGM scenario from the spacetime origin in NLSUSY to (Yukawa-)Higgs origin in LSUSY. We can perform the same arguments for $D = 4$ N -extended SUSY by using the chiral representations (for fermions), where the induced global $SU(N)$ internal symmetry are formulated manifestly and compactly as shown for $D = 4$ $N = 2$ [7]. The large mass scales and the compact (broken) gauge d.o.f. necessary for constructing the realistic model will appear through the linearization of $T_{\mu\nu}(e, \psi)$ with the large N which contains the mass scale Λ^{-1} and the higher order terms with ψ . NLSUSY GR with extra spacetime dimensions equipped with the Big Decay and the spontaneous compactification is interesting and in the same scope.

References

- [1] K. Shima, *Phys. Lett.* **B501** (2001) 237.
K. Shima, *European Phys. J.* **C7** (1999) 341.
- [2] K. Shima and M. Tsuda, *PoS HEP2005* (2006) 011.
- [3] D.V. Volkov and V.P. Akulov, *Phys. Lett.* **B46** (1973) 109.
- [4] K. Shima and M. Tsuda, *Phys. Lett.* **B507** (2001) 260.
- [5] K. Shima and M. Tsuda, *Class. and Quantum. Grav.* **19** (2002) 5101.
K. Shima, M. Tsuda and M. Sawaguchi, *Int. J. Mod. Phys.* **E13** (2004) 539.
- [6] E.A. Ivanov and A.A. Kapustnikov, *J. Phys.* **A11** (1978) 2375.
M. Roček, *Phys. Rev. Lett.* **41** (1978) 451.
T. Uematsu and C.K. Zachos, *Nucl. Phys.* **B201** (1982) 250.
K. Shima, Y. Tanii and M. Tsuda, *Phys. Lett.* **B525** (2002) 183.
- [7] K. Shima, Y. Tanii and M. Tsuda, *Phys. Lett.* **B546** (2002) 162.
- [8] J. Wess and J. Bagger, *Supersymmetry and Supergravity* (Princeton University Press, Princeton, New Jersey, 1992).
- [9] K. Shima, *Phys. Rev.* **D20** (1979) 574.
- [10] K. Shima and M. Tsuda, hep-th/0611051.
- [11] T. Yanagida, *Workshop on Grand Unified Theories*, KEK report 79-18 (1979).
M. Gell-Mann, P. Ramond, R. Slank, *Supergravity* (North Holland, Amsterdam, 1979).