

U SC -91/009

# Supersymmetric Gelfand-Dickey Algebra

K. Huitu and D. Nemeschansky

Department of Physics  
University of Southern California  
Los Angeles, CA 90089

We study the classical version of supersymmetric  $W$ -algebras. Using the second Gelfand-Dickey Hamiltonian structure we work out in detail  $W_2$  and  $W_3$ -algebras.

7/91

---

Supported by a grant from the Finnish Cultural Foundation

Supported by the Department of Energy grant DE-FG 03-84ER-40168

## 1. Introduction

Recently  $W$ -algebras have attracted a lot of attention. They play an important role both in integrable systems and conformal field theories [1], [2]. The Virasoro algebra and its  $W$ -algebra generalization appear naturally in integrable systems [3] of KdV-type [4], [5], [6]. The  $W$ -algebras were originally introduced by Zamolodchikov [7]. In addition to the Virasoro algebra generated by the stress tensor, the  $W$ -algebras contain higher integer spin currents. The supersymmetric versions of these algebras contain in addition to the integer spin currents also integer fermionic currents. A lot of work has been done in trying to find  $W$ -algebras with  $N = 1$  supersymmetry [8], [9], [10].

For a string theory application it is well known that models with  $N = 2$  supersymmetry are the most desirable since they give rise to  $N = 1$  supersymmetry in space-time. The  $W$ -algebras with  $N = 2$  supersymmetry were studied in [11] through Hamiltonian reduction of non-compact Lie algebras. In this paper we consider also supersymmetric  $N = 2$   $W$ -algebras. We consider superdifferential operators whose coefficients can be identified with conformal fields. Using the second Gelfand-Dickey Hamiltonian structure [12] we generate the classical version of extended supersymmetric  $W$ -algebra.

The paper is organized as follows. In section two we study the algebra of pseudodifferential operators. The minimal  $N = 2$  algebras are considered in section three. In section four we extend our analysis to  $W_3$ -algebras. Chapter five contains our conclusions.

## 2. Supersymmetric Pseudodifferential Operators

We start our analysis by considering the algebra of supersymmetric pseudodifferential operators [13]. Throughout the paper we use a  $N = 1$  supersymmetric description. With this in mind we introduce an anticommuting coordinate  $\theta$ , with  $\theta^2 = 0$ . We work in two-dimensional superspace and use complex coordinates  $z = (z; \theta)$ . The coordinate  $z$  is even and the anticommuting coordinate  $\theta$  is odd. In general we set  $X = 0$  (respectively 1) if  $X$  is even (respectively odd). The superderivative  $D$  is the square root of the ordinary derivative

$$D = \frac{\partial}{\partial z} + \frac{\partial}{\partial \theta} ; \quad (2.1)$$

and it acts as an odd superderivative

$$\begin{aligned} D u &= \alpha + 1 \\ D(uv) &= (D u)v + (-1)^{\alpha} uD v; \end{aligned} \quad (2.2)$$

The supercommutator is defined by

$$[X; Y] = XY - (-)^{XY} YX \quad (2.3)$$

The differential operators that we consider have the form

$$L = u_n(z; )D^n + u_{n-1}(z; )D^{n-1} + \dots + (z; )D + u_0(z; ) : \quad (2.4)$$

The multiplication rule is provided by the Leibniz rule,

$$\begin{aligned} D^n u &= \sum_{k=0}^n (-1)^{nk} (D^{n-k} u) D^k ; \\ X u_j D^j \cdot X v_i D^i &= \sum_{k=0}^{\min(j,i)} (-1)^{jk} u_j (D^{j-k} v_i) D^{k+i} ; \end{aligned} \quad (2.5)$$

Above we have used form multiplication to distinguish  $D^n u$  from  $(D^n u)$ . The coefficients  $\binom{j}{k}$  in eq. (2.5) are the superbinomial coefficients,

$$\binom{j}{k} = \begin{cases} 0 & \text{for } k > j \text{ and for } (j;k) \equiv (0;1) \pmod{2} \\ \binom{j+2}{k+2} & \text{for } k \leq j, (j;k) \not\equiv (0;1) \pmod{2} \end{cases}, \quad (2.6)$$

where  $\binom{j}{k}$  is the ordinary binomial coefficient.

Next let us introduce the ring of formal pseudodifferential operators

$$L = u_n(z; )D^n + u_{n-1}(z; )D^{n-1} + \dots + (z; )D + u_0(z; ) + \sum_{k=1}^{\infty} u_k(z; )D^k : \quad (2.7)$$

We set

$$L_+ = L; \quad L_- = \sum_{k=1}^{\infty} u_k(z; )D^k; \quad \text{res}L = u_0(z; ) ; \quad (2.8)$$

where  $\text{res}$  stands for the residue. The multiplication rule of formal pseudodifferential operators is also given by the Leibniz rule. We can introduce a bracket for the pseudodifferential operators

$$[L_1; L_2] = L_1 L_2 - (-)^{L_1 L_2} L_2 L_1 : \quad (2.9)$$

From the definition of the bracket we see that the commutator  $[L_1; L_2]$  of two pseudodifferential operators  $L_1$  and  $L_2$  with degrees  $n_1$  and  $n_2$  respectively has degree  $n_1 + n_2 - 1$ . The pseudodifferential operators with negative degree form an algebra with respect to the

bracket  $[ ; ]$ . This algebra is known as the Volterra algebra. Let  $X$  be an element of the Volterra algebra,

$$X = \sum_{i=1}^{\infty} D^{-i} x_i(z; ) : \quad (2.10)$$

Later when we use (2.10) only a finite number of  $x_i(z; )$ 's are non-zero. To each element  $X$  of the Volterra algebra we can associate a differential operator  $L$  via the mapping

$$(L; X) = \text{Tr}(LX) ; \quad (2.11)$$

where

$$\text{Tr}(LX) = \int_Z \text{res}(LX) dz d\bar{z} : \quad (2.12)$$

Let us consider differential operators with fixed degree  $n > 0$ . Without loss of generality we can set  $u_n(z; ) = 1$ ,

$$L = D^n + u_{n-1}(z; )D^{n-1} + \dots + (z; )D + u_0(z; ) : \quad (2.13)$$

For any functional  $f[u_0; \dots; u_{n-1}]$ , let  $X_f$  be the formal sum

$$X_f = \sum_{i=1}^{n-1} D^{-i} x_i(z; ) \quad \text{with } x_i = \frac{f}{u_{i-1}} : \quad (2.14)$$

Then for any pair of functionals  $f$  and  $g$  we introduce the Poisson bracket

$$\{f; g\} = \int_Z \text{res}(V_{X_f}(L) - X_g) dz d\bar{z} ; \quad (2.15)$$

where

$$V_{X_f}(L) = L(X_f - L) - (L - X_f)_+ - L : \quad (2.16)$$

The Poisson bracket defined by equation (2.15) is called the Gelfand-Dickey bracket of the second kind [12], [14]. In the next section we work out Gelfand-Dickey bracket for differential operators of degree three and five.

For the rest of the paper we restrict ourselves to differential operators of odd degree. These give rise to extended supersymmetric algebras. They act on functions  $f(z; )$ . By a change of the function

$$\begin{aligned} f(z; ) &= \exp \left( \frac{2}{n} \int_z^{Z_1} u_{n-1}(z_1; ) dz_1 \right) f(z; ) \quad n \text{ even} \\ f(z; ) &= \exp \left( \frac{2}{n-1} \int_{Z_1}^{Z_2} \int_{z_1}^{Z_2} u_{n-1}(z_1; ) dz_1 dz^0 \right) f(z; ) \quad n \text{ odd} : \end{aligned} \quad (2.17)$$

we can eliminate the coefficient  $u_{n-1}$  from the expansion of the differential operator (2.4). Therefore the differential operator  $L$  has the form

$$L = D^n + u_{n-2}(z; )D^{n-2} + \dots + (z; )D + u_0 : \quad (2.18)$$

Next let us consider the behavior of the differential operator under a superconformal transformation. A super analytic map  $z \mapsto z(z) = (\bar{z}(z; ); \tilde{z}(z; ))$  transforms the superderivative according to, e.g. [15]

$$D = (\bar{D} \tilde{z})\bar{D} + (\bar{D} z - \tilde{D} \tilde{z})\tilde{D}^2 : \quad (2.19)$$

A super analytic map is called a superconformal transformation when the superderivative transforms homogeneously,

$$D = (\bar{D} \tilde{z})\bar{D} : \quad (2.20)$$

From (2.19) we see that for a superconformal transformation

$$\bar{D} z - \tilde{D} \tilde{z} = 0 : \quad (2.21)$$

The superconformal tensor field  $f(z)$  is defined by the condition that  $f(z)d^{2h}$  is superconformally covariant. The conformal dimension of the field  $f$  is  $h$ . From the transformation law (2.20) we see that a superdifferential  $dz$  transforms as follows,

$$dz = (\bar{D} \tilde{z})dz \text{ or } \frac{dz}{dz} = \bar{D} \tilde{z} : \quad (2.22)$$

This means that the field  $f(z; )$  with conformal dimension  $h$  has the following transformation law

$$f(z) = f(\bar{z})(\bar{D} \tilde{z})^{2h} : \quad (2.23)$$

The superconformal tensors are the analogs of ordinary tensor fields  $f$  with conformal weight  $h$ , for which  $f(z) = f(\bar{z})(dz/dz)^h$ .

If we denote the space of functions with conformal dimension  $h$  by  $F_h$  one can show that the differential operator  $L$  maps the space  $F_{\frac{n-1}{2}}$  into the space  $F_{\frac{n+1}{2}}$ . Therefore we have

$$\tilde{L} = (\bar{D} \tilde{z})^{\frac{n-1}{2}} L (\bar{D} \tilde{z})^{\frac{n+1}{2}} : \quad (2.24)$$

This then allows to determine the transformation properties of the coefficients  $u_i(z; )$ . In the next sections we work out the transformation properties of these functions when the differential operator has degree three and five.

In eq. (2.18) we have set  $u_{n-1}(z; )$  to zero. From eq. (2.14) we see that the coefficient  $x_n(z; )$  of  $X_f$  is undetermined. This problem can be solved by demanding that  $\text{res}[L; X_f]$  vanishes. It turns out that this is equivalent to demanding that the coefficient of the term with degree  $n-1$  in  $V_{X_f}(L)$  vanishes. This concludes our general discussion.

$$3. L = D^3 + u_1 D + u_0$$

In this section we work out the details when the differential operator  $L$  is of degree three,

$$L = D^3 + u_1(z; )D + u_0(z; ) : \quad (3:1)$$

Following our general discussion let us consider

$$X_f = D^{-1} - \frac{f}{u_0} + D^{-2} - \frac{f}{u_1} + D^{-3}x_3 ; \quad (3:2)$$

where  $x_3$  is determined by the requirement that  $\text{res}[L; X_f] = 0$ : An explicit calculation then gives the condition

$$D x_3 = (D^2 \frac{f}{u_1}) + (D^3 \frac{f}{u_0}) - (D \frac{f}{u_0}) u_1 - (D u_1) \frac{f}{u_2} : \quad (3:3)$$

Using this condition we then calculate  $V_x(L)$  and we find

$$\begin{aligned} V_x(L) = & 2(D^3 \frac{f}{u_1}) + (D^4 \frac{f}{u_0}) + 2u_0 \frac{f}{u_1} + u_0(D \frac{f}{u_0}) - (D u_1) \frac{f}{u_1} \\ & (D^2 u_1) \frac{f}{u_0} - (D^2 \frac{f}{u_0}) u_1 - D + (-1)^{\frac{f}{u_0}} - (D^4 \frac{f}{u_1}) - (D^2 u_0) \frac{f}{u_0} \\ & u_0(D \frac{f}{u_1}) - 2u_0(D^2 \frac{f}{u_0}) + \frac{f}{u_1}(D u_0) + u_1(D^2 \frac{f}{u_1}) : \end{aligned} \quad (3:4)$$

The Gelfand-Dickey algebra of the second kind then has the form ,

$$\begin{aligned} ff(z; )g(z^0; )^0g &= \text{res}(V_{X_f}(L)X_g) \\ &= 2(D @_z \frac{f}{u_1}) + (@_z^2 \frac{f}{u_0}) + 2u_0 \frac{f}{u_1} + u_0(D \frac{f}{u_0}) \\ &\quad (D u_1) \frac{f}{u_1} - (D^2 u_1) \frac{f}{u_0} - @_z \frac{f}{u_0} u_1 - \frac{g}{u_1} (-1)^{\frac{g}{u_1}} \\ &+ @_z^2 \frac{f}{u_1} u_0 - D \frac{f}{u_1} + \frac{f}{u_1}(D u_0) + u_1 @_z \frac{f}{u_1} \\ &- (@_z u_0) \frac{f}{u_0} - 2u_0 @_z \frac{f}{u_0} - \frac{g}{u_0} (-1)^{\frac{g}{u_0} + \frac{g}{u_1}} \end{aligned} \quad (3:5)$$

Before we calculate the Poisson bracket of the fields  $u(z; )$  and  $u_0(z; )$  let us look how they transform under a superconformal transformation. From equation (2.24) we have

$$D^3 + u_1 D + u_0 = (D \tilde{\gamma})^2 (D^3 + u_1 D + u_0) (D \tilde{\gamma})^{-1} ; \quad (3:6)$$

where

$$\begin{aligned} u_1 &= \alpha_1 (\mathcal{D}^\sim)^2 \\ u_0 &= \alpha_0 (\mathcal{D}^\sim)^3 + \alpha_1 (\mathcal{D}^\sim) (\mathcal{D}^{2\sim}) + S ; \end{aligned} \quad (3.7)$$

The super Schwarzian derivative  $S$  of eq. (3.7) is given by

$$S(z; z) = \frac{\mathcal{D}^{4\sim}}{\mathcal{D}^\sim} - 2 \frac{\mathcal{D}^{3\sim} \mathcal{D}^{2\sim}}{\mathcal{D}^\sim \mathcal{D}^\sim} : \quad (3.8)$$

The above equations show that  $u_1(z; )$  transforms as a field with conformal dimension one. Therefore we can identify it with

$$J(z; ) = J(z) + G^2(z) ; \quad (3.9)$$

where  $G^2(z)$  is the superpartner of  $J(z)$ . The field  $u_0(z; )$  does not have the correct transformation property for a field with conformal dimension  $3=2$ . The second term is the one that causes us trouble. The Schwarzian derivative reflects the fact that there is a central term in the corresponding operator product expansion. We can eliminate the extra term from the transformation law by considering a linear combination of  $u_0(z; )$  and  $\mathcal{D} u_1(z; )$ . We find that

$$\hat{u}_0(z; ) = u_0(z; ) - \frac{1}{2} \mathcal{D} u_1(z; ) \quad (3.10)$$

has the correct transformation property. Therefore we can identify it with the stress tensor

$$T(z; ) = G^1(z) + T(z) : \quad (3.11)$$

Having identified the correct fields in our theory we now calculate their Poisson brackets. From the Poisson bracket of  $u_1(z; )$  with itself we can read off the Poisson brackets  $fJ(z); J(z^0)g$ ,  $fJ(z); G^2(z^0)g$  and  $fG^2(z); G^2(z^0)g$ . We find

$$\begin{aligned} fJ(z); J(z^0)g &= 2\partial_{z^0}(z - z^0) \\ fJ(z); G^2(z^0)g &= -2G^1(z)(z - z^0) \\ fG^2(z); G^2(z^0)g &= -2\partial_{z^0}(z - z^0) - 2T(z)(z - z^0) : \end{aligned} \quad (3.12)$$

From the Poisson bracket  $\hat{u}_0(z; )$  with  $u_1(z; )$  we obtain four new Poisson brackets  $fJ(z); G^1(z^0)g$ ,  $fT(z); J(z^0)g$ ,  $fG^1(z); G^2(z^0)g$  and  $fT(z); G^2(z^0)g$ . After some algebra we get

$$\begin{aligned} fJ(z); G^1(z^0)g &= -\frac{1}{2}G^2(z)(z - z^0) \\ fT(z); J(z^0)g &= \partial_{z^0}J(z - z^0) + J(z^0)\partial_{z^0}(z - z^0) \\ fG^1(z); G^2(z^0)g &= \frac{1}{2}\partial_{z^0}J(z - z^0) + J(z^0)\partial_{z^0}(z - z^0) \\ fT(z); G^2(z^0)g &= \frac{3}{2}G^2(z^0)\partial_{z^0}(z - z^0) + \partial_{z^0}G^2(z^0)(z - z^0) : \end{aligned} \quad (3.13)$$

We still need one more Poisson bracket. The Poisson bracket of the field  $\hat{u}(z; \ )$  with itself will give us the remaining unknown brackets of the component fields,  $fG^1(z); G^1(z^0)g$ ,  $fT(z); T(z^0)g$  and  $fT(z); G^1(z^0)g$ . They have the form,

$$\begin{aligned} fG^1(z); G^1(z^0)g &= \frac{1}{2}\partial_{z^0}(z - z^0) + \frac{1}{2}T(z^0)(z - z^0) \\ fT(z); T(z^0)g &= \frac{1}{2}\partial_{z^0}^3(z - z^0) + 2T(z^0)\partial_{z^0}(z - z^0) + \partial_{z^0}T(z^0)(z - z^0) \\ fT(z); G^1(z^0)g &= \frac{3}{2}G^1(z^0)\partial_{z^0}(z - z^0) + \partial_{z^0}G^1(z^0)(z - z^0) : \end{aligned} \quad (3.14)$$

The fields  $J(z); G^1(z); G^2(z)$  and  $T(z)$  form an  $N = 2$  supermultiplet. Furthermore the Poisson brackets that we have written down above are precisely the brackets of an  $N = 2$  supersymmetric algebra. We can summarize these Poisson brackets in a compact form using the  $N = 1$  superfields introduced above,

$$fJ(z; \ ); J(z^0; \ )g = -2(D^{03}) - 2T(z^0; \ ) \quad (3.15)$$

$$fT(z; \ ); J(z^0; \ )g = J(z^0; \ )D^{02} - \frac{1}{2}(D^0J)D^0 + (D^{02}J) \quad (3.16)$$

$$\begin{aligned} fT(z; \ ); T(z^0; \ )g &= \frac{1}{2}(D^{05}) - \frac{3}{2}T(z^0; \ )D^{02} \\ &\quad \frac{1}{2}(D^0T)D^0 - (D^{02}T) \end{aligned} \quad (3.17)$$

where

$$= ( \quad ^0)(z - z^0) \quad (3.18)$$

is the  $N = 1$  supersymmetric delta function and

$$D^0 = \frac{\partial}{\partial z^0} + \partial_{z^0} \quad (3.19)$$

In the next section we consider  $N = 2$  supersymmetric algebras with an extended symmetry.

#### 4. W<sub>3</sub>-algebra, $L = D^5 + u_3D^3 + u_2D^2 + u_1D + u_0$

In this section we use the Gel'fand-Dickey algebra to construct the classical version of the  $N = 2$  supersymmetric W<sub>3</sub>-algebra. The differential operator in this case is of order five,

$$L = D^5 + u_3(z; \ )D^3 + u_2(z; \ )D^2 + u_1(z; \ )D + u_0(z; \ ) : \quad (4.1)$$

Before we work out the Poisson bracket of the differential fields let us first identify the fields that transform covariantly. From the general discussion of section two we see that the differential operator of order five satisfies

$$D^5 + \alpha_3 D^3 + \alpha_2 D^2 + \alpha_1 D + \alpha_0 = (D^\gamma)^4 (D^5 + u_3 D^3 + u_2 D^2 + u_1 D + u_0) (D^\gamma)^3 : (4.2)$$

Comparing both sides in eq. (4.1) we obtain the following transformation laws,

$$\begin{aligned} u_3 &= \alpha_3 (D^\gamma)^2 \\ u_2 &= \alpha_2 (D^\gamma)^3 + \alpha_3 (D^\gamma) (D^2 \gamma) + 3S(z; z) \\ u_1 &= \alpha_1 (D^\gamma)^4 - \alpha_2 (D^\gamma)^2 (D^2 \gamma) + \alpha_3 (D^\gamma) (D^3 \gamma) + (D S(z; z)) \\ u_0 &= \alpha_0 (D^\gamma)^5 + 2\alpha_1 (D^\gamma)^3 (D^2 \gamma) + 2\alpha_2 (D^\gamma)^2 (D^3 \gamma) \\ &\quad + 2\alpha_3 (D^\gamma) (D^4 \gamma) - (D^2 \gamma) (D^3 \gamma) + 2(D^2 S(z; z)) \end{aligned} \quad (4.3)$$

As in our discussion in the previous sections only the field  $u_3$  transforms covariantly. It can be again identified with a super field with conformal dimension one, as we did in section two. The field with conformal dimension 3=2 is as before  $\hat{u}_2(z; ) = u_2 - 1/2(D u_3)$ . This can then be identified with the stress tensor  $T(z; )$ . It is easy to see that to construct a field with conformal dimension two we need a linear combination of  $u_1$ ,  $D u_2$  and  $D^2 u_3$ . The term  $(u_3(z; ))^2$  transforms by itself as a superconformal field with dimension 2. It turns out that we need to add also a term  $-2/9 u_3^2$  in order to cancel the unwanted terms in the brackets. We define

$$\hat{u}_1(z; ) = u_1(z; ) - \frac{1}{3} D u_2(z; ) - \frac{1}{3} (D^2 u_3(z; )) - \frac{2}{9} (u_3(z; ))^2 \quad (4.4)$$

We can identify this with the super field  $S(z; )$  with conformal dimension two

$$S(z; ) = S(z) + F^2(z) ; \quad (4.5)$$

where  $S(z)$  is the lowest component of an  $N = 2$  supermultiplet. It has conformal dimension two and  $F^2(z)$  is its superpartner with conformal dimension 5=2. To construct the highest component of the supermultiplet we need to construct a field that transforms with weight 5=2,

$$\begin{aligned} \hat{u}_0(z; ) &= u_0(z; ) - \frac{1}{2} (D u_1(z; )) - \frac{1}{2} (D^2 u_2(z; )) \\ &\quad + \frac{1}{6} (D^3 u_3(z; )) - \frac{4}{9} u_3(z; ) \hat{u}_2(z; ) : \end{aligned} \quad (4.6)$$

It is now clear that we can identify this with super field  $W(z; \bar{z})$  with conformal dimension  $5=2$ ,

$$W(z; \bar{z}) = F^1(z) + W(z); \quad (4.7)$$

where  $W(z)$  is a bosonic field with conformal dimension three and  $F^1$  is its superpartner with conformal dimension  $5=2$ . The fields that we have introduced above:  $S(z)$ ,  $F^1(z)$ ,  $F^2(z)$  and  $W(z)$  form an  $N = 2$  supermultiplet. The Poisson brackets of these fields are determined by eq. (2.15). For the differential operator of order five it is not very hard to see that  $V_{X_f}(L)$  is differential operator of order three,

$$V_{X_f}(L) = aD^3 + bD^2 + cD + e; \quad (4.8)$$

The coefficients  $a, b, c, e$  can be determined after a long and tedious calculation. The first coefficient  $a$  contains 27 differential terms. In the second term we find 22 differential terms. When we carry out the expansion we find 68 terms in  $c$  and 56 terms in  $e$ . The explicit form of these terms is not very illuminating. They, however allow us to determine the Poisson bracket of the supersymmetric  $W_3$  algebra, which we write down below.

Let us start by writing down the Poisson brackets that involve the  $U(1)$  current  $J(z; \bar{z})$  and its  $N = 2$  superpartner the stress tensor  $T(z; \bar{z})$

$$fJ(z; \bar{z}); J(z^0; \bar{z}^0)g = -6(D^0)^3 - 2T(z^0; \bar{z}^0) \quad (4.9)$$

$$fT(z; \bar{z}); J(z^0; \bar{z}^0)g = -J(z^0; \bar{z}^0)(D^0)^2 - \frac{1}{2}(D^0 J)(D^0) + (D^0)^2 J \quad (4.10)$$

$$fS(z; \bar{z}); J(z^0; \bar{z}^0)g = -2W(z^0; \bar{z}^0) \quad (4.11)$$

$$fW(z; \bar{z}); J(z^0; \bar{z}^0)g = 2S(z^0; \bar{z}^0)(D^0)^2 - \frac{1}{2}(D^0 S)(D^0) + 2(D^0)^2 S \quad (4.12)$$

$$\begin{aligned} fT(z; \bar{z}); T(z^0; \bar{z}^0)g = & -\frac{3}{2}(D^0)^5 - \frac{3}{2}T(z^0; \bar{z}^0)(D^0)^2 \\ & - \frac{1}{2}(D^0 T)(D^0) - (D^0)^2 T \end{aligned} \quad (4.13)$$

$$fS(z; \bar{z}); T(z^0; \bar{z}^0)g = -2S(z^0; \bar{z}^0)(D^0)^2 + \frac{1}{2}(D^0 S)(D^0) - \frac{3}{2}(D^0)^2 S \quad (4.14)$$

$$fW(z; \bar{z}); T(z^0; \bar{z}^0)g = -\frac{5}{2}W(z^0; \bar{z}^0)(D^0)^2 - \frac{1}{2}(D^0 W)(D^0) - 2(D^0)^2 W \quad (4.15)$$

The Poisson brackets involving only the super field  $S(z; \theta)$  and its  $N = 2$  superpartner  $W(z; \theta)$  are more complicated. After a long calculation we find

$$\begin{aligned}
& fS(z; \theta; S(z^0; \theta)g \\
&= + \frac{2}{3}(\mathbb{D}^{07}) + \frac{10}{3}S(z^0; \theta)(\mathbb{D}^{03}) + \frac{5}{3}(\mathbb{D}^0S)(\mathbb{D}^{02}) + \frac{5}{3}(\mathbb{D}^{02}S)(\mathbb{D}^0) \\
&+ (\mathbb{D}^{03}S) + \frac{4}{3}T(z^0; \theta)(\mathbb{D}^{04}) + \frac{8}{9}(\mathbb{D}^0T)(\mathbb{D}^{03}) + \frac{16}{9}(\mathbb{D}^{02}T)(\mathbb{D}^{02}) \\
&+ \frac{4}{9}(\mathbb{D}^{03}T)(\mathbb{D}^0) + \frac{2}{3}(\mathbb{D}^{04}T) - \frac{2}{27}(J(z^0; \theta))^2(\mathbb{D}^{03}) - \frac{1}{27}(\mathbb{D}^0(J^2))(\mathbb{D}^{02}) \\
&- \frac{1}{27}(\mathbb{D}^{02}(J^2))(\mathbb{D}^0) - \frac{1}{54}(\mathbb{D}^{03}(J^2)) - \frac{1}{54}(\mathbb{D}^0J)(\mathbb{D}^{02}J) - \frac{2}{9}W(z^0; \theta)J(z^0; \theta) \\
&+ 2T(z^0; \theta)S(z^0; \theta) + \frac{2}{3}T(z^0; \theta)(\mathbb{D}^0T) - \frac{4}{81}(J(z^0; \theta))^2T(z^0; \theta)
\end{aligned} \tag{4.16}$$

For the Poisson bracket of  $S(z; \theta)$  with  $W(z; \theta)$  we get

$$\begin{aligned}
& fW(z; \theta; S(z^0; \theta)g = \\
& \frac{2}{9}J(z^0; \theta)(\mathbb{D}^{06}) + \frac{1}{3}(\mathbb{D}^0J)(\mathbb{D}^{05}) - \frac{2}{3}(\mathbb{D}^{02}J)(\mathbb{D}^{04}) + \frac{4}{9}(\mathbb{D}^{03}J)(\mathbb{D}^{03}) \\
& \frac{2}{3}(\mathbb{D}^{04}J)(\mathbb{D}^{02}) + \frac{1}{6}(\mathbb{D}^{05}J)(\mathbb{D}^0) - \frac{2}{9}(\mathbb{D}^{06}J) + \frac{5}{3}W(z^0; \theta)(\mathbb{D}^{03}) \\
& (\mathbb{D}^0W)(\mathbb{D}^{02}) - \frac{2}{3}(\mathbb{D}^{03}W) + (\mathbb{D}^{02}W)(\mathbb{D}^0) - \frac{2}{27}T(z^0; \theta)J(z^0; \theta)(\mathbb{D}^{03}) \\
& \frac{5}{9}T(z^0; \theta)(\mathbb{D}^0J)(\mathbb{D}^{02}) - \frac{8}{27}(\mathbb{D}^0T)J(z^0; \theta)(\mathbb{D}^{02}) - \frac{1}{18}T(z^0; \theta)(\mathbb{D}^{02}J)(\mathbb{D}^0) \\
& \frac{1}{27}(\mathbb{D}^{02}T)J(z^0; \theta)(\mathbb{D}^0) - \frac{4}{27}(\mathbb{D}^{03}T)J(z^0; \theta) - \frac{10}{27}T(z^0; \theta)(\mathbb{D}^{03}J) \\
& \frac{8}{27}(\mathbb{D}^0T)(\mathbb{D}^{02}J) - \frac{10}{27}(\mathbb{D}^{02}T)(\mathbb{D}^0J) + \frac{1}{6}(\mathbb{D}^0T)(\mathbb{D}^0J)(\mathbb{D}^0) \\
& \frac{1}{18}J(z^0; \theta)(\mathbb{D}^0S)(\mathbb{D}^0) - \frac{7}{9}J(z^0; \theta)S(z^0; \theta)(\mathbb{D}^{02}) + \frac{1}{2}(\mathbb{D}^0J)S(z^0; \theta)(\mathbb{D}^0) \\
& + \frac{5}{18}(\mathbb{D}^0J)(\mathbb{D}^0S) - \frac{8}{9}(\mathbb{D}^{02}J)S(z^0; \theta) - \frac{1}{3}J(z^0; \theta)(\mathbb{D}^{02}S) \\
& + \frac{10}{9}W(z^0; \theta)T(z^0; \theta) - \frac{1}{81}(J(z^0; \theta))^2(\mathbb{D}^0J)(\mathbb{D}^0) + \frac{2}{81}(J(z^0; \theta))^3(\mathbb{D}^{02}) \\
& + \frac{4}{81}(J(z^0; \theta))^2(\mathbb{D}^{02}J)
\end{aligned} \tag{4.17}$$

There remains one more Poisson bracket to complete the supersymmetric  $W$ -algebra. A

long and tedious calculation gives the Poisson bracket of the super field  $W(z; \bar{z})$  with itself,

$$\begin{aligned}
& f_W(z; \bar{z}) ; W(z^0; \bar{z}^0) g = \\
& + \frac{1}{6} (\mathcal{D}^{09}) + \frac{5}{6} S(z^0; \bar{z}^0) (\mathcal{D}^{05}) + \frac{5}{12} (\mathcal{D}^0 S) (\mathcal{D}^{04}) + \frac{5}{6} (\mathcal{D}^{02} S) (\mathcal{D}^{03}) \\
& + \frac{1}{2} (\mathcal{D}^{03} S) (\mathcal{D}^{02}) + \frac{1}{4} (\mathcal{D}^{04} S) (\mathcal{D}^0) + \frac{1}{6} (\mathcal{D}^{05} S) - \frac{1}{81} (\mathcal{D}^0 T) (J(z^0; \bar{z}^0))^2 (\mathcal{D}^0) \\
& + \frac{2}{81} T(z^0; \bar{z}^0) J(z^0; \bar{z}^0) (\mathcal{D}^0 J) (\mathcal{D}^0) - \frac{7}{81} T(z^0; \bar{z}^0) (J(z^0; \bar{z}^0))^2 (\mathcal{D}^{02}) \\
& - \frac{4}{81} (\mathcal{D}^{02} T) (J(z^0; \bar{z}^0))^2 - \frac{8}{81} T(z^0; \bar{z}^0) (\mathcal{D}^{02} J) J(z^0; \bar{z}^0) + \frac{5}{9} T(z^0; \bar{z}^0) (\mathcal{D}^{06}) \\
& + \frac{5}{9} (\mathcal{D}^0 T) (\mathcal{D}^{05}) + \frac{10}{9} (\mathcal{D}^{02} T) (\mathcal{D}^{04}) + \frac{5}{9} (\mathcal{D}^{03} T) (\mathcal{D}^{03}) + \frac{5}{6} (\mathcal{D}^{04} T) (\mathcal{D}^{02}) \\
& + \frac{1}{6} (\mathcal{D}^{05} T) (\mathcal{D}^0) + \frac{2}{9} (\mathcal{D}^{06} T) - \frac{1}{54} (J(z^0; \bar{z}^0))^2 (\mathcal{D}^{05}) \\
& - \frac{1}{54} J(z^0; \bar{z}^0) (\mathcal{D}^0 J) (\mathcal{D}^{04}) - \frac{1}{27} J(z^0; \bar{z}^0) (\mathcal{D}^{02} J) (\mathcal{D}^{03}) \\
& + \frac{7}{108} J(z^0; \bar{z}^0) (\mathcal{D}^{03} J) (\mathcal{D}^{02}) - \frac{1}{108} J(z^0; \bar{z}^0) (\mathcal{D}^{04} J) (\mathcal{D}^0) \\
& + \frac{1}{27} J(z^0; \bar{z}^0) (\mathcal{D}^{05} J) - \frac{11}{72} (\mathcal{D}^0 J) (\mathcal{D}^{02} J) (\mathcal{D}^{02}) - \frac{1}{72} (\mathcal{D}^{02} J)^2 (\mathcal{D}^0) \\
& - \frac{1}{54} (\mathcal{D}^{02} J) (\mathcal{D}^{03} J) - \frac{5}{54} (\mathcal{D}^0 J) (\mathcal{D}^{04} J) + \frac{1}{24} (\mathcal{D}^0 J) (\mathcal{D}^{03} J) (\mathcal{D}^0) \\
& + \frac{1}{6} (\mathcal{D}^0 T)^2 (\mathcal{D}^0) - \frac{1}{54} T(z^0; \bar{z}^0) (\mathcal{D}^{02} T) (\mathcal{D}^0) + \frac{55}{54} T(z^0; \bar{z}^0) (\mathcal{D}^0 T) (\mathcal{D}^{02}) \\
& + \frac{14}{27} T(z^0; \bar{z}^0) (\mathcal{D}^{03} T) + \frac{2}{3} (\mathcal{D}^0 T) (\mathcal{D}^{02} T) + \frac{23}{18} S(z^0; \bar{z}^0) T(z^0; \bar{z}^0) (\mathcal{D}^{02}) \\
& - \frac{1}{18} (\mathcal{D}^0 S) T(z^0; \bar{z}^0) (\mathcal{D}^0) + \frac{1}{2} S(z^0; \bar{z}^0) (\mathcal{D}^0 T) (\mathcal{D}^0) + \frac{11}{18} (\mathcal{D}^{02} S) T(z^0; \bar{z}^0) \\
& + \frac{8}{9} S(z^0; \bar{z}^0) (\mathcal{D}^{02} T) + \frac{5}{18} (\mathcal{D}^0 S) (\mathcal{D}^0 T) + \frac{13}{18} J(z^0; \bar{z}^0) W(z^0; \bar{z}^0) (\mathcal{D}^{02}) \\
& - \frac{1}{18} J(z^0; \bar{z}^0) (\mathcal{D}^0 W) (\mathcal{D}^0) + \frac{1}{2} (\mathcal{D}^0 J) W(z^0; \bar{z}^0) (\mathcal{D}^0) - \frac{5}{18} (\mathcal{D}^0 J) (\mathcal{D}^0 W) \\
& + \frac{11}{18} (\mathcal{D}^{02} J) W(z^0; \bar{z}^0) + \frac{1}{3} J(z^0; \bar{z}^0) (\mathcal{D}^{02} W)
\end{aligned} \tag{4.18}$$

This completes our analysis of the  $N = 2$  supersymmetric  $W$ -algebra.

## 5. Conclusion

In this paper we have shown how to derive in a systematic way the classical version of the  $N = 2$  supersymmetric  $W$ -algebra. As an example of our technique we first considered the minimal  $N = 2$  model. Then we analyzed the  $N = 2$  version of the  $W_3$ -algebra. We

saw that to obtain the Poisson brackets required a lot of algebra. It is clear that our method generates the general  $W$ -algebra although the explicit form of the algebra may be very complicated.

It would be interesting to study the integrable perturbations that correspond to the differential operator (2.13). For the minimal models this has been carried out by Mathieu and Walton [16]. They found three integrable  $N = 2$  Korteweg-de-Vries equations. The generalization to  $W$ -algebras is not known. We hope to return to this question in future work.

After finishing this work we received a preprint [17], where similar results were obtained.

## R eferences

- [1] V .A Fateev and S .L .Lykyanov, Int. J. Mod. Phys. A 3 (1988) 507.
- [2] F .Bais, P .Bouwknegt, M .Surridge and K .Schoutens, Nucl. Phys. B 304 (1988) 348.
- [3] M .Jimbo and T .M iwa, Integrable systems in statistical mechanics, eds. G .D 'Ariano et.al. (World Scientific, Singapore 1985).
- [4] V .Drinfeld and V .Sokolov, Jour. Sov. Math. 30 (1985) 1975.
- [5] I .Bakas, Phys. Lett. 213B (1988) 313.
- [6] T .Inami, Y .Matsuo and I .Yamanaka, Phys. Lett. 215B (1988) 701.
- [7] A .Zamolodchikov, Teor. Mat. Fiz. 65 (1985) 347.
- [8] S .Komata, K .Mohri, H .Nohara, Tokyo preprint UT-Komaba 90-21 (July 1990).
- [9] JM .Figueroa-O'Farrill, E .Ram os, preprints K U L-T F-91/6 (March 1991), K U L-T F-91/4 (April 1991).
- [10] H .Lu, C .Pope, L .Romans and X .Shen and X .Wang, Phys. Lett. B , to appear.
- [11] D .Nevezhansky and S .Yankielowicz, USC-005-91-preprint, (1991).
- [12] I .Gelfand and L .Dickey, Preprint 136, IPM SSSR, Moscow 1988.
- [13] Yu .I .Manin, A .O .Radul, Commun. Math. Phys. 98 (1985) 65-77.
- [14] IM .Gelfand, I .Dorfman, Funct. Anal. Appl. 15 (1981) 173.
- [15] D .Friedan, in Superstrings, Santa Barbara workshop 1985, eds. M .Green, D .Gross
- [16] P .Mathieu and M .Walton, Phys. Lett. 254B (1991) 106.
- [17] T .Inami, H .Kanno, preprint YITP/K-928