

Correlation functions in super Liouville theory

E . Abdalla¹, M . C B . Abdalla²,

D . D alm azí², Koji Harada^{1(a)}

¹Instituto de Física, Univ. São Paulo, CP 20516, São Paulo, Brazil

²Instituto de Física Teórica, UNESP, Rua Pamplona 145,
CEP 01405, São Paulo, Brazil

We calculate three- and four-point functions in super Liouville theory coupled to super Coulomb gas on world sheets with spherical topology. We first integrate over the zero mode and assume that a parameter takes an integer value. After calculating the amplitudes, we formally continue the parameter to an arbitrary real number. Remarkably the result is completely parallel to the bosonic case, the amplitudes being of the same form as those of the bosonic case.

The matrix model definition of 2D-gravity has been proving to be very powerful in calculating correlation functions [1], although it seems difficult to generalize the results to supersymmetric theories. On the other hand, in the continuum approach (Liouville theory) [2{5] it is difficult to calculate correlation functions, while its supersymmetric generalization (super Liouville theory) [6{8] is well known. Recently, however, several authors [9{13] have exactly calculated correlation functions in the continuum approach to conformal matter fields coupled to 2D-gravity. (See also Ref. [14]). They have used a technique based on the integration over the Liouville zero mode, and their results agree with those obtained earlier in the discrete approach (matrix models). It is thus very urgent to extend the continuum method [15] to the supersymmetric case, i.e., superconformal matter fields coupled to 2D-supergravity.

The aim of this Letter is to calculate the three- and four-point functions in super Liouville theory coupled to superconformal matter with the central charge $\hat{c} < 1$, represented as super Coulomb gas [16]. Our approach is close to that of Di Francesco and Kutasov [10]. The result is remarkable and is very parallel to the bosonic case; it amounts to a redefinition of the cosmological constant and of the primary superelds, resulting the same amplitudes as those of the bosonic theory.

The relevant framework has been given by Distler, Bouček and Kawai [8]. With a translation invariant measure, the total action is given by $S = S_{SL} + S_M$,

$$S_{SL} = \frac{1}{4} \int^Z d^2 z \hat{E} \left(\frac{1}{2} \hat{D}_{SL} \hat{D}_{SL} - Q \hat{Y}_{SL} e^{+ \hat{S}_{SL}} \right); \quad (1)$$

$$S_M = \frac{1}{4} \int^Z d^2 z \hat{E} \left(\frac{1}{2} \hat{D}_M \hat{D}_M + 2i \epsilon_0 \hat{Y}_M \right); \quad (2)$$

where S_{SL} , S_M , are super Liouville and matter superelds respectively. (See Refs. [8,17]). The matter sector has the central charge $\hat{c}_m = 1 - 8 \frac{2}{\alpha'}$. The parameters Q and ϵ_0 are given by

$$Q = 2 \frac{q}{1 + \frac{2}{\alpha'}}; \quad \epsilon_0 = \frac{Q}{2} \frac{1}{2} \frac{q}{Q^2 - 4} = \frac{Q}{2} \quad \text{job}; \quad (3)$$

The (gravitationally dressed) primary supereld e_{NS} is given by

$$e_{NS}(z; k) = d^2 z \hat{E} e^{ik_M(z)} e^{(k)_S L(z)}; \quad (4)$$

$$(k) = \frac{1}{2} Q + j_0 : \quad (5)$$

Note that the expressions for (k) and (k) are the same in terms of Q and j_0 as those of the bosonic theory.

Screening charges in the matter sector are of the form $d^2 z e^{id_M(z)}$, where d are the two solutions of $\frac{1}{2}d(d - 2j_0) = \frac{1}{2}$. In this Letter, however, we will concentrate on the case without screening charges. The case with screening charges, N (4)-point functions and the inclusion of the Ramond sector will be discussed elsewhere.

We shall calculate three-point functions of the primary field e_{NS} on world sheets with spherical topology (without screening charges), that is,

$$\left[\prod_{i=1}^* Y^3 Z e_{NS}(z_i; k_i) \right] + \left[\prod_{i=1}^Z D_E^{-S_L}] D_E^{-M} \right] \left[\prod_{i=1}^Y Y^3 e_{NS}(z_i; k_i) e^S \right] : \quad (6)$$

Our first step is to integrate over the zero modes,

$$\begin{aligned} & \left[\prod_{i=1}^* Y^3 Z e_{NS}(z_i; k_i) \right] + \left[\prod_{i=1}^X X^3 k_i \right] h ! \\ A(k_1; k_2; k_3) &= (s) \left(\frac{i}{2} \right)^3 \left(\frac{j}{2} \right)^s \left[\prod_{i=1}^* Y^3 d^2 z_i e^{ik_i M(z_i)} e^{i S_L(z_i)} \right] \left[\prod_{i=1}^Z d^2 z e^{+ S_L(z)} \right]^s ; \end{aligned} \quad (7)$$

where h_0 denotes the expectation value evaluated in the free theory ($= 0$) and we have absorbed the factor $[_+ (-2)^3]^1$ into the normalization of the path integral. The parameter s is defined as

$$s = \frac{1}{+} Q + \left[\prod_{i=1}^X X^3 \right] : \quad (8)$$

In general, s can take any real value and there is no obvious way of calculating the path-integral. However, if we assume that s is a non-negative integer [8{13], this is just a free-field correlator. Under this assumption, we evaluate the path-integral, and formally continue s to non-integer values. For s non-negative integer,

$$\begin{aligned}
A(k_1; k_2; k_3) &= (s) \left(\frac{i}{2} \right)^3 \frac{i^s}{2^s} \\
&= (s) \left(\frac{i}{2} \right)^3 \frac{i^s}{2^s} Z_{ij}^s + \sum_{i=1}^s \sum_{j=1}^s \frac{i^s}{2^s} Z_{ij}^s \\
&= (s) \left(\frac{i}{2} \right)^3 \frac{i^s}{2^s} Z_{ij}^s + \sum_{i=1}^s \sum_{j=1}^s \frac{i^s}{2^s} Z_{ij}^s
\end{aligned}$$

We have divided by the $S^1_{L_2}$ volume by setting $z_1 = 0; z_2 = 1; z_3 = 1; \tilde{z}_2 = \tilde{z}_3 = 0$ and $\tilde{z}_1 = \tilde{z}$. The integral is a supersymmetric generalization of (B.9) of Ref. [18]. Alternatively, using $S_L = + + +$, we can write

$$A(k_1; k_2; k_3) = \left(\frac{s}{2} \right)^3 \frac{\frac{1}{2}^2}{\frac{1}{2}^2} \frac{z}{z} \frac{y^s}{y^s} d^2 z_i \frac{y^s}{y^s} \frac{j_i j^2 + j_j}{j_i j^2 + j_j} \frac{z_i j^2 + z_j j^2}{z_i j^2 + z_j j^2} \frac{j_i}{j_i} \frac{j_j}{j_j} \frac{D}{(0)} \frac{(z_1)}{(z_s)} \frac{E}{s(z_0)} : \quad (10)$$

This is non-vanishing only for s odd; we thus write $s = 2m + 1$. One may evaluate h^{+-+} and h^{++0} independently. Since the rest of the integrand is symmetric, one may write the result in a simple form by relabelling coordinates:

$$A(k_1; k_2; k_3) = i \left(\frac{1}{2}\right)^3 (-s)(s+1) \frac{1}{\frac{s}{2}^+} \frac{!}{I^m(;;)} \quad (11)$$

where

$$\text{and } = +_1, = +_2, = \frac{1}{2} +_2.$$

We now have to calculate $I^m(; ;)$. First of all, we observe that $I^m(; ;)$ should be symmetric in α and β : $I^m(; ;) = I^m(; ;)$. It is easy to check that when $m = 0$. The large- α and large- β behaviors are consistent with it, and it is physically natural because

the amplitude should be symmetric under the exchange of two external momenta. Thus $I^m(\ ; \ ; \)$ exhibits the following symmetry

$$I^m(\ ; \ ; \) = I^m(1 \ m \ ; \ ; \) \quad (13)$$

Thus we may write $I^m(\ ; \ ; \)$ in the following way

$$\begin{aligned} I^m(\ ; \ ; \) &= C_m(\ ; \ ; \) \sum_{i=0}^{2m} (1 + \dots + 2i) (1 + \dots + 2i) (\dots + (2i - 4m)) \\ &\quad \sum_{i=1}^{2m} \left(\frac{1}{2} + \dots + (2i - 1) \right) \left(\frac{1}{2} + \dots + (2i - 1) \right) \left(\frac{1}{2} + \dots + (2i - 4m - 1) \right) \end{aligned} \quad (14)$$

where $C_m(\ ; \ ; \)$ has the same symmetries as $I^m(\ ; \ ; \)$, and where $(x) = (1 \ x)$.

By looking at the large- behavior:

$$I^m(\ ; \ ; \) \sim 2m \cdot 2(2m+1) \cdot 4m \cdot (2m+1) \ ; \quad (15)$$

one can confirm that $C_m(\ ; \ ; \)$ is, as a function of , bounded as $j j! \ 1$ and analytic.

This means that $C_m(\ ; \ ; \)$ is independent of , and by symmetry, of as well; $C_m = C_m(\)$.

It is hard to calculate $C_m(\)$. For this purpose, it is useful to consider the simpler integral:

$$\begin{aligned} J^m(\ ; \ ; \ ; \) &= \int_0^{\infty} d^2 \int_0^{\infty} d^2 \int_0^{\infty} j_1^2 j_2^2 j_3^2 j_4^2 \dots j_i^2 j_{i+1}^2 \dots j_{2m+1}^2 \\ &\quad j_1^4 j_2^4 j_3^4 \dots j_i^4 \dots : \end{aligned} \quad (16)$$

By using similar arguments, one may obtain

$$\begin{aligned} &J^m(\ ; \ ; \ ; \) \\ &= C_m(\ ; \) \sum_{i=0}^{m-1} (1 + \dots + 2i) (1 + \dots + 2i) (1 + \dots + (2i - 4m + 2)) \\ &\quad \sum_{i=1}^{m-1} (1 + \dots + (2i - 1)) (1 + \dots + (2i - 1)) (1 + \dots + (2i - 4m + 2)) : \end{aligned} \quad (17)$$

Again, it is very difficult to calculate $C_m(\cdot)$. Unfortunately we could not get it in a rigorous way. A series of trials and errors, however, led us to the following form;

$$C_m(\cdot) = \frac{2^m}{2^m} m! [(\cdot + \cdot)]^{2^m} \prod_{i=1}^{2^m} (1 + 2(\cdot + i)) (1 + \cdot + (2i - 1)) : \quad (18)$$

This is consistent with $C_1(\cdot) = \frac{2^2}{2^1} [(\cdot + \cdot)]^2 (1 + \cdot + \cdot) (1 + 2(\cdot + \cdot))$ and the two other (calculable) cases $= 0$ and $= 0$ (up to symmetry factors). It is very difficult to get anything else consistent with these constraints. And a posteriori it seems to be correct since it gives a physically reasonable result. Let us assume that (18) is correct and see its consequences.

The two integrals are related by

$$I^m(\cdot; \cdot) = \frac{1}{2^m m!} (1 + \cdot) (1 + \cdot) (\cdot) J^m(2; \cdot; 1=2; \cdot) : \quad (19)$$

Therefore $C_m(\cdot) = (-2^m m!) C_m(1=2; \cdot) (\frac{1}{2} \cdot \cdot) (\frac{1}{2} + (2m + 1) \cdot)$. If we substitute (18) we get

$$C_m(\cdot) = \frac{2^m + 1}{2^{2^m}} \frac{1}{2} \prod_{i=1}^{2^m} (2i) \prod_{i=0}^{2^m} \frac{1}{2} + (2i + 1) : \quad (20)$$

Now we are ready to write down the amplitude. Without loss of generality, we can choose $k_1; k_3 = 0; k_2 = 0$. By using (5), (8) and $\sum_{i=1}^3 k_i = 2_0$, one gets

$$\begin{aligned} &\stackrel{8}{\approx} (1 - s) (0 > 0) \\ &\stackrel{\infty}{\approx} \frac{1}{2} - s (0 < 0) : \end{aligned}$$

It is easily seen that, for $0 > 0$, $A = 0$ identically, as in the bosonic theory. For $0 < 0$, there are many cancellations, leading to

$$\begin{aligned} &I^m(\cdot; \cdot) \\ &= C_m(\cdot) \frac{1}{2} (2i + 1) \prod_{i=1}^{2^m} (2i) (1 + \cdot + 2m) \frac{1}{2} + \\ &= (-1)^{m+1} \frac{1}{(m!)^2} \frac{1}{2} \prod_{i=0}^{2^m} (4)^{2^m} (1 + \cdot + 2m) \frac{1}{2} + : \quad (21) \end{aligned}$$

We finally obtain the three-point function:

$$A(k_1; k_2, k_3) = \left(\frac{i}{2}\right)^3 \frac{1}{2} \frac{1}{2}^s \frac{1}{2} \frac{s}{2} (1 + \dots + 2m) \frac{1}{2} + \\ = \frac{1}{2} \frac{1}{2}^s Y^3 \left(\frac{i}{2}\right) \frac{1}{2} [1 + \frac{2}{j} k_j^2] : \quad (22)$$

By redefining the cosmological constant and the primary supereld e_{NS} as

$$! \frac{2}{\frac{1}{2}} ; e_{NS}(k_j) ! \frac{1}{\left(\frac{i}{2}\right) \frac{1}{2} [1 + \frac{2}{j} k_j^2]} e_{NS}(k_j) ; \quad (23)$$

we get our main result

$$A(k_1; k_2, k_3) = : \quad (24)$$

Remarkably, this amplitude is of the same form as the bosonic one [10].

It is natural to expect that this feature continues to be true for N (> 4)-point functions.

In fact, for $k_1, k_2, k_3 \neq 0; k_4 \neq 0$ (and without screening charges), the four point function turns out to be

$$A(k_1; k_2, k_3, k_4) = (s+1)^{-s}; \quad (25)$$

with the same redefinition of the cosmological constant and the primary superelds. In order to get the amplitude for general k_i , one may argue, as in Ref. [10], that non-analyticity comes entirely from massless intermediate states and one may calculate the amplitude by using the analyticity of the one particle irreducible (1PI) correlators. After setting $\epsilon = 1$, we obtain the four-point function for all k_i :

$$A(k_1; k_2, k_3, k_4) = [k_1 + k_2 \rightarrow j k_1 + k_3 \rightarrow j k_1 + k_4 \rightarrow j] A_{1PI}; \quad (26)$$

with $A_{1PI} = \frac{1}{2} (1 + \epsilon^2)$. Compare with Eq. (37) in Ref. [10]. The analogy to the bosonic case is obvious. A detailed account will appear elsewhere.

The close connection to the bosonic amplitudes was suggested from super-KP systems [19], and more recently, from supermatrix models [20]. We think, however, that our demonstration is more direct.

In conclusion, we calculated the three- and four-point functions of super Liouville theory coupled to super Coulomb gas (without screening charges) on a sphere and found that they

are essentially the same as those of the usual Liouville theory, obtained in Ref. [11]. As a by-product we get the supersymmetric generalization of (B.9) formula of Ref. [18] ($s = 2m + 1$),

$$\begin{aligned}
& \frac{1}{s!} \sum_{i=1}^{\infty} d^2 z_i d^2 \sim \sum_{i=1}^{\infty} \bar{z}_i + \sum_{i,j=1}^{\infty} \bar{z}_i \bar{z}_j z_i \bar{z}_j \bar{z}_{ij}^4 \\
& = (-1)^m \sum_{i=1}^{2m+1} \frac{1}{2} \sum_{i=1}^{2m+1} \frac{1}{2} (2i) \sum_{i=0}^{2m+1} \frac{1}{2} + (2i+1) \\
& \quad \sum_{i=0}^{2m+1} (1 + \dots + 2i) (1 + \dots + 2i) (\dots + (2i - 4m)) \\
& \quad \sum_{i=1}^{2m+1} \frac{1}{2} + (2i-1) \quad \frac{1}{2} + \dots + (2i-1) \quad \frac{1}{2} + (2i-4m-1) : \\
& \tag{27}
\end{aligned}$$

A C K N O W L E D G M E N T S

K.H. would like to thank the members of Instituto de Física Teórica, UNESP, for their hospitality extended to him and FAPESP (# 90/1799-9) for the financial support. He also acknowledges communications on 2D gravity with Y.Tanii and with N.Ishibashi. D.D. thanks FAPESP (# 90/2246-3) for financial support. Work of E.A. and M.C.B.A. is partially supported by CNPq.

(a) Address after October 1991, Department of Physics, Kyushu University, Fukuoka 812, Japan.

REFERENCES

- [1] E . B rezin and V . A . K azakov, Phys. Lett. B 236, 144 (1990); M . R . D ouglas and S . H . Shenker, Nucl. Phys. B 335, 635 (1990); D . J . G ross and A . A . M igdal, Phys. Rev. Lett. 64, 127 (1990).
- [2] A . M . Polyakov, Phys. Lett. B 103, 207 (1981).
- [3] T . L . C urtright and C . B . Thom, Phys. Rev. Lett. 48, 1309 (1982); E . B raaten, T . L . Curtright and C . B . Thom, Phys. Lett. B 118, 115 (1982); Ann. Phys. 147, 365 (1983); E . B raaten, T . L . C urtright, G . G andour and C . B . Thom, Phys. Rev. Lett. 51, 19 (1983); Ann. Phys. 153, 147 (1984); J . L . G ervais and A . N eveu, Nucl. Phys. B 199, 50 (1982); B 209, 125 (1982); B 224, 329 (1983); B 238, 123, 396 (1984); E . D 'H oker and R . Jackiw, Phys. Rev. D 26, 3517 (1982); T . Y oneya, Phys. Lett. B 148, 111 (1984).
- [4] F . D avid, M od. Phys. Lett. A 3, 1651 (1988); J . D istler and H . K awai, Nucl. Phys. B 321, 509 (1989).
- [5] N . Seiberg, Lecture at 1990 Yukawa Int. Sem . C omm on Trends in M ath. and Quantum Field Theory, and C argese meeting Random Surfaces, Quantum Gravity and Strings, May 27, June 2, 1990; J . Polchinski, Strings '90 Conference, College Station, TX , Mar 12-17, 1990, Nucl. Phys. B 357 241 (1991).
- [6] A . M . Polyakov Phys. Lett. B 103 , 211 (1981).
- [7] J . F . A rvis, Nucl. Phys. B 212, 151 (1983); B 218, 309 (1983); O . B abelon, Nucl. Phys. B 258, 680 (1985); Phys. Lett. 141B , 353 (1984); T . C urtright and G . G andour, Phys. Lett. B 136, 50 (1984).
- [8] J . D istler, Z . H lousek and H . K awai, Int. J. Mod. Phys. A 5, 391 (1990).
- [9] M . G oulian and M . Li, Phys. Rev. Lett. 66, 2051 (1991).
- [10] P . D iFrancesco and D . K utasov, Phys. Lett. B 261, 385 (1991).

- [11] V.L.S. Dotsenko, Paris VI preprint PAR-LP THE 91-18 (1991).
- [12] Y.K. Itazawa, Harvard Univ. preprint HUTP-91/A 013 (1991).
- [13] N. Sakai and Y. Tanii, Tokyo Institute of Technology preprint TIT/HEP-169 (STUPP-91-116) (1991).
- [14] A. Gupta, S.P. Trivedi and M.B.W.isse, Nucl. Phys. B 340, 475 (1990).
- [15] (See also) M. Bershadsky and I.R. Klebanov, Phys. Rev. Lett. 65, 3088 (1990); Harvard Univ. preprint HUTP-91/A 002 (PUPT-1236) (1991); A.M. Polyakov, Mod. Phys. Lett. A 6, 635 (1991).
- [16] M.A. Bershadsky, V.G. Knizhnik and M.G. Teitelman, Phys. Lett. B 151, 131 (1985); D. Friedan, Z. Qiu and S. Shenker, Phys. Lett. B 151, 37 (1985).
- [17] E. Martinec, Phys. Rev. D 28, 2604 (1983).
- [18] V.L.S. Dotsenko and V. Fateev, Nucl. Phys. B 240, 312 (1984); B 251, 691 (1985).
- [19] P.D. Francesco, J. Distler and D. Kutasov, Mod. Phys. Lett. A 5, 2135 (1990).
- [20] L. Alvarez-Gaume and J.L. Marquez, Mod. Phys. Lett. A 6, 2039 (1991).