

NBI-HE-93-07  
January 1993  
PRELIMINARY  
hep-th/9302007

# VECTOR INDUCED LATTICE GAUGE THEORIES

Igor Pesando<sup>1</sup>

The Niels Bohr Institute  
University of Copenhagen  
Blegdamsvej 17, DK-2100 Copenhagen Ø  
Denmark

## Abstract

We consider vector induced lattice gauge theories, in particular we consider the QED induced and we show that at negative temperature corresponds to the dimer problem while at positive temperature it describes a gas of branched polymers with loops. We argue that these models are critical at  $D_{cr} = 6$  for  $N \neq \infty$  and they are *not* critical for  $N = \infty$ .

---

<sup>1</sup>E-mail PESANDO@NBIVAX.NBI.DK, 22105::PESANDO, 39163::LPESANDO

In this letter we would like to show how it is possible to induce a lattice gauge theory with group  $SU(N)$  starting from a lattice gauge theory for  $SU(N)$  coupled to vector in the fundamental irrep. Differently from ([1, 2]), we would like also to consider the relationship among these models, the dimer problem, branched polymers with loops and "induced QCD" ([3]).

We start examining the easiest example, the QED induced, at negative temperature. To this purpose let us consider the following action defined on a lattice  $\mathcal{G}$ <sup>1</sup>:

$$Z_{U(1)}^{(-)} = \int \prod_{links} dU_{ij} \int \prod_{l \in \mathcal{G}} d\epsilon_l d\bar{\epsilon}_l \exp \left( -\mu \sum_i \bar{\epsilon}_i \epsilon_i + K \sum_{i,j} A_{ij} U_{ij} \bar{\epsilon}_i \epsilon_j \right) \quad (1)$$

where  $A_{ij}$  is the adjacency matrix of the lattice  $\mathcal{G}$ ,  $\bar{\epsilon}_i = (\epsilon_i)^*$  are complex grassman variables and  $U_{ij} = e^{i\theta_{ij}} = U_{ji}^\dagger$  are elements of  $U(1)$ . This action can be regarded as a lattice gauge theory for  $U(1)$  with infinite coupling constant  $g = \infty$  coupled to a scalar ghost and because of that it describes a non unitary theory. The hamiltonian can be rewritten as

$$H_{U(1)}^{(-)} = \sum_{ij} \bar{\epsilon}_i (\mu \delta_{ij} - K A_{ij} U_{ij}) \epsilon_j \quad (2)$$

and it can be easily verified to be hermitian. We want to demonstrate that the partition function (1) is the generating function for the the dimer problem, more explicitly:

$$Z_{U(1)}^{(-)} = (-\mu)^N \sum_D N(D) \left( -\frac{K^2}{\mu^2} \right)^D \quad (3)$$

where  $D$  is the number of dimers and  $N(D)$  is the number of possible configurations with  $D$  dimers ( $N$  is the number of lattice sites).

To prove this assertion we consider the high temperature expansion (HTE) of the partition function (1); to this purpose we give the following graphical picture to the "interaction terms":

$$\begin{aligned} \overline{i} \longrightarrow j &\implies K \bar{\epsilon}_i U_{ij} \epsilon_j \\ \overline{i} \longleftarrow j &\implies -K \epsilon_i U_{ij}^\dagger \bar{\epsilon}_j \end{aligned}$$

---

<sup>1</sup> We use  $i, j, \dots$  to indicate sites (vertices) of the lattice;  $A, B, \dots$  to indicate rows and colons of group elements  $U$ ;  $\alpha, \beta, \dots$  to indicate the generation of the replica .

Now if a term of the HTE gives a non vanishing contribution to the partition function  $Z$ , it must have an equal number of  $U$  and  $U^\dagger$  on each link, but this number is limited (in this peculiar case) to 2 since we have grassman variables sitting in the vertices. Hence the fundamental unit (dimer) of our expansion is:

$$K \bar{\epsilon}_i U_{ij} \epsilon_j \cdot K \bar{\epsilon}_j U_{ji}^\dagger \epsilon_i = -K^2 \bar{\epsilon}_i \epsilon_i \bar{\epsilon}_j \epsilon_j \quad (4)$$

which can be given a graphical picture as follows:

$$\begin{array}{c} i \quad j \\ \hline \leftarrow \quad \rightarrow \end{array} \quad \equiv \quad \begin{array}{c} i \quad j \\ \hline \rightarrow \quad \leftarrow \end{array} \quad \Longrightarrow \quad -K^2 \bar{\epsilon}_i \epsilon_i \bar{\epsilon}_j \epsilon_j$$

Moreover we cannot put two dimers on the lattice sharing a vertex because  $(\bar{\epsilon}_i \epsilon_i)^2 = 0$ , hence we get that the HTE yields just the dimer recovering problem. Now we consider the corresponding induced QED. We find easily that

$$\begin{aligned} -H_{QED} &= N \log(\mu) - \sum_{\text{oriented } \Gamma} \frac{1}{l(\Gamma)} \left( \frac{K}{\mu} \right)^{l(\Gamma)} U_\Gamma \\ &= N \log(\mu) - 2 \sum_{\Gamma} \frac{1}{l(\Gamma)} \left( \frac{K}{\mu} \right)^{l(\Gamma)} \text{Re}(U_\Gamma) \end{aligned} \quad (5)$$

where  $\Gamma$  is a generic closed random walk and  $U_\Gamma$  is the product of the group element along the oriented path. Obviously only the non backtracking  $\Gamma$  will give a non trivial result since  $U(1)$  is an abelian group. Moreover the theory is at negative temperature reflecting the non unitarity of the initial model, nevertheless this model describes the dimer covering problem with negative weight that is known ([4]) to be equivalent to the Lee-Yang edge singularity and hence it is critical at  $D_{cr} = 6$ . We want to stress that the same result could have been reached with the abelian  $\mathbb{Z}_2$  group, i.e. with  $U_{ij} = \pm 1$  in (1). These results show that the induced  $\mathbb{Z}_2$  model and the induced QED are equivalent each other but they are *not* equivalent to the usual models at the criticality. As last observation we want to notice that in the QED case setting  $\mu = 0$  we get the close packing dimer problem that it is known to be equivalent to Ising and hence in this case the theory has  $D_{cr} = 4$  ([5]).

In a complete similar way we introduce the induced QED at positive temperature. It is given by the partition function

$$Z_{U(1)} = \int \prod_{links} dU_{ij} \int \prod_{l \in \mathcal{G}} d\phi_l d\phi_l^* e^{-\mu \sum_i \phi_i^* \phi_i + K \sum_{i,j} A_{ij} U_{ij} \phi_i^* \phi_j} \quad (6)$$

where  $\phi_i$  are complex scalar defined on the vertices of  $\mathcal{G}$ . Its HTE yields

$$Z = (w_0)^N \sum_{gas \text{ of } BP} \tilde{N}(B, \{V_f\}) (K^2)^B \prod_{f=1}^{f=q} \left(\frac{w_f}{w_0}\right)^{V_f} \quad (7)$$


where the sum is extended over a gas of branched polymers with  $B$  bonds,  $V_f$   $f$ -branching units and  $\tilde{N}(B, \{V_f\})$  is the weighted sum of the number of such configurations. Explicitly  $\tilde{N}(B, \{V_f\})$  is given by

$$\tilde{N}(B, \{V_f\}) = \sum_{\substack{\Gamma \text{ with } B \text{ bonds} \\ \text{and } V_f \text{ } f \text{ units}}} \prod_{bond \ b \in \Gamma} \left(\frac{1}{n_b!}\right)^2 \quad (8)$$

where  $n_b$  is the occupation number of the bond  $b$  belonging to a gas  $\Gamma$  of branched polymers with loops (i.e.  $\Gamma$  can be made of many disconnected pieces). Besides the statistical weights of the  $f$ -branching units are given by:

$$w_f = \frac{\pi f!}{\mu^{f+1}} \quad (9)$$

The formula (7) can be demonstrated as in the previous case of ghosts: also in this case a non vanishing contribution to HTE has on each link an equal number of  $U$  and  $U^\dagger$ , in such a way that a bond of a branched polymer (a monomer) is composed by one  $U$  and one  $U^\dagger$  bond, exactly as shown by the following figure



In this way each monomer, i.e. polymer link, yields a  $\phi\phi^\star$  contribution to each of its endpoints, so that a  $f$ -branching vertex contributes as

$$w_f = \int d\phi d\phi^\star \exp(-\mu\phi\phi^\star)(\phi\phi^\star)^f = \frac{\pi f!}{\mu^{f+1}} \quad (10)$$

Moreover each link occupied  $n$  times yields a factor  $(\frac{1}{n!})^2$  instead of  $\frac{1}{n!}$  because its compositeness.

It is now useful to make a number of comments. Firstly we notice that the this induced QED is also critical at  $D_{cr} = 6$ . This is easily demonstrated as (see also (17)):

$$\begin{aligned} Z_{U(1)} &\propto \int \prod_{links} dU_{ij} \det(\mu\delta_{ij} - K A_{ij}U_{ij})^{-1} = \lim_{n \rightarrow -1} \int \prod_{links} dU_{ij} (\det(\mu\delta_{ij} - K A_{ij}U_{ij}))^n \\ &= \lim_{n \rightarrow -1} \int \prod_{links} dU_{ij} \int \prod_{l \in \mathcal{G}} d\epsilon_{l\alpha} d\bar{\epsilon}_{l\alpha} e^{\sum_{\alpha=1}^n \left( -\mu \sum_i \bar{\epsilon}_{i\alpha} \epsilon_{i\alpha} + K \sum_{i,j} A_{ij} U_{ij} \bar{\epsilon}_{i\alpha} \epsilon_{j\alpha} \right)} \end{aligned} \quad (11)$$

and relying on the fact that the critical dimension is independent of the number of components of the order parameter.

Secondly the difference between branched polymers with loops and random walks is *both* in the silhouette of the terms of the HTE ( due to the appearance of odd branching vertices) and in the weights used for  $\tilde{N}$  ( $\frac{1}{n_b!}$  for RW,  $(\frac{1}{n_b!})^2$  for BP).

Now we would like also to consider the more general case of the  $SU(N)$  or  $U(N)$  induced by the the following partition function:

$$Z_{SU(N)} = \int \prod_{links} dU_{ij} \int \prod_{l \in \mathcal{G}} d\phi_{lA} d\phi_{lA}^\star e^{-\mu N \sum_i \phi_{iA}^\star \phi_{iA} + N K \sum_{i,j} A_{ij} U_{ij,AB} \phi_{iA}^\star \phi_{jB}} \quad (12)$$

where  $\phi_i \equiv (\phi_{iA}) = (\phi_{i1} \dots \phi_{iN})$  are complex scalars in the fundamental irrep of  $SU(N)$  defined on every vertex  $i \in \mathcal{G}$ . The HTE of these models yield graphs with the same silhouettes of the  $U(1)$  case but with different weights. To get these weights we integrate over the gauge variables firstly, to this purpose we use the fact that

$$f_{SU(N)}(M, M^\dagger) = e^{N g_{SU(N)}(M, M^\dagger)} = \int_{SU(N)} dU_{ij} \exp(N \text{tr}(M^\dagger U + U^\dagger M))$$

is a function of  $tr(MM^\dagger)^p$  and of  $detM$  (in the case of  $SU(N)$ ) and that in our case  $M_{ij} = \phi_i \otimes \phi_j^\dagger$  to deduce that  $tr(MM^\dagger)_{ij}^p = (\phi^\dagger \phi)_i^p (\phi^\dagger \phi)_j^p$  and  $detM = 0$ , so that<sup>2</sup>

$$f_{SU(N)}(\phi_i \otimes \phi_j^\dagger, \phi_j \otimes \phi_i^\dagger) = e^{N g_{SU(N)}(\phi_i^\dagger \phi_i \times \phi_j^\dagger \phi_j)} = \sum_p c_p(N) (\phi^\dagger \phi)_i^p (\phi^\dagger \phi)_j^p \quad (13)$$

Hence we can rewrite (12) as

$$Z_{SU(N)} = \int \prod_{l \in \mathcal{G}} d\phi_{lA} d\phi_{lA}^* e^{-N\mu \sum_i \phi_i^\dagger \phi_i} \prod_{links \ ij} \left( \sum_p c_p(N) K^{2p} (\phi^\dagger \phi)_i^p (\phi^\dagger \phi)_j^p \right) \quad (14)$$

Similarly to the  $U(1)$  case we associate to each polymer link a factor  $\phi^\dagger \phi$  for each of its endpoints, so that now the weight for a f-branching vertex becomes

$$w_f = \int \prod_A d\phi_A d\phi_A^* \exp(-\mu N \phi^\dagger \phi) (\phi^\dagger \phi)^f = \frac{\pi^N (N+f)!}{N! (\mu N)^{N+f+1}} \quad (15)$$

meanwhile the occupation weights can be deduced from (13) and yield for  $\tilde{N}(B, \{V_f\})$  the expansion

$$\tilde{N}(B, \{V_f\}) = \sum_{\substack{\Gamma \text{ with } B \text{ bonds} \\ \text{and } V_f \text{ f units}}} \prod_{bond \ b \in \Gamma} c_{n_b}(N) \quad (16)$$

where in particular we have  $c_n(1) = (\frac{1}{n!})^2$ .

It is important to notice that also these models are critical at  $D_{cr} = 6$  because they are in the same universality class of branched polymers with loops and both odd and even vertices. In order to show this point also from a more mathematical point of view, we change coordinates in (14) and we introduce  $\exp \varphi_i = \phi_i^\dagger \phi_i$  ( $\varphi \in ]-\infty, +\infty[$ ) along with the radial coordinates, so that (14) becomes

$$Z_{SU(N)} = const \int \prod_{l \in \mathcal{G}} d\varphi_l \exp \left( N \sum_i (\varphi_i - \mu e^{\varphi_i}) + N \sum_{\langle ij \rangle} g_{SU(N)}(e^{\varphi_i + \varphi_j}) \right) \quad (17)$$

---

<sup>2</sup>It is easy to show that using the normalization  $vol(U(N)) = vol(SU(N))$  we have  $f_{U(N)}(\phi_i, \phi_j) = f_{SU(N)}(\phi_i, \phi_j)$ . Moreover this implies that using  $N > 1$  scalars we cannot induce  $U(N)$  but only  $SU(N)$ .

If now we expand the action, we shift  $\varphi$  as the shifted  $\varphi$  has null expectation value and we throw away the irrelevant term we recover a  $\varphi^3$  action that is critical at  $D_{cr} = 6$ . Obviously this is not a rigorous proof since we should check that the irrelevant operators are really irrelevant at the fixed point ([6]), nevertheless up to now this approach did not fail to get the upper critical dimension.

Extending the known result valid for ordinary lattice gauge theories for which both  $SU(N)$  and  $SU(N)/\mathbb{Z}_N$  have the same upper critical dimension, here we would deduce that both the vector induced lattice gauge theories and the adjoint induced theories ("induced QCD") at finite  $N$  share the same upper critical dimension  $D_{cr} = 6$ , nevertheless this seems not to be true in this case since the induced QCD seems to be critical at  $D_{cr} = 4$  ([7]).

Now we would solve exactly this model in the large  $N$  limit. To this purpose we compute exactly (13) in the large  $N$  limit. Using the observation that  $f_{SU(N)}$  depends only on  $x = \phi_i^\dagger \phi_i \cdot \phi_j^\dagger \phi_j$  we get easily the equation

$$\frac{x}{N} \frac{d^2 g}{dx^2} + \left(\frac{dg}{dx}\right)^2 + \frac{dg}{dx} = 1 \quad (18)$$

that can be solved easily ([8]) as:

$$g(x) = \sqrt{1+4x} - 1 - \log(1 + \sqrt{1+4x}) + \log 2 \quad (19)$$

Plugging this solution into (14) and changing coordinates from  $\phi_{iA}, \phi_{iA}^*$  to  $R_i = \phi_i^\dagger \phi_i$  and angular coordinates, we get

$$Z_{SU(N)} = \text{const} \int_0^\infty \prod_{i \in \mathcal{G}} dR_i e^N \left[ \sum_i \left(1 - \frac{1}{N}\right) \log(R_i) - \mu \sum_i R_i + \sum_{\langle ij \rangle} \sqrt{1+4K^2 R_i R_j} - \log(1 + \sqrt{1+4K^2 R_i R_j}) \right] \quad (20)$$

Now we can use the usual saddle point method. If we look for a translational invariant solution we get

$$\begin{aligned} KR &= \frac{-\frac{\mu}{K}(D-1) + \sqrt{\left(\frac{\mu}{K}\right)^2 - 4(2D-1)}}{\left(\frac{\mu}{K}\right)^2 - 4D^2} && \text{for } \frac{\mu}{K} > 2D \text{ and } D \geq 0 \\ KR &= \frac{2D-1}{4D(D-1)} && \text{for } D > 1 \end{aligned} \quad (21)$$

while in the range  $D < \frac{\mu}{J} < 2D$  the saddle point is unstable. The explicit form of the free energy density is then ( $\mathcal{V}$  is the volume of the lattice)

$$F_{SU(\infty)} = \frac{\log Z_{SU(N)}}{N\mathcal{V}} = \log(KR) - \frac{\mu}{J}R + D\sqrt{1 + 4K^2R^2} - D\log(1 + \sqrt{1 + 4K^2R^2}) - \log(K) \quad (22)$$

that does not show any transition. It can be shown that  $F_{SU(\infty)}^{\text{fer}} = -F_{SU(\infty)}^{\text{bos}}$ .

In conclusion we show that the  $N = \infty$  theories (both bosonic and fermionic) are "trivial" since they have no critical dimension and can be solved exactly by the saddle point; on the contrary the theories at  $N \neq \infty$  are critical so the role of coupling constant is played by  $\frac{1}{N}$  but the  $N = \infty$  theory is not able to catch all the features of the  $N \neq \infty$  theories, exactly as the gaussian approximation has not all the characteristics of  $\phi^4$  theory. Perhaps the same conclusion can be reached for the  $SU(N)$  induced by up to  $N - 1$  scalars since in this range we are in the strong coupling regime of Brezin-Gross formula ([8]), nevertheless a different regime should show up starting at  $N$  scalars (or fermions) since it appears both the strong and weak coupling regime ([2]).

**Acknowledgement** We thank for useful discussion D. Boulatov, M. Caselle, A. D'Adda, P. Di Vecchia, F. Gliozzi and M. Rasetti.

## References

- [1] M. Bander, Phys. Lett. **126B** (1983) 463  
H.W. Hamber, Phys. Lett. **126B** (1983) 471
- [2] I.Ya. Aref'eva, "A simple model inducing QCD", SMI-1-93  
I.Ya. Aref'eva, "Large-N Quenching in the Kazakov-Migdal model", SMI-12-92 (both available on hep-th)
- [3] V.A. Kazakov and A.A. Migdal, "Induced QCD at Large  $N$ ", preprint PUPT - 1322, LPTENS -92/15, May, 1992.  
A.A. Migdal, "Exact Solution of Induced Lattice Gauge Theory at large  $N$ ", preprint PUPT - 1323, Revised, June, 1992;
- [4] G.A. Baker and P. Moussa, J. Appl. Phys. **49** (1978) 1360  
D. Kurze and M.E. Fisher, Phys. Rev **B20** (1979) 2785  
Y. Shapir, J. Phys. **A15** (1982) L433



- [5] M.E Fisher, J. Mod. Phys. **7** (1966) 1776  
M. Rasetti and T. Regge, Riv. Nuovo Cimento **4** (1981) 1
- [6] F. Fucito and G. Parisi, J. Phys. **A14** (1981) L507
- [7] M. Caselle, A. D’Adda and S. Panzeri, “The Kazakov-Migdal model as a high temperature lattice gauge theory”, DFTT 71/92, December 1992
- [8] E. Brezin and D.J. Gross, Phys. Lett. **97B** (1980) 120