

CBPF-NF-054/94

DFTT-045/94

# The Quantum Algebraic Structure of the Twisted XXZ Chain

M. R. Monteiro<sup>1a</sup>, I. Roditi<sup>1b</sup>, L. M. C. S. Rodrigues<sup>1c</sup> and S. Sciuto<sup>2d</sup>

<sup>1</sup>Centro Brasileiro de Pesquisas Físicas - CBPF

Rua Dr. Xavier Sigaud, 150

22290-180 Rio de Janeiro, RJ, Brasil

<sup>2</sup>Dipartimento di Fisica Teorica dell'Università di Torino

and Sezione di Torino dell'INFN, Via P. Giuria 1, I-10125, Torino, Italy

## Abstract

We consider the Quantum Inverse Scattering Method with a new R-matrix depending on two parameters  $q$  and  $t$ . We find that the underlying algebraic structure is the two-parameter deformed algebra  $SU_{q,t}(2)$  enlarged by introducing an element belonging to the centre. The corresponding Hamiltonian describes the spin-1/2 XXZ model with twisted periodic boundary conditions.

---

e-mail addresses:

(a) mmont@cbpfsul.cat.cbpf.br

(b) roditi@cbpfsul.cat.cbpf.br

(c) ligia@brlnccbitnet

(d) sciuto@to.infn.it

Many completely integrable one-dimensional quantum models have been treated [1] by the Quantum Inverse Scattering Method (QISM) [2, 3] which, among other achievements, led to the discovery of Quantum Groups [4] independently from Drinfeld and Jimbo [5]. In this letter we use this method to describe the relation between the spin-1/2 XXZ chain with twisted periodic boundary conditions and the two-parameter deformed algebra  $SU_{qt}(2)$  enlarged by introducing an element belonging to the centre, showing that the second parameter of the deformation,  $t$ , is linked to the twist.

The QISM introduces an auxiliary problem with the help of the so-called Lax operator. In our model this operator is

$$L_n(\lambda; t) = \begin{pmatrix} 0 & 1 \\ t^{Z_n} S_n^3 \operatorname{sh}[(\lambda + iS_n^3)] & iS_n \sin \frac{\lambda}{A} \\ iS_n^+ \sin \frac{\lambda}{A} & t^{Z_n} S_n^3 \operatorname{sh}[(\lambda - iS_n^3)] \end{pmatrix}; \quad (1)$$

where  $S_n$  and  $Z_n$  are operators defined on the  $n$ -th vectorial space of the periodic  $(S_{N+1} = S_1; Z_{N+1} = Z_1)$  chain, which in the fundamental representation are given by

$$Z_n = \frac{1}{2} 1I_n; S_n = \frac{1}{2} \tilde{\sigma}_n; \quad (2)$$

where  $\tilde{\sigma}$  are the Pauli matrices and  $1I$  is the identity operator.

The R-matrix associated to the Lax operator (1) is

$$R(\lambda, \mu; t) = \begin{pmatrix} 0 & 1 \\ a(\lambda, \mu) & 0 & 0 & 0 \\ 0 & c^0(\lambda, \mu) & b(\lambda, \mu) & 0 \\ 0 & b(\lambda, \mu) & c^0(\mu, \lambda) & 0 \\ 0 & 0 & 0 & a(\mu, \lambda) \end{pmatrix}; \quad (3)$$

where

$$\begin{aligned} a(\lambda, \mu) &= \operatorname{sh}[(\lambda + \mu)] \\ b(\lambda, \mu) &= i \sin \frac{\lambda - \mu}{A} \\ c^0(\lambda, \mu) &= t c(\lambda, \mu) \\ c^0(\lambda, \mu) &= t^{-1} c(\mu, \lambda) \end{aligned} \quad (4)$$

and

$$c(\lambda) = \text{sh } \lambda; \quad (5)$$

clearly,  $R(\lambda) = R(\lambda; t=1)$  is the appropriate matrix for the XXZ model [6]. It is easy to check that the matrix  $R(\lambda; t)$  (eqs. (3-5)) satisfies the Yang-Baxter equation [7, 8]

$$R_{12}(\lambda_{12}; t) R_{13}(\lambda_{13}; t) R_{23}(\lambda_{23}; t) = R_{23}(\lambda_{23}; t) R_{13}(\lambda_{13}; t) R_{12}(\lambda_{12}; t) \quad (6)$$

and that  $L_n(\lambda; t)$  (eq. (1)) obeys the Fundamental Commutation Relations (FCR)

$$R_{12}(\lambda_{12}; t) L_n^1(\lambda_1; t) L_n^2(\lambda_2; t) = L_n^2(\lambda_2; t) L_n^1(\lambda_1; t) R_{12}(\lambda_{12}; t); \quad (7)$$

In (6) and (7),  $\lambda_{ij} = \lambda_i - \lambda_j$  and

$$\begin{aligned} R_{12} &= \sum_i a_i \lambda_i - \lambda_i \mathbb{I} \quad ; \quad R_{13} = \sum_i a_i \mathbb{I} - b_i \lambda_i; \\ R_{23} &= \sum_i \lambda_i - a_i \lambda_i - b_i \lambda_i; \end{aligned} \quad (8)$$

with the  $R(\lambda; t)$  matrix written as

$$R(\lambda; t) = \sum_i a_i \lambda_i - b_i \lambda_i \quad (9)$$

and the upper indices in eq. (7) follow

$$L^1 = L - \mathbb{I} \mathbb{I} \quad ; \quad L^2 = \mathbb{I} \mathbb{I} - L; \quad (10)$$

We notice that the  $R$ -matrix is defined on the tensor product of two auxiliary spaces  $\mathbb{C}^2 \otimes \mathbb{C}^2$  and the  $L$ -matrix is defined on the tensor product of the auxiliary space  $\mathbb{C}^2$  and the internal space  $\mathbb{C}^d$ , with  $d$  the dimension of the representation of the associated algebra satisfied by the operators in the elements of the matrix  $L$ .

The reason for  $R(\lambda; t)$  to satisfy eq. (6) is that it can be written in terms of  $R(\lambda)$  as<sup>1</sup>

$$R_{12}(\lambda; t) = g^1 (g^2)^{-1} R_{12}(\lambda) g^1 (g^2)^{-1}; \quad (11)$$

---

<sup>1</sup>Equivalently one could write

$$g^1 (g^2)^{-1} = t^{S^3 - Z} Z^{-1} S^3; \quad ;$$

with  $S^3$  and  $Z$  given by eq. (2). This form is more appropriate if one wishes to compare the algebraic structure here presented with ref. [9]. In a forthcoming paper we shall discuss this subject, as well as the relationship of our approach with the one in ref. [10].

where

$$g^1 = g \otimes 1 \quad ; \quad g^2 = 1 \otimes g \quad ; \quad g = t^{\frac{1}{2}S^3} \quad (12)$$

and

$$[g^1 g^2; R_{12}(\lambda)] = 0: \quad (13)$$

Moreover, eq. (7) follows from eq. (6), because (for  $d=2$ )

$$L_n(\lambda; t) = R_{on}(\lambda - \frac{i}{2}; t); \quad (14)$$

where  $\lambda$  labels the auxiliary space. We also observe that

$$R(0; t) = P; \quad (15)$$

where  $P$  is the permutation operator on the tensor product of the two spaces where the  $R$ -matrix is defined.

According to the standard procedure of the QISM, eq. (7) allows one to build an infinite set of commuting operators

$$F(\lambda; t) = \text{Tr}[L_N(\lambda; t) \otimes_2 (L; t) L_1(\lambda; t)]; \quad (16)$$

where both the matrix product and the trace are performed in the auxiliary space.

In our case the Bethe Ansatz equations for the fundamental representation are given by

$$\frac{(\lambda)}{(\lambda)}^N = t^N \prod_{j=1}^M \left( \frac{a(\lambda)}{a(\lambda)} \frac{c(\lambda)}{c(\lambda)} \right); \quad \lambda = 1; \quad ; M = N; \quad (17)$$

with

$$\begin{aligned} a(\lambda) &= \text{sh} \left( \lambda + \frac{i}{2} \right) \\ c(\lambda) &= \text{sh} \left( \lambda - \frac{i}{2} \right); \end{aligned} \quad (18)$$

explicitly showing the contribution due to the parameter  $t$ , as  $a$ ,  $b$ ,  $c$  and  $d$  are the same functions appearing in the XXZ model.

In order to show the algebraic structure underlying the  $R$  and  $L$  matrices defined in eqs. (1) and (3-5) we perform a suitable similarity transformation [3] on (6) and (7) which permits us to have the following decomposition:

$$\tilde{L}_n(\cdot; t) = \frac{1}{2} (e^{-L_+} - e^{-L_-}) \quad (19)$$

$$\tilde{R}(\cdot; t) = e^{-L_+} R_+ - e^{-L_-} R_- ;$$

where

$$L_+ = \begin{pmatrix} 0 & 1 \\ q^{s^3} t^z s^3 & S \\ 0 & q^{s^3} t^z s^3 \end{pmatrix} \begin{matrix} C \\ C \\ A \end{matrix} \quad (20)$$

$$L_- = \begin{pmatrix} 0 & 1 \\ q^{s^3} t^z s^3 & 0 \\ S^+ & q^{s^3} t^z s^3 \end{pmatrix} \begin{matrix} C \\ C \\ A \end{matrix}$$

and

$$R_+ = \begin{pmatrix} q & 0 & 0 & 0 \\ 0 & t & 0 & 0 \\ 0 & 0 & t^{-1} & 0 \\ 0 & 0 & 0 & q \end{pmatrix} ; \quad (21)$$

with  $\cdot = q - q^{-1}$  and  $R_- = P R_+^{-1} P$ .

Substituting eq. (19) in the Y-B equation (6) and in the FCR, eq. (7), one gets the following independent equations:

$$R_+ L_-^1 L_-^2 = L_-^2 L_-^1 R_+ \quad (n = 1) \quad (22)$$

$$R_+ L_+^1 L_-^2 = L_-^2 L_+^1 R_+ ;$$

which imply that the operators in the entries of  $L$  must satisfy

$$[s^3; z] = [s; z] = 0 \quad (23a)$$

$$[s^3; s] = s \quad (23b)$$

$$t^{-1} s^+ s - t s s^+ = t^{2s^3} [s^3]_t ; \quad (23c)$$

where  $[x]_q = (q^x - q^{-x})/(q - q^{-1})$  with  $q = \exp(i\theta)$ . Eqs. (23) are the commutation relations of the two-parametric deformed  $SU(2)$  [9, 11, 12] with  $Z$ , an element of the center of the resulting algebra. The coproduct is obtained by considering the product of two  $L_n$  acting on two internal spaces and we find:

$$S^3 = S^3 \otimes 1 + 1 \otimes S^3 \quad (24a)$$

$$Z = Z \otimes 1 + 1 \otimes Z \quad (24b)$$

$$S^\pm = q^{S^3} t^{\pm Z} S^\pm \otimes S^\pm + S^\pm \otimes q^{\mp S^3} t^{\mp Z} S^\pm; \quad (24c)$$

The coproduct (24c) is related to the one in ref. [12] by a similarity transformation generated by the operator  $t^{S^3 \otimes Z - Z \otimes S^3}$ .

Following the QISM, a local Hamiltonian can be written as

$$H = \sum_{i=1}^N \frac{\partial}{\partial t} \log F(i; t) \quad (25)$$

and thanks to eqs. (14-16), it becomes for the fundamental representation of the algebra

$$H = \sum_{i=1}^N H_{i,i+1} \quad (N+1 \rightarrow 1) \quad (26)$$

$$H_{i,i+1} = \frac{J \sin \theta}{i} \frac{\partial}{\partial t} \hat{R}_{i,i+1}(i; t) \big|_{j=0};$$

where  $\hat{R} = P R$  and  $R(i; t)$  is given by eqs. (3-5). In the above equation  $R_{i,i+1}(i; t)$  acts on the two internal spaces  $(i; i+1)$  instead of acting on two auxiliary spaces.

Substituting  $R(i; t)$  given by eqs. (3-5) in eq. (26), apart from an additive constant, we get

$$H = \sum_{i=1}^N \left[ 2t^{-1} \sigma_i^x \sigma_{i+1}^x + 2t \sigma_i^y \sigma_{i+1}^y + \frac{q + q^{-1}}{2} \sigma_i^z \sigma_{i+1}^z \right]; \quad (27)$$

where  $\sigma = (\sigma^x, \sigma^y, \sigma^z)$ .

Such a Hamiltonian is very similar to the XXZ model with periodic boundary conditions but for each pair of sites  $(i; i+1)$ , the site  $(i+1)$  is rotated of an angle  $\theta$  ( $t = e^{i\theta}$ ) in the  $x-y$  plane with respect to the site  $i$ .

The similarity transformation generated by  $\exp \left( i \sum_{n=1}^N (\sigma_n^x - 1) \right)$  takes the Hamiltonian (eq. (27)) to

$$H = \frac{J}{2} \sum_{n=1}^N \left( \sigma_n^x \sigma_{n+1}^x + \sigma_n^y \sigma_{n+1}^y + \cos \theta \sigma_n^z \sigma_{n+1}^z + \right. \\ \left. \cos \theta \sigma_N^z \sigma_1^z + 2t \sum_{n=1}^N \sigma_n^+ \sigma_{n+1}^- + 2t^N \sigma_N^+ \sigma_1^- \right); \quad (28)$$

which is the well-known [13] Hamiltonian for the XXZ chain with twisted periodic boundary conditions.

It is amusing to observe that, thanks to eq. (13) and following the procedure of ref. [10], the Hamiltonian (eq. (27)) could also be obtained from the R-matrix of the XXZ model  $R(\lambda) = R(\lambda; t=1)$ , using  $L_n^0(\lambda; t) = t^{S^3} L_n(\lambda; t=1) \in L_n(\lambda; t)$ . Conversely, the untwisted XXZ model can be built from  $R(\lambda; t)$  (eqs. (3-5)) and  $L_n^0(\lambda; t) = t^{S^3} L_n(\lambda; t)$ . All these topics will be discussed in detail in a forthcoming paper.

Finally, we would like to point out that by introducing a central element  $Z$  which enlarges the  $SU_{q,t}(2)$  algebra, we make appear the underlying algebraic structure of the so-called twisted XXZ model.

## Acknowledgements

S. Sciuto thanks CBPF for its kind hospitality and CNPq for financial support. I. Roditi thanks D. A. Iltshuler for useful discussions.

## References

- [1] P. Kulish and E. Sklyanin, Lect. Notes in Phys. 151 (1982) 61 (and ref. therein);  
N. Andrei, "Integrable Models in Condensed Matter Physics", ICTP Summer Course,  
Trieste, 1992 (and ref. therein);  
V. Korepin and F. Eiler, "Exactly Solvable Models of Strongly Correlated Electrons"  
(and ref. therein), cond-mat/9408092.
- [2] E. Sklyanin, L. Takhtajan and L. Faddeev, Teor. Matem. Fiz. 40 (1979) 194;  
L. Takhtajan and L. Faddeev, Russian Math. Surveys 34 (1979) 11.
- [3] L. Faddeev, in "New Problems, Methods and Techniques in Quantum Field Theory  
and Statistical Mechanics", edited by M. Rasetti, World Scientific Pub. (1990);  
"Algebraic Aspects of Bethe-Ansatz", preprint ITP-SB-94-11.
- [4] P. Kulish and N. Reshetikhin, Zap. Nauch. Seminarov LOMI 101 (1981) 101; J. Sov.  
Math. 23 (1983) 2435.
- [5] V. Drinfeld, Sov. Math. Dokl. 32 (1985) 254;  
M. Jimbo, Lett. Math. Phys. 10 (1985) 63; 11 (1986) 247.
- [6] E. Lieb, Phys. Rev. Lett. 18 (1967) 692; Phys. Rev. 162 (1968) 162; Phys. Rev. Lett.  
18 (1967) 1046; Phys. Rev. Lett. 19 (1967) 108.
- [7] R. Baxter, Ann. of Phys. 70 (1972) 193; Ann. of Phys. 70 (1972) 323.
- [8] C. N. Yang, Phys. Rev. Lett. 19 (1967) 1312.
- [9] N. Reshetikhin, Lett. Math. Phys. 20 (1990) 331.
- [10] H. J. de Vega, Nucl. Phys. B 240 (1984) 495; Int. J. Mod. Phys. A 4 (1989) 2371.
- [11] A. Sudbery, J. Phys. A 23 (1990) L697; M. Takeuchi, Proc. Jap. Acad. 66 (1990)  
112.



[12] C .Burdik and L .H lavaty, J.Phys. A 24 (1991) L165.

[13] F .A lcaraz, U .G rim m and V .R ittenberg, Nucl.Phys. B 316 (1989) 735;  
V .Pasquier and H .Sakur, Nucl.Phys. B 330 (1990) 523.