

ABSENCE OF CLASSICAL AND QUANTUM MIXING

L.L. Salcedo

Departamento de Física Moderna, Universidad de Granada, E-18071 Granada, Spain

It is shown, under mild assumptions, that classical degrees of freedom dynamically coupled to quantum ones do not inherit their quantum fluctuations. It is further shown that, if the assumptions are strengthened by imposing the existence of a canonical structure, only purely classical or purely quantum dynamics are allowed.

1. There is consensus among physicists that Quantum Mechanics is the correct description of Nature, at least within the range of presently observable scales. Nevertheless, some systems are routinely described using a classical, and thus approximated, dynamics. This can be so either for simplicity or due to the lack of a consistent quantum theory. Einstein's General Relativity is an example of the latter. In an excellent speculative paper [1], Boucher and Traschen consider several physical systems which require a mixed description in terms of quantum and classical degrees of freedom, mutually interacting. A good example is provided by early universe physics, where fully quantum matter fields are coupled to classical gravitational fields. The traditional approach to this problem has been to couple the gravitational fields to the expectation values of the quantum energy-momentum tensor, see e.g. Ref. [2]. This kind of approach has been criticized [1], on the grounds that the classical fields evolve deterministically, hence, the quantum fluctuations in these fields, induced by their coupling to the quantum fields, are missed.

This criticism (as well as presumably the challenge it presents) has led to look for a mathematically consistent description of semi-quantized systems, i.e., mixed classical-quantum systems [3,4,6]. These systems are considered by themselves, that is, not as the limit of a fully quantum theory. The fact that the classical description is just an approximation is disregarded in this context, since the purpose is to define a mathematical structure with some physical input.

In this letter it is shown that in fact there are severe obstructions to construct such a description and, if it exists at all, it will not enjoy the elegant mathematical structures common to classical or quantum mechanics. Since presently there is no widely accepted definition of what is meant by a semi-quantized system, and in order not to discard potentially interesting choices, we should rely on properties as general as possible, which must hold, in particular, for the purely classical and purely quantum cases. Throughout, the degrees of freedom will be bosonic, though this assumption does not seem to be essential for the arguments.

2. It is assumed (i) that the set of observables forms an associative algebra A over the field of complex numbers. Let A_L be the algebra spanned by the coordinates q_i , the conjugate momenta p_i , $i = 1, \dots, N$ and the identity E , as generators, i.e., the set of formal series of ordered products of them. Then the physical algebra A is defined as the quotient algebra of A_L modulo some identities among the generators (e.g., commutation relations). These identities characterize the algebra and are to be specified. A will be a non-commutative algebra in general. By definition, $AE = EA = A$ for every observable A . As a consequence, E is the only element with this property. In classical mechanics A is just the set of complex functions in phase space and E is the unity function. In quantum mechanics we have the algebra of operators in the Hilbert space of the system. Here, the word observable is being used in a slightly wider sense than usual, since it includes non-real functions and non-Hermitian operators as well. This axiom is also present in Refs. [1,4],

A second axiom (ii) refers to the time evolution of the observables (Heisenberg picture). Namely, there is a family U of evolution operators $U(t_1; t_2)$ in A such that $U(t_1; t_1)$ is the identity operator and $U(t_1; t_2)U(t_2; t_3) = U(t_1; t_3)$. Furthermore, the evolution preserves the algebraic structure, that is, if two observables $A_{1,2}(t_0)$ evolve to $A_{1,2}(t)$, and $c_{1,2}$ are constant complex numbers, $c_1 A_1(t_0) + c_2 A_2(t_0)$ evolves to $c_1 A_1(t) + c_2 A_2(t)$, and $A_1(t_0)A_2(t_0)$ evolves to $A_1(t)A_2(t)$. In other words, time evolution forms a groupoid of algebra automorphisms of A . Certainly, this axiom holds both in classical and in quantum mechanics, and it is hard to imagine an interesting formulation which would violate it. Moreover, the endomorphism property follows from the Schrodinger picture, since there the observables do not evolve, and hence the algebraic structure is trivially preserved. Note that U gives only the dynamic time dependence of observables, and also that, in principle, the operator $U(t_1; t_2)$ does not correspond to an algebra element (e.g., in the purely classical case). On the other hand, it is not assumed that the system is conservative. There can be time dependent external fields which break invariance under time translations. Similarly, time reversal invariance is not required.

Some relevant conclusions can be extracted from these two axioms. If a set of elements generates the algebra, this property is maintained through time evolution. This follows from time evolution being an automorphism. The

observable E is time independent: for any observable A , the relation $E A = A E$ evolves to $E(t)A(t) = A(t)E(t)$ and thus $E(t) = E$, using that $A(t)$ is an arbitrary observable, since time evolution is a bijection and A was arbitrary. Another consequence is that commutation relations of the form $[A(t_0); B(t_0)] = cE$ are also preserved, since they evolve to $[A(t); B(t)] = cE$. In particular, if two observables commute at any given time they do so at any time.

A last axiom (iii) is needed referring to the commutation relations. The classical dynamics is characterized by commuting coordinates and momenta which evolve according to Hamilton's equations. On the other hand, the quantum dynamics satisfies the canonical commutation relations and the Heisenberg evolution equation, $dA/dt = i[H(t); A]$. For the semi-quantum dynamics it is postulated that the classical commutation relations hold among the classical generators and similarly for the quantum sector. Furthermore, the generators of the classical sector commute with those of the quantum one. In other words, the commutation relations are as follows:

$$[q_i; q_j] = [p_i; p_j] = 0; \quad [q_i; p_j] = i \delta_{ij} E; \quad i, j = 1; \dots; N; \quad (1)$$

where δ_{ij} is zero if i is the label of a classical degree of freedom, and unity (or \hbar) if it labels a quantum one. These are the defining identities of the algebra of the semi-quantized system.

This axiom can be justified as follows. Certainly, eqs. (1) are natural if the semi-quantized system consists of a classical sector and a quantum sector without any interaction among them; this is a physical assumption. Since both in classical and quantum dynamics the commutation relations are unaffected by the choice of the interaction, one should expect that this is true as well in the semi-quantized case, and hence eqs. (1) follow. Another argument can be given by introducing a second physical assumption, namely, that the coupling among the two sectors can be switched on and off by playing with suitable time dependent coupling constants. Now, we can imagine starting with an uncoupled system, which satisfies the relations (1), switching on the interaction to end up with any given fully coupled system. Since the commutation relations are preserved by time evolution (even for non conservative dynamics) eqs. (1) will hold too in an arbitrary coupled semi-quantized system. We think that these considerations make axiom (iii) inescapable.

Now, from the previous considerations, a quite strong result can be derived, namely, the subalgebra spanned by the classical sector is invariant under time evolution. To simplify the notation, let us consider a system with just two degrees of freedom, one of them, $(q_1; p_1)$, quantum and the other $(q_2; p_2)$ classical, i.e., $\epsilon_1 = 1$ and $\epsilon_2 = 0$ in eqs. (1). Further, the coordinates and momenta at $t = t_0$ are denoted by q_i and p_i , respectively. For any time t , the set $\{E; q_1(t); p_1(t); q_2(t); p_2(t)\}$ generates the whole algebra A , and $q_2(t)$ commutes with all of them from eqs. (1), thus $q_2(t)$ commutes with all the algebra elements and, in particular, with q_1 and p_1 , and the same holds for $p_2(t)$. On the other hand, again using the commutation relations, every $A \in A$ is uniquely characterized by a set of coefficients $\alpha_{k'm'n}$, $k, m, n = 0; 1; \dots$, as $A = \sum_{k'm'n} \alpha_{k'm'n} q_1^k p_1^m q_2^n p_2^n E$. It is immediate to see that any element commuting with q_1 cannot contain p_1 and vice versa. Therefore, $q_2(t)$ must be of the form $\sum_{m,n} c_{m,n}(t) q_2^m p_2^n E$, and similarly $p_2(t)$. In other words, q_2 and p_2 are commuting objects which evolve by themselves following classical trajectories, without fluctuations. On the other hand, $q_1(t)$ and $p_1(t)$ may depend on $q_2(t)$ and $p_2(t)$ which, in this regard, behave as external sources.

One realization of the above picture is the traditional approach to semi-quantization, namely, the quantum degrees of freedom move in the presence of the classical background. On the other hand, the classical degrees of freedom are coupled to the expectation values of the quantum variables. For instance, if the system consists of two coupled harmonic oscillators, the equations of motion take the following form:

$$\frac{dq_1(t)}{dt} = \frac{p_1(t)}{m_1}; \quad \frac{dp_1(t)}{dt} = -m_1 \omega_1^2 q_1(t) - g(t) q_2(t); \quad (2a)$$

$$\frac{dq_2(t)}{dt} = \frac{p_2(t)}{m_2}; \quad \frac{dp_2(t)}{dt} = -m_2 \omega_2^2 q_2(t) - g(t) \langle q_1(t) \rangle_{\psi_1}; \quad (2b)$$

Here, ψ_1 is the state of the quantum sector in Heisenberg picture, i.e., a certain time independent wavefunction in the Hilbert space of q_1 and p_1 . Technically, our axioms apply here by considering $\langle q_1 \rangle_{\psi_1}$ and $\langle p_1 \rangle_{\psi_1}$ as fixed parameters, that is, independent of $q_{1,2}$ and $p_{1,2}$. Indeed, we can take expectation values of eqs. (2a) in ψ_1 , and solve the resulting system for $q_2(t)$ and $p_2(t)$; they will depend dynamically on q_2, p_2 and t (as well as on the fixed parameters $\langle q_1 \rangle_{\psi_1}$ and $\langle p_1 \rangle_{\psi_1}$). Substituting the classical solution in eqs. (2a), $q_1(t)$ and $p_1(t)$ are obtained as functions of q_1, p_1, q_2, p_2 and t . Afterwards, to extract meaningful physical results, one must choose precisely ψ_1 as the state of the quantum sector in Heisenberg picture, but this is not required by our axioms. It is immediate to check that eqs. (2) preserve the commutation relations (1): $q_2(t)$ and $p_2(t)$ are just ordinary functions and hence are commuting objects; $q_1(t)$ and $p_1(t)$ describe a purely quantum harmonic oscillator coupled to an applied external force $-g(t)q_2(t)$.

If one insisted in keeping the operator $q_1(t)$ in eq. (2b), instead of its expectation value, i.e.,

$$\frac{dp_2(t)}{dt} = -m_2 \omega_2^2 q_2(t) - g(t) q_1(t); \quad (3)$$

a violation of the commutation relations would result. For instance, assuming that eqs. (1) hold at $t=0$, and, there, treating $q_{1,2}$ as free parameters, one would find $[p_1(t); p_2(t)] = (\omega_1 - \omega_2)ig(0)tE + O(t^2)$, which only vanishes if either $g=0$ and thus the two subsystems are decoupled, or else if $\omega_1 = \omega_2$, i.e., the purely classical case if they vanish, or the purely quantum case, if they do not. Similarly, $[q_1(t); p_2(t)]$ would break down at $O(t^2)$.

3. The canonical structure of both classical and quantum mechanics (Poisson bracket and commutator, respectively) has been invoked in the literature [1,4], as a guiding principle to define semi-quantized theories. From this point of view, it is of interest to consider whether there exist canonical structures interpolating between the quantum and the classical limits.

Let us then study which new constraints are found if, in addition to previous assumptions (i-iii), a canonical structure is present. For convenience, the relations (1) are rewritten in the form:

$$[\xi_i; \xi_j] = E_{ij}; \quad \xi_i = 1; \dots; 2N; \quad (4)$$

where the single symbol ξ_i has been introduced to denote both q_i and p_i , and E_{ij} is an antisymmetric tensor with complex components.

The canonical structure is introduced by three new postulates. First, there exists (iv) a Lie bracket $(;)$ in A , which generates the (infinitesimal) canonical transformations by $A \cdot B = (A; B)$, $A; B \in A$, and in particular time evolution is a canonical transformation

$$\frac{dA(t)}{dt} = (H(t); A(t)); \quad (5)$$

where $H(t) \in A$ is the Hamiltonian of the system. Second, it is assumed (v) that the canonical transformations are algebra automorphisms. Of course, axioms (iv) and (v) imply (ii). And third, the following canonical relations are assumed (vi):

$$(q_i; q_j) = 0; \quad (p_i; p_j) = 0; \quad (q_i; p_j) = \delta_{ij}; \quad i, j = 1; \dots; 2N; \quad (6)$$

where δ_{ij} is the usual symplectic matrix, namely, zero for $(q; q)$ or $(p; p)$ and δ_{ij} for $(q_i; p_j)$, common to classical and quantum mechanics.

Since $(;)$ is a Lie bracket, it is bilinear, antisymmetric and satisfies Jacobi's identity. This is a consistency requirement among canonical transformations, which guarantees that $A(B; C) = (A; B; C) + (B; A; C)$, i.e., the bracket itself is invariant. In particular, the relationship $(A; B)(t) = (A(t); B(t))$ will be consistent with the equations of motion. The endomorphism property of the canonical transformations implies that the Lie bracket is a derivation in A , i.e., it satisfies the product (Leibniz) rule: $(A; BC) = (A; B)C + B(A; C)$. From here it is immediate to deduce that E is invariant under canonical transformations, that is, $A \cdot E = (A; E) = 0$. A consequence of the two previous observations is that the canonical relations between the ξ_i are preserved by canonical transformations.

As noted in [7] (see also [8]), the brackets defined in Refs. [1] or [4] are not derivations, thus the algebraic structure among observables is not preserved under time evolution (thus, violating axiom (ii)) and this seems unphysical. Also they are not Lie brackets, since they fail to satisfy Jacobi's identity. As a consequence, the canonical relations are not preserved either. Actually, the bracket defined in [4] is not even antisymmetric, hence, in general, the energy is not conserved even by time independent Hamiltonians, and hermiticity is broken by the dynamic evolution [7,8].

We still have to determine in which cases the canonical structure is consistent with axioms (i-iii). Using only linearity, antisymmetry, the product rule and the canonical relations (6), the bracket of every two observables can be worked out and $(;)$ becomes completely determined. Hence, it can be checked whether it admits the commutation relations (4). Indeed, arbitrary canonical transformations of both sides in (4) must coincide. For any $\xi_i; \xi_j = 1; \dots; 2N$, we find:

$$0 = (E_{ij}; E) = (E_{ij}; [q; p]) = (E_{ij}; q) \cdot (E_{ij}; p) + (E_{ij}; p) \cdot (E_{ij}; q) \quad (7)$$

The last line follows from repeatedly applying the product rule. Contracting this equation with δ_{ij} , the inverse matrix of E_{ij} , one concludes that consistency is only achieved if $\delta_{ij} = i$ for some i . In fact, from eqs. (1),

all the α_i are equal to α . In other words, there can be just one sector. Furthermore, the commutator of every two observables can also be worked out using eqs. (4); it follows that $[A;B] = i\alpha(A;B)$, for arbitrary $A;B$. There are only two possibilities: first that α is non vanishing. In this case, we end up with the usual purely quantum dynamics. Second, if α vanishes, all variables are commuting. Moreover, since the bracket is completely determined, it coincides with the Poisson bracket. That is, the dynamics is purely classical.

Note that this result is consistent with that found regarding eq. (3), namely, the canonical evolution generated by an arbitrary quadratic Hamiltonian does not preserve the semi quantized commutation relations (1).

The previous result means that there are no quotient algebras of A_L of the form (4), and supporting a canonical structure, which mix the classical and the quantum cases. In passing, it can be proven [9], that the bracket defined in A_L by using only linearity, antisymmetry, Leibniz rule and the canonical relations (6), satisfies Jacobi's identity as a byproduct, and thus this will be true as well for any of the quotient algebras considered here if and only if the bracket preserves the characteristic identities of that quotient algebra.

4. We conclude that, assumptions (i-iii) prevent the classical sector from inheriting quantum fluctuations and further, assumptions (i-vi) actually discard any non-trivial semi quantized theory. Note that further details on how to actually extract physical information from the observables (e.g., expectations values in the quantum case) are not required to reach the previous conclusion. We comment that the approach in Ref. [5], based directly on time ordered vacuum expectation values, is also flawed since it breaks physical positivity of the expectation values. It is entirely possible that there is no non-trivial (or at least elegant) semi quantization scheme, since, after all, such a concept is not presently known to be physically required.

The author thanks J. Caro and C. Garcia-Recio for useful criticism and for providing the proof in Ref. [9], and also J. Kruytzer for calling my attention to Refs. [6,8]. This work was partially supported by the Spanish DGIYT under contract PB92-0927 and the Junta de Andalucía.

-
- [1] W. Boucher and J. Traschen, Phys. Rev. D 37 3522 (1988).
 - [2] R. H. Brandenberger, Rev. Mod. Phys. 57, 1 (1985).
 - [3] I. V. Aleksandrov, Z. Naturforsch. 36A, 902 (1981).
 - [4] A. Anderson, Phys. Rev. Lett. 74, 621 (1995).
 - [5] L. L. Salcedo, Granada Univ. preprint UG-DFM-28/94, bull. board hep-th/9410007.
 - [6] L. Diosi, bull. board quant-ph/9503023.
 - [7] L. Diosi, Comment on 'Quantum Backreaction on 'Classical' Variables', submitted to Phys. Rev. Lett.
 - [8] A. Anderson, Reply to Comment on 'Quantum Backreaction on 'Classical' Variables', submitted to Phys. Rev. Lett.
 - [9] J. Caro, private communication.