

NSF-ITP-95-129
 CPTH-S379.1095
 hep-th/9510094

NATURAL GAUGE-COUPLING UNIFICATION AT THE STRING SCALE

C. Bachas^{a,b}, C. Fabre^b and T. Yanagida^{a,c}

^a Institute for Theoretical Physics, University of California, Santa
 Barbara, CA 93106, USA

^b Centre de Physique Theorique, Ecole Polytechnique, 91128 Palaiseau,
 FRANCE

^c Department of Physics, University of Tokyo, Tokyo 113, JAPAN

ABSTRACT

We argue that moduli in the adjoint representation of the standard model gauge group are a natural feature of superstring models, and that they can account for the apparent discrepancy between the string and unification scales.

NSF-ITP-95-129
 CPTH-S379.1095
 October 1995

An important hint in favour of supersymmetric GUTs is the unification of gauge couplings extrapolated from their low-energy experimental values under the assumptions of minimal particle content and superpartner thresholds at 1 TeV [1]. The scale at which the couplings meet ($M_{\text{GUT}}^{\text{minimal}} \approx 2 \times 10^{16} \text{ GeV}$) lies, however, one order of magnitude below the heterotic string scale ($M_{\text{str}} \approx 0.5 g_{\text{str}}^{-2} \approx 10^{18} \text{ GeV}$) [2], suggesting that there should exist some string states (beyond those of the MSSM) which are significantly lighter than M_{str} . It has been proposed that these light states can be exotic vector-like quarks and leptons with non-conventional assignment of hypercharge [5], or else the extra gauge bosons and Higgses of a unifying symmetry broken somewhere below M_{str} [6, 7]. In this note we would like to point out a natural alternative: a color-SU(3) octet and a color-neutral triplet of weak-SU(2), both having zero hypercharge. As we will argue (a) these appear in many string models as continuous moduli which is why they can remain light naturally, and (b) they push the unification scale up to M_{str} when their masses lie in the best-motivated intermediate range $M_{\text{Planck}}^{2/3} m_{\text{susy}}^{1/3} \approx 10^{13} \text{ GeV}$. Furthermore contrary to exotic stable remnants they present no danger for cosmology.

The one-loop running coupling constants in the presence of an adjoint scalar multiplet read [8]:

$$\begin{aligned} \frac{2}{1(\gamma)} - \frac{2}{1(m_Z)} &= \left(\frac{1}{2} + \frac{2}{3}N_g\right) \log\left(\frac{m_{\text{susy}}}{m_Z}\right) - \left(2N_g + \frac{3}{5}\right) \log\left(\frac{m_{(3)}}{m_Z}\right) \\ \frac{2}{2(\gamma)} - \frac{2}{2(m_Z)} &= \left(\frac{4}{3} + \frac{2}{3}N_g + \frac{5}{6}\right) \log\left(\frac{m_{\text{susy}}}{m_Z}\right) + \\ &\quad + (6 - 2N_g - 1) \log\left(\frac{m_{(3)}}{m_Z}\right) - 2 \log\left(\frac{m_{(3)}}{m_{(3)}}\right) \end{aligned}$$

Note however that their mass can be pushed up if they are very numerous [3]. They may even be unnecessary if one allows a non-standard hypercharge normalization and an α_3 at the higher end of its allowed range [4].

and

$$\frac{2}{3}(\gamma) - \frac{2}{3}(\gamma_Z) = (2 + \frac{2}{3}N_g) \log\left(\frac{m_{\text{susy}}}{m_Z}\right) + (9 - 2N_g) \log\left(\frac{m_{(8)}}{m_Z}\right) - 3 \log\left(\frac{m_{(3)}}{m_{(8)}}\right)$$

where $m_{(8)}$ and $m_{(3)}$ are the masses of the color octet and weak triplet, respectively. Using as input the experimental values of the couplings at m_Z (with $\gamma = 0.116 \pm 0.005$) [9] and assuming a usual $m_{\text{susy}} \sim 1 \text{ TeV}$, one can calculate from the above equations the required masses of these extra particles such that couplings unify at the string scale. Taking $M_{\text{str}} \sim 6 \cdot 10^{17} \text{ GeV}$, corresponding to a $k = 2$ Kac-Moody level,^y we find: $m_{(3)} \sim 5.3 \cdot 10^{12} \text{ GeV}$ and $m_{(8)} \sim 2.8 \cdot 10^{12} \text{ GeV}$. Surprisingly enough these masses are not only close to each other, but also of the order of magnitude one would expect if supersymmetry breaking were induced by condensing gauginos [10]. Our underlying assumptions are of course (i) standard hypercharge normalization as in the usual $SU(5)$ embedding, and (ii) zero vevs for the adjoint moduli, consistent with an unbroken standard model gauge group.

Such adjoint moduli are an ingredient of all the recently constructed models based on a group structure $G = G$ [11, 12, 13]. They are the relics after truncation of extended supersymmetric vector multiplets, which is why their potential stays flat [14]. To illustrate this point explicitly consider a $N = 2$ supersymmetric pure gauge theory with gauge group $SO(2n)$. We may define in this theory two parity operations, $(\gamma)^A$ and P , where A counts the number of vector indices under the first factor in the embedding $SO(2n) = SO(n)_A \times SO(n)_B$, while P breaks down a $N = 2$ vector multiplet into an even vector and an odd scalar superfield of $N = 1$. Imposing the combined parity projection $(\gamma)^A P = +1$ leads to a truncated $N = 1$ supersymmetric theory with gauge group $SO(n)_A \times SO(n)_B$ and with extra chiral multiplets in the representation $(n; n)$. A heritage of the extended supersymmetry is that these latter have a flat potential along directions lying inside the Cartan subalgebra of the original $SO(2n)$. By turning on a

^yAdjoint scalars require that k be at least equal to two. Higher values push the string scale up and would demand somewhat lighter intermediate masses.

non-vanishing vev for the trace of the $(n;n)$ matrix, we can break the gauge symmetry to the diagonal subgroup $SO(n)_{\text{diag}}$. As a minute's thought will convince the reader, the resulting theory still has $SO(n)_{\text{diag}}$ -adjoint moduli. Note that the level of the corresponding algebra is the sum of the levels of the two group factors, and hence it is at least equal to two. Note also that in string theory a Z_2 truncation gives rise to new (twisted) states. These however only appear in pairs, and thus do not spoil the flat directions as long as their expectation values vanish.

The original motivation [6, 7] for constructing $G \times G$ string models was to allow minimal unification by enlarging the symmetry at a scale significantly lower than M_{str} . As we have just argued the existence of adjoint moduli turns this motivation around on its head: it invalidates the desert hypothesis and, in the absence of other light states, can render premature unification unnecessary. Our argument also carries over to those string GUT models, constructed directly at $k > 1$ [12, 15], which are just stringy realizations of the last $G \times G \rightarrow H_{\text{diag}}$ breaking step [6]. More generally, it is hard to imagine how a unifying gauge symmetry can break at a scale an order of magnitude below M_{str} if not along some flat directions which would leave behind light extra matter populating the desert. Thus it might be better motivated (and safer with respect to proton decay) to search for models whose observable light states include only a $k > 1$ MSSM and adjoint moduli.

Acknowledgments

This work was supported by the NSF under grant no. PHY 94-07194, and by EEC grants CHRX-CT 93-0340 and SC1-CT 92-0792. We thank our colleagues at the ITP workshop for enlightening discussions, and in particular K. Dienes and P. Langacker for a careful reading of the manuscript.

References

- [1] S. Dimopoulos and H. Georgi, Nucl. Phys. B 193 (1981) 150; S. Dimopoulos, S. Raby and F. Wilczek, Phys. Rev. D 24 (1981) 1681;

- L. Ibanez and G. G. Ross, Phys.Lett. 105B (1981)439; N. Sakai, Z.Phys. C 11 (1981)153; J. Ellis, S. Kelley and D. V. Nanopoulos, Phys.Lett.B 249 (1990) 441; U. Amaldi, W. de Boer and H. Furstenau, Phys.Lett. B 260 (1991)447; P. Langacker and M. X. Luo, Phys.Rev. D 44 (1991)817; for a recent review see P. Langacker, hep-ph/9411247.
- [2] P. Ginsparg, Phys.Lett. 197B (1987)139; V. Kaplunovsky, Nucl.Phys. B 307 (1988)145; erratum ibid B 382 (1992)436.
- [3] For a recent discussion see for instance H. P. Nilles and S. Stieberger, hep-th/9510009, and references therein.
- [4] L. Ibanez, Phys.Lett. 318B (1993)73.
- [5] For a recent systematic discussion and extensive references see K. Dienes and A. Faraggi, Phys.Rev.Lett. 75 (1995)2646, hep-th/9505018; hep-th/9505046.
- [6] A. Font, L. Ibanez and F. Quevedo, Nucl. Phys.B 345 (1990)389; D. Lewellen, Nucl.Phys. B 337 (1990)61.
- [7] R. Barbieri, G. Dvali and A. Stumia, Phys.Lett. 333B (1994)79 and Nucl.Phys. B 435 (1995)102.
- [8] J. Hisano, H. Murayama and T. Yanagida, Phys.Rev.Lett. 69 (1992)1014.
- [9] J. Erler and P. Langacker, Phys.Rev. D 52 (1995)441.
- [10] J.-P. Derendinger, L. Ibanez and H. P. Nilles, Phys.Lett. B 155 (1985)65; M. Dine, R. Rohm, N. Seiberg and E. Witten, Phys.Lett. B 156 (1985)55.
- [11] A. A. Maslikov, S. M. Sergeev and G. G. Volkov, Phys.Lett. B 328 (1994)319; Phys.Rev. D 50 (1994)319; BUTP-94-24 preprint.
- [12] A. Aldazabal, A. Font, L. Ibanez and A. Uranga, hep-th/9410206 and hep-th/9508033.

- [13] D .F innell, hep-th/9508073.
- [14] I.Antoniadis, C .Bachas and C .K ounnas, NuclPhys. B 289 (1987)87.
- [15] S.Chaudhuri, S.W .Chung, G .Hockney and J.Lykken,
hep-ph/9501361; G .C leaver, hep-th/9506006.