arXiv:hep-th/9704181v1 25 Apr 1997

q-Functional Wick's theorems for particles with exotic statistics

K.N.Ilinski [1,2,3] *, G.V.Kalinin [1] † and A.S.Stepanenko [1,2,4] ‡

[1] *IPhys Group, CAPE, St.-Petersburg State University, 14-th line of V.I. 29, St-Petersburg, 199178, Russian Federation*

[2] *School of Physics and Space Research, University of Birmingham, Edgbaston B15 2TT, Birmingham, United Kingdom*

[3] *Theoretical Department, Institute of Spectroscopy, Troitsk, Moscow region, 142092, Russian Federation*

[4] *Theoretical Department, St.-Petersburg Nuclear Physics Institute, Gatchina, St.-Petersburg, 188350, Russian Federation*

Abstract

In the paper we begin a description of functional methods of quantum field theory for systems of interacting q-particles. These particles obey exotic statistics and are the q-generalization of the colored particles which appear in many problems of condensed matter physics, magnetism and quantum optics. Motivated by the general ideas of standard field theory we prove the q-functional analogues of Hori's formulation of Wick's theorems for the different ordered q-particle creation and annihilation operators. *The formulae have the same formal expressions as fermionic and bosonic ones but differ by a nature of fields.* This allows us to derive the perturbation series for the theory and develop analogues of standard quantum field theory constructions in q-functional form.

*E-mail: kni@th.ph.bham.ac.uk
†E-mail: kalinin@snoopy.phys.spbu.ru
‡E-mail: ass@th.ph.bham.ac.uk

1 Introduction

Last decade many mathematical structures were deformed and gained the subscript q in their notations. In this way remarkable mathematical objects such as non-commutative geometries [1], quantum groups (q-groups) [2] and their representations on quantum vector spaces [3] have arisen. These objects led to the investigations of the quantum group gauge theory [4], q-deformed Schrodinger equation [5] and classical and quantum dynamics on q-deformed phase spaces [6]. In Refs. [7, 8] q-deformed spaces were considered as graded-commutative algebras. On this basis the classical and quantum dynamics on q-deformed spaces were proposed in close analogy with the case of $\mathbf{Z}_2$-commutative spaces (Grassmann algebras) [8].

After quantization the particles satisfy the q-deformed commutation relations similar to the commutation relations of the colored particles [9]:

$$\begin{array}{l} a_k a_j \mp q a_j a_k = 0 \ , \quad a_k a_j^+ \mp q^{-1} a_j^+ a_k = 0 \ , \quad q = e^{i\alpha} \ , \quad 1 \leq k < j \leq n \ , \\ a_k a_k^+ \mp a_k^+ a_k = 1 \qquad \text{and for } q\text{-deformed fermions} \quad [a_k^+]^2 = [a_k]^2 = 0 \ . \end{array} \tag{1}$$

where the upper sign corresponds to the quantization on q-deformed common space and the lower one corresponds to the quantization on q-deformed Grassmann algebra. These particles were called q-particles to distinguish them from the famous q-oscillators which a number of papers are devoted to. These q-particles are the only objects of our consideration.

It is surprising that q-particles are interesting not only from the mathematical point of view but also from the physical one. Here we do not mean the appearance of q-particles in the parastatistics [7, 8, 10], q-extended Supersymmetry [11], Parasupersymmetry [11, 12] and other similar problems because they are rather mathematical applications. By physical applications we mean firstly solid state physics and quantum optics where q-particles appear in a natural way as well as the theory of magnetics where Paulions originally appeared and play a central role.

In solid state physics, anyons (particles with exotic braiding statistics) are important in some attempts to understand the physical features of planar systems [13]. The main physical interest in anyon systems is their possible connection with some effects in two dimensional condensed matter physics in particular in the quantum Hall effect [14] and high temperature superconductivity [15] (in the framework of the investigations of the t-J and Hubbard models).

In contrast to the previous example where anyons serve as auxiliary objects for the construction of one of possible scenarios there is a wide field in the quantum nonlinear optics in which q-particles are the main components. This is a theory of the collective behavior of excitons with small radius (Frenkel Excitons and Charge-Transfer Excitons (CTE)) [16]. The studies investigate possibilities of formation of the Frenkel biexcitons and the observation of phase transitions in exciton systems in molecular crystals (Bose- Einstein condensation of excitons [17], structural phase transitions in crystals with high excitonic concentrations, dielectric-metal phase transition in a system of interacting CTE [18] and others). Strictly speaking excitons are not particles. They are quasiparticles describing molecular excitations

and are of great importance in the analysis of nonlinear optical processes which accompany propagation of high-intensity light fluxes whose frequencies are in the range of the exciton absorption bands [19]. Moreover excitons obey exotic statistics (Pauli statistics) [20] coinciding with q-particles statistics for $q = -1$. The general case of $q = e^{i\alpha}$ arises if we try to take into account phenomenologically some nonlinear effects (such as the difference in the creation time of molecular excitations for different types of molecules). This effect can be modeled by the change of the Paulion commutation relations to those of the q-particles using the method developed in [21].

Surprisingly, even the investigation of the behavior of low dimensional exciton systems is meaningful. For example the exact solutions for 1-dimensional Paulion chains [22] caused great advances in the theory of the so called J-aggregates, *i.e.* molecular aggregates with unusually sharp absorption band ([23] and Refs. therein). The investigations of exciton systems on interfaces closely connect with the successes of contemporary technology . All these show that q-particles find deep applications in modern physical theories and motivate our objective to derive the appropriate field theoretical technique for them.

The paper is organized as follows. In the next section we motivate the necessity for the introduction of q-ordered objects (such as the normal product) and introduce useful notation. Then in the sections 3 and 4 we prove Wick's theorems in q-functional form for the simple product and q-symmetrical products of q-operators (q-symmetrical and q-chronological products). This allows us to find in section 5 the Wick's theorems for the q-operator functionals. The last section contains the conclusions and some remarks.

2 Motivation and notations

In this section we motivate the necessity to introducing q-objects (such as N-, T-products of q-operators and others) which we study in the following sections. We also define "universal" notations following [24] that allow us to prove Wick's theorems for the cases of q-operators with internal degree of freedom (spin, colors) in the unified way.

As in the usual Fermi-Bose situation in q-particle physics we are interested in two kinds of physical quantities. The first one is generally speaking the class of equilibrium thermodynamical characteristics calculated via the partition function of the system. The second one is a set of correlators which are important in describing the kinetics.

For clarity we consider a system of q-particles with creation (annihilation) operators obeying the commutation relations (1). The equilibrium thermodynamics of the model is described by the statistical operator $\rho = e^{-\beta H}$ and the partition function $Z = \mathrm{Tr}\rho$. In close analogy with undeformed many-particle quantum theory there are several approaches for calculating Z. The comparative convenience of each of them is defined by the specific features of the system in question. The most straightforward way is to present ρ by the series

$$\rho = \sum_{n=0}^{\infty} \frac{(-\beta)^n}{n!} H^n$$

and to find matrix elements of the operators H^n in the occupation number basis $\{|\varphi_{i_1,i_2,\ldots,i_m}\rangle\}$ $= \{a^+_{i_1}a^+_{i_2}\ldots a^+_{i_m}|0\rangle\}$ and then to try to calculate the sum of all diagonal matrix elements in this basis. Usually it is difficult enough to calculate the sum in a closed form.

To escape this problem one can introduce the basis coherent states corresponding to creation (annihilation) operators (1) [25]. These states use elements of the graded-commutative algebra $HQ^{0|L}$ (or $HQ^{L|0}$ in the bosonic case) as parameters. By proceeding this way we need to calculate matrix elements of the operator in the Bargmann-Fock representation. This means we need to find a representation of the operator in N-ordering form in which all creation operators are on the right-hand side with respect to all annihilation operators in all monomial terms. This can be done using the Wick's theorem for q-operators in the same way that it was done for the usual Fermi-Bose statistics [24]. The next section is devoted to the formulation and proof of the theorem.

It is a very natural idea to use the objects of the non-commutative algebra for the investigations of systems with quantum particles obeying some exotic statistics. The first example of its useful application is in fermionic physics. Moreover all functional methods of quantum bosonic field theory can be modified for the treatment of quantum fermionic fields using Grassmann variables instead of the usual complex numbers. This allows us to hope that the methods can be adopted for particles with exotic statistics (q-statistics) if permutation relations for the classical analogues are defined according to the rules of the statistics. In this way we can obtain a self-consistent scenario for building the quantum field theory (QFT) of particles with exotic statistics.

In this paper we consider creation and annihilation operators obeying the following commutation relations which generalize (1):

$$\left.\begin{array}{l} \hat{a}(\mathbf{x})\hat{a}(\mathbf{y}) - \kappa q(\mathbf{x},\mathbf{y})\hat{a}(\mathbf{y})\hat{a}(\mathbf{x}) = 0 \ , \\ \hat{a}^\dagger(\mathbf{x})\hat{a}^\dagger(\mathbf{y}) - \kappa q(\mathbf{x},\mathbf{y})\hat{a}^\dagger(\mathbf{y})\hat{a}^\dagger(\mathbf{x}) = 0 \ , \\ \hat{a}(\mathbf{x})\hat{a}^\dagger(\mathbf{y}) - \kappa q^*(\mathbf{x},\mathbf{y})\hat{a}^\dagger(\mathbf{y})\hat{a}(\mathbf{x}) = \delta(\mathbf{x},\mathbf{y}) \ , \\ [\hat{a}(\mathbf{x})]^2 = [\hat{a}^\dagger(\mathbf{x})]^2 = 0 \ , \quad \text{for } q\text{-fermions} \ , \end{array}\right\} \tag{2}$$

This form allows us to consider continuous indices as well as discrete ones. $\mathbf{x}$, $\mathbf{y}$ are in general D-dimensional vectors (or/and D-dimensional multi-indices for a lattice) which describe external and internal degree of freedom. The function $\delta(\mathbf{x},\mathbf{y})$ is δ-function $\delta(\mathbf{x}-\mathbf{y})$ for a continuous space and Kronecker δ-function for a lattice or internal color indices. The statistical factor $q(\mathbf{x},\mathbf{y})$ possesses the following property generalized Paulionic and Anyonic property:

$$q(\mathbf{x},\mathbf{y}) = q^{-1}(\mathbf{y},\mathbf{x}) = q^*(\mathbf{y},\mathbf{x}) \ , \qquad q(\mathbf{x},\mathbf{x}) = 1 \ . \tag{3}$$

Finally κ serves to unify formulae for deformed bosonic and deformed fermionic cases. As usual it has the form:

$$\kappa = \left\{ \begin{array}{ll} +1 \ , & \text{for } q\text{-bosons} \\ -1 \ , & \text{for } q\text{-fermions} \ . \end{array}\right. \tag{4}$$

Hereafter letters with hats denote operators and those without hats denote the corresponding classical variables. For operator algebra (2) the corresponding classical variables satisfy the

following permutation relations:

$$\left.\begin{array}{l} a(\mathbf{x})a(\mathbf{y}) - \kappa q(\mathbf{x},\mathbf{y})a(\mathbf{y})a(\mathbf{x}) = 0 \ , \\ a^\dagger(\mathbf{x})a^\dagger(\mathbf{y}) - \kappa q(\mathbf{x},\mathbf{y})a^\dagger(\mathbf{y})a^\dagger(\mathbf{x}) = 0 \ , \\ a(\mathbf{x})a^\dagger(\mathbf{y}) - \kappa q^*(\mathbf{x},\mathbf{y})a^\dagger(\mathbf{y})a(\mathbf{x}) = 0 \ , \\ [a(\mathbf{x})]^2 = [a^\dagger(\mathbf{x})]^2 = 0 \ , \quad \text{for } q\text{-fermions} \ , \end{array}\right\} \tag{5}$$

Let us introduce "universal" notations to escape repetitions and enable us formulate statements in unified form for different cases. Following [24] we collect the creation and annihilation operators into the single vector "field" operator:

$$\hat{\varphi}(x) = \left(\begin{array}{c} \hat{a}^\dagger(\mathbf{x}) \\ \hat{a}(\mathbf{x}) \end{array} \right) \ , \quad x \equiv (s, \mathbf{x}) \ , \tag{6}$$

such that the additional vector index s indicates a type of operator ($\hat{a}^\dagger$ (when $s = 1$) or $\hat{a}$ (when $s = 2$)). It is convenient to represent quantities by vectors or matrices with respect to the index s.

Using this notation the commutation relations (2) can be rewritten in the following form:

$$\left.\begin{array}{l} \hat{\varphi}(x)\hat{\varphi}(y) - \kappa Q(x,y)\hat{\varphi}(y)\hat{\varphi}(x) = u(x,y) \ , \\ [\hat{\varphi}(x)]^2 = 0 \ , \quad \text{for } q\text{-fermions} \ , \end{array}\right\} \tag{7}$$

where the statistical matrix $Q(x, y)$ is defined by statistical factor $q(\mathbf{x}, \mathbf{y})$:

$$Q(x,y) = \left(\begin{array}{cc} q(\mathbf{x},\mathbf{y}) & q^*(\mathbf{x},\mathbf{y}) \\ q^*(\mathbf{x},\mathbf{y}) & q(\mathbf{x},\mathbf{y}) \end{array} \right) \ , \tag{8a}$$

and the quantum deformation parameter matrix is given by the equality:

$$u(x,y) = \delta(\mathbf{x},\mathbf{y}) \left(\begin{array}{cc} 0 & -\kappa \\ 1 & 0 \end{array} \right) \ . \tag{8b}$$

The row number of the matrices from (8a) and (8b) corresponds to s_x and the column number to s_y. We can now associate classical analogues — classical vector "field" variables – with the corresponding vector "field" operator. For the classical variables we get the permutation relations in the following "universal" form:

$$\left.\begin{array}{l} \varphi(x)\varphi(y) - \kappa Q(x,y)\varphi(y)\varphi(x) = 0 \ , \\ [\varphi(x)]^2 = 0 \ , \quad \text{for } q\text{-fermions} \ , \end{array}\right\} \tag{9}$$

which is defined by the statistical matrix $Q(x, y)$ from (8a).

To formulate Wick's theorems in the functional form we need to introduce left ($\frac{\vec{\partial}}{\partial\varphi(x)}$) and right ($\frac{\overleftarrow{\partial}}{\partial\varphi(x)}$) functional derivatives on the algebra (9). We do as was done for the Grassmann algebra case: to act by the left (right) derivative on an expression we must bring the corresponding variable to the extremely left (right) position in accordance with the permutation relations and cancel it. This leads to the following commutation relations for the left derivative:

$$\frac{\vec{\partial}}{\partial\varphi(x)}\varphi(y) - \kappa Q^*(x,y)\varphi(y)\frac{\vec{\partial}}{\partial\varphi(x)} = \delta(x,y) \ , \tag{10}$$

where $\delta(x,y) = \delta_{s_x,s_y}\delta(\mathbf{x},\mathbf{y})$.

To construct Hori's functional expressions we will also need extended algebras which will contain several copies of the algebra of our vector "field" variables as sub-algebras. The mutual permutation relations for the sub-algebras we define to keep the graded-commutative structure. For example, the pair of fields $\varphi_1(x), \varphi_2(x)$ like (9) from different sub-algebras commute with one another as follows:

$$\varphi_1(x)\varphi_2(y) - \kappa Q(x,y)\varphi_2(y)\varphi_1(x) = 0 \ . \tag{11}$$

We are now ready to proceed with Wick's theorems for the operators (7).

3 Wick's theorem for the simple product of q-operators

In this section we formulate and prove the analog of Hori's formula [26] which gives the functional form of the Wick's theorem for the normal form of a simple product of the creation and annihilation (or vector "field") operators. *The formal expression does not depend on the statistics of the fields.* It shows that the exotic statistics can be taken into account by changing the permutation relations for the classical analogues according to the rules of the statistics and keeping the formal expression for the Hori's formulae.

As was said above we define the normal form $N(A)$ of the monomial operator A as an expression where all creation operators are placed on the left hand side in respect to the annihilation operators (using permutation relations (5)). Then the definition is generalized for an arbitrary polynomial operator by the linearity. The definition is standard [24].

As usual we can do this in the vector "field" form (in the "universal" notations) where the final expressions only contain the normal contractions $n(x,y)$. For the pair of "field" operators the normal product is defined by the following relation:

$$\mathrm{N}[\hat{\varphi}(x)\hat{\varphi}(y)] = \hat{\varphi}(x)\hat{\varphi}(y) - n(x,y) \ , \tag{12}$$

For system like (2) the normal contraction hence has the form:

$$n(x,y) = \delta(\mathbf{x},\mathbf{y}) \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} \tag{13}$$

We note that the normal contraction has δ-functional (or δ-symbol) character with respect to the indices. In the following we will frequently use this fact. More precisely we will use the fact that the product $\varphi(x)\varphi(y)$ is C-number function when $n(x,y) \neq 0$.

We now prove the following theorem which reduces the simple operator product to the normal form:

Theorem 1

$$\hat{\varphi}(x_1)\ldots\hat{\varphi}(x_n) = \mathrm{N}\left[\prod_{i<k}\left(1 + \frac{\overleftarrow{\partial}}{\partial\varphi_i} n \frac{\overleftarrow{\partial}}{\partial\varphi_k}\right)\varphi_1(x_1)\ldots\varphi_n(x_n)\Big|_{\ldots}\right] \tag{14}$$

where

1. the variables $\varphi_1, \ldots, \varphi_n$ form the extended algebra (9) following eq.(11),

2. The symbol $|_{\ldots}$ means substitution $\varphi_1 = \ldots = \varphi_n = \hat{\varphi}$,

3. Substituting the differential forms into eq.(14) in expanded form we have

$$\frac{\partial}{\partial\varphi_i} n \frac{\partial}{\partial\varphi_k} = \sum_{x,x'} \frac{\partial}{\partial\varphi_i(x)} n(x,x') \frac{\partial}{\partial\varphi_k(x')} \ . \tag{15}$$

(here the symbol $\sum_x$ means summation over discrete variables and integration over continuous ones).

Proof. The strategy of the proof coincides with that of the proof of the Fermi-Bose Hori's formulae [24] and is fulfilled by induction. Indeed, for the particular cases $n = 1, 2$ the statement can be proved immediately:

$$\left.\begin{array}{l} \hat{\varphi}(x) = \mathrm{N}[\hat{\varphi}(x)] \ , \\ \hat{\varphi}(x_1)\hat{\varphi}(x_2) = \mathrm{N}[\hat{\varphi}(x_1)\hat{\varphi}(x_2) + n(x_1,x_2)] \ , \end{array}\right\} \tag{16}$$

by the application of the definitions of the normal form and the normal contraction (12,13).

Let us now assume that eq.(14) is true for any $n \leq N$ and consider the product of $N+1$ operators $\hat{\varphi}(x_1)\ldots\hat{\varphi}(x_{N+1})$.

1. If the field $\hat{\varphi}(x_{N+1})$ contains only the second component $\hat{a}(\mathbf{x}_{N+1})$ then

$$\hat{\varphi}(x_1)\ldots\hat{\varphi}(x_{N+1}) = \hat{\varphi}(x_1)\ldots\hat{\varphi}(x_N)\hat{a}(\mathbf{x}_{N+1}) \tag{17}$$

By the inductive assumption the first N multipliers in the RHS of eq.(17) can be mapped to the normal form by the following reduction operator $\overleftarrow{\mathcal{P}}_N$:

$$\overleftarrow{\mathcal{P}}_N = \prod_{1\leq i<k\leq N} \left(1 + \frac{\overleftarrow{\partial}}{\partial\varphi_i} n \frac{\overleftarrow{\partial}}{\partial\varphi_k}\right) \ . \tag{18}$$

Then the full expression (17) also takes the normal form. It is not difficult to see that due to the structure of normal contraction (13) the differential expression (15) in (14) does not contain the derivative on $\hat{\varphi}^{(2)}(x_{N+1}) = \hat{a}(\mathbf{x}_{N+1})$. Hence the terms with derivative $\frac{\overleftarrow{\partial}}{\partial\varphi_{N+1}}$ do not contribute in this case. So we can substitute $\overleftarrow{\mathcal{P}}_N$ by $\overleftarrow{\mathcal{P}}_{N+1}$ what finishes the proof for this case.

2. Let us now consider the case of $\hat{\varphi}(x_{N+1}) = \hat{a}^\dagger(\mathbf{x}_{N+1})$. Using N times the commutation relations (7) we obtain the following relation:

$$\begin{array}{rcl} \hat{\varphi}(x_1)\ldots\hat{\varphi}(x_{N+1}) & = & Q\hat{a}^\dagger(\mathbf{x}_{N+1})\hat{\varphi}(x_1)\ldots\hat{\varphi}(x_N) + \\ & + & \displaystyle\sum_{k=1}^{N} Q_k n(x_k, x_{N+1})[\hat{\varphi}(x_1)\ldots\hat{\varphi}(x_N)]_k \ , \end{array} \tag{19}$$

where $[\hat{\varphi}(x_1)\ldots\hat{\varphi}(x_N)]_k$ means product $\hat{\varphi}(x_1)\ldots\hat{\varphi}(x_N)$ without multiplier $\hat{\varphi}(x_k)$ and the following notations

$$Q = \kappa^N Q(x_1, x_{N+1})\ldots Q(x_N, x_{N+1}) \ ,$$

$$Q_k = \kappa^{N-k} Q(x_{k+1}, x_{N+1}) \dots Q(x_N, x_{N+1})$$

were introduced. The multipliers Q and Q_k reflect the q-deformation of the statistics and are due to the commutation relations (7). The first summand in the RHS of eq.(19) can be rewritten as

$$\mathrm{N}[Q\hat{a}^\dagger(\mathbf{x}_{N+1}) \cdot \overleftarrow{\mathcal{P}}_N \varphi_1 \dots \varphi_N|_{\dots}] \tag{20}$$

Let us now note that under the sign of the normal products (20) the operator $\hat{a}^\dagger(\mathbf{x}_{N+1})$ can be substituted by the corresponding classical variable. Then if we move the variable $a^\dagger(\mathbf{x}_{N+1})$ to the original (right) position into expression (20) we gain the statistical multiplier Q^{-1} which is exactly cancels the multiplier Q. This is because the reduction operator $\overleftarrow{\mathcal{P}}_N$ commutes with $a^\dagger(\mathbf{x}_{N+1})$ due to the δ-character of the normal contraction and the permutation relations for the classical analogues give the same statistical phases as the quantum ones. So

$$\mathrm{N}\left[Q\hat{a}^\dagger(\mathbf{x}_{N+1})\overleftarrow{\mathcal{P}}_N \varphi_1(x_1) \dots \varphi_N(x_N)\Big|_{\dots}\right] = \mathrm{N}\left[\overleftarrow{\mathcal{P}}_N \varphi_1 \dots \varphi_{N+1}\Big|_{\dots}\right] \ . \tag{21}$$

Now consider the second summand in RHS of (19). By applying the inductive assumption and substituting normal contraction by the expression in parentheses we get:

$$\mathrm{N}\left[\sum_{k=1}^{N} Q_k \left(\frac{\overleftarrow{\partial}}{\partial\varphi_k} n \frac{\overleftarrow{\partial}}{\partial\varphi_{N+1}} \varphi_k \varphi_{N+1}\right) \overleftarrow{\mathcal{P}}_N [\varphi(x_1) \dots \varphi(x_N)]_k\right] \ .$$

Product $[\varphi(x_1) \dots \varphi(x_N)]_k$ does not contain fields φ_k, φ_{N+1}. Using arguments similar to those for the derivation of (21) we can move them to there original position and get

$$\mathrm{N}\left[\sum_{k=1}^{N} \frac{\overleftarrow{\partial}}{\partial\varphi_k} n \frac{\overleftarrow{\partial}}{\partial\varphi_{N+1}} \overleftarrow{\mathcal{P}}_N \varphi_1 \dots \varphi_{N+1}\Big|_{\dots}\right] \ . \tag{22}$$

Collecting together eqs.(19), (21) and (22) we get

$$\hat{\varphi}(x_1) \dots \hat{\varphi}(x_{N+1}) = \mathrm{N}\left[\overleftarrow{\mathcal{P}}_{N+1} \varphi_1 \dots \varphi_{N+1}\Big|_{\dots}\right] \ . \tag{23}$$

Hence the proof of Theorem 1 is completed. □

Corollary 1.1 Due to the linearity of the expression (14) on each variable φ_i the formulae (14) can be rewritten in the following form:

$$\hat{\varphi}(x_1) \dots \hat{\varphi}(x_n) = \mathrm{N}\left[\exp\left[\sum_{i<k} \frac{\overleftarrow{\partial}}{\partial\varphi_i} n \frac{\overleftarrow{\partial}}{\partial\varphi_k}\right] \varphi_1(x_1) \dots \varphi_n(x_n)\Big|_{\dots}\right] \tag{24}$$

Below we will use this relation to find the compact functional form for the Wick's theorems for the symmetrical and chronological products.

Corollary 1.2 Formulae (14, 24) hold true if we substitute right derivatives by left ones, sign "$<$" in product limits by "$>$" and normal contraction n by n^{T}.

Let us note here that the formal form of the statement of Theorem 1 does not depend on any statistics (*i.e.* is the same as in fermionic and bosonic cases) and the commutation factors are "hidden" in the nature of the classical variables. It is useful to remember this when studying all other theorems of the paper where it is also true.

4 Wick's theorem for q-symmetrical products

In this section we consider other kinds of products, in particular, q-symmetrical and q-chronological. Such products naturally play the role of the usual symmetrical for bosons (antisymmetrical for fermions) and chronological products in the framework of the standard quantum field theory. Moreover they are interesting by themselves because they keep the symmetry properties of the q-operators.

In close analogy with the undeformed cases we define q-Symmetrical product (Sym_q-product) of q-operators in the following way

$$\mathrm{Sym}_q[\hat{\varphi}(x_1)\dots\hat{\varphi}(x_n)] = \frac{1}{n!}\sum_P Q_P P[\hat{\varphi}(x_1)\dots\hat{\varphi}(x_n)] \ . \tag{25}$$

The sum is taken over all $n!$ permutations P of the q-operators $\hat{\varphi}$ with corresponding statistical phases Q_P. These Q_P are the factors arising from permutation of operators from the original order to the order P. We assume that under the permutation q-operators $\hat{\varphi}$ are replaced by the corresponding classical variables $\hat{\varphi} \to \varphi$ and we do not pick up the expressions due to the RHS of the eq.(7). In other words Q_P is defined from the relation

$$\varphi(x_1)\dots\varphi_n(x_n) = Q_P P[\varphi(x_1)\dots\varphi_n(x_n)] \ . \tag{26}$$

We now consider the q-chronological product. We assume that q-operators $\hat{\varphi}$ depend also on time t and define q-chronological product (or T_q-product) of the q-operators by the equation (all times t_i are assumed to be different):

$$\mathrm{T}_q[\hat{\varphi}(x_1)\dots\hat{\varphi}(x_n)] = \sum_P Q_P P[\theta(1\dots n)\hat{\varphi}(x_1)\dots\hat{\varphi}(x_n)] \ , \tag{27}$$

where the n-point θ-function is given by the relation:

$$\theta(1\dots n) \equiv \prod_{k=1}^{n-1}\theta(t_k - t_{k+1}) \ . \tag{28}$$

The summation in eq.(27) is taken over all simultaneous permutations of the operators $\hat{\varphi}(x_i)$ and the corresponding times t_i in θ-function.

If all or any part of the arguments t_i of the operators are equal then the T_q-product is not rigorously defined. We need to complete the definition. We define T_q-product under equal times as the Sym_q-product. An action of Sym_q- and T_q-products on zeroth power of operators is defined as usual by the equalities: $\mathrm{Sym}_q[1] = \mathrm{T}_q[1] = 1$.

We would like to note that Sym_q and T_q (similar to N) are not true linear operator on the space of the field operators: an operator equality $\hat{F}_1 = \hat{F}_2$ is not followed by the equalities $\mathrm{T}_q\hat{F}_1 = \mathrm{T}_q\hat{F}_2$ or $\mathrm{Sym}_q\hat{F}_1 = \mathrm{Sym}_q\hat{F}_2$ (because the field operators in the argument of operations $\mathrm{T}_q, \mathrm{Sym}_q, \mathrm{N}$ behave themselves as the classical variables).

We call a product *q-symmetric* if under permutation multipliers (q-operators) in the product it behaves in the same manner as if the q-operators would be classical variables, *i.e.* only the corresponding statistical phase appears (26). This definition naturally generalizes

the notion of (anti)symmetry of the field operator products in the Bose (Fermi) case. For example, the normal product is q-symmetric

$$\mathrm{N}[\hat{\varphi}(x_1)\dots\hat{\varphi}(x_n)] = Q_P \mathrm{N}[P\{\hat{\varphi}(x_1)\dots\hat{\varphi}(x_n)\}] \ , \tag{29}$$

It is also not difficult to check that the Sym_q- and T_q-products introduced above are q-symmetric:

$$\begin{aligned} \mathrm{Sym}_q[\hat{\varphi}(x_1)\dots\hat{\varphi}(x_n)] &= Q_P \mathrm{Sym}_q[P\{\hat{\varphi}(x_1)\dots\hat{\varphi}(x_n)\}] \ , \\ \mathrm{T}_q[\hat{\varphi}(x_1)\dots\hat{\varphi}(x_n)] &= Q_P \mathrm{T}_q[P\{\hat{\varphi}(x_1)\dots\hat{\varphi}(x_n)\}] \ . \end{aligned} \tag{30}$$

To formulate the functional form of the Wick's theorem for the above defined products we need to introduce the q-chronological contraction $\Delta_q(x,y)$ and the q-symmetrical contraction (or the q-symmetric part of the normal contraction n) n_s^q. They are defined by the relations

$$\mathrm{T}_q[\hat{\varphi}(x)\hat{\varphi}(y)] = \mathrm{N}[\hat{\varphi}(x)\hat{\varphi}(y)] + \Delta_q(x,y) \ , \tag{31}$$

$$\mathrm{Sym}_q[\hat{\varphi}(x_1)\hat{\varphi}(x_2)] = \mathrm{N}[\hat{\varphi}(x_1)\hat{\varphi}(x_2)] + n_s^q(x_1,x_2) \ . \tag{32}$$

Using definitions (12, 25, 27) we get expressions for the contraction via normal one (13):

$$\Delta_q = \theta(12)n + \kappa Q\theta(21)n^{\mathrm{T}} = \theta(12)n(x_1,x_2) + \kappa Q(x_1,x_2)\theta(21)n(x_2,x_1) \tag{33}$$

$$n_s^q = \frac{1}{2}[n + \kappa Q n^{\mathrm{T}}] = \frac{1}{2}[n(x_1,x_2) + \kappa Q(x_1,x_2)n(x_2,x_1)] \tag{34}$$

Due to our definition of the q-chronological product on equal times we have

$$\Delta_q(x_1,x_2)|_{t_1=t_2} = n_s^q(x_1,x_2)|_{t_1=t_2} \ . \tag{35}$$

We wish to note that the q-chronological contraction (similar to n_s^q) in contrast to the normal contraction n has the property of q-symmetry, *i.e.* $\Delta_q = \kappa Q \Delta_q^{\mathrm{T}}$.

We are now ready to formulate the Wick's theorem for q-symmetric products which gives Sym_q-product and T_q-product in the normal form.

Theorem 2

$$\mathrm{Sym}_q[\hat{\varphi}(x_1)\dots\hat{\varphi}(x_n)] = \mathrm{N}\left[\exp\left(\frac{1}{2}\frac{\overleftarrow{\partial}}{\partial\varphi}n\frac{\overleftarrow{\partial}}{\partial\varphi}\right)\varphi(x_1)\dots\varphi(x_n)\Big|_{\varphi=\hat{\varphi}}\right] \tag{36}$$

$$\mathrm{T}_q[\hat{\varphi}(x_1)\dots\hat{\varphi}(x_n)] = \mathrm{N}\left[\exp\left(\frac{\overleftarrow{\partial}}{\partial\varphi}\Delta_q\frac{\overleftarrow{\partial}}{\partial\varphi}\right)\varphi(x_1)\dots\varphi(x_n)\Big|_{\varphi=\hat{\varphi}}\right] \tag{37}$$

The proof can be performed in close analogy with the undeformed case [24] and the proof of Theorem 1.

Corollary 2.1 The normal contraction n in eq.(36) can be replaced by n_s^q from (34) due to the fact that the kernel of the differential operation is automatically symmetrized:

$$\frac{\partial}{\partial\varphi_i}n\frac{\partial}{\partial\varphi_k} = \frac{\partial}{\partial\varphi_k}\kappa Q n^{\mathrm{T}}\frac{\partial}{\partial\varphi_i} \ , \qquad \frac{\partial}{\partial\varphi}n\frac{\partial}{\partial\varphi} = \frac{\partial}{\partial\varphi}n_s^q\frac{\partial}{\partial\varphi} \ .$$

5 Wick's theorems for q-operator functionals

To describe the Green function technique and perturbation theory for the systems of the q-deformed particles we also need the rules for finding the normal form not only of the products of the operators but the whole functionals as well. A q-operator expression $F(\hat{\varphi})$ is said to be *q-operator functional* if it has the following form:

$$F(\hat{\varphi}) = \sum_{n=0}^{\infty} \int \dots \int \mathrm{d}x_1 \dots \mathrm{d}x_n \; F_n(x_1, \dots, x_n) \hat{\varphi}(x_1) \dots \hat{\varphi}(x_n) \; . \tag{38}$$

The operator functional is completely defined by the set of its (may be generalized and singular) coefficient functions $F_n(x_1 \dots x_n)$. We will call an operator functional *q-symmetrical* if all its coefficient functions satisfy the following relations:

$$\begin{aligned} F_n(\dots x_i \dots x_k \dots) &= Q_{ik}^{-1} F_n(\dots x_k \dots x_i \dots) \\ \dots \varphi(x_i) \dots \varphi(x_k) \dots &= Q_{ik} (\dots \varphi(x_k) \dots \varphi(x_i) \dots) \end{aligned} \tag{39}$$

Classical functionals (which are obtained from the operator one by replacing $\hat{\varphi}$ by φ) correspond to the q-symmetrical operator functional. q-Symmetrical coefficient functions are uniquely determined by the classical functional. Let's note also that any q-symmetrical functional possesses the property $F(\hat{\varphi}) = \mathrm{Sym}_q F(\hat{\varphi})$. We call an operator functional $F(\hat{\varphi})$ as *an operator functional in normal form* if it possesses the property $F(\hat{\varphi}) = \mathrm{N}[F(\hat{\varphi})]$

We emphasize that operator functionals are defined just by their coefficient functions (not by operator $F(\hat{\varphi})$). The functions F_n determine operator $F(\hat{\varphi})$ uniquely but the opposite statement is not true in general.

The formulae (36) and (37) are generalized directly to operator functionals due to "universality" of the reduction operation (we assume that operator functionals do not contain time derivatives, though all considerations can be extended to this case). Thus we obtain the following rules for reducing operator functionals to the normal form which we collect into Theorem 3:

Theorem 3

$$\mathrm{Sym}_q F(\hat{\varphi}) = \mathrm{N} \exp \left(\frac{1}{2} \frac{\overleftarrow{\partial}}{\partial \varphi} n \frac{\overleftarrow{\partial}}{\partial \varphi} \right) F(\varphi) \Big|_{\varphi = \hat{\varphi}} \tag{40}$$

$$\mathrm{T}_q F(\hat{\varphi}) = \mathrm{N} \exp \left(\frac{\overleftarrow{\partial}}{\partial \varphi} \Delta_q \frac{\overleftarrow{\partial}}{\partial \varphi} \right) F(\varphi) \Big|_{\varphi = \hat{\varphi}} \tag{41}$$

Corollary 3.1 From these various formulae of the inverse and combined transformations can be easily found. For example

$$\mathrm{N} F(\hat{\varphi}) = \mathrm{Sym}_q \exp \left(-\frac{1}{2} \frac{\overleftarrow{\partial}}{\partial \varphi} n \frac{\overleftarrow{\partial}}{\partial \varphi} \right) F(\varphi) \Big|_{\varphi = \hat{\varphi}} \tag{42a}$$

$$\mathrm{T}_q F(\hat{\varphi}) = \mathrm{Sym}_q \exp\left(\frac{1}{2}\frac{\overleftarrow{\partial}}{\partial\varphi}(\Delta_q - n)\frac{\overleftarrow{\partial}}{\partial\varphi}\right) F(\varphi)\Big|_{\varphi=\hat{\varphi}} \tag{42b}$$

We proceed with the consideration of the normal form of a product of the operator functionals. We formulate the Wick's theorem for product of operator functionals *in normal form*:

Theorem 4

$$\prod_{k=1}^{n}\left[F^{(k)}(\hat{\varphi})\right] = \mathrm{N}\left\{\exp\left[\sum_{i<k}\frac{\overleftarrow{\partial}}{\partial\varphi_i}n\frac{\overleftarrow{\partial}}{\partial\varphi_k}\right]\prod_{k=1}^{n}F^{(k)}(\varphi_k)\Big|_{\dots}\right\} \tag{43}$$

From now on the non-commuting multipliers are believed to be ordered in accordance with increasing index:

$$\prod_{k=1}^{n}\left[F^{(k)}(\hat{\varphi})\right] \equiv F^{(1)}(\hat{\varphi})\cdot\ldots\cdot F^{(n)}(\hat{\varphi})\ . \tag{44}$$

Corollary 4.1 Starting from the basic Wick's theorem (14) and arguing as was done in deriving eq.(36) one can obtain the following rule of reduction of a product of *q-symmetrical functionals* to the normal form:

$$\prod_{k=1}^{n}\left[\mathrm{Sym}_q F^{(k)}(\hat{\varphi})\right] = \mathrm{N}\left\{\exp\left[\frac{1}{2}\sum_i\frac{\overleftarrow{\partial}}{\partial\varphi_i}n\frac{\overleftarrow{\partial}}{\partial\varphi_i} + \sum_{i<k}\frac{\overleftarrow{\partial}}{\partial\varphi_i}n\frac{\overleftarrow{\partial}}{\partial\varphi_k}\right]\prod_{k=1}^{n}F^{(k)}(\varphi_k)\Big|_{\dots}\right\} \tag{45}$$

Corollary 4.2 The eqs.(45) are obviously generalized to the case when some multipliers in the LHS are given in the N-form (not Sym_q-form):

$$\prod_{k=1}^{n}\left[\mathcal{A}F^{(k)}(\hat{\varphi})\right] = \mathrm{N}\left\{\exp\left[\frac{1}{2}\sum_{\mathrm{Sym}_q}\frac{\overleftarrow{\partial}}{\partial\varphi_i}n\frac{\overleftarrow{\partial}}{\partial\varphi_i} + \sum_{i<k}\frac{\overleftarrow{\partial}}{\partial\varphi_i}n\frac{\overleftarrow{\partial}}{\partial\varphi_k}\right]\prod_{k=1}^{n}F^{(k)}(\varphi_k)\Big|_{\dots}\right\}\ . \tag{46}$$

Here $\mathcal{A}$ denote Sym_q or N and summation in the diagonal terms of the quadratic form is only over functionals that stand in Sym_q-form.

If a product of operator functionals stands under the common sign of some q-symmetrical product (Sym_q or T_q) it is automatically q-symmetrized and one can use the usual formulae (40) and (41). For example,

$$\mathrm{T}_q\left[\prod_{k=1}^{n}F^{(k)}(\hat{\varphi})\right] = \mathrm{N}\exp\left(\frac{\overleftarrow{\partial}}{\partial\varphi}\Delta_q\frac{\overleftarrow{\partial}}{\partial\varphi}\right)\prod_{k=1}^{n}F^{(k)}(\varphi_k)\Big|_{\varphi=\hat{\varphi}} \tag{47}$$

In conclusion we note that it is possible to complete the definition of the T_q-product not via the Sym_q-product but by using the N-form. This is equivalent to the definition of the θ-function entering into the chronological contraction under coincided times as follows: $\theta(t_{12})|_{t_1=t_2} = 0$. It is not difficult to check that the formula (37) remain true under this convention. The same can be said about formulae (41) but the classical functional entering into this formulae should represent N-form of the corresponding q-operator functional, *i.e.* $F(\hat{\varphi}) = \mathrm{N}F(\hat{\varphi})$.

6 Conclusion.

In this paper we have formulated Wick's theorems for general q-operators and q-operator functionals. The formal expression does not depend on the statistics of the fields and the statistical aspect is revealed only when the nature of the algebra of the corresponding classical variables is defined. This implies that the functional approach to the Wick's theorems using the analog of the Hori's formulae is the most natural one. Moreover, the theorems of the paper show that it is possible to take into account the exotic statistics keeping the formal expression for the Hori's formulae and changing the permutation relations for the classical analogues according to the rules of the statistics. This fact will allow us to derive the perturbation series for the theory and develop analogous constructions of standard quantum field theory in q-functional form in a straightforward way.

Acknowledgments.

We want to thank V.M.Agranovich for the drawing of our attention to the problems of quantum optics where q-particles find deep applications. We are also grateful to A.S.Vasiliev for interesting discussions. This work was supported by Grant of the Russian Fund of Fundamental Investigations N 95-01-00548 and UK EPSRC grants GR/L29156 and GR/K68356.

References

[1] A.Connes, *Non-commutative differential geometry*, Publ. Math. I.H.E.S. N62(1986);

[2] L.D. Faddeev, N.Yu. Reshetikhin and L.A. Takhtadzhyan , *Algebra and Analysis*, **1** (1987) 178;
S. Majid, *Int. J. of Mod. Phys.* **A 5** N1, (1990) 1;

[3] Yu.I.Manin, *Quantum groups and non-commutative geometry*, Preprint Montreal University, CRM-1561,1988;

[4] I.Ya.Aref'eva and I.V.Volovich, *Mod.Phys.Lett.* **A 6**(1991) 893;
A.P.Isaev and Z.Popowicz, *Phys.Lett.* **B 281** (1992) 271;
L.Castellani, *Phys.Lett.* **B 292** (1992) 93;
D.Bernard, *Suppl.Progr.Theor.Phys.* **102** (1992) 49;
M.Hirayama, *Progr.Theor.Phys.* **88** (1992) 111;

[5] J. Wess and B. Zumino, *Nucl.Phys.(Proc.Suppl.)* **B 18** (1990) 302;
U.Carow-Watamura, M.Schlieker and F.Watamura, *Z.Phys. C-Particles and Fields*, **49** (1991) 439;

[6] I.Ya.Aref'eva and I.V.Volovich, *Phys.Lett.* **B 268** (1991) 179;
J.Rembielinski and K.A.Smolinski, *Non-commutative quantum dynamics*, preprint KFT UL 3/93 (1993) ;
M.R.Ubriaco, *Mod.Phys.Lett.* **A 8** N 28 (1993) 89;

[7] W.Marcinek *J.Math.Phys.* **33**, (1992) 1631.

[8] N.V.Borisov, K.N.Ilinski, V.M.Uzdin , *Phys.Lett.* **A 169** (1992) 427

[9] Y.Ohnuki and S.Kamefuchi, *J.Math.Phys.* **21**, (1980) 601;
W.Marcinek: *On color quantization: relations and parastatistics.*, preprint ITP UWr 799/92;
W.Marcinek: *On unital braidings and quantization.*, preprint IFT UWr 847/93;

[10] A.T.Filippov, a.o., *Mod.Phys.Lett.* A 7 (1992) 2129;

[11] K.N.Ilinski, V.M.Uzdin , *Mod.Phys Lett.* **A 8** N 28 (1993), 2657;

[12] J.Beckers, N.Debergh,*J.Phys.***A 24**, (1991) L597;

[13] E.Fradkin, *Field Theories of Condensed Matter Systems*, Addison-Wesley, Reading, MA, USA 1991;
R.Prange and S.Girvin, *The quantum Hall Effect*, Springer-Verlag Inc., New York, 1987;

[14] S.Girvin and A.MacDonald, *Phys. Rev. Lett.* **58** (1987) 1252;

[15] *Fractional Statistics and Anyon Superconductivity*, edited by F. Wilczek, World Scientific Publishing Co., Singapore, 1990;

[16] V.M.Agranovich, M.D.Galanin, *Electronic excitation energy transfer in condensed matter.*, North-Holland Publishing Company, Amsterdam 1982, 381 p.;
Modern problem in condensed matter sciences. v.2. Excitons., ed. E.I.Rashba and M.D.Sturge, North-Holland Publishing Company, Amsterdam 1982, 866 p.;

[17] V.M.Agranovich and B.S.Toshich, Zh.ETF **53**, 149 (1967);

[18] V.M.Agranovich, K.N.Ilinski, Phys.Lett. **A 191**, 309 (1994) .

[19] V.M.Agranovich, *Theory of excitons.*,("Nauka",Moscow) 1968, 384 p. (in Russian);

[20] V.M.Agranovich, Zh.ETF **37**, 430 (1959);

[21] K.N.Ilinski, V.M.Uzdin, *Phys. Lett.***A 174**, (1993), 179-181;

[22] E.Lieb, T.Schults, D.Mattis, Ann.Phys.(N.Y.) **16**, 407, (1961);
D.B.Chesnut, A.Suna, J.Chem.Phys. **39**, 146 (1963);

[23] F.S.Spano, J.Knoester, in *Advances in magnetic and optical resonance* **18**, 117 (1994);

[24] A.N.Vasiliev, *Functional Methods in Quantum Field Theory and Statistics*, Leningrad Univ. Press, Leningrad, 1976 (in Russian);

[25] K.N.Ilinski, A.S.Stepanenko: "q-Functional integrals in anyonic physics.", preprint PNPI TH-2-1994 1944;
K.N.Ilinski, A.S.Stepanenko: "q-Functional integrals in anyonic physics.", Phys. Lett. **A 187** 1 (1994);

[26] S.Hori, Progr.Theor.Phys., v.7, 578 (1952).