

Multi-spin correlation functions for the \mathbb{Z} -Invariant Ising model

J.R. Reyes Martínez

Institut für Theoretische Physik
Freie Universität Berlin
Alte Allee 14, 14195 Berlin
martinez@physik.fu-berlin.de

Abstract

Continuing our work [12], where an explicit formula for the two-point functions of the two dimensional \mathbb{Z} -invariant Ising model were found. I obtain here different results for the higher correlation functions and several consistency checks are done.

1 Introduction

In this paper I want to apply the vertex operator method (VOM) for the calculation of different correlations functions of the two dimensional Z -invariant Ising model (IM). Z -invariance is the property that allows to introduce inhomogeneities in the lattice in conformity with the integrability of the model. In particular when we put all inhomogeneities equal zero we obtain a homogeneous Ising model in a square lattice. Also by choosing appropriately the inhomogeneity parameters it is possible to obtain other regular lattice configurations, like a triangle, checkerboard and others (see [1] for examples). The VOM was introduced by the Kyoto group in the case of the six-vertex model [7] and then extended to a variety of other integrable lattice models. The novelty of this approach is the treatment from the beginning of the lattice model in the thermodynamic limit, this is made possible because of two fundamental concepts: the corner transfer matrix (CTM) of Baxter [2] and the lattice vertex operator (VO) introduced by the Kyoto group, see [8] for a review. Whereas the CTM allows the calculation of the one-point functions the VO's permit in connection with the CTM the calculation of higher correlation functions (CF). Physically they correspond to inserting a semi-infinite inhomogeneity line at bulk of the lattice. This inhomogeneity is described by a spectral parameter that enters the parameterization of the Boltzmann's weights of the model such that they satisfy some integrability conditions (in the case of the IM they are called the star-triangle relations), which imply that the diagonal transfer matrices of the model commute and also allow the possibility of choosing different spectral parameters in the definition of the usual transfer matrices [2].

In the paper [12] the author obtained the two-point functions in the general Z -invariant IM whose vertex operator algebra was developed in [4]. These were given in forms of products of well known elliptic functions over the inhomogeneity parameters. Because of its importance for the calculation of the higher correlation functions I review this result in Sect. 3 (see also Sect. 4.1) and in the Appendix I give explicit details of the calculations. After reviewing some important definitions in Sect. 2, then in Sect. 3.3 I give a general integral formula for all higher correlations with an even number of spins. Particular cases of this formula are treated next. Section 4 contains our principal results, the first part reviews the general two-point functions as already mentioned, in Sect. 4.2 the homogeneous CF with adjacent diagonal spins are found in the form of generalized Toeplitz determinants and in Sect. 4.3 I find general homogeneous diagonal four-point functions, obtaining a general determinant expression. In Sect. 5 I consider finally some particular cases of three-point functions. A conclusion section ends the paper.

2 Fundamental definitions

2.1 Elliptic parameterization

Let us begin by introducing a number of definitions for the Z -invariant IM; the model is defined initially on a square lattice rotated by 45 degrees, the spin variables taking values $+1$ or -1 live on the vertices of the lattice the coupling being between nearest neighbors. Assume first two different interaction energies (measured in kT units) with $K > 0$ in the NW-SE direction and $L > 0$ in the NO-SW direction between two spins as in the figure 1




Figure 1: Boltzmann Weights for the IM

In the figure the two lines indicate one horizontal u_H and one vertical u_V spectral parameters, their difference $u = u_H - u_V$ is used to code the couplings energies under the critical temperature by mean of Jacobian elliptic functions,

$$\begin{aligned} \sinh(2K) &= i \operatorname{sn} iu; & \cosh(2K) &= \operatorname{cn} iu; \\ \sinh(2L) &= ik^{-1} \operatorname{ns} iu; & \cosh(2L) &= ik^{-1} \operatorname{ds} iu; \end{aligned} \quad (1)$$

where $\operatorname{sn} u$ is a Jacobian elliptic function with half-periods I, iI^0 , k is the modulus, $0 < u < I^0$ (see e.g. [2]), and Glashier's notation for the Jacobian elliptic functions is used, $\operatorname{ns} u \equiv 1/\operatorname{sn} u$, $\operatorname{ds} u \equiv \operatorname{dn} u/\operatorname{sn} u$. These parameterization already introduced by Onsager is a consequence of the integrability conditions, in the case of the IM the so called star-triangle relations, which imply as well known, that diagonal matrices commute when the difference of the spectral parameters from site to site remain constant (see pag. 374 in Baxter's book). The module $k = (\sinh(2K) \sinh(2L))^{-1}$ does not depend on the spectral parameter and tends to 1 when the system goes critical. It is possible to write a similar parameterization for temperatures higher than the critical but it will not be necessary due to the duality between order and disorder operators to be introduced below, which imply that the disorder CF under the critical temperature correspond to the order CF above it.

2.2 Vertex Operator Algebra

I review now the fundamental results of the paper [4], where the vertex operator algebra for the IM was found. Dividing the diagonal lattice in four equal quadrants to apply Baxter's CTM method we have two cases according to whether the central point belongs or not to the lattice; this defines two CTM's Fig.2 that act on the two sectors of the model, NS and R, respectively (in analogy to conformal field theory). Diagonalising the adjoint action of the CTM Hamiltonians we can describe the two sectors as Fock spaces (called H^{NS} and H^R) of two irreducible representations of free fermions (apart from a different normalization). We have:

$$[D^{NS}; r] = 2r \quad \text{for } r \in \mathbb{Z} + \frac{1}{2} \quad \text{for the NS sector,} \quad [D^R; r] = 2r \quad \text{for } r \in \mathbb{Z} \quad \text{for the R sector}$$

where $r \in \mathbb{Z} + \frac{1}{2}$ for the NS sector, $r \in \mathbb{Z}$ for the R sector and D^{NS}, D^R are the respective CTM hamiltonians.

There are two vertex operators (VOs in the following) depending on which sector they intertwine:

$$\begin{aligned} R_{NS}(\epsilon) &: H^{NS} \rightarrow H^R; \\ NS_R(\epsilon) &: H^R \rightarrow H^{NS}; \end{aligned}$$




Figure 2: The CTMs for the IM

The Vertex Operators (VO) satisfy the following elegant intertwining relations with the fermions

$$\begin{aligned} (\psi)^R(z) &= NS(z) f(z^2) (\psi); \\ (\psi)^{NS}(z) &= f(z^2) R(z) (\psi); \end{aligned} \quad (2)$$

$$\begin{aligned} (\psi) &= i_1^{NS}(\epsilon) (\psi); \\ (\psi) &= R_1(\epsilon) (\psi); \end{aligned} \quad (3)$$

Here we use the simplified notation of [4] $R_{NS}(\epsilon) = (\epsilon)$, being equal to +1 or -1 distinguishes the two sub-sectors (with even or odd parity) of the NS sector and $R_{NS}(\epsilon) = (\epsilon)$, also we define $[z] := f(z) = \prod_{n \in \mathbb{Z}} (1 - z^{2n+1})$ with $z = \exp(i v = I)$ and we have introduced the generating functions

$$NS(z) = \sum_{r \in \mathbb{Z} + \frac{1}{2}} NS_r z^r; \quad R(z) = \sum_{s \in \mathbb{Z}} R_s z^s; \quad (4)$$

and

$$i_1^{NS}(\epsilon) = \frac{dz}{2 - iz} [z]_{NS}^{NS}(z^2); \quad R_1(\epsilon) = \frac{dz}{2 - iz} [z]_R^R(z^2); \quad (5)$$

where

$$[z]_{NS} = \prod_{r \in \mathbb{Z} + \frac{1}{2}} \frac{2Ik}{kI} \epsilon v = \prod_{r \in \mathbb{Z} + \frac{1}{2}} \frac{2}{kI} \sum_{r \in \mathbb{Z} + \frac{1}{2}} z^r;$$

$$[z]_R = \frac{r}{2I} \text{dnv} = \frac{r}{I} \frac{X}{s^2 Z} s^1 z^s; \\ x_r = x^{2r} + x^{-2r}; \quad x = \exp\left(\frac{I^0}{2I}\right);$$

By crossing symmetry i.e. $u \rightarrow u - I^0$ or in the multiplicative language $x \rightarrow x^{-1}$ we can obtain (using also the transformation $f(x^2 z) = 1/f(z)$) similar relations with the parameter of the VO replaced with $-r$ but in the calculation of the CF in the next sections I will need only the expressions above. Finally the VO's satisfy the following unitary relations:

$$X (x) () = g^R \text{id}_{H^R}; \quad (6)$$

$$(x) () = g^{NS} \text{id}_{H^{NS}}; \quad (7)$$

where the scalars g^R and g^{NS} can be given as infinite products in the variable x but as we will not need them for the calculation of the CF we leave them undefined, similarly the expectation values of the VO's in the respective fermion basis are known but will not be needed (see [4] for their explicit values).

3 Even correlation functions

In this section I will give a general multi-integral expression for the CF with an even number of spins. The case two spin operators was considered in [12] where the integration was carried explicitly. This result will be remembered in the Sect. 4.1. We start with an easy example and proceed in order of difficulty.

3.1 Nearest diagonal neighbors CF

Now I review the calculation of the two-point functions. For simplicity let us consider first the case for nearest diagonal order-order and disorder-disorder CF Fig. 3. In terms of the VO




Figure 3: Order-Order and Disorder-Disorder CF for nearest diagonal neighbors

they are respectively

$$P_0 = \frac{\text{tr}_{H^{NS}} (x^{2D^{NS}} (x_2) (x_1) (1) (2))}{g^{NS} g^R \text{tr}_{H^{NS}} (x^{2D^{NS}})}; \quad (8)$$

$$\frac{P \quad \text{tr}_{\mathbb{H}^R} \quad \mathbf{x}^{2D^R} \quad (\mathbf{x}_2) \quad (\mathbf{x}_1) \quad (1) \quad (2)}{g^N g^R \text{tr}_{\mathbb{H}^R} (\mathbf{x}^{2D^R})} : \quad (9)$$

Consider as an example (9); the reader can look at App. B in [4] for an explicit determination of (8). We transform the four VO under the trace symbol in terms of fermions in the following way

$$\begin{aligned}
X & \quad (\mathbf{x}_2) \quad (\mathbf{x}_1) \quad (1) \quad (2) = \\
X & \quad (\mathbf{x}_2) \quad (\mathbf{x}_1) \quad (1) \quad (2) = \\
X & \quad {}^R_1(1) \quad (\mathbf{x}_2) i_1^{NS}(1) \quad (\mathbf{x}_1) \quad (1) \quad (2) = \\
& \quad I \\
& \quad {}^R_1(2) \quad [w_1]_{NS} \quad {}^R(w_1) [w_1 2] \quad X \quad (\mathbf{x}_2) \quad (\mathbf{x}_1) \quad (1) \quad (2) = \\
& \quad I^{w_1} \quad I \\
ig^{NS} g^R & \quad [w_2 2]_R \quad {}^R(w_2) \quad [w_1 1]_{NS} [w_1 2]^R(w_1) = \\
& \quad I^{w_2} I \quad w_1 \\
ig^{NS} g^R & \quad [w_1 2]_R [w_1 1]_{NS} [w_1 2]^R(w_2)^R(w_1); \tag{10} \\
& \quad w_2 \quad w_1
\end{aligned}$$

where we changed the variables to $w_i = z_i = \frac{2}{i}$ and used the notation $\frac{H}{i} = \frac{H}{2} \frac{dw_i}{iw_i}$, $[w_i] = [w_i \frac{2}{i}]$. Following substitutions were then made, in the second line the identity $2 = 1$, in the third line eqs. (3), the intertwining relation (2) and the second eq. in (5) in the fourth line and finally the unitarity conditions for the VO's.

It only remain to take the trace of the fermion operators, which we state as a simple lemma,

$$\frac{\text{tr}_{\mathbf{H}^R}(\mathbf{x}^{2D^R} \mathbf{w}_2 \mathbf{w}_1)}{\text{tr}_{\mathbf{H}^R}(\mathbf{x}^{2D^R})} = \mathbf{w}_2 \mathbf{w}_1 : \mathbf{x}^{2D^R} \quad (11)$$

With the aid of this formula and the relation $\chi^{-2} z_R[z] = i[z]_{N_S}$ we are left with an integral expression for the disorder-disorder CF,

$$\mathbb{I}_{\mathbb{W}}^{[1]_{\mathbb{N}_S} [2]_{\mathbb{N}_S}}: \quad (12)$$

Remembering that $w = \exp(i v/I)$ and $\bar{w} = \exp(-i v/I)$ the last expression can be written as:

$$\frac{k^Z}{k} \frac{d\sigma}{d\Omega}(\theta_1) \frac{d\sigma}{d\Omega}(\theta_2) = \frac{2I}{k} \frac{k^0}{k} \frac{1}{H(\theta_2)} \frac{1}{H(\theta_1)};$$

where H , θ , H_1 and θ_1 (see e.g.[6]) are the Jacobi theta functions and $\eta := \eta_1 - \eta_2$. The prime denotes the derivative under the argument. This integration formula appears in other disguise in the Appendix, see eq. (52).

3.2 Two-point functions with an arbitrary number of inhomogeneities

Let us call the general order-order and disorder-disorder CF by $g_{2n}(\epsilon_1; \epsilon_2; \dots; \epsilon_{2n})$ and $g_{2n}(\epsilon_1; \epsilon_2; \dots; \epsilon_{2n})$ respectively (when we don't want to emphasize the spectral parameters g_{2n} and g_{2n} simply), where $\epsilon_j = \exp(\epsilon_j = 2I)$, then we have as simple generalizations of the trace formulas from Sect. 3.1:

$$g_{2n}(\epsilon_1; \epsilon_2; \dots; \epsilon_{2n}) = \frac{\prod_{i=1}^n \text{tr}_{NS} x^{2D^{NS}} F(\epsilon_i; \epsilon_1; \epsilon_2; \dots; \epsilon_{2n})}{(g^{NS} g^R)^n \text{tr}_{NS} x^{2D^{NS}}}; \quad (13)$$

$$F(\epsilon_i; \epsilon_1; \epsilon_2; \dots; \epsilon_{2n}) = \sum_{\epsilon_1, \dots, \epsilon_{2n}} (\epsilon_{2n})_{\epsilon_1} (\epsilon_{2n-1})_{\epsilon_2} \dots (\epsilon_2)_{\epsilon_{2n-1}} (\epsilon_1)_{\epsilon_{2n}} = \sum_{\epsilon_1, \dots, \epsilon_{2n}} (\epsilon_{2n})_{\epsilon_1} (\epsilon_{2n-1})_{\epsilon_2} \dots (\epsilon_2)_{\epsilon_{2n-1}} (\epsilon_1)_{\epsilon_{2n}};$$

where the subscript ϵ_i in the sum above means the sum for all $1 \leq i \leq n-1$. Similarly for the disorder-disorder case:

$$g_{2n}(\epsilon_1; \epsilon_2; \dots; \epsilon_{2n}) = \frac{\prod_{i=1}^n \text{tr}_R x^{2D^R} G(\epsilon_i; \epsilon_1; \epsilon_2; \dots; \epsilon_{2n})}{(g^{NS} g^R)^n \text{tr}_R x^{2D^R}}; \quad (14)$$

$$G(\epsilon_i; \epsilon_1; \epsilon_2; \dots; \epsilon_{2n}) = \sum_{\epsilon_1, \dots, \epsilon_{2n}} (\epsilon_{2n})_{\epsilon_1} (\epsilon_{2n-1})_{\epsilon_2} \dots (\epsilon_2)_{\epsilon_{2n-1}} (\epsilon_1)_{\epsilon_{2n}} = \sum_{\epsilon_1, \dots, \epsilon_{2n}} (\epsilon_{2n})_{\epsilon_1} (\epsilon_{2n-1})_{\epsilon_2} \dots (\epsilon_2)_{\epsilon_{2n-1}} (\epsilon_1)_{\epsilon_{2n}};$$

with $1 \leq i \leq n$. Consider the disorder CF (the other case is similar, see [12] for the differences). We see that if we begin in the middle of the operator expression $G(\epsilon_i; \epsilon_1; \epsilon_2; \dots; \epsilon_{2n})$ we have the same product of four vertex operators (10) appearing before, thus we can substitute them by a pair of fermion operators and move them to the left by means of the intertwining property (2). The result is, apart from a product of elliptic functions depending on the spectral parameters, of the same form but with four vertex operators less as before. Trading in this way each time four vertex operators by two fermions operators and using the following antisymmetry property of the fermion algebra under the integrations, i.e.,

$$\int \int [w_1]_R [w_i]_{NS} [w_i]_R [w_1]_R (w_1; w_i) = \int \int [w_1]_R [w_i]_{NS} [w_i]_R (x^2 w_1 = w_i) + \int \int [w_1]_R [w_i]_{NS} [w_i]_R (x^2 w_1 = w_i) = 0;$$

one arrives at the expression (with the fermions ordered from left to right according to increasing indexes)

$$(ig^{NS} g^R)^n \prod_{i=1}^n \int [w_1]_{NS} [w_2]_R \dots [w_{2n}]_{NS} [w_{2n}]_R \prod_{1 \leq i < j \leq 2n} [w_i]_R [w_j]_R (w_i): \quad (15)$$

The corresponding formula for the order-order CF is obtained by simply interchanging the NS with R representation and vice-versa. We continue with the analysis of this expression in the Section 4.

3.3 Higher even correlation functions

By the same method of calculation of the two-point functions explained above the generalization to arbitrary higher even CF is straightforward. We consider the CF with only order operators the purely disorder case can be obtained in the same way as before interchanging representations. Let us call the general $2n$ -CF G_{2n} . Define first some parameters:

$2n$ spins σ_i for i between 1 and $2n$ at arbitrary positions in a diagonal ordered from left to right;

$2N$ spectral parameters λ_i for i between 1 and $2N$, N being a positive integer;

$2n$ integers a_1, a_2, \dots, a_n with $a_1 = 1 < a_2 < a_3 < \dots < a_n \leq 2N$, such that for odd i the spectral parameter λ_i is adjacent to the right of the spin σ_i and when i is even λ_i is adjacent to the left of the spin σ_i ;

A set W of $L := \sum_{i=1}^n (a_{2i} - a_{2i-1} + 1)$ complex parameters w_j , note that L is an even integer, ordered in the following way:

$$w_{a_1}, w_{a_1+1}, \dots, w_{a_2}, w_{a_3}, w_{a_3+1}, \dots, w_{a_4}, \dots, w_{a_{2n-1}}, w_{a_{2n-1}+1}, \dots, w_{a_{2n}}.$$

Consider the following example for a 6-point CF: In the picture we have that: $n = 3, N = 7$,




Figure 4: An example of a 6-point spin-spin correlation function.

$a_1 = 1, a_2 = 2, a_3 = 7, a_4 = 10, a_5 = 13$, and $a_6 = 14$. The set W of complex variables w_j is: $w_1, w_2, w_7, w_8, w_9, w_{10}, w_{13}, w_{14}$.

The $2n$ -CF G_{2n} can now be written as:

$$G_{2n} = \left(i \right)^{\frac{L}{2}} \frac{\prod_{j=1}^n \prod_{k=1}^{a_{2i}-a_{2i-1}+1} \langle \sigma_i \sigma_k \rangle}{\prod_{j=1}^n \prod_{k=1}^{a_{2i}-a_{2i-1}+1} \langle \sigma_i \sigma_k \rangle}; \quad G_{fw_j}; \quad F_{fw_j}; \quad (16)$$

where F_{fw_j} is a 'temperature' state:

$$F_{fw_j} := \frac{\text{tr}_{H^{NS}} x^{2D^{NS}} Q_{w_j, 2W}^{NS}(w_j)}{\text{tr}_{H^{NS}} x^{2D^{NS}}}; \quad (17)$$

and is known, see Sect. 4.1, to be a sum of products of Dirac-delta functions.

The function $G_{fw_j}; \sigma_i$ is given by the product of n blocks g_i , where

$$g_i = \prod_{a_{2i-1} < r < a_{2i}} [w_r r]_k \prod_{a_{2i-1} < s < a_{2i}} [w_s s]_{NS} \prod_{a_{2i-1} < k < a_{2i}} [w_k j]; \quad (18)$$

in the above expression the parameters r, s, k , and j are positive integers. The prime in the products means that we increment the parameters r and s in steps of 2, i.e. the r runs over odd and the s even integers respectively.

In the example presented before we have three blocks g_1, g_2 , and g_3 :

$$\begin{aligned} g_1 &= [w_1 1]_R [w_2 2]_{NS} \prod_{j=1}^{Y^2-Y^4} [w_j k] \\ g_2 &= [w_7 7]_R [w_8 8]_{NS} [w_9 9]_R [w_{10} 10]_{NS} \prod_{j=7}^{Y^0-Y^4} [w_j k] \\ g_3 &= [w_{13} 13]_R [w_{14} 14]_{NS} [w_{13} 14]: \end{aligned}$$

And $G = g_1 g_2 g_3$.

The next step in the calculation is easy, namely to integrate the delta functions in eq. (16). The number of integrals is reduced by a half and the final result is a clumsy sum of integrals expressions. Perhaps some symmetrization procedure can give more adequate results as we will show in the next section for the two-point functions.

4 Explicit results for the even correlation functions

In this section we review the general integration of the two-point functions announced in [12], details of the calculation are relegated to the appendix. Also some examples of higher homogeneous nearest neighbors diagonal CF are given in the form of determinants, which generalize the well-known results for the diagonal two-point functions. Finally we treat the diagonal homogeneous four-point functions for arbitrary spins locations.

4.1 The general two-point functions

To integrate the expression found in Sect. 3.2 we need the trace of the fermions operators in (15) that is given by a generalization of the lemma used before, i.e.

$$\frac{\text{tr}_{H^R} (x^{2D^R} (z_1) (z_2) \dots (z_{2n-1}) (z_{2n}))}{\text{tr}_{H^R} (x^{2D^R})} = \sum_p \prod_p (p_1 p_2) (p_3 p_4) \dots (p_{2n-1} p_{2n});$$

with the notation $\prod_p (ij) = \prod (x^2 w_i = w_j)$. Here p_1, \dots, p_{2n} is some permutation of the numbers $1, 2, \dots, 2n$, \prod_p is a summation over all $(2n-1)!!$ permutations which satisfy the restrictions $p_{2m-1} < p_{2m}$ for $1 < m < n$ and $p_{2m-1} < p_{2m+1}$ for $1 < m < n-1$ and \prod_p is the parity of the permutation. There is a similar expression for the NS case (note that the integrations in this case are well defined although the function $\prod^{NS}(ij)$ has half-integral powers in the integration variable z).

Now changing the spectral parameters to their additive version i.e. interchanging λ_i with μ_i (remember that $\lambda_i = \exp(\mu_i/2I)$) and using some identities for the product of theta functions (see the appendix for the technicalities) we arrive at the following simple

expressions for the general two-point functions

$$g_{2n}^i = \frac{P \text{fa} \text{an} (h; (i_1 \dots i_j))}{P \text{fa} \text{an} \frac{1}{k} \text{sn}(i_1 \dots i_j)}; \quad (19)$$

$$h(i) := \frac{2I}{k} \frac{k^0 \frac{1}{2}}{(i)} \frac{1^0(i)}{(i)}; \quad h(i) := \frac{2I}{(k^0)^{\frac{1}{2}}} \frac{H_1^0(i)}{(i)}; \quad (20)$$

Expression (19) is symmetric in all spectral parameters and satisfies the recurrence relation of Baxter¹

$$g_{2n}^i(i_1, i_2, \dots, i_{2n-1} + I^0) = g_{2n-2}^i(i_1, i_2, \dots, i_{2n-2});$$

$$g_2^i(i_1 + I^0) = 1;$$

To obtain the homogeneous correlations one has to take the limit of the expression (19) when all spectral parameters i are equal which requires a generalization of the L'Hôpital's rule for determinants already well known (see e.g. [1] or [16]),

$$g_{\text{hom}, 2n}^i = \frac{\det h^{(2j+2k-3)}(0)}{\det h^{(2j+2k-3)}(0)g}; \quad (21)$$

In the last formula $h^n(0)$ means the n 'th derivative of the function h evaluated at the point 0, $h(i) = \frac{1}{k} \text{sn}(i)$ and $h; ,$ were already defined in (20). The determinants have rank n and their elements at the row j and column k are given by $h^{(2j+2k-3)}(0)$. In the mathematical literature they are called persymmetric Wronskians (a terminology due to Sylvester). Because the denominator in (21) can be expressed as a product one can calculate explicitly its homogeneous limit:

$$\det \text{sn}^{(2j+2k-3)}(0) = k^{n^2-n} \prod_{i=1}^{2n-1} (i!); \quad (22)$$

therefore²

$$g_{\text{hom}, 2n}^i = k^{n^2-n} \prod_{i=1}^{2n-1} (i!)^{-1} \det \frac{1}{k} h^{(2j+2k-3)}(0); \quad (23)$$

This expressions are useful for studying correlations for small separations, as one only has to take derivatives of known elliptic functions. Several checks of eq. (23) with the results in [5] were done.

4.2 Homogeneous even-point functions for adjacent spins on a diagonal

By adjacent spins I mean nearest diagonal neighbors. I discuss the 4-point spin function and then the 6-point spin function the generalization afterward will be easy.

¹The integral formula conjectured by Baxter [3] for the general two-point function can also be derived see [12].

²Yamada has obtained a similar expression by different methods in [16].

In the the case of the four-point CF G is the product of two blocks g_1 and g_2 and the temperature state $F(w_i)$ is

$$F(w_i) = \int_{-1}^1 x^2 \frac{w_1}{w_2} \int_{-1}^1 x^2 \frac{w_5}{w_6} \int_{-1}^1 x^2 \frac{w_1}{w_5} \int_{-1}^1 x^2 \frac{w_2}{w_6} + \int_{-1}^1 x^2 \frac{w_1}{w_6} \int_{-1}^1 x^2 \frac{w_2}{w_5}$$

The functions g_1 and g_2 are

$$g_1 = [w_{11}]_R [w_{12}] [w_{13}] [w_{14}] [w_{15}] [w_{16}] \\ [w_{22}]_{NS} [w_{23}] [w_{24}] [w_{25}] [w_{26}]$$

$$g_2 = [w_{55}]_R [w_{56}] \\ [w_{66}]_{NS}$$

Doing the integration for the variables half of the variables w_i in (16) gives

$$G_4 = \int_{-1}^1 \int_{-1}^1 [w_{11}]_R [w_{22}]_R [w_{55}]_R [w_{66}]_R \\ \int_{-1}^1 \int_{-1}^1 [w_{11}]_R [w_{12}] [w_{13}] [w_{14}] [w_{15}]_{NS} [w_{22}]_{NS} [w_{23}] [w_{24}] [w_{25}] [w_{26}]_R + \\ \int_{-1}^1 \int_{-1}^1 [w_{11}]_R [w_{12}] [w_{13}] [w_{14}] [w_{15}] [w_{16}]_R [w_{22}]_{NS} [w_{23}] [w_{24}] [w_{25}]_{NS} : \quad (24)$$

Equation (24) is the inhomogeneous 4-point function. Now I restrict to the homogeneous 4-point function i.e., take all spectral parameters w_i equal to 1.

Let us change eq. (24) to the additive variables v_i ($w_i = \exp(i v_i)$), but first I introduce the following notations:

$$a_n := \frac{k^n}{2\pi} \int_{-1}^1 \int_{-1}^1 dv (dn v)^2 (sn v)^{2n} \quad \text{for } n \in \mathbb{Z} : \quad (25)$$

Note that for $n < 0$, a_n can be put in a different form: by the shifting $v \rightarrow v + i\pi^0$ in such a way that the poles of integrand remain outside the contour of integration and because the integrand is an even function the limits of integration can be leave as in (25), namely

$$a_{-n} := \frac{k^n}{2\pi} \int_{-1}^1 \int_{-1}^1 dv (cn v)^2 (sn v)^{2(n-1)} \quad \text{for } n = 1, 2, \dots$$

Returning to the homogeneous 4-point CF eq. (24), inserting the definitions for the different factors in the additive variables

$$G_4 = \frac{1}{2\pi} \int_{-1}^1 \int_{-1}^1 dv (dn v)^2 \int_{-1}^1 \int_{-1}^1 dv k^2 dn v cn v (sn v)^3 + \\ + \frac{1}{2\pi} \int_{-1}^1 \int_{-1}^1 dv k^2 (dn v)^2 (sn v)^4 \int_{-1}^1 \int_{-1}^1 dv k^2 (cn v)^2 (sn v)^2 ;$$

where the integrals go from -1 to 1 , the second term vanishes and with the above notation we have

$$G_4 = a_0^2 - a_2 a_{-2} : \quad (26)$$

Proceeding in the same way the 6-point CF is given by

$$G_6 = a_0^3 - 2a_0 a_2 a_4 + a_2^2 a_4 + a_4 a_2^2 - a_0 a_4 a_4 = \begin{pmatrix} a_0 & a_2 & a_4 \\ a_2 & a_0 & a_2 \\ a_4 & a_2 & a_0 \end{pmatrix} : \quad (27)$$

The general expression for G_{2n} can be induced to be:

$$G_{2n} = \begin{pmatrix} a_0 & a_2 & \dots & a_{2(n-1)} \\ a_2 & a_0 & \dots & a_{2(n-2)} \\ \vdots & \vdots & \ddots & \vdots \\ a_{2(n-1)} & a_{2(n-2)} & \dots & a_0 \end{pmatrix} : \quad (28)$$

Remark. The definitions of the coefficients of the above matrices is equivalent to those given on [10] pag. 199. For temperatures below the critical temperature, when $0 < k < 1$ they are given by

$$a_n = \frac{1}{2} \int_0^{2\pi} d\theta e^{in\theta} \frac{k^{-1} e^{i\theta/2}}{k^{-1} e^{i\theta}} ; \quad (29)$$

deforming the contour of integration from the unit circle $j = 1$ ($\theta = e^{i\theta}$) to the real axis one rewrites (29) as

$$a_n = \frac{1}{2} \int_0^k \frac{d\theta}{\theta} \frac{1 - k}{k^{-1} - 1} ;$$

substituting $\theta = k(\sin u)^2$ one sees that the definition (29) coincides with (25).

In a similar manner the Toeplitz determinant for the two-point function (see e.g. [10]) can be derived by the method presented above, giving a strong support of the vertex operator method.

4.3 Homogeneous diagonal Four-point functions

We label the four spins from left to right as s_1, s_2, s_3, s_4 and let the distance between the s_1 and s_2 be r and between s_3 and s_4 be s , finally let the distance between s_2 and s_3 be p , where the distances are measured in diagonal steps of course. Then by the same procedure as before, we have the following general result.

The CF can be written as a determinant G of order $r+s$ whose elements on the principal diagonal are all equal to a_0 and whose elements a_n above and below the diagonal are negative and positive sub-indices respectively. To give its form decompose G in four matrix blocks:

$$G = \begin{pmatrix} A_1 & B_1 \\ B_2 & A_2 \end{pmatrix} : \quad (30)$$

A_1 is the same Toeplitz matrix of order r for the two-point function $h_{s_1 s_2}$ i.e.

$$A_1 = \begin{pmatrix} a_0 & a_1 & \dots & a_{r-1} \\ a_1 & a_0 & \dots & a_{r-2} \\ \vdots & \vdots & \ddots & \vdots \\ a_{r-1} & a_{r-2} & \dots & a_0 \end{pmatrix} :$$

A_2 is also a Toeplitz matrix but now of order s . B_1 is a $r \times s$ matrix with all its a_n coefficients negative whose element in the r st row and at the s column is equal to $a_{-(r+s+p)+1}$ and with coefficient sub-indices growing to the left and to the bottom of the matrix

$$B_1 = \begin{pmatrix} a_{-(r+p)} & a_{-(r+p)-1} & \dots & a_{-(r+s+p)+1} \\ a_{-(r+p)+1} & a_{-(r+p)} & \dots & a_{-(r+s+p)+2} \\ \vdots & \vdots & \ddots & \vdots \\ a_{-p-1} & a_{-p-2} & \dots & a_{-p} \end{pmatrix}$$

Finally the matrix B_2 is the transpose of B_1 but with all its coefficients positive.

Let us give some examples for illustration. Consider the following configuration where each circle represents a point in the diagonal:

$$\begin{matrix} 1 & 2 & 3 & 4 \end{matrix}$$

The correlation function for this configuration is (with $r = 2$, $p = 3$ and $s = 1$):

$$\begin{pmatrix} a_0 & a_{-1} & a_{-5} \\ a_{-1} & a_0 & a_{-4} \\ a_{-5} & a_{-4} & a_0 \end{pmatrix}$$

An interesting example is given by the following picture

$$\begin{matrix} 1 & 2 & 3 & 4 \end{matrix}$$

in the figure we suppose that the distance between 2 and 3 is n (i.e. $p = n$). Then the CF is

$$\begin{pmatrix} a_0 & a_{-n-1} \\ a_{-n-1} & a_0 \end{pmatrix};$$

this coincides with the value found in [14] using combinatorial methods.

5 Odd correlation functions

In this section we want to give some examples of odd-point spin CF (there are not odd disorder CF under the critical temperature). As example of higher odd-point functions I treat some particular cases of the three spin CF.

Consider first the three spins in adjacent diagonal positions see Fig. 5. Let us call the 3-




Figure 5: Adjacent 3-point correlation function.

point CF $C_1(i_1; i_2; i_3; i_4)$ (the choice of the index will be clear below). Its explicit expression is given by

$$C_1 = \frac{1}{N} \sum_i \text{tr}_{H^{NS}} \mathbf{x}^{2D^{NS}} \begin{matrix} 3 & 2 & 1 \\ (x_4) & (x_3) & (x_2) & (x_1) \end{matrix} \begin{matrix} 1 & 2 & 3 & 4 \\ (1) & (2) & (3) & (4) \end{matrix}; \quad (31)$$

where the normalization factor N is $(g^R g^{NS})^2 \text{tr}_{H^{NS}} (\mathbf{x}^{2D^{NS}})$. Now using eqs. (2) and (3) and starting at the middle of the operator expression in (31). After using the unitarity conditions for the VOs we arrive at

$$\sum_i \begin{matrix} 3 & 2 & 1 \\ (x_4) & (x_3) & (x_2) & (x_1) \end{matrix} = g^{NS} (\text{id}_{H^{NS};+} - \text{id}_{H^{NS};-}) = g^{NS} \begin{matrix} NS \\ 0 \end{matrix};$$

where the unitarity relation (7) was employed and the fermion operator $\begin{matrix} NS \\ 0 \end{matrix}$ anticommutes with all the fermions in the NS representation and has eigenvalue +1 in even subspace and -1 at the odd subspace. Collecting all the terms we have

$$C_1 = \sum_{w_1, w_2} g(w_1; w_2) F(w_1; w_2); \quad (32)$$

$$g(w_1; w_2) = [w_1 1]_R [w_1 2] [w_1 3] [w_1 4] [w_2 2]_{NS} [w_2 3] [w_2 4];$$

and

$$F(w_1; w_2) = \frac{\text{tr}_{H^{NS}} \mathbf{x}^{2D^{NS}} \begin{matrix} NS \\ 0 \end{matrix} (w_1) \begin{matrix} NS \\ 0 \end{matrix} (w_2)}{\text{tr}_{H^{NS}} \mathbf{x}^{2D^{NS}}}; \quad (33)$$

Evaluating the expression (33) gives

$$F(w_1; w_2) = M \begin{matrix} NS \\ 0 \end{matrix} \left(\mathbf{x}^2 \frac{w_1}{w_2} \right) + \frac{2kI}{i} \text{sn}(v_1 - v_2); \quad (34)$$

M being the magnetization³

$$M = \frac{\text{tr}_{H^{NS}} \mathbf{x}^{2D^{NS}} \begin{matrix} NS \\ 0 \end{matrix}}{\text{tr}_{H^{NS}} \mathbf{x}^{2D^{NS}}} = (1 - k^2)^{1/8};$$

Carrying out the integrations explicitly the three-point function is

$$C_1 = M \sum_n \prod_{1 \leq i < j \leq 4} (\text{sn}(i_{ij}))^{-1} \begin{matrix} \text{dn}(i_{24}) \text{sn}(i_{14}) [\text{sn}(i_{23}) - \text{cn}(i_{12}) \text{sn}(i_{13}) + \text{sn}(i_{12}) \text{cn}(i_{13})] \\ \text{dn}(i_{23}) \text{sn}(i_{13}) [\text{sn}(i_{24}) - \text{cn}(i_{12}) \text{sn}(i_{14}) + \text{sn}(i_{12}) \text{cn}(i_{14})] \\ + \text{sn}(i_{12}) [\text{sn}(i_{34}) - \text{cn}(i_{13}) \text{sn}(i_{14}) + \text{sn}(i_{13}) \text{cn}(i_{14})] \end{matrix}; \quad (35)$$

³The operator $\begin{matrix} NS \\ 0 \end{matrix}$ represents the central spin matrix $\frac{z}{1}$ under the Jordan-Wigner transformation, see [4].

To make explicit the symmetries of this function, we employ the identities, $\text{cn}(u-v) = \text{cn}u\text{cn}v + \text{sn}u\text{sn}v\text{dn}(u-v)$ and $\text{dn}u\text{sn}(u-v) = -\text{cn}u\text{sn}v + \text{sn}u\text{cn}v\text{dn}(u-v)$, obtaining

$$C_1 = M \sum_{1 \leq i < j \leq 4} (\text{sn}(i_{ij}))^{-1} \text{sn}(i_{12})\text{sn}(i_{34}) + \text{cn}(i_{13})\text{cn}(i_{24}) - \text{cn}(i_{14})\text{cn}(i_{23}) + \text{sn}(i_{14})\text{sn}(i_{23})[\text{dn}(i_{13}) + \text{dn}(i_{24})] - \text{sn}(i_{13})\text{sn}(i_{24})[\text{dn}(i_{14}) + \text{dn}(i_{23})] ; \quad (36)$$

eq. (36) has now symmetries under the transpositions $1 \leftrightarrow 2$ or $3 \leftrightarrow 4$ and the geometrical symmetry under the double transposition $1 \leftrightarrow 4$ and $2 \leftrightarrow 3$ see Fig. 5; note that the above expression is not invariant under the sole transposition $2 \leftrightarrow 3$ or $1 \leftrightarrow 4$.

Taking all the i in (35) or (36) equal and using (22) we obtain the homogeneous diagonal three-point function⁴

$$C_1^{\text{hom}} = M^{-1} \frac{k^2}{4} ; \quad (37)$$

Let us investigate the case that the third spin goes to infinity the other spins remaining adjacent diagonal neighbors in the homogeneous case Fig. 6. By the cluster property we expect that the CF tends to the product of spin expectation value times the two-point function as the spin $_3$ tends to infinity, this limit will be another check for the VOM. The



Figure 6: 3-point function with one spin at arbitrary distance.

double integral expression for the homogeneous case is then

$$C_n^{\text{hom}} = M \frac{k^{2n+2}}{2} \int_I^Z \int_I^Z dv_1 dv_2 \text{dn } v_1 \text{sn}^{2n+1} v_1 \text{cn } v_2 \text{sn}^{2n} v_2 - \text{sn}(v_1 - v_2) + \frac{i}{k} \text{NS}(v_1 - v_2 + i\Gamma^0) ; \quad (38)$$

where $n = -1$ corresponds to the case of adjacent diagonal spins. The integration of the delta function gives

$$\frac{1}{2} \int_I^Z \int_I^Z dv \text{dn}^2 v = \frac{2}{-E(k)} ; \quad (39)$$

where $E(k)$ is the complete elliptic integral of second kind. The integration of the first term in (38) could be done in principle by residues but the most simple way seems to be to use the addition formula for the function $\text{sn}(v_1 - v_2)$,

$$\text{sn}(v_1 - v_2) = \frac{\text{sn } v_1 \text{cn } v_2 \text{dn } v_2 - \text{sn } v_2 \text{cn } v_1 \text{dn } v_1}{1 - k^2 \text{sn}^2 v_1 \text{sn}^2 v_2} ;$$

substituting this formula we see that only the first factor is not zero on symmetry grounds

$$I_n = \frac{k^{2n+2}}{2} \int_I^Z \int_I^Z dv_1 dv_2 \frac{\text{sn}^{2n+2} v_1 \text{sn}^{2n} v_2 \text{cn}^2 v_2 \text{dn } v_1 \text{dn } v_2}{1 - k^2 \text{sn}^2 v_1 \text{sn}^2 v_2} ; \quad (40)$$

⁴This limit can also be done, of course, in the integral expression first and then integrate by residues.

Define the following integrals

$$J_n = \int_I dv \, dn \, v \, \text{sn}^{2n} v; \quad (41)$$

by differentiation of the expression $\int_I dv \, \text{sn}^{2n} v$ it can be proved that the integrals J_n satisfy the following recursion relations

$$J_{n+1} = \frac{2n+1}{2n+2} J_n;$$

In the interval $I = [v_1; v_2]$ we can expand the denominator in (40)

$$(1 - k^2 \text{sn}^2 v_1 \text{sn}^2 v_2)^{-1} = \sum_{j=0}^{\infty} k^{2j} \text{sn}^{2j} v_1 \text{sn}^{2j} v_2;$$

putting this expression under the integrations and using the definitions of J_n one arrives to the following series (interchange of the series with the integrals is allowed due to the uniform convergence of the series in the interval of integration)

$$I_n = \sum_{j=0}^{\infty} \frac{k^{2n+2}}{2} \sum_{j=0}^{\infty} k^{2j} J_{n+j+1} (J_{n+j} - J_{n+j+1});$$

using the recursion relations and changing the index of summation one has

$$I_n = \frac{1}{2} \sum_{j=n+1}^{\infty} \frac{k^{2j} J_j^2}{2j-1};$$

now because $J_0 = 1$ by the recursion relations

$$J_j = \frac{(2j-1)!!}{(2j)!!};$$

finally

$$I_n = \sum_{j=n+1}^{\infty} k^{2j} \frac{(2j-1)!!^2}{(2j)!!^2} \frac{1}{2j-1};$$

by the series definition of the elliptic integral $E(k)$ (see e.g. [6])

$$I_0 = 1 - \frac{1}{2} E(k)$$

we also have that

$$I_n - I_0 = \sum_{j=1}^n k^{2j} \frac{(2j-1)!!^2}{(2j)!!^2} \frac{1}{2j-1};$$

therefore the final expression for the 3-point function G_n is

$$C_n^{\text{hom}} = M^{-1} \sum_{j=1}^n k^{2j} \frac{(2j-1)!!^2}{(2j)!!^2} \frac{1}{2j-1}; \quad (42)$$

with $n \rightarrow \infty$ G_n tends to $M^{-1} E(k)$ i.e. the product of the magnetization with the two-point function with as we expect.

6 Conclusion

We have seen in the calculation of different CF how intimate is the connection of the IM with the theory of elliptic functions and how the vertex operator method allows to make this connection very directly. In this way several scattered results about the CF of the Ising model in the literature could be treated in a general way. Because the method is not totally rigorous, specially in the definition of the CTM, several consistency checks were done: comparison with results already known by other methods or physical reasonable consequences.

We have only treated purely order or disorder CF but with the same methods exposed in this article one can write immediately integral expressions for the mixed CF. A more difficult problem is to obtain a complete integration of the higher CF as was done for the two-point functions or alternatively to obtain possible equations satisfied by the CF like the ones found in [9] and references therein. Also the generalized Toeplitz determinants found in Sect. 4.2 and 4.3 could be used to see what kind of nonlinear differential equations the CF satisfy in the same way as was done in [13].

Acknowledgments Supported in part by DFG Sfb 288 "Differentialgeometrie und Quantenphysik" and by research funds of the Freie Universität Berlin.

A Integration of the two-point correlation function

Here I give the details of the explicit integration of order-order correlation function. The disorder-disorder case is analogous and I will indicate in passing the modification needed.

I begin remembering the expression obtained for the disorder-disorder CF on page 6 equation (15), that I write now in the following way

$$g_{2n} = \int \prod_{i=1}^{2n} dw_i G_{2n}^R fw_i g = F_{2n}^R fw_i g; \quad (43)$$

where

$$F_{2n}^R fw_i g = \sum_p \int \prod_{i=1}^{2n} dw_i \exp \left(\sum_{i,j=1}^{2n} p_{ij} w_i w_j \right); \quad (44)$$

with the same notation as in Sect. 4.1, and

$$G_{2n}^R fw_i g = \int \prod_{i=1}^{2n} dw_i \exp \left(\sum_{i,j=1}^{2n} w_i w_j \right); \quad (45)$$

As I showed in [12], before inserting this formula under the integral expression it is more convenient to antisymmetrize the factor multiplying the fermions in all the w_i variables (under the integrals the fermions can be anticommutated as already observed), for this we change the spectral parameters and integration variables to their additive version,

$$\begin{aligned} \int \prod_{i=1}^{2n} dw_i &= \int \prod_{i=1}^{2n} \frac{dw_i}{2\pi i}; \\ \int \prod_{i=1}^{2n} dw_i \exp \left(\sum_{i,j=1}^{2n} w_i w_j \right) &= \int \prod_{i=1}^{2n} \frac{dw_i}{2\pi i} \exp \left(\sum_{i,j=1}^{2n} w_i w_j \right); \\ \int \prod_{i=1}^{2n} dw_i \exp \left(\sum_{i,j=1}^{2n} w_i w_j \right) &= \int \prod_{i=1}^{2n} \frac{dw_i}{2\pi i} \exp \left(\sum_{i,j=1}^{2n} w_i w_j \right); \\ \int \prod_{i=1}^{2n} dw_i \exp \left(\sum_{i,j=1}^{2n} w_i w_j \right) &= \int \prod_{i=1}^{2n} \frac{dw_i}{2\pi i} \exp \left(\sum_{i,j=1}^{2n} w_i w_j \right); \end{aligned}$$

The result of the antisymmetrization is

$$\frac{\prod_{i=1}^n H_1(v_{2i-1} - i_{2i-1}) \prod_{i=1}^n H_1(v_{2i} - i_{2i})}{\prod_{i=1}^n \prod_{j=1}^n (v_i - i_j)} + \text{perm} = \frac{\prod_{i=1}^n \prod_{j=1}^n (i_i - i_j) H(v_i - v_j) (H_1(0) - 1(0))^n}{H_1(0)^{2n} \prod_{i=1}^n \prod_{j=1}^n (v_i - i_j)} \prod_{i=1}^n X^{2n}_{v_i - i} \prod_{r=1}^n X^{2n}_r ; \quad (46)$$

where by perm we mean the antisymmetrization in all v_i . The same formula is valid under the interchange of v_i with H_1 . This identity is proved showing first that both sides transform identically under the periods $2I$ and $2iI^0$ in all v_i , and then through an inductive process it is possible show that the residues of the simple poles at e.g. $v_1 = i_1 + iI^0$ are identical for the two sides. As a simple consequence of Liouville's theorem the rhs can only differ by an additive constant, which is proved to be zero by comparing the two sides at e.g. $v_{2n} = i_{2n}$. Therefore we can now write

$$G_{2n}^R f v_i g = \frac{c}{(2n)!} \frac{\prod_{i=1}^n \prod_{j=1}^n (i_i - i_j) H(v_{ij})}{H_1(0)^{2n} \prod_{i,j=1}^n (v_i - i_j)} \prod_{i=1}^n X^{2n}_{v_i - i} \prod_{r=1}^n X^{2n}_r ; \quad (47)$$

$c := (ik^0 H_1(0) - 1(0))^n$ is a constant factor; $i_{ij} = i_i - i_j$ and $v_{ij} = v_i - v_j$.

Comparing the rhs. of the above equation with the rhs. of the following classical result⁵ (see e.g. App. A in [16])

$$\det_{1 \leq i < j \leq 2n} \frac{(v_i - i_j) H(0)^0}{(v_i - i_j) H_1(0)} = \frac{(H^0(0))^{2n}}{H_1(0)^{2n}} \frac{\prod_{i=1}^n \prod_{j=1}^n H(v_{ij}) H(i_{ij})}{\prod_{i,j=1}^n (v_i - i_j)} \prod_{i=1}^n X^{2n}_{v_i - i} ;$$

we see that by multiplication of some factors dependent on the variables v_i one can write

$$G_{2n}^R f v_i g = B(i) \frac{1}{(2n)!} \det_{1 \leq i < j \leq 2n} \hat{h}(v_i - i_j) \quad (48)$$

with

$$\hat{h}(v) := \frac{(v) H(0)^0}{(v) H_1(0)} = \text{dn } v ; \quad (49)$$

In the order case one has instead $\hat{h}(v) := \text{kc}n v$.

The function $B(i)$ is:

$$B(i) := \frac{c}{(H^0(0))^{2n}} \prod_{i < j} \frac{(i_i - i_j)}{H(i_{ij})} = \frac{1}{k^{2n}} \prod_{i < j} \frac{p_{-i_j}}{k^{2n} i_{ij}} ; \quad (50)$$

Inserting eqs. (48), (49) and (50) under the integration symbols gives (I consider now simultaneously the order case also)

$$g_{2n}^i = B(i) \int_{i=1}^n \int_{I} dv_i \frac{1}{(2n)!} \prod_{p=1}^n \hat{h}(v_j - i_{pj}) \prod_{q=1}^n (v_{q1} - v_{q2} + iI^0) \prod_{q=1}^n (v_{q1} - v_{q2n} + iI^0) ; \quad (51)$$

⁵This formula is incorrectly given in reference [12].

Now each of the $(2n-1)!!$ terms in the sum over the delta functions gives the same result when integrated over their respective variables v_i , because both factors under the integrals are totally antisymmetric in the v_i 's, therefore it is only necessary to integrate over the first term in the second sum i.e., $(v_1 - v_2 + iI^0)(v_3 - v_4 + iI^0) \dots (v_{2n-1} - v_{2n} + iI^0)$ multiplied by the constant $(2n-1)!! = (2n)! = 1 = (2^n n!)$. Integrating out the deltas one is left with (after renaming the integration variables)

$$g_{2n}^i = B(i) \int_{-I}^I \prod_{i=1}^{2n} dv_i \frac{1}{2^n n!} \prod_{j=1}^{2n} \hat{h}(v_j - i_{P(j)});$$

$$\prod_{i < j} \frac{1}{k \sin i_{ij}} \prod_{i=1}^{2n} \hat{h}(i_{P(i)} - i_{P(2)}); \hat{h}(i_{P(2n-1)} - i_{P(2n)});$$

where $\hat{h}(i) = \int_{-I}^I dv \hat{h}(v - i)$ and the following definition was introduced:

$$\hat{h}(i_i - i_j) = \int_{-I}^I \frac{1}{k} dv \hat{h}(v - i_i) \hat{h}(v - i_j):$$

For the integration of the last formula we use (see [15])

$$\int_{-I}^I du \operatorname{cn}(u + ia) \operatorname{cn}(u + ib) = 2I [\operatorname{cn}(ia - ib) + ik^{-1} \operatorname{cn}(ia - ib - iI^0) \operatorname{zn}(ia - ib)] \quad (52)$$

$$\int_{-I}^I du \operatorname{dn}(u + ia) \operatorname{dn}(u + ib) = 2I [\operatorname{dn}(ia - ib) + idn(ia - ib - iI^0) \operatorname{zn}(ia - ib)]: \quad (53)$$

Equations (52) and (53) can be shown under the condition

$$I^0 < a; b < I^0$$

for real numbers a and b . $\operatorname{zn} u = \operatorname{zn}^0(u) = \operatorname{zn}(u)$ is the Jacobi zeta function. Employing these results we have by analytic continuation,

$$\hat{h}(i_i - i_j) = \frac{2I}{k} [\operatorname{dn} i_{ij} \operatorname{sn} i_{ij} - \operatorname{cn} i_{ij} \operatorname{zn} i_{ij}] + \frac{1}{k} \operatorname{cn} i_{ij} \quad (54)$$

$$\hat{h}(i_i - i_j) = \frac{2I}{k} [k^2 \operatorname{cn} i_{ij} \operatorname{sn} i_{ij} - \operatorname{dn} i_{ij} \operatorname{zn} i_{ij}] + \frac{1}{k} \operatorname{dn} i_{ij}: \quad (55)$$

The terms in square brackets in eqs. (54) and (55) are odd under interchanging i with j ; the last terms are even. Inserting these expressions under the formula for g_{2n}^i only the odd terms survive. Now the sum in the eqs. for g_{2n}^i is an equivalent way of writing a pfaffian, thus

$$g_{2n}^i = \prod_{i < j} \frac{1}{k \sin i_{ij}} \operatorname{Pfaffian} [\hat{h}(i_i - i_j)]$$

Using the equivalence of the zeta function with the logarithmic derivative of the $\zeta(u)$ function one sees that the factors in square brackets are equivalent to the expressions on page 9, eqs. (20). Using now the identity

$$\operatorname{Pfaffian} \frac{1}{k \sin(i_i - i_j)} = \prod_{1 \leq i < j \leq 2n} \frac{1}{k \sin(i_i - i_j)}:$$

proved by use of Liouville's theorem, see e.g. [11], eq. (5.8). This proves finally eq. (19).

References

- [1] Au-Yang, H., Perk, J., *Physica* 144A (1987) 44.
- [2] Baxter, R. *Exactly solved models in statistical mechanics* (Academic Press, London, 1982)
- [3] Baxter, R. *Phil. Trans. Roy. Soc. A* 289 (1978) 315.
- [4] Foda, O., Jimbo, M., Miwa, T., Mukai, T. and Nakayashiki, A., *J. Math. Phys.* 35 (1994) 13, hep-th/9305100
- [5] Gosh, R., Shrock, R., *Phys. Rev. B* 30 (1984) 3790.
- [6] Gradshteyn, I., Ryzhik, I., *Table of Integrals, series and products* (Academic Press, New York, 1965)
- [7] Jimbo, M., Mukai, K., Miwa, T., and Nakayashiki, A., *Phys. Lett. A* 168 (1992) 256, hep-th/9205055
- [8] Jimbo, M., Miwa, T., *Algebraic Analysis of Solvable Lattice Models* CBMS 85 (Amer. Math. Soc., 1993)
- [9] McCoy, B. in, *Important developments in Soliton theory*, A.S. Fokas and V.E. Zakharov (Eds.) (Springer-Verlag, 1993)
- [10] McCoy, B., Wu, T.T., *The two-dimensional Ising model* (Harvard University Press, 1973)
- [11] Palmer, J., Tracy, C., *Adv. in Appl. Math.* 2 (1981) 329.
- [12] Reyes Martinez, J., *Phys. Lett. A* 277 (1997) 203, hep-th/9609135
- [13] Sato, M., Jimbo, M., and Miwa, T., *Publ. RIMS, Kyoto Univ.* 15 (1979) 577, 871
- [14] Stephenson, J., *J. Math. Phys.* 7 (1966) 1123.
- [15] Yamada, K., *Prog. Theor. Phys.* 72 (1984) 922.
- [16] Yamada, K., *Prog. Theor. Phys.* 76 (1986) 602.