

Exact static soliton solutions of $3 + 1$ dimensional integrable theory with nonzero Hopf numbers

H. Aratyn¹, L.A. Ferreira² and A.H. Zimerman²

¹Department of Physics
University of Illinois at Chicago
845 W. Taylor St., Chicago, IL 60607-7059

²Instituto de Física Teórica - IFT/UNESP
Rua Pamplona 145
01405-900, São Paulo - SP, Brazil

Abstract

In this paper we construct explicitly an infinite number of Hopfions (static, soliton solutions with non-zero Hopf topological charges) within the recently proposed $3 + 1$ -dimensional, integrable and relativistically invariant field theory.

Two integers label the family of Hopfions we have found. Their product is equal to the Hopf charge which provides a lower bound to the soliton's finite energy. The Hopfions are constructed explicitly in terms of the toroidal coordinates and shown to have a form of linked closed vortices.

Recent numerical studies of the Faddeev-Skyrme modified $O(3)$ sigma-model [1] support existence of static toroidal solutions stabilized by their non-zero Hopf numbers [2]. The emerging string-like structures are quite intriguing and may find applications in various physical models of condensed matter physics and gauge field theory.

In [3] we have introduced the three-dimensional field model which falls into a class of higher dimensional integrable models from the point of view of the generalized zero-curvature approach [4]. The question posed in [3] was whether this form of integrability is linked to the existence of soliton solutions as it is expected from the study of two-dimensional integrable models. Our analysis of the model in [3] has indeed revealed one non-trivial soliton solution described by a standard Hopf map of unit Hopf index. To fully establish connection between integrability and soliton solutions would require finding other topological solitons with arbitrary topological charges. This is accomplished in this letter. The equations of motion of the model are solved in toroidal coordinates and the space of solutions is found to be represented by a family of maps $\mathbb{R}^3 \rightarrow \mathbb{R}^2$ labeled by two integers. The integers count number of times the map winds around two independent angular directions.

The model under consideration is described by the Lagrangean density

$$\mathcal{L} = -\eta_0 \left(H_{\mu\nu}^2 \right)^{\frac{3}{4}} \quad (1)$$

where $\eta_0 = \pm 1$, determines the choice of the signature of the Minkowski metric, $g_{\mu\nu} = \eta_0 \text{diag}(1, -1, -1, -1)$ and the field-tensor $H_{\mu\nu}$ is defined in terms of the three component, unit vector field $\vec{n} \in S^2$ as

$$H_{\mu\nu} \equiv \vec{n} \cdot (\partial_\mu \vec{n} \times \partial_\nu \vec{n}) . \quad (2)$$

The action in (1) is $O(3)$ and Poincare invariant. The value 3/4 of the power of $H_{\mu\nu}^2$ in (1) is such that the theory circumvents the usual obstacle of Derrick's scaling argument against existence of stable solitons. We are interested in the boundary condition $\vec{n} = (0, 0, 1)$ at spatial infinity. This condition compactifies effectively the Euclidean space \mathbb{R}^3 to the three-sphere S^3 . Accordingly, \vec{n} becomes a map: $S^3 \rightarrow S^2$. Due to $\pi_3(S^2) = \mathbb{Z}$, the field configurations fall into disjoint classes characterized by the value of the Hopf invariant Q_H .

Using stereographic projection of S^2 :

$$\vec{n} = \frac{1}{1 + |u|^2} \left(u + u^*, -i(u - u^*), |u|^2 - 1 \right) \quad (3)$$

one obtains

$$H_{\mu\nu} = \frac{-2i}{(1 + |u|^2)^2} (\partial_\mu u \partial_\nu u^* - \partial_\nu u \partial_\mu u^*) \quad (4)$$

where u is a complex scalar field.

In terms of the vector quantities

$$K_\mu = \frac{i}{2} (1 + |u|^2)^2 H_{\mu\nu} \partial^\nu u = (\partial^\nu u^* \partial_\nu u) \partial_\mu u - (\partial_\nu u)^2 \partial_\mu u^* \quad (5)$$

and

$$\mathcal{K}_\mu \equiv \frac{(K \partial u^*)^{-\frac{1}{4}} K_\mu}{1 + |u|^2} \quad (6)$$

the corresponding equations of motion can be rewritten compactly as

$$\partial^\mu \mathcal{K}_\mu = 0 \quad (7)$$

The quantity (5) automatically satisfies the relations

$$K^\mu \partial_\mu u = 0 \quad \text{Im}(K^\mu \partial_\mu u^*) = 0 \quad (8)$$

which play a crucial role in establishing integrability of the model. Indeed, using relations (8) and the equations of motion (7) one obtains an infinite number of conserved currents given by

$$J_\mu \equiv \mathcal{K}_\mu \frac{\delta G}{\delta u} - \mathcal{K}_\mu^* \frac{\delta G}{\delta u^*} \quad (9)$$

with G being any functional of u and u^* only (no derivatives). In reference [3] we have analyzed the integrability properties of this theory using the generalized version of the zero curvature condition [4]. The equations of motion (7) can be represented as $\partial^\mu B_\mu + [A^\mu, B_\mu] = 0$, with A^μ being a flat $SU(2)$ connection, and B_μ being an operator living in any integer spin representation of $SU(2)$ (for more details see [3, 4]). The integrability properties emerge due to the fact that B_μ can be put in any integer spin representation, and that is a direct consequence of (8) [4, 5]. The question we address here is whether the above notion of integrability implies existence of the infinite many soliton solutions.

We shall look for time-independent solutions to equations of motion (7) of the type¹:

$$u(\eta, \xi, \varphi) \equiv f(\eta) e^{i(m\xi + n\varphi)} \quad (10)$$

where we used toroidal coordinates on \mathbb{R}^3 :

$$\begin{aligned} x &= aq^{-1} \sinh \eta \cos \varphi, & y &= aq^{-1} \sinh \eta \sin \varphi \\ z &= aq^{-1} \sin \xi; & a &> 0; & q &= \cosh \eta - \cos \xi \end{aligned} \quad (11)$$

¹We shall take m and n integers in order for u to be single valued.

The angles ξ, φ both vary from 0 to 2π and η varies from 0 to ∞ . The surfaces of constant η are toroids that circle the z -axis, ξ =constant are spheres and φ =constant are half-planes. The corresponding gradient of u becomes in toroidal coordinates:

$$\vec{\nabla}u = (q/a)e^{i(m\xi+n\varphi)} \left(f'(\eta)\hat{e}_\eta + imf(\eta)\hat{e}_\xi + inf(\eta)\frac{\hat{e}_\varphi}{\sinh \eta} \right) \quad (12)$$

where we introduced the unit vectors spanning the orthogonal toroidal coordinate system having properties $\hat{e}_i \cdot \hat{e}_j = \delta_{ij}$, $i, j \equiv \eta, \xi, \varphi$.

In terms of the vector-field $\vec{\mathcal{K}} \equiv \left(\vec{K} \cdot \vec{\nabla}u^* \right)^{-1/4} \vec{K}/(|u|^2+1)$ the equations of motion (7) become, in the static case, $\vec{\nabla} \cdot \vec{\mathcal{K}} = 0$, i.e. $\vec{\mathcal{K}}$ is solenoidal. Using (10) and (12) we obtain for the components of $\vec{\mathcal{K}}$:

$$\mathcal{K}_\eta = \mathcal{K}_0 \left(m^2 + \frac{n^2}{(\sinh \eta)^2} \right) f ; \quad \mathcal{K}_\xi = im\mathcal{K}_0 f' ; \quad \mathcal{K}_\varphi = in\mathcal{K}_0 \frac{f'}{\sinh \eta} \quad (13)$$

with

$$\mathcal{K}_0 \equiv \sqrt{2}(q/a)^2 e^{i(m\xi+n\varphi)} \left(m^2 + \frac{n^2}{(\sinh \eta)^2} \right)^{-1/4} \frac{f^{1/2} f'^{1/2}}{(f^2 + 1)} \quad (14)$$

where the components of the vector field are defined according to $\vec{V} = \hat{e}_\eta V_\eta + \hat{e}_\xi V_\xi + \hat{e}_\varphi V_\varphi$.

Plugging the components of $\vec{\mathcal{K}}$ into the expression for divergence of the vector field in the toroidal coordinates

$$\vec{\nabla} \cdot \vec{V} = \frac{q}{a} \left(\frac{\partial V_\eta}{\partial \eta} + \frac{\partial V_\xi}{\partial \xi} + \frac{1}{\sinh \eta} \frac{\partial V_\varphi}{\partial \varphi} - 2V_\xi \frac{\sin \xi}{q} + V_\eta \left(\frac{\cosh \eta}{\sinh \eta} - \frac{2 \sinh \eta}{q} \right) \right) \quad (15)$$

and using

$$\frac{\partial q^2}{\partial \eta} = q^2 \frac{2 \sinh \eta}{q} ; \quad \frac{\partial q^2}{\partial \xi} = q^2 \frac{2 \sin \xi}{q} \quad (16)$$

we arrive at:

$$\frac{\partial}{\partial \eta} \ln \frac{f f'}{(f^2 + 1)^2} = - \frac{2m^2(\sinh \eta)^2 - n^2 \cosh \eta}{m^2(\sinh \eta)^2 + n^2} \frac{1}{\sinh \eta} \quad (17)$$

We take $m^2 > n^2$ in eq.(17). The integration yields:

$$\frac{1}{f^2 + 1} = \frac{2k_1}{|m|(m^2 - n^2)} \frac{\cosh \eta}{\left(\frac{n^2 - m^2}{m^2} + \cosh^2 \eta \right)^{1/2}} + k_2 \quad (18)$$

with k_2 being the integration constant of the last integration. Imposing boundary conditions:

$$\vec{n} \rightarrow (0, 0, 1) \quad \text{or} \quad |u| \rightarrow \infty \quad \text{or} \quad f \rightarrow \infty \quad \text{as} \quad \eta \rightarrow 0 \quad (19)$$

$$\vec{n} \rightarrow (0, 0, -1) \quad \text{or} \quad u \rightarrow 0 \quad \text{or} \quad f \rightarrow 0 \quad \text{as} \quad \eta \rightarrow \infty \quad (20)$$

one gets

$$f^2 = \frac{\cosh \eta - \sqrt{n^2/m^2 + \sinh^2 \eta}}{\sqrt{1 + m^2/n^2 \sinh^2 \eta} - \cosh \eta} \quad (21)$$

One observes that f depends only on the ratio m^2/n^2 , and that $f^2 \geq 0$ for any value of m^2/n^2 . Note, that taking the limit $m \rightarrow n$ in (21) and using L'Hospital rule yields:

$$\lim_{m \rightarrow n} f^2 = \frac{1}{\sinh^2 \eta} \quad (22)$$

which gives for $m = n$ a solution:

$$u = \frac{e^{im(\xi+\varphi)}}{\sinh \eta} \quad (23)$$

In the special case of $m = n = 1$ eq.(23) reproduces the standard soliton solution [6, 3]. Expression (23) can be written as a composite of the Hopf map together with the stereographic map $: \mathbb{R}^3 \rightarrow S^3$ of degree 1.

A formula:

$$E \equiv \int d^3x \Theta_{00} = 8^{\frac{3}{4}} \int d^3x \frac{(K_i \partial^i u^*)^{\frac{3}{4}}}{(1 + |u|^2)^3} \quad (24)$$

describes energy of the static configuration [3]. Inserting our solution into (24) we obtain an expression for the energy $E = E_{m,n}$ of the soliton configuration

$$E_{m,n} = (2\pi)^2 8 \cdot 2^{3/4} \int_0^\infty \frac{d\eta \sinh \eta}{(1 + f^2)^3} \left(m^2 + \frac{n^2}{(\sinh \eta)^2} \right)^{3/4} f^{3/2} f'^{3/2} \quad (25)$$

which after the η -integration yields:

$$E_{m,n} = (2\pi)^2 4 \cdot 2^{1/4} \sqrt{|n||m|(|n| + |m|)} \quad (26)$$

We now turn to calculation of the Hopf numbers. Define functions $\Phi_i, i = 1, \dots, 4$ as follows:

$$\Phi \begin{Bmatrix} 1 \\ 2 \end{Bmatrix} = \left(\frac{\cosh \eta - (n^2/m^2 + \sinh^2 \eta)^{1/2}}{(|m/n| - 1) (n^2/m^2 + \sinh^2 \eta)^{1/2}} \right)^{1/2} \begin{Bmatrix} \cos m\xi \\ \sin m\xi \end{Bmatrix} \quad (27)$$

$$\Phi \begin{Bmatrix} 3 \\ 4 \end{Bmatrix} = \left(\frac{(1 + (m^2/n^2) \sinh^2 \eta)^{1/2} - \cosh \eta}{(|m/n| - 1) (n^2/m^2 + \sinh^2 \eta)^{1/2}} \right)^{1/2} \begin{Bmatrix} \cos n\varphi \\ -\sin n\varphi \end{Bmatrix} \quad (28)$$

They provide parametrization of S^3 and satisfy :

$$u = \frac{Z_1}{Z_2} = \frac{\Phi_1 + i\Phi_2}{\Phi_3 + i\Phi_4} \quad |Z_1|^2 + |Z_2|^2 = 1 \quad (29)$$

or equivalently, for \vec{n} defined from u via relation (3) :

$$n_i = Z^\dagger \sigma_i Z \quad ; \quad Z^\dagger = (Z_1^*, Z_2^*) \quad ; \quad Z = \begin{pmatrix} Z_1 \\ Z_2 \end{pmatrix} \quad (30)$$

Let, furthermore

$$A_i = \frac{i}{2} (Z^\dagger \partial_i Z - \partial_i Z^\dagger Z) \quad (31)$$

be a vector potential for the two-form $H_{ij} = \partial_i A_j - \partial_j A_i$. In terms of the toroidal coordinates the components of \vec{A} are

$$A_\eta = q \partial_\eta (\Phi_3 \Phi_4) \quad ; \quad A_\xi = -mq (\Phi_1^2 + \Phi_2^2) \quad ; \quad A_\varphi = -\frac{nq}{\sinh \eta} (\Phi_3^2 + \Phi_4^2) \quad (32)$$

Next, we calculate the vector function $\vec{B} = \vec{\nabla} \times \vec{A}$ (such that $B_i = \epsilon_{ijk} H_{jk}/2$), using expression for the curl operator in toroidal coordinates and plugging it into the Hopf index defined as :

$$Q_H = \frac{1}{4\pi^2} \int d^3x \vec{A} \cdot \vec{B} \quad (33)$$

for which we now find

$$Q_H = \frac{nm}{2} \left(\left(\Phi_1^2 + \Phi_2^2 \right)^2 \Big|_0^\infty - \left(\Phi_3^2 + \Phi_4^2 \right)^2 \Big|_0^\infty \right) = -nm \quad (34)$$

It is always possible to choose the Hopf index such that it is positive $Q_H \geq 0$. This amounts to the right choice of orientation, which determines the sign of Q_H .

Note, that the following inequality:

$$|m| + |n| \geq 2\sqrt{|n||m|} \quad (35)$$

always holds. Hence

$$\sqrt{|n||m|(|n| + |m|)} \geq \sqrt{2}(|n||m|)^{3/4} \rightarrow E_{m,n} \geq (2\pi)^2 4 \cdot 2^{3/4} |Q_H|^{3/4} \quad (36)$$

which agrees with the lower bound result by [7] for the Faddeev-Skyrme model.

Notice that the solution obtained above implies that the vector field \vec{n} (see (3)) is given by

$$n_1 = \frac{2f}{f^2 + 1} \cos(m\xi + n\varphi) \quad ; \quad n_2 = \frac{2f}{f^2 + 1} \sin(m\xi + n\varphi) \quad ; \quad n_3 = \frac{f^2 - 1}{f^2 + 1} \quad (37)$$

Since f is a function of η only, one observes that the surfaces of constant n_3 are tori. In addition, on those surfaces, the lines of constant n_1 and n_2 wind around the tori with frequencies in the ξ and φ -directions given by m and n , respectively. One can check that n_3 falls monotonically from $n_3 = 1$ at $\eta = 0$ to $n_3 = -1$ at $\eta = \infty$. In addition, the bigger the ratio m/n is, the faster it performs that flip. Therefore the size of our soliton decreases with the increase of m/n .

The Hopf index of our solution can alternatively be calculated in the following way. The solution provides a mapping of the spatial \mathbb{R}^3 into S^3 defined by $\sum_{i=1}^4 \Phi_i^2 = 1$. Then, (3) and (29) provide the Hopf map $S^3 \rightarrow S^2$. The Hopf index is given by the linking number of the pre-images, under the Hopf map, of any two points of S^2 . Consider the point $\vec{n} = (0, 0, -1)$ which corresponds in \mathbb{R}^3 to $\eta \rightarrow \infty$. For this value of η , Φ_1, Φ_2 go to zero and $\Phi_3 = \cos n\varphi$ and $\Phi_4 = -\sin n\varphi$. On S^3 we find the circle $\Phi_3^2 + \Phi_4^2 = \cos^2 n\varphi + \sin^2 n\varphi = 1$ of radius 1 and as φ varies between 0 and 2π the preimage wraps $|n|$ times around the φ direction. Similarly the preimage of $\vec{n} = (0, 0, 1)$ corresponds to $\eta = 0$. For this value of η , Φ_3, Φ_4 go to zero and $\Phi_1 = \cos m\xi$ and $\Phi_2 = \sin m\xi$. On S^3 we find the circle $\Phi_1^2 + \Phi_2^2 = \cos^2 m\xi + \sin^2 m\xi = 1$ of radius 1 and as ξ varies between 0 and 2π the preimage wraps $|m|$ times around the ξ direction. Since these two circles intersect the linking number is $|nm|$. That is indeed the Hopf number calculated above.

Finally, let us mention that other models circumventing Derrick's theorem have been proposed previously in the literature [8, 6]. A common feature is that, like the model considered in the present paper, the Lagrangians are non-polynomial functions of the fields and their derivatives. The corresponding solutions have been constructed and in the special case of [6] a soliton with the unit Hopf charge was obtained. To our knowledge, an infinite number of soliton solutions to the field theoretical equations of motion with the Hopf charges bigger than unity were not obtained previously in the literature in an exact form.

Acknowledgements

LAF and AHZ are partially supported by CNPq (Brazil).

References

- [1] L. Faddeev, in *40 Years in Mathematical Physics*, (World Scientific, 1995).

- [2] J. Gladikowski and M. Hellmund, *Phys. Rev.* **D56** (1997) 5194, hep-th/9609035; L. Faddeev and A. Niemi, *Nature* **387** (1997) 58, hep-th/9610193; R.A. Battye, P.M. Sutcliffe, *Phys. Rev. Lett.* **81** (1998) 4798 and hep-th/9811077; J. Hietarinta and P. Salo, *Phys. Lett.* **451B** (1999) 60, hep-th/9811053
- [3] H. Aratyn, L. A. Ferreira and A.H. Zimerman, *Toroidal solitons in $3 + 1$ dimensional integrable theories*, hep-th/9902142, to appear in *Physics Letters* **B**.
- [4] O. Alvarez, L. A. Ferreira and J. Sánchez Guillén, *Nucl. Phys.* **B529** (1998) 689, hep-th/9710147.
- [5] L. A. Ferreira and E. E. Leite, *Integrable theories in any dimension and homogeneous spaces*, IFT-P/067/98, hep-th/9810067, to appear in *Nuclear Physics* **B**.
- [6] D.A. Nicole, *J. Phys.* **G4** (1978) 1363.
- [7] A. F. Vakulenko and L.V. Kapitansky, *Sov. Phys. Dokl.* **24** (1979) 432
- [8] S. Deser, M.J. Duff and C.J. Isham, *Nucl. Phys.* **B114** (1976) 29.