

# Maximal superintegrability on N-dimensional curved spaces

Angel Ballesteros<sup>†</sup>, Francisco J. Herranz<sup>†</sup>,  
Mariano Santander<sup>‡</sup> and Teresa Sanz-Gil<sup>‡</sup>

<sup>†</sup> *Departamento de Física, Facultad de Ciencias,  
Universidad de Burgos, 09001 Burgos, Spain*

<sup>‡</sup> *Departamento de Física Teórica, Facultad de Ciencias,  
Universidad de Valladolid, 47011 Valladolid, Spain*

## Abstract

A unified algebraic construction of the classical Smorodinsky–Winternitz systems on the  $ND$  sphere, Euclidean and hyperbolic spaces through the Lie groups  $SO(N+1)$ ,  $ISO(N)$ , and  $SO(N,1)$  is presented. Firstly, general expressions for the Hamiltonian and its integrals of motion are given in a linear ambient space  $\mathbb{R}^{N+1}$ , and secondly they are expressed in terms of two geodesic coordinate systems on the  $ND$  spaces themselves, with an explicit dependence on the curvature as a parameter. On the sphere, the potential is interpreted as a superposition of  $N+1$  oscillators. Furthermore each Lie algebra generator provides an integral of motion and a set of  $2N-1$  functionally independent ones are explicitly given. In this way the maximal superintegrability of the  $ND$  Euclidean Smorodinsky–Winternitz system is shown for any value of the curvature.

Superintegrable systems on the two- and three-dimensional (3D) Euclidean spaces have been classified in [1, 2], and also extended to the 2D and 3D spheres [3] as well as to the hyperbolic spaces [4, 5]. Recent classifications of superintegrable systems for these 2D Riemannian spaces can be found in [6, 7, 8]. In the 2D sphere there are two (maximal) superintegrable potentials: the harmonic oscillator ( $\tan^2 r$ ) with ‘centrifugal terms’ and the Kepler or Coulomb potential ( $1/\tan r$ ) with some ‘additional’ terms. The former is the version with non-zero curvature of the Smorodinsky–Winternitz (SW) system [9, 10, 11, 12]. Both potentials  $\tan^2 r$  and  $1/\tan r$  on the  $ND$  sphere have been studied in quantum mechanics in [13, 14, 15], and have been mutually related in [16].

The SW Hamiltonian on the  $ND$  Euclidean space is given by

$$\mathcal{H} = \frac{1}{2} \sum_{i=1}^N \left( p_i^2 + 2\beta_0 q_i^2 + \frac{2\beta_i}{q_i^2} \right) \quad (1)$$

The following functions are integrals of motion for (1) ( $i < j$ ;  $i, j = 1, \dots, N$ ):

$$I_{0i} = \tilde{P}_i^2 + 2\beta_0 q_i^2 + 2\frac{\beta_i}{q_i^2} \quad \text{with} \quad \tilde{P}_i = p_i \quad (2)$$

$$I_{ij} = \tilde{J}_{ij}^2 + 2\beta_i \frac{q_j^2}{q_i^2} + 2\beta_j \frac{q_i^2}{q_j^2} \quad \text{with} \quad \tilde{J}_{ij} = q_i p_j - q_j p_i. \quad (3)$$

The set (2) comes from the separability of the Hamiltonian  $2\mathcal{H} = \sum_i I_{0i}$ , while (3) are just the square of the components of the angular momentum tensor plus some additional terms. The functions  $\tilde{P}_i, \tilde{J}_{ij}$  close the commutation relations of the Euclidean algebra  $iso(N)$  with respect to the canonical Lie–Poisson bracket:

$$\{f, g\} = \sum_{i=1}^N \left( \frac{\partial f}{\partial q_i} \frac{\partial g}{\partial p_i} - \frac{\partial g}{\partial q_i} \frac{\partial f}{\partial p_i} \right). \quad (4)$$

Our aim is to construct, simultaneously, the non-zero curvature version of (1) on the three classical Riemannian spaces with constant curvature in arbitrary dimension, as well as to prove its maximal superintegrability, from a group theoretical standpoint.

Let  $so_\kappa(N+1)$  be the real Lie algebra of the Lie group  $SO_\kappa(N+1)$  with generators  $\{J_{0i} \equiv P_i, J_{ij}\}$  ( $i, j = 1, \dots, N$ ;  $i < j$ ) and non-vanishing commutation relations given by

$$\begin{aligned} [J_{ij}, J_{ik}] &= J_{jk} & [J_{ij}, J_{jk}] &= -J_{ik} & [J_{ik}, J_{jk}] &= J_{ij} \\ [J_{ij}, P_i] &= P_j & [J_{ij}, P_j] &= -P_i & [P_i, P_j] &= \kappa J_{ij} \end{aligned} \quad (5)$$

with  $i < j < k$ . If we consider the following Cartan decomposition of  $so_\kappa(N+1)$ :

$$so_\kappa(N+1) = h \oplus p \quad h = \langle J_{ij} \rangle \quad p = \langle P_i \rangle \quad (6)$$

where  $h$  is the Lie algebra of  $H \simeq SO(N)$ , we obtain a family of  $ND$  symmetric homogeneous spaces  $S_{[\kappa]}^N = SO_\kappa(N+1)/SO(N)$  parametrized by  $\kappa$ , which turns out to be the constant sectional curvature of the space. Thus  $J_{ij}$  leave a point  $\mathcal{O}$  invariant by acting as rotations, while  $P_i$  generate translations that move  $\mathcal{O}$  along  $N$  basic

geodesics  $l_i$  orthogonal at  $\mathcal{O}$ . For  $\kappa >, =, < 0$ ,  $S_{[\kappa]}^N$  reproduces the sphere  $\mathbf{S}^N = SO(N+1)/SO(N)$ , Euclidean  $\mathbf{E}^N = ISO(N)/SO(N)$  and hyperbolic  $\mathbf{H}^N = SO(N,1)/SO(N)$  spaces, respectively. The case  $\kappa = 0$  is the contraction around  $\mathcal{O}$ :  $\mathbf{S}^N \rightarrow \mathbf{E}^N \leftarrow \mathbf{H}^N$ .

The vector representation of  $so_\kappa(N+1)$  is given by  $(N+1) \times (N+1)$  real matrices:

$$P_i = -\kappa e_{0i} + e_{i0} \quad J_{ij} = -e_{ij} + e_{ji} \quad (7)$$

where  $e_{ij}$  is the matrix with entries  $(e_{ij})_m^l = \delta_i^l \delta_j^m$ . Any generator  $X$  of  $so_\kappa(N+1)$  fulfils

$$X^T \Lambda + \Lambda X = 0 \quad \Lambda = e_{00} + \kappa \sum_{i=1}^N e_{ii} = \text{diag}(1, \kappa, \dots, \kappa) \quad (8)$$

so that any element  $G \in SO_\kappa(N+1)$  verifies  $G^T \Lambda G = \Lambda$ . In this way,  $SO_\kappa(N+1)$  is a group of linear transformations in an ambient space  $\mathbb{R}^{N+1}$ , with Weierstrass coordinates  $\mathbf{x} = (x_0, x_1, \dots, x_N)$ , acting as the group of isometries of the bilinear form  $\Lambda$  via matrix multiplication. The Lie group  $H \simeq SO(N) = \langle J_{ij} \rangle$  is the isotopy subgroup of the origin  $\mathcal{O} = (1, 0, \dots, 0) \in \mathbb{R}^{N+1}$ . The space  $S_{[\kappa]}^N$  is identified with the orbit of  $\mathcal{O}$ , which is contained in the ‘sphere’  $\Sigma$ :

$$\Sigma \equiv x_0^2 + \kappa \sum_{i=1}^N x_i^2 = 1 \quad (9)$$

and the metric on  $S_{[\kappa]}^N$  comes from the flat ambient metric in  $\mathbb{R}^{N+1}$  in the form:

$$ds^2 = \frac{1}{\kappa} \left( dx_0^2 + \kappa \sum_{i=1}^N dx_i^2 \right) \Big|_\Sigma. \quad (10)$$

A point  $Q \in S_{[\kappa]}^N$  with Weierstrass coordinates  $\mathbf{x}$  can be reached in different ways starting from  $\mathcal{O}$  through the action of  $N$  one-parametric subgroups of  $SO_\kappa(N+1)$ :

$$\begin{aligned} \mathbf{x} &= \exp(a_1 P_1) \exp(a_2 P_2) \dots \exp(a_{N-1} P_{N-1}) \exp(a_N P_N) \mathcal{O} \\ &= \exp(\theta_N J_{N-1 N}) \exp(\theta_{N-1} J_{N-2 N-1}) \dots \exp(\theta_2 J_{12}) \exp(r P_1) \mathcal{O}. \end{aligned} \quad (11)$$

The canonical parameters involved are intrinsic quantities on  $S_{[\kappa]}^N$ , called geodesic parallel  $a = (a_1, \dots, a_N)$  and geodesic polar  $\theta = (r, \theta_2, \dots, \theta_N)$  coordinates of the point  $\mathbf{x}$ :

$$\begin{aligned} x_0 &= \prod_{s=1}^N C_\kappa(a_s) = C_\kappa(r) \\ x_1 &= S_\kappa(a_1) \prod_{s=2}^N C_\kappa(a_s) = S_\kappa(r) \cos \theta_2 \\ x_i &= S_\kappa(a_i) \prod_{s=i+1}^N C_\kappa(a_s) = S_\kappa(r) \prod_{s=2}^i \sin \theta_s \cos \theta_{i+1} \\ x_N &= S_\kappa(a_N) = S_\kappa(r) \prod_{s=2}^N \sin \theta_s. \end{aligned} \quad (12)$$

where the curvature-dependent functions  $C_\kappa(x)$  and  $S_\kappa(x)$  are defined by [17, 18]:

$$C_\kappa(x) = \begin{cases} \cos \sqrt{\kappa} x & \kappa > 0 \\ 1 & \kappa = 0 \\ \cosh \sqrt{-\kappa} x & \kappa < 0 \end{cases} \quad S_\kappa(x) = \begin{cases} \frac{1}{\sqrt{\kappa}} \sin \sqrt{\kappa} x & \kappa > 0 \\ x & \kappa = 0 \\ \frac{1}{\sqrt{-\kappa}} \sinh \sqrt{-\kappa} x & \kappa < 0 \end{cases} \quad (13)$$

The  $\kappa$ -tangent is defined by  $T_\kappa(x) = S_\kappa(x)/C_\kappa(x)$ ; its contraction  $\kappa = 0$  is  $T_0(x) = x$ .

Each parallel coordinate  $a_i$ , associated to  $P_i$ , has dimensions of *length*:  $a_1$  is the distance between  $\mathcal{O}$  and a point  $Q_1$ , measured along the basic geodesic  $l_1$ ;  $a_2$  is the distance between  $Q_1$  and another point  $Q_2$ , measured along a geodesic  $l'_2$  through  $Q_1$  and orthogonal to  $l_1$  (and ‘parallel’ in the sense of parallel transport to  $l_2$ ) and so on, up to reaching  $Q$  [18]. On the other hand, the first polar coordinate  $r$ , associated to  $P_1$ , has dimensions of *length* and is the distance between  $\mathcal{O}$  and  $Q$  measured along the geodesic  $l$  joining both points. The remaining  $\theta_i$ , associated to  $J_{i-1}i$ , are ordinary *angles*, the polar angles of  $l$  relative to the reference flag at  $\mathcal{O}$  spanned by  $\{l_1\}, \{l_1, l_2\}, \dots$ . On the sphere  $\mathbf{S}^N$  with positive curvature  $\kappa = 1/R^2$ , the usual spherical coordinates, all of which are angles, differ from ours [19] only in the first coordinate, which conventionally is taken as the dimensionless quantity  $r/R$  (see for instance [20]). When  $\kappa = 0$  we recover directly the Cartesian and polar coordinates on  $\mathbf{E}^N$ .

Next, by introducing (12) in (10), we obtain the metric in  $S_{[\kappa]}^N$ :

$$\begin{aligned} ds^2 &= \sum_{i=1}^{N-1} \left( \prod_{s=i+1}^N C_\kappa^2(a_s) \right) da_i^2 + da_N^2 \\ &= dr^2 + S_\kappa^2(r) \left( d\theta_2^2 + \sum_{i=3}^N \left( \prod_{s=2}^{i-1} \sin^2 \theta_s \right) d\theta_i^2 \right) \end{aligned} \quad (14)$$

which provides the kinetic energy  $\mathcal{T}$  in terms of the velocities ( $\dot{q} = \dot{a}, \dot{\theta}$ ), that is, the Lagrangian  $\mathcal{L} \equiv \mathcal{T}$  of a geodesic motion on  $S_{[\kappa]}^N$ . If we introduce the canonical momenta  $p = \partial\mathcal{L}/\partial\dot{q}$  ( $p = p, \pi$ ), we obtain the free Hamiltonian  $\mathcal{H} \equiv \mathcal{T}$  on  $S_{[\kappa]}^N$ :

$$\begin{aligned} \mathcal{T} &= \frac{1}{2} \left( \sum_{i=1}^{N-1} \frac{p_i^2}{\prod_{s=i+1}^N C_\kappa^2(a_s)} + p_N^2 \right) \\ &= \frac{1}{2} \left( \pi_1^2 + \frac{\pi_2^2}{S_\kappa^2(r)} + \sum_{i=3}^N \frac{\pi_i^2}{S_\kappa^2(r) \prod_{s=2}^{i-1} \sin^2 \theta_s} \right). \end{aligned} \quad (15)$$

An  $N$ -particle realization of  $so_\kappa(N+1)$  in the phase space is obtained by starting from the following expressions in terms of Weierstrass coordinates:

$$\tilde{P}_i(x(q), \dot{x}(q, p)) = x_0 \dot{x}_i - x_i \dot{x}_0 \quad \tilde{J}_{ij}(x(q), \dot{x}(q, p)) = x_i \dot{x}_j - x_j \dot{x}_i \quad (16)$$

and expressing everything either in parallel  $(a, p)$  or polar  $(\theta, \pi)$  canonical coordinates and momenta. In geodesic parallel coordinates we obtain that  $(i, j = 1, \dots, N)$ :

$$\tilde{P}_i = \prod_{k=1}^i C_\kappa(a_k) C_\kappa(a_i) p_i + \kappa S_\kappa(a_i) \sum_{s=1}^i S_\kappa(a_s) \frac{\prod_{m=1}^s C_\kappa(a_m)}{\prod_{l=s}^i C_\kappa(a_l)} p_s$$

$$\begin{aligned}\tilde{J}_{ij} = & S_\kappa(a_i) C_\kappa(a_j) \prod_{s=i+1}^j C_\kappa(a_s) p_j - \frac{C_\kappa(a_i) S_\kappa(a_j)}{\prod_{k=i+1}^j C_\kappa(a_k)} p_i \\ & + \kappa S_\kappa(a_i) S_\kappa(a_j) \sum_{s=i+1}^j S_\kappa(a_s) \frac{\prod_{m=i+1}^s C_\kappa(a_m)}{\prod_{l=s}^j C_\kappa(a_l)} p_s\end{aligned}\quad (17)$$

while in geodesic polar coordinates the same quantities read  $(i, j = 1, \dots, N-1)$ :

$$\begin{aligned}\tilde{P}_i = & \frac{\prod_{k=2}^{i+1} \sin \theta_k}{\tan \theta_{i+1}} \pi_1 + \sum_{s=2}^{i+1} \frac{\prod_{m=s}^{i+1} \sin \theta_m \cos \theta_s \pi_s}{T_\kappa(r) \tan \theta_{i+1} \prod_{l=2}^s \sin \theta_l} - \frac{\pi_{i+1}}{T_\kappa(r) \prod_{l=2}^{i+1} \sin \theta_l} \\ \tilde{P}_N = & \prod_{k=2}^N \sin \theta_k \pi_1 + \sum_{s=2}^N \frac{\prod_{m=s}^N \sin \theta_m \cos \theta_s}{T_\kappa(r) \prod_{l=2}^s \sin \theta_l} \pi_s \\ \tilde{J}_{ij} = & \sin \theta_{i+1} \cos \theta_{j+1} \prod_{k=i+1}^j \sin \theta_k \pi_{i+1} - \frac{\cos \theta_{i+1} \sin \theta_{j+1}}{\prod_{l=i+1}^j \sin \theta_l} \pi_{j+1} \\ & + \cos \theta_{i+1} \cos \theta_{j+1} \sum_{s=i+1}^j \frac{\prod_{m=s}^j \sin \theta_m \cos \theta_s}{\prod_{l=i+1}^s \sin \theta_l} \pi_s \\ \tilde{J}_{iN} = & \sin \theta_{i+1} \prod_{k=i+1}^N \sin \theta_k \pi_{i+1} + \cos \theta_{i+1} \sum_{s=i+1}^N \frac{\prod_{m=s}^N \sin \theta_m \cos \theta_s}{\prod_{l=i+1}^s \sin \theta_l} \pi_s.\end{aligned}\quad (18)$$

Both sets of generators (17) and (18) fulfil the commutation rules (5) with respect to the canonical Poisson bracket. The kinetic energy is related to the second-order Casimir of  $so_\kappa(N+1)$  through

$$2\mathcal{T} = \tilde{\mathcal{C}} = \sum_{i=1}^N \tilde{P}_i^2 + \kappa \sum_{i,j=1}^N \tilde{J}_{ij}^2 \quad (19)$$

so that any generator Poisson-commutes with  $\mathcal{T}$ . The geodesic motion is maximally superintegrable and its integrals of motion come from any function of the Lie generators.

Now the crucial problem is to find potentials  $\mathcal{U}(q)$  that can be added to  $\mathcal{T}$  in such a manner that the new Hamiltonian  $\mathcal{H} = \mathcal{T} + \mathcal{U}$  preserves the maximal superintegrability. This requires to add ‘some’ terms to ‘some’ functions of the generators in order to ensure their involutivity with respect to  $\mathcal{H}$ . By taking into account the results given in [6] for  $\mathbf{S}^2$  and  $\mathbf{H}^2$ , we propose the following generalization of the SW potential (1) to the space  $S_{[\kappa]}^N$ :

$$\begin{aligned}\mathcal{U} = & \beta_0 \frac{\sum_{s=1}^N x_s^2}{x_0^2} + \sum_{i=1}^N \frac{\beta_i}{x_i^2} \\ = & \beta_0 \sum_{i=1}^N \frac{S_\kappa^2(a_i)}{\prod_{s=1}^i C_\kappa^2(a_s)} + \sum_{i=1}^{N-1} \frac{\beta_i}{S_\kappa^2(a_i) \prod_{s=i+1}^N C_\kappa^2(a_s)} + \frac{\beta_N}{S_\kappa^2(a_N)} \\ = & \beta_0 T_\kappa^2(r) + \frac{1}{S_\kappa^2(r)} \left( \frac{\beta_1}{\cos^2 \theta_2} + \sum_{i=2}^{N-1} \frac{\beta_i}{\cos^2 \theta_{i+1} \prod_{s=2}^i \sin^2 \theta_s} + \frac{\beta_N}{\prod_{s=2}^N \sin^2 \theta_s} \right).\end{aligned}\quad (20)$$

On the sphere  $\mathbf{S}^N$  with  $\kappa > 0$ , this can be interpreted as the joint potential due to a superposition of  $N + 1$  harmonic oscillators whose centers are placed at  $N + 1$  points on  $\mathbf{S}^N$  mutually separated a quadrant (a distance  $\pi/2\sqrt{\kappa}$ , which for  $\kappa = 1$  is  $\pi/2$ ); on  $\mathbf{S}^2$  these would be placed at the three vertices of a sphere's octant [21]. Explicitly, if we take  $\kappa = 1$  and consider the polar coordinate  $r$  together with  $N$  geodesic distances  $r_i$  ( $i = 1, \dots, N$ ) such that  $x_0 = \cos r$ ,  $x_i = \cos r_i$ , the potential (20) turns out to be

$$\mathcal{U} = \beta_0 \tan^2 r + \sum_{i=1}^N \frac{\beta_i}{\cos^2 r_i} = \beta_0 \tan^2 r + \sum_{i=1}^N \beta_i \tan^2 r_i + \sum_{i=1}^N \beta_i. \quad (21)$$

The first term is  $\beta_0 \tan^2 r$ , where  $r$  is the distance from the particle and the origin  $\mathcal{O}$  along the geodesic  $l$ ; this is the spherical Higgs potential with center at  $\mathcal{O}$  where the 0-th coordinate axis  $x_0$  in the ambient space intersects the sphere. Each of the  $N$  remaining terms (apparently very different in (20)),  $\beta_i \tan^2 r_i$ , is written in terms of the spherical distance  $r_i$  to the point where the  $i$ -th coordinate axis  $x_i$  intersects the sphere. Under the contraction  $\kappa = 0$ ,  $\mathbf{S}^N \rightarrow \mathbf{E}^N$ , the first term gives rise to the ‘flat’ harmonic oscillator  $r^2 = \sum_i a_i^2$ , while the  $N$  remaining oscillators (whose centers would be now ‘at infinity’) leave the ‘centrifugal’ barriers  $\beta_i/a_i^2$  as their imprints.

Let us consider the following functions  $I_{ij}$  ( $i < j$ ;  $i, j = 0, 1, \dots, N$ ):

$$I_{ij} = (x_i \dot{x}_j - x_j \dot{x}_i)^2 + 2\beta_i \frac{x_j^2}{x_i^2} + 2\beta_j \frac{x_i^2}{x_j^2} \quad (22)$$

which are quadratic in the momenta through the square of the generators. In parallel coordinates with the phase space realization (17), they turn out to be

$$\begin{aligned} I_{0i} &= \tilde{P}_i^2 + 2\beta_0 \frac{S_\kappa^2(a_i)}{\prod_{s=1}^i C_\kappa^2(a_s)} + 2\beta_i \frac{\prod_{s=1}^i C_\kappa^2(a_s)}{S_\kappa^2(a_i)} \\ I_{ij} &= \tilde{J}_{ij}^2 + 2\beta_i \frac{S_\kappa^2(a_j)}{S_\kappa^2(a_i) \prod_{s=i+1}^j C_\kappa^2(a_s)} + 2\beta_j \frac{S_\kappa^2(a_i) \prod_{s=i+1}^j C_\kappa^2(a_s)}{S_\kappa^2(a_j)}. \end{aligned} \quad (23)$$

Likewise these can be written in geodesic polar coordinates. Hereafter we consider the Hamiltonian  $\mathcal{H} = \mathcal{T} + \mathcal{U}$  with  $\mathcal{T}$  and  $\mathcal{U}$  given in (15) and (20). Notice that the analogous property to (19) is given by

$$2\mathcal{H} = \sum_{i=1}^N I_{0i} + \kappa \sum_{i,j=1}^N I_{ij} + 2\kappa \sum_{i=1}^N \beta_i. \quad (24)$$

When  $\kappa = 0$ , the expressions (17), (23) and (24) reduce to (1)–(3). Next it can be proven that:

**Proposition 1.** *The  $N(N + 1)/2$  functions (23) are integrals of the motion for  $\mathcal{H}$ .*

Let us choose the following subsets  $Q^{(k)}$  and  $Q_{(k)}$  of  $N - 1$  integrals ( $k = 2, \dots, N$ ):

$$Q^{(k)} = \sum_{i,j=1}^k I_{ij} \quad Q_{(k)} = \sum_{i,j=N-k+1}^N I_{ij} \quad (25)$$

where  $Q^{(N)} \equiv Q_{(N)}$ . The maximal superintegrability of  $\mathcal{H}$  is characterized as follows.

**Theorem 2.** (i) The  $N$  functions  $\{Q^{(2)}, \dots, Q^{(N)}, \mathcal{H}\}$  are mutually in involution. The same property holds for the set  $\{Q_{(2)}, \dots, Q_{(N)}, \mathcal{H}\}$ .

(ii) The  $2N-1$  functions  $\{Q^{(2)}, \dots, Q^{(N-1)}, Q^{(N)} \equiv Q_{(N)}, Q_{(N-1)}, \dots, Q_{(2)}, I_{0i}, \mathcal{H}\}$  (with  $i$  fixed) are functionally independent, thus  $\mathcal{H}$  is maximally superintegrable.

The set  $Q^{(k)}$  can be associated to a sequence of orthogonal subalgebras within  $\mathfrak{h} = \mathfrak{so}(N) = \langle J_{ij} \rangle$ , the generators of which determine the terms quadratic in the momenta in the integrals  $I_{ij}$  starting ‘upwards’ from  $\langle J_{12} \rangle = \mathfrak{so}(2)$ :

$$\begin{array}{ccccccc} Q^{(2)} & \subset & Q^{(3)} & \subset & \dots & \subset & Q^{(k)} & \subset & \dots & \subset & Q^{(N-1)} & \subset & Q^{(N)} \\ \mathfrak{so}(2) & \subset & \mathfrak{so}(3) & \subset & \dots & \subset & \mathfrak{so}(k) & \subset & \dots & \subset & \mathfrak{so}(N-1) & \subset & \mathfrak{so}(N) \end{array}$$

with a similar embedding for  $Q_{(k)}$  but starting ‘backwards’ from  $\langle J_{N-1N} \rangle = \mathfrak{so}(2)$ . In fact, the SW system on  $\mathbf{E}^N$  can be constructed from a coalgebra approach [22] by means of  $N$  copies of  $\mathfrak{sl}(2, \mathbb{R})$ . When  $\kappa = 0$ , each  $Q^{(k)}$  (or  $Q_{(k)}$ ) is related to the  $k$ -th order coproduct of the Casimir of  $\mathfrak{sl}(2, \mathbb{R})$  [23]. In this sense, the results of theorem 2 show that the set of integrals ensuring the maximal superintegrability of the ‘flat’ SW system coming from a  $\mathfrak{sl}(2, \mathbb{R})$ -coalgebra also holds for any curvature.

Explicit proofs and details for this algebraic construction—which could also be applied to the ND Kepler potential—will be given elsewhere. Furthermore, the consideration of a second contraction parameter  $\kappa_2$ , that determines the signature of the metric [17, 18], would allow one to obtain superintegrable systems on different spacetimes.

## Acknowledgments

This work was partially supported by the Ministerio de Ciencia y Tecnología, Spain (Projects BFM2000-1055 and BFM2002-03773). The authors are also grateful to G.S. Pogosyan for helpful discussions

## References

- [1] Evans N W 1990 *Phys. Rev. A* **41** 5666
- [2] Kalnins E G, Williams G C, Miller W Jr and Pogosyan G S 1999 *J. Math. Phys.* **40** 708
- [3] Grosche C, Pogosyan G S and Sissakian A N 1995 *Fortschr. Phys.* **43** 523
- [4] Kalnins E G, Miller W Jr and Pogosyan G S 1997 *J. Math. Phys.* **38** 5416
- [5] Grosche C, Pogosyan G S and Sissakian A N 1997 *Phys. Part. Nuclei* **28** 486
- [6] Rañada M F and Santander M 1999 *J. Math. Phys.* **40** 5026
- [7] Kalnins E G, Miller W Jr and Pogosyan G S 2000 *J. Phys. A: Math. Gen.* **33** 6791

- [8] Kalnins E G, Kress J M, Pogosyan G S and Miller W Jr 2001 *J. Phys. A: Math. Gen.* **34** 4705
- [9] Fris J, Mandrosov V, Smorodinsky Ya A, Uhler M and Winternitz P 1965 *Phys. Lett.* **16** 354
- [10] Evans N W 1990 *Phys. Lett. A* **147** 483
- [11] Evans N W 1991 *J. Math. Phys.* **32** 3369
- [12] Grosche C, Pogosyan G S and Sissakian A N 1995 *Fortschr. Phys.* **43** 453
- [13] Schrödinger E 1940 *Proc. R. Ir. Acad. A* **46** 9
- [14] Higgs P W 1979 *J. Phys. A: Math. Gen.* **12** 309
- [15] Leemon H I 1979 *J. Phys. A: Math. Gen.* **12** 489
- [16] Kalnins E G, Miller W Jr and Pogosyan G S 2002 *Phys. Atom. Nucl.* **65** 1086
- [17] Herranz F J, Ortega R and Santander M 2000 *J. Phys. A: Math. Gen.* **33** 4525
- [18] Herranz F J and Santander M 2002 *J. Phys. A: Math. Gen.* **35** 6601
- [19] Rañada M F and Santander M 2002 *J. Math. Phys.* **43** 431
- [20] Izmet'sev A A, Pogosyan G S, Sissakian A N and Winternitz P 1999 *J. Math. Phys.* **40** 1549
- [21] Rañada M F, Santander M and Sanz-Gil T 2002 Superintegrable potentials and the superposition of Higgs oscillators on the sphere  $S^2$  (Warszawa: Banach Center Publications) to be published
- [22] Ballesteros A and Ragnisco O 1998 *J. Phys. A: Math. Gen.* **31** 3791
- [23] Ballesteros A and Herranz F J 1999 *J. Phys. A: Math. Gen.* **32** 8851