## Topological Phenomena in the Real Periodic Sine-Gordon Theory

P.G. Grinevich<sup>1</sup>
Landau Institute for Theoretical Physics
of the Russian Academy of Sciences,
Moscow 117940 Kosygin Street 2,
emailpgg@landau.ac.ru

S.P.Novikov<sup>2</sup>
IPST, University of Maryland-College Park,
M.D. 20742-2431
and Landau Institute for Theoretical Physics, Moscow,
e-mail novikov@ipst.umd.edu

1 Introduction. A lgebro-G eom etrical Solutions and the R eality R eductions. The topological charge.

The algebro-geom etrical (or nite-gap) solutions play a very important role in the modern theory of soliton equations with periodic and quasiperiodic boundary conditions. The main property characterizing these solutions is the following: the wave function of the auxiliary linear operator is merom orphic in spectral parameter on the nite part of an algebraic Riemann surface. It allows to apply the powerful tools of classical algebraic geometry, and to write explicit representations for solutions and wave functions in terms of Riemann—functions of several variables. Usually the algebro-geometrical solutions are dense in the space of the periodic ones, and if the nonlinearity tends to zero, they degenerate to nite Fourier sums. Therefore their role in the soliton theory is rather similar to the role of nite Fourier series for the linear PDE's.

The algebro-geom etrical solutions of solition equations were st introduced in the paper [1] in 1974 dedicated to the integration of the K orteveg-

 $<sup>^{1}</sup>$ The authorwas partially supported by the RFBR G rant N o 01-01-00874-A. Numerical simulations were partially performed on a computer, donated to P.G. Grinevich by the Humboldt Foundation.

<sup>&</sup>lt;sup>2</sup>The author was partially supported by NSF Grant DMS 0072700

de V ries (K dV) equation with the periodic boundary conditions. Real non-singular periodic algebro-geom etrical K dV solutions can be characterised by the following property: the number of gaps (forbidden bands or zones) in the spectrum of the auxiliary Schrodinger operator acting in the H ilbert Space  $L_2$  (R) is nite. This characterization is not valid for other systems (for example the spectrum of non-selfad joint auxiliary operator for the self-focusing N onlinear Schrodinger equation (N LS) has no gaps for any regular potential), but neverthereless the algebro-geom etrical solutions are called nite-gap.

The complete theory of algebro-geom etrical K dV solutions involving the spectral theory of periodic (quasiperiodic) nite-gap Schrodinger operators on the line and its uni cation with algebraic geometry and analysis on the Riem ann surfaces, tim e dynamics and algebro-geometric ham iltonian aspects was constructed in the works [2, 3, 4, 5, 6, 7] (see in the survey article [8]) and in the works [9, 10]. Complex nite-gap solutions of the NLS and Sine-Gordon (SG) equations were constructed in [11, 12]. Finite-gap solutions of the 2+1 K adom tsev-Petviashvili (KP) eqation were consturcted in [13] on a basis of a purely algebraic formulation of the nite-gap approach developed in this paper. The algebro-geom etric spectral theory of the stationary periodic 2D Schrödinger operator was developed in the works [14, 15]. In these papers it was shown, that the proper analog of the nite-gap constraint in the 2dim ensional case is the following: the restriction of the Bloch variety to one energy level is algebraic, and the spectral data is \collected" at this energy level only. Such potentials can be treated as \integrable" at one energy. It means, that the wave functions for this energy and arbitrary complex quaism om enta can be written explicitly in terms of Riemann -functions, but the inform ation about the spectral properties at other energies is very restricted.

The algebro-geom etrical approach allows to obtain explicit compact formulas for solutions. But the —fucntions of several variables are very complicated from the analytic point of view; therefore the existence of such formulas does not lead sometimes to the simple solution of the problem of selecting physically orgeom etrically meaningful solutions among generic complex ones. For example, it very easy to describe reality conditions for the KdV, defocusing NLS or Sinh-Gordon equation

$$\frac{e^2u}{a \ a} = 4 \sinh u ( ; ):$$

(in all these cases the auxiliary linear operators are self-adjoint). However,

the reality reductions for the self-focusing NLS or the Sine-G ordon equation found in [16] are rather nontrivial. It was shown in [15] that analogous reductions are responsible for the absence of magnetic eld it the theory of the 2D Schrodinger operator. A good review of reality reductions can be found if [17].

The Sine-Gordon equation

$$u_{tt} u_{xx} + \sin u (x;t) = 0$$
 (1)

derived in the X IX century geometry in the light-cone representation

$$u = 4 \sin u; \quad u = u(;);$$
 (2)

w here

$$x = 2(+); t = 2(); 0 = 20x + 20t; 0 = 20x - 20t; (3)$$

is one the most important systems of the soliton theory. One it's distinguished properties is the following. Usually if we x the spectral curve (or equivalently all local conservation laws), then the level set is a connected Jacoby torus. It is true for K dV, defocusing and self-focusing NLS, the both real reductions of the KP equation. But in the Sine-G ordon case the level set is a union of  $2^m$  real Jacoby tori, where m denotes the number of pairs of negative branch points. As a corollary, the result of averaging a generic functional depends not only on the spectral curve, but also on the component. This property is also essential if we calculate the action-angle variables (this problem was studied in [18, 19]).

The averaged densities of conservation laws arose naturally as ham iltonians of the W hitham equations. These equations describe the evolution of parametrs of the asymptotic solutions in the form of a slow ly varying N - phase wavetrains. The W hitham equations for the Sine-G ordon were derived in the paper [20], but the connectivity components of the level sets were not discussed. Let us point out, that the W hitham equations are hyperbolic only in the \stable" case, when all branch point are negative and the number of components is always greater than 1.

It is possible to extract from the calculations of [20], that for all classical local conservation laws except the topological charge the result of averaging is the same for all components, but this property was never pointed out explicitly. In Section 4 we give a simple explanation of it.

Remark. The number of realtoriwas calculated in [16], their characterization in terms of the complex Jacoby torus was obtained in [21], see also [22]. An elementary description of these toriwas obtained by the authors in [23, 24], see Section 2 below.

A nother interesting property of the Sine-G ordon equation is the following. In the theory of constant negative curvature surfaces  $u(\cdot;\cdot)$  denotes the angle between asymptotic lines. In the theory of Josephson jucntions u has the meaning of a phase. In both situations it is dened up to an additive constant c=2 n. Therefore it is natural to call a solution u(x;t) space-periodic w if the period u if expfiu u is called u opological charge, the real number u is called u opological charge, the real number u is called u opological charge. These quantities are the u ost stable conservation laws surviving all real periodic nonintegrable perturbations. The density of topological charge can be naturally extended to all nonsingular quasiperiodic solutions using the form u la

$$n = \lim_{T \mid 1} [u(x + T) u(x)] = 2 T$$
:

One of the most natural problems is the following: How to calculate the topological charge of real periodic SG solutions in terms of the Spectral Data: the Riemann surface and the divisor? It turned out, that nobody succeeded to extract the answer from the explicit description of the real components described in terms of the Jacoby torus and —functional formulas, found in the works [21, 22]. An attempt has been made almost 20 years ago in the work [18] to solve this problem. For this purpose the so-called \A lgebro-Topological" approach was developed in [18], and an interesting formula has been proposed. However, as it was pointed out in [19], the idea of the proof presented in [18] works only if the spectral curve is su ciently close to a degenerate one. A complete solution using a new development of the main idea of the work [18] was obtained only recently by the authors in the papers [23] (for the so-called stable curves) and [24] for generic ones. Our proof is based on a new elective description of the real components in terms of divisors. We present a summary of these results in Section 3.

Remark. Formally we can de ne topological charge for any non-singular complex solution. But if the solution has poles, this quantity became ill-de ned, and no natural non-singularity condition in terms of the spectral data for complex potentials is known. It explains why only real solutions are discussed.

#### 2 Finite-gap Sine-Gordon solutions.

The \soliton-type" solutions of Sine-G ordon were found already in the X IX century using substitution discovered by Bianchi and S. Lie. Now this method is called \Backlund transform ation".

In the early 1970's G. Lamb found out (see [25]), that the the Sine-Gordon equation is analogous to KdV in the following sence: the so-called \inverse scattering method" can be applied to it. The modern approach was started in the work [26]. The critical point of this approach is the following zerocurvature representation for SG

$$_{x} = \frac{1}{4} (U + V) ; \quad _{t} = \frac{1}{4} (U \quad V) ; \quad (4)$$

w here

$$U = U (;x;t) = \begin{array}{ccc} i(u_x + u_t) & 1 & & \\ & i(u_x + u_t) & ; & & \\ V = V (;x;t) = & 0 & \frac{1}{2}e^{iu} & \\ e^{iu} & 0 : & & \\ \end{array}$$
(5)

To construct generic complex nite-gap SG solutions assume, that we have the following spectral data:

- 1. A nonsingular hyperelliptic Riemann surface  $[^2 = R()]$ , where R() =  $(E_k)$ , such that  $E_0 = 0$  and  $E_i \in E_j$ ; i; j = 0; 1; ...; 2g. It has exactly 2g + 2 branch points  $E_0 = 0$ ,  $E_1$ , ...,  $E_{2q}$ , 1; the genus of is equal to g. A point 2 is by de nition a pair of complex numbers = (;) such, that  $^2 = R()$ .
- 2. A divisor D of degree g, i.e. set (or form al sum ) of g points D = 1 + :::+ g.

For generic data , D there exists an unique two-component \Baker-A khiezer" vector-function (;x;t) such that

1. For xed (x;t) the function (;x;t) is merom orphic in the variable outside the points 0, 1 and has at most 1-st order poles at the divisor points k, k = 1; ...; q.

2. (;x;t) has essential singularities at the points 0, 1 with the following asymptotics:

$$(;x;t) = p^{1+o(1)} + o(1) = e^{i^{p} - x + t}$$
 as ! 1; (7)

$$(;x;t) = p_{1}^{-1}(x;t) + o(1)$$
  $e^{\frac{p^{\frac{1}{2}}x-t}{4}}$  as ! 0; (8)

where  $_{1}(x;t)$ ,  $_{2}(x;t)$  are some functions of the variables x;t.

P Denote the divisor of zeroes of the st component  $_1$  by D (x;t) =  $_j$   $_j$  (x;t). The vector-function (;x;t) satisfies to the zero-curvature equations (4) with potential given by the form ula

$$u(x;t) = i \ln \frac{2(x;t)}{1(x;t)};$$
 (9)

and the function u(x;t) solves the SG equation (2). In terms of the divisor of zeroes D (x;t) the potenital u(x;t) can be easily written explicitly:

$$e^{iu (x,t)} = \frac{\int_{j=1}^{j=1} x}{\sum_{j=1}^{j=1} E_{j}}$$
(10)

The x and t dynam ics of the divisor D (x;t) can be described it terms of D ubrovin equations (see [24]). From (10) it follows, that the potential u(x;t) is singular at the point  $(x_0;t_0)$  if and only if one of the points  $k(x_0;t_0)$  coincides with 0 or 1 . Collisions of 2 or more divisor points result in singularities in the solutions of D ubrovin equations, but all symmetric combinations of the divisor points including the potential u(x;t) remain non-singular.

R em ark. The rst component  $_1$  of the Baker-Akhiezer vector-function is a partial case of the general scalar two-point Baker-Akhiezer functions satisfying to the second order linear Schrodinger equation  $L_{1-1}=0$  invented in the work [14]. Here we have

$$L_1 = \frac{e^2}{e^2} + A(;)\frac{e}{e} + W(;);$$

A=0 log  $_1$ . In this special case our surface is hyperelliptic, the selected points coincide with 0;1 and corresponding local parameters are chosen as  $_1;1=1$ . The second function  $_2$  satisfies to similar equation  $L_2$   $_2=0$  where the operator  $L_2$  is obtained from  $L_1$  by the so-called Laplace Transform ations and vice versa. It is known that cyclic Laplace Chains of the length 2 lead to the SG equation in the general complex case (see the work [27]).

Lem m a 1 Assume, that the spectral data (;D), D =  $_1$  + :::;  $_g$  satisfy the following reality constraints:

1. The set of branch points of the R iem ann surface contains real points or complex conjugate pairs of points only, and all real branch points are nonpositive. Without loss of generality we may assume, that the rst 2m+1 of them  $E_0=0$ ,  $E_1$ ,  $E_2$ ,...,  $E_{2m}$  are real and  $0>E_1>E_2>\dots$ :: >  $E_{2m}$ , and for k>m we have  $E_{2k}=\overline{E_{2k-1}}$ , Im  $E_{2k}$  60. Then the spectral curve is real and admits a  $Z_2$   $Z_2$ -group of involutions generated by the standard holomorphic involution

$$: ! : (; ) = (; );$$
 (11)

transposing the sheets, and by the antiholom orphic involution

$$: ! : (;) = (;);$$
 (12)

(com plex con jugation).

2. There exists a merom ophic dierential with two simple poles at the points 0;1 such, that the zeroes of are located at the points D + D. We shall call such divisors admissible. Without loss of generality we can normalize to be real: = .

Then the potential u(x;t) is real and non-singular.

Remark The constraints on the spectral curve we found in [12]. The characterisation of admissible divisors and the proof of non-singularity were obatined in the work [16].

It is easy to check (see, for example [24])

Lem m a 2 If D is adm issible divisor, then the divisor of zeroes D  $(x;t) = {}_{1}(x;t) + :::+ {}_{q}(x;t)$  is adm issible for all x;t

Assume, that we have a real merom orphic dierential with exactly 2 simple poles at the points 0;1. We can associate an admissible divisor D to it if and only if 2g zeroes of can be can be presented as an union of 2 subsets D and D<sub>1</sub> such, that D<sub>1</sub> = D. It is possible if and only if all real roots of have even multiplicity.

De nition 1 Assume that the spectral curve satisfy the reality condinitions formulated above. A realmerom orphic dierential on with exactly 2 simple poles at the points 0;1 is called admissible if the multiplicity of all real roots is even.

W ithout loss of generality we may assume that the residued of at the points 0 and 1 are equal to +1 and 1 respectively. Then we have

$$= 1 \frac{P_{g 1}()}{R()^{1=2}} \frac{d}{2}; \qquad (13)$$

where  $P_{g-1}$  ( ) is a polynom ial of degree at most g-1 with real coe cients. Let us call the polynom ial  $P_{g-1}$  ( ) adm is sible if is an adm is sible dierential.

Denote the map associating an admissible polynomial with an admissible divisor by :D ! P $_{g-1}$ (). The inverse map  $^{-1}$  is multivalued. If P $_{g-1}$ () is an admissible polynomial, then the equation

$$= P_{g 1}()$$
  
R() =  $^{2}$ ; (14)

has 2g + 1 roots. One of them is the point (0;0), other 2g roots form g pairs. To de ne an admissible divisor we have to choose one point from each pair: therefore we have at most  $2^g$  possibilities depending on the number of real points. In the generic case we have no real roots, so the number is equal to  $2^g$  in this case.

The description of connected components suggested by the authors in [23, 24] is based on the following simple observation:

Consider a pair of functions of the real variable:

$$f() = \frac{p}{R()}$$
: (15)

We assume, that the functions f () are defined only for such 6 0 that R () 0, i.e. these functions are real-valued. Let us draw the graphs of these functions  $y = f_+$  (), y = f (), and ll in the following domains:

< 
$$0; y^2 < \frac{R()}{2};$$
  
>  $0; y^2 > \frac{R()}{2}:$  (16)




Fig 1.

Lem m a 3 The polynom ial  $P_{g-1}($  ) is admissible if and only if the graph of  $P_{g-1}($  ) does not cross the black open domains (but it can touch their boundaries).

Now we can associate with each admissible polynomial a Topological Type, i.e. a collection of m numbers  $s_k = 1$ , k = 1;::; m de ned by the following rule:

$$s_k = 1 \text{ if } P_{g-1}()$$
  $f_+() \text{ as } E_{2k}$   $E_{2k-1}$ ,  $s_k = 1 \text{ if } P_{g-1}()$   $f_-() \text{ as } E_{2k}$   $E_{2k-1}$ 

U sing sim ple algebraic estimates (see [24]) it is easy to check:

Lem m a 4 Each topological type is presented by the convex subset in the space of polynom ials  $P_{g-1}$ . Each of these  $2^m$  connected components is non-empty and depends continuously on the branch points  $E_{1}, \ldots, E_{2g}$ .

The x and t dynam ics of adm issible divisors m ay be rather non-trivial (a number of numeric experiments are discussed in [22]). In contrast with the self-a joint case (K dV, defocusing NLS, sinh-G ordon) the adm issible position of a given divisor point depends on the position of other ones (the adm issibility constraints are non-local). The trajectory of a single divisor point m ay be not well-de ned because divisor points m ay collide and a small variation of parameters near the collision point results in a bifurcation of trajectories. If the potential u (x;t) is periodic with the period T, the corresponding divisor is generically periodic only modulo permutations. But the following properties can be easily proved using the characterization formulated above.

Lem m a 5 P rojections of the points of adm issible divisors to the -plane could not lie in the segments  $\mathbb{E}_1$ ;  $\mathbb{E}_0$ ],  $\mathbb{E}_3$ ;  $\mathbb{E}_2$ ],..., (1;  $\mathbb{E}_{2m}$ ]. Moreover, it is possible to choose an open neighborhood of these segments such, that the projections of these points could not lie in it.

Lem m a 6 Assume, that the projection of a point  $_s = (_s;_s)$  of an adm issible divisor to the -plane lie in the segments  $\mathbb{E}_{2k}$ ;  $\mathbb{E}_{2k}$   $_1$ ]; then  $_s < 0$  if  $s_k = 1$  and  $_s > 0$  if  $s_k = 1$ .

## 3 The topological charge of real solutions

It is well-known (see, for example [21, 22]) that after the Abel transform the x-dynamics of Sine-G ordon corresponds to the motion along the straight line

$$X' = X_0 + UX; (17)$$

in the Jacoby torus, where  $\tilde{\textbf{U}}$  denotes the norm alized vector of b-periods of the quasim omentum dierential dp.

$$U^{k} = \frac{1}{2} \operatorname{dp}; \tag{18}$$

Let us recall, that by de nition dp is a merom orphic di erential with zero a-periods and two second order poles at 0, 1 such, that

$$\frac{1}{8} + 0 = 0 \text{ das } ! 1;$$

$$dp = 0 \text{ das } ! 0:$$

$$\frac{1}{8} + 0 = 0 \text{ das } ! 0:$$

The real part of the Jacoby torus is isom or phic to the factor  $R^n = Z^n$ , therefore we can apply the following simple analytic lem m a:

Lem m a 7 Letu (X), X 2  $R^n$  be a sm ooth function in  $R^n$  such, that  $\exp(iu(X))$  is single-valued on the torus  $R^n = Z^n$ , i.e.  $\exp(iu(X + N)) = \exp(iu(X))$  for any integer vector N. Denote by u(x) restriction of u(X) to the straight line  $X = X_0 + x$  V. Then the density of topological charge  $n = \lim_{T \ge 1} [u(x + T) u(x)] = 2$  T is well-de ned; it does not depend on the point  $X_0$  and can be expressed by the following form u(x):

$$n = \sum_{k=1}^{X^n} n_k U^k; \qquad (20)$$

where  $U = (U^1; U^2; ...; U^n)$ , and  $n_k$  are topological charges along the basic cycles  $A_k$ , k = 1; ...; n:

$$u(X^{1};X^{2};...;X^{k} + 1;...;X^{n}) \quad u(X^{1};X^{2};...;X^{k};...;X^{n}) = 2 n_{k};$$
(21)

The main step is the calculation of the the charges along the basic cycles  $A_k$ . To obtain a convenient answer the choice of canonical basis is very important. For stable surfaces (surfaces without complex branch points) the proper system of a-cycles was constructed in [18].





Fig 2.

We work with the following system of cuts  $[E_1; E_0]$ ,  $[E_3; E_2]$ ,...,  $(1; E_{2g}]$ . Let us denote the sheet containing the line 2R, > 0 > 0 by  $G_+$  and the second sheet by G (see Fig 2).

For the generic real curves with 2m negative branch points  $E_1; \ldots; E_{2m}$  and g m complex adjoint pairs  $E_{2j-1}; E_{2j} = E_{2j-1}$ ,  $j = m+1; \ldots; g$  we choose rst m a-cycles exactly as in the stable case. The last g m a-cycles should be chosen as coverings on over the path on the -plane connecting the points  $E_{2j-1}$  and  $E_{2j}$  for j > m. All these path should not meet each other. All of them should cross positive part of the line > 0 in one point  $_j$  (i.e. in two points of R iem ann surface) such that  $_{m+1} < _{m+2} < \ldots < _{g}$ . This basis depends on the order of the complex conjugate pairs only (see Fig 3).



We assume also that the cycles  $b_1$ ,  $b_2$ , ...,  $b_m$  are the ovals lying over the segments  $E_2$ ;  $E_1$ ,  $E_4$ ;  $E_3$ ,  $E_{2m}$ ;  $E_{2m-1}$ ] respectively.

For the complex conjugate pairs of branch points  $E_{2j}$ ;  $E_{2j-1}$ ; j > m; we make the cuts connecting each of these points with if or if such that the cuts do not intersect the real line. As before we denote the sheet containing the line  $2 R_* > 0 > 0$  by  $G_+$  and the second sheet by  $G_-$ . The orientation of the cycles  $b_{1}$ , l = 1; :::; modified with the standard orientation

of the negative sem i-line R  $\,$  at the sheet G  $_{+}$  ; they are opposite at the sheet G  $\,$  .

Consider a basic cycle  $A_k$  on the real component of Jacoby torus, represented by the closed curve  $B_k$  on this torus. The image of this cycle in the surface (i.e. image of it under the inverse Abelmap) is a closed oriented curve  $C_k$  (may be consisting from several connected components) homological to the cycle  $a_k$  2  $H_1$  (; Z). From Lemma 5 it follows, that this curve does not touch the closed segments on the real line [1;  $E_{2m}$ ]; :::;  $E_3$ ;  $E_2$ ];  $E_1$ ; 0], and if it crosses the negative semi-line at the inteval  $E_{21}$ ;  $E_{21}$ , then it happens in the sheet  $G_+$  if (1)  $E_{2m}$  in the sheet  $E_{2m}$  if (1)  $E_{2m}$  is given by

$$n_k = C_k \quad R ; \qquad (22)$$

where  $\mathcal{C}_k$  is the projection of  $C_k$  to the -plane, and denotes the intersection index of curves on the -plane. Taking into account that any intercetion with the negative sem i-line is the intersection with one of the b-cycles we obtain:

Lem m a 8 The topological charge along the cycle  $A_k$  on the real torus in the Jacobian variety is equal to the intersection index of the corresponding curve  $C_k$  with the Counting Cycle of the real component on the Riemann surface

$$n_k = C_k$$

$$(1)^{1} s_1 b_1 :$$

$$(23)$$

Let us mention that the cycle C  $_k$  is homotopic to  $a_k$  on  $% A_k$  . Therefore C  $_k$   $% A_k=0$  , and

$$n_k = (1)^{k-1} s_k$$
 for  $k = m$   
 $n_k = 0$  for  $k > m$ : (24)

Combining (20) with (24), we obtain our main result:

Theorem 1 For any real solution the density of topological charge along the variable x is given by the form u la

$$n = \frac{1}{2} \sum_{k=1}^{X^{n}} (1)^{k-1} s_{k} U^{k};$$
 (25)

where the components  $U^k$  of the vector  $\tilde{U}$  are dened by (18).

# 4 A veraging of the local conservation laws and topological types

For a generic functional

$$F [u] = f(e^{iu}; e^{-iu}; u_x; u_{xx}; ...; u_t; u_{tx}; u_{txx}; ...;) dx$$
 (26)

the result of averaging over a real com ponet of the Jacoby torus depends on the topological type. For example by avearging of  $u_x$  we get the topological charge. But if we average the Sine-G ordon H am iltonian

H [u] = 
$$\frac{u_t^2}{2} + \frac{u_x^2}{2} + (1 \cos(u)) dx$$
 (27)

the result is the same for all components. Therefore it is natural to formulate the following:

P roblem. How to characterize the functional such, that the averaging does not depend on the topological type?

A complete classication seems to be a complicated problem. But it is rather easy to formulate a natural su cient condition.

If  $f(e^{iu}; e^{-iu}; u_x; u_{xx}; \dots; u_t; u_{tx}; u_{txx}; \dots)$  is a polynom ial of its arguments then it can be written as a symmetric rational function of the divisor coordinates  $f = f'(\ _1(x;t); \dots; \ _q(x;t); \ _1(x;t); \dots; \ _q(x;t))$ .

Lem m a 9 Consider the following form on the  $\left| \frac{z}{z} \right|$ 

If we x all divisor points except  $_1$  we obtain a merom orphic diemtial on with poles only at 0 and 1 . Assume that the residues of this dierential are equal to 0 identically in  $_2, \ldots, _g$ . Then for a given spectral curve the averaging does not depend on the topological type

Proof. Following [20] we can calculate the averaging of F [u] using the formula:

$$\overline{F[u]} = C \qquad \cdots \qquad W_f \qquad (29)$$

$$a_1 \qquad a_g$$

where C is the normalization constant, and all cycles  $a_k$  have the proper topological type. The last condition was not discussed in [20]. The deformations of the integration cycles do not a ect the integral if we do not cross 0. If the condition of Lemma 9 is full led, then the we can move the integration path through 0 and \reach" any topological type without changing the integral.

As a corollary we im mediately get the following statement:

Theorem 2 Assume that the functional F[u] is invariant with respect to the following symmetries: f(u): f

Im portant exam ple. Consider the densities of \higher SG ham iltonians" de ned as odd expansion coe cients of the function

$$\frac{x (;x;t)}{(;x;t)} \tag{30}$$

at the points 0 and 1 . Direct calculation shows, that

$$\frac{x}{1} = \frac{1}{4} - x + \frac{1}{4}$$
 (31)

w here

$$-- = i \frac{+ Q ()}{( (x;t)) ::: ( (x;t))};$$
 (32)

$$- = i \frac{[ + Q ()][ _{1}(x;t); :::( _{g}(x;t))]}{( _{1}(x;t)) :::( _{g}(x;t))^{p} \frac{\overline{E_{1} :::E_{2g}}}{\overline{E_{1} :::E_{2g}}};$$
 (33)

Q (), Q () are polynomials of degree g 1 determined by the following conditions:

$$Q (_{k}(x;t)) = _{k}(x;t); Q (_{k}(x;t)) = \frac{_{k}(x;t)}{_{k}(x;t)}:$$
(34)

We see that the odd part of  $x^{-1}$  depends on  $x^{-1}$  but not  $x^{-1}$ . Therefore the averaging of the \higher SG ham iltonians" does not depend on the topological type.

#### R eferences

- [1] Novikov, SP. The periodic problem for the Korteweg-de Vries equation. I. Funkts. Anal. iego Pril. 8 (1974), no. 3, 54 (66.; English translation: Funct. Anal. Appl. 8 (1974), 236 (246.
- [2] Dubrovin, B.A.; Novikov, S.P. Periodic and conditionally periodic analogs of the many-soliton solutions of the Korteweg-de Vries equation. Zh. Eksper. Teoret. Fiz. 67 (1974), no. 6, 2131 (2144; English translation: Sov. Phys-JETP 40 (1974), 1058 (1063.
- [3] Dubrovin, B.A.; Novikov, S.P.A periodicity problem for the Korteweg-de Vries and Sturm-Liouville equations. Their connections with algebraic geometry. Dokl. Akad. Nauk SSSR 219 (1974), no. 3, 531 (534; English translation: Sov. Math. Dokl. 15 (1974), 1597 (1601.
- [4] Dubrovin, B A. Periodic problem for the Kortweg-de Vries equation in the class of nite-band potentials. Funkts. Anal. i ego Pril. 9 (1975), no. 3, 41 (51; English translation: Funct. Anal. Appl. 9 (1975), 215 (223.
- [5] Its, A.R.; Matveev, V.B. Hilloperators with nitely many gaps. Funkts. Anal. i ego Pril. 9 (1975), no. 1, 69{70; English translation: Funct. Anal. Appl. 9 (1975), 65{66.
- [6] Lax, P.D. Periodic solutions of the KdV equation. Comm. Pure Appl. Math. 28 1975, 141 (188.
- [7] McKean, HP.; van Moerbeke, P. The spectrum of Hill's equation. Invent. Math. 30 (1975), no. 3, 217{274.
- [8] Dubrovin, B A; M atveev, V B; N ovikov, S P. N on-linear equations of K orteweg-de V ries type, nite-zone linear operators and A belian varieties. U spekhi M at. N auk 31 (1976), no. 1, 55{136; English translation: Russ. M ath. Surveys 31 (1976), no. 1, 59{146.
- [9] Flaschka, H.; M. cLaughlin, D. W. Canonically conjugate variables for the Korteweg-de Viries equation and the Toda lattice with periodic boundary conditions. Progr. Theoret. Phys. 55 (1976), no. 2, 438 (456.
- [10] Novikov, S.P.; Veselov, A.P.On Poisson brackets compatible with algebraic geometry and Korteweg-de Vries dynamics on the set of nite-zone

- potentials. Dokl. A kad. Nauk SSSR 266 (1982), no. 3, 533 (537; English translation: Sov. Math. Dokl. 26 (1982), 357 (362.
- [11] K ozel, V A.; K otlyarov, V P.A lm ost periodic solutions of the equation  $u_{\rm tt} = u_{\rm xx} + \sin u = 0$ . (Russian) Dokl. A kad. N auk U krain. SSR Ser. A, (1976), no. 10, 878 [881.
- [12] Its, A.R.; Kotlyarov, V.P. Explicit formulas for solutions of the Nonlinear Schrodinger equation. (Russian) Dokl. Akad. Nauk Ukrain. SSR Ser. A, (1976), no. 11, 965{968;
- [13] K richever, IM. An algebraic-geometric construction of the Zakharov-Shabat equations and their periodic solutions. Dokl. A kad. Nauk SSSR 227 (1976), no. 2, 291 (294; English translation: Sov. Math. Dokl. 17 (1976), 394 (397.
- [14] Dubrovin, B A.; Krichever, IM.; Novikov, S.P. The Schrodinger equation in a periodic eld and Riemann surfaces. Dokl. Akad. Nauk SSSR 229 (1976), no. 1, 15{18; English translation: Sov. Math. Dokl. 17 (1976), 947{951.
- [15] Novikov, S.P.; Veselov, A.P. Finite-zone, two-dimensional Schrodinger operators. Potential operators. Dokl. Akad. Nauk SSSR 279 (1984), no. 4, 784{788. English translation: Sov. Math. Dokl. 30 (1984), 705{708; Finite-zone, two-dimensional potential Schrodinger operators. Explicit formulas and evolution equations. Dokl. Akad. Nauk SSSR 279 (1984), no. 1, 20{24; English translation: Sov. Math. Dokl., 30 (1984), 588{591.
- [16] Cherednik, I.V. Reality conditions in \ nite-zone" integration. Dokl. A kad. Nauk SSSR 252 (1980), no. 5, 1104{1108; English translation: Sov. Phys. Dokl. 25 (1980), 450{452.
- [17] Natanzon, S.M. Moduli of real algebraic surfaces, and their superanalogues. Dierentials, spinors and Jacobians of real curves. Uspekhi Mat. Nauk 54 (1999), no. 6, 3{60; English translation: Russ. Math. Surveys 54 (1999), no. 6, 1091{1147.
- [18] Dubrovin, B A ; Novikov, S.P. A lgebro-geometrical Poisson brackets for real nite-zone solutions of the Sine-Gordon equation and the nonlinear

- Schrodinger equation. Dokl. A kad. Nauk SSSR 267 (1982), no. 6, 1295 (1300; English translation: Sov. Math. Dokl. 26 (1982), no. 3, 760 (765.
- [19] Novikov, S.P.A lgebrotopological approach to the reality problem s.Real action variables in the theory of nite-zone solutions of the Sine-Gordon equation. Zap. Nauchn. Sem. LOM I: Di erential geometry, Lie groups and mechanics. VI.133 (1984), 177{196; English Translation: J. Soviet Math. 31 (1985), no. 6, 3373{3387
- [20] Ercolani, N.; Forest M. G.; M. CLaughlin D. W.; M. ontgom ery R. Ham iltonian structure for them odulation equations of a Sine-Gordon wavetrain. Duke M. athematical Journal 55 (1987), no 4, 949 (983.
- [21] Dubrovin, B A.; Natanzon, S M. Real two-zone solutions of the sine-Gordon equation. Funkts. Anal. i ego Pril. 16 (1982), no. 1, 27{43; English translation: Funct. Anal. Appl. 16 (1982), 21{33.
- [22] Ercolani, N.M.; Forest, M.G. The geometry of real Sine-Gordon wave-trains. Comm. Math. Phys. 99 (1985), no. 1, 1 (49.
- [23] Grinevich, P.G.; Novikov, S.P. Real nite-gap Sine-Gordon solutions: a formula for topological charge. (in Russian) Uspekhi. Mat. Nauk 56 (2001), no. 5, 181{182.; English translation: Russ. Math. Surveys 56 (2001), no. 5, 980{981.
- [24] Grinevich, P.G.; Novikov, S.P. Topological Charge of the real periodic nite-gap Sine-Gordon solutions. To appear in Commun. Pure. Appl. Math..
- [25] Lamb, G. L. Analytical description of ultrashort optical pulse propagation in a resonant medium. Rev. Mod. Phys. 43 (1971), 99{124.
- [26] Ablow itz, M. J.; Kaup, D. J.; Newell, A. S.; Segur, H. Method for solving the Sine-Gordon equation. Phys. Rev. Lett. 30 (1973), no. 25.1262 {1264.
- [27] Novikov, S.P.; Veselov, A.P. Exactly solvable two-dimensional Schrödinger operators and Laplace transformations. In: Solitons, geometry, and topology: on the crossroad. Am er. Math. Soc. Transl. Ser. 2, 179 (1997), 109{132.