

INTEGRALS INVOLVING FOUR MACDONALD FUNCTIONS AND THEIR RELATION TO $7(3)=2$

Cyril Furtlehner¹ and Stephane Ouvry²

¹ INRIA, Preval, Domaine de Voluceau-Rocquencourt, B.P. 105, 78153 Le Chesnay Cedex,
France

² LPTMS, Bat 100, Université Paris Sud, 91405 Orsay, France

A family of multiple integrals over four variables is rewritten in terms of a family of simple integrals involving the product of four modified Bessel (Macdonald functions). The latter are shown to be related to $7(3)=2$. A generalization to $2n$ integration variables is given which yields only at odd arguments.

1 Introduction

Consider

$$I_{1100}(\lambda) = \int_0^\infty d_1 \int_0^\infty d_2 \int_0^\infty d_3 \int_0^\infty d_4 \frac{(a+c)(b+d)}{abc + bcd + cda + dab}$$

where $a = \lambda_1 \lambda_2$; $b = \lambda_2 \lambda_3$; $c = \lambda_3 \lambda_4$; $d = \lambda_4 \lambda_1$. One has

$$I_{1100}(\lambda) = \int_0^\infty da \int_0^\infty db \int_0^\infty dc \frac{d(a+c)(b+d)}{abc + bcd + cda + dab}$$

where $d = (a+b+c)$ is understood¹. By successive integration on c ; b ; a one has obtained [1]

$$I_{1100}(\lambda) = \frac{1 + \tilde{\gamma}(3)}{16}$$

where $\tilde{\gamma}(3) = 7(3)=2$. Rewrite $I_{1100}(\lambda)$ differently

$$I_{1100}(\lambda) = \int_{a,b,c,d=0}^\infty da db dc dd \frac{d(a+c)(b+d)}{abc + bcd + cda + dab} \quad (a+b+c+d)$$

¹ same as

$$I_{1100}(\lambda) = \int_0^\infty da db dc \frac{d(a+c)(b+d)}{abc + bcd + cda + dab}$$

and also

$$I_{1100}(\cdot) = \sum_{a,b,c,d=0}^{\infty} da db dc dd \frac{ab}{abc + bcd + cda + dab} (a + b + c + d)$$

Laplace transform

$$I_{1100}(s) = \sum_{d=0}^{\infty} d e^{-s} \frac{I_{1100}(\cdot)}{s} = \frac{1}{s^3} \sum_{a,b,c,d=0}^{\infty} da db dc dd \frac{ab}{abc + bcd + cda + dab} e^{-(a+b+c+d)}$$

such that $I_{1100}(s) = 2I_{1100}(\cdot = 1) = s^3$. Fix $s = 1$ and denote $I_{1100}(s = 1) = I_{1100}$. One has

$$I_{1100} = \sum_{a,b,c,d=0}^{\infty} da db dc dd \sum_{t=0}^{\infty} dt \frac{1}{t!} e^{-(a+b+c+d) - t(\frac{1}{a} + \frac{1}{b} + \frac{1}{c} + \frac{1}{d})}$$

Change variable $u = 2^{\frac{p}{t}}$

$$I_{1100} = 2^4 \sum_{u=0}^{\infty} \frac{udu}{2} \left(\frac{u}{2}\right)^2 K_1(u)^2 K_0(u)^2$$

where the $K_{\cdot}(u)$'s are modified Bessel functions also called MacDonald functions [2]. Therefore

$$I_{1100} = 2^4 \sum_{u=0}^{\infty} \frac{udu}{2} \left(\frac{u}{2}\right)^2 K_1(u)^2 K_0(u)^2 = 2 \frac{1 + \sqrt{3}}{16} \quad (1)$$

2 Generalization

To generalize (1), introduce for $n_a, n_b, n_c, n_d \in \mathbb{N}$

$$I_{n_a n_b n_c n_d} = \sum_{a,b,c,d=0}^{\infty} da db dc dd \frac{a^{n_a} b^{n_b} c^{n_c} d^{n_d}}{abc + bcd + cda + dab} e^{-(a+b+c+d)} \quad (2)$$

and rewrite it as²

$$I_{n_a n_b n_c n_d} = 2^4 \sum_{u=0}^{\infty} \frac{udu}{2} \left(\frac{u}{2}\right)^{n_a + n_b + n_c + n_d} K_{n_a}(u) K_{n_b}(u) K_{n_c}(u) K_{n_d}(u)$$

²This is also formally true for $n_a, n_b, n_c, n_d \in \mathbb{Z}$ but (2) diverges as soon as $n_a < 0$ or $n_b < 0$ or $n_c < 0$ or $n_d < 0$. For $n_a, n_b, n_c, n_d \in \mathbb{N}$ $m \geq 1$; $m \geq 1$

$$I_{n_a n_b n_c n_d}^m = \sum_{a,b,c,d=0}^{\infty} da db dc dd \frac{a^{n_a} b^{n_b} c^{n_c} d^{n_d}}{(abc + bcd + cda + dab)^m} e^{-(a+b+c+d)}$$

One has

$$I_{n_a n_b n_c n_d} = u_{n_a n_b n_c n_d} + v_{n_a n_b n_c n_d} \quad (3)$$

where $u_{n_a n_b n_c n_d}, v_{n_a n_b n_c n_d}$ are positive or negative rational numbers.

-Proof: Since

$$K_{n_a}(u) = 2(n_a - 1) \frac{K_{n_a-1}(u)}{u} + K_{n_a-2}(u)$$

then if $n_a > 0$ even/odd

$$\begin{aligned} \frac{u^{n_a}}{2^{n_a}} K_{n_a}(u) &= \frac{u^{n_a}}{2^{n_a}} K_{0=1}(u) + \frac{u^{n_a-1}}{2} K_{1=0}(u) + c_{n_a-2} u^{n_a-2} K_{0=1}(u) + \dots \\ &+ c_1 u K_1(u) \end{aligned}$$

where $c_{n_a-2}, \dots, c_1$ are rational numbers. It follows that (3) is true if

$$p_n(0000) = \int_0^1 du u^{n+1} K_0(u)^4 \quad n=0 \quad (4)$$

$$p_n(0011) = \int_0^1 du u^{n+1} K_0(u)^2 K_1(u)^2 \quad n=2$$

$$p_n(1111) = \int_0^1 du u^{n+1} K_1(u)^4 \quad n=4 \quad (5)$$

(n even)

rewrites as

$$\frac{1}{(m)} 2^4 \int_0^1 \frac{u du}{2} \left(\frac{u}{2}\right)^{n_a+n_b+n_c+n_d-4(m-1)} K_{n_a-(m-1)}(u) K_{n_b-(m-1)}(u) K_{n_c-(m-1)}(u) K_{n_d-(m-1)}(u)$$

The considerations that follow for $I_{n_a n_b n_c n_d}$ (ie. the case $m = 1$) generalizes to $I_{n_a n_b n_c n_d}^m$ form 1 implying that the latter can also be built from $p_n(0000)$ and $p_n(1111)$ defined in (4) and (5). Therefore $I_{n_a n_b n_c n_d}^m$ is of the form (3). Note that if one introduces back s in $I_{n_a n_b n_c n_d}^m$

$$I_{n_a n_b n_c n_d}^m(s) = \int_0^1 da db dc dd \frac{a^{n_a} b^{n_b} c^{n_c} d^{n_d}}{(abc + bcd + cda + dab)^m} e^{-s(a+b+c+d)}$$

and differentiates n times with respect to s , one obtains with $s = 1$

$$I_{n_a n_b n_c n_d}^{m,n} = \int_0^1 da db dc dd (a+b+c+d)^n \frac{a^{n_a} b^{n_b} c^{n_c} d^{n_d}}{(abc + bcd + cda + dab)^m} e^{-(a+b+c+d)}$$

again of the form (3).

and

$$i_n(0001) = \int_0^{Z_1} du u^{n+1} K_0(u)^3 K_1(u) \quad n-1$$

$$i_n(0111) = \int_0^{Z_1} du u^{n+1} K_0(u) K_1(u)^3 \quad n-3$$

(n odd)

are again³ of the form (3). To show this, integrate by parts (use $dK_0(u) = du = K_1(u)$ and $d(uK_1(u)) = du = uK_0(u)$) to obtain

$$p_n(0000) = \frac{4}{n+2} i_{n+1}(0001) \quad n-0$$

$$p_n(0011) = \frac{2}{n} (i_{n+1}(0001) + i_{n+1}(0111)) \quad n-2$$

$$p_n(1111) = \frac{4}{n-2} i_{n+1}(0111) \quad n-4$$

which implies

$$2np_n(0011) = (n+2)p_n(0000) + (n-2)p_n(1111) \quad n-4$$

It remains to be found $p_n(0000)$ and $p_n(1111)$: integration by parts gives

$$i_n(0111) = \frac{1}{n-1} ((p_{n+1}(1111) + 3p_{n+1}(0011)) \quad n-3$$

$$i_n(0001) = \frac{1}{n+1} ((p_{n+1}(0000) + 3p_{n+1}(0011)) \quad n-1$$

thus the recurrence relation acting in a 2 dimensional vector space

$$\begin{pmatrix} p_{n+2}(0000) \\ p_{n+2}(1111) \end{pmatrix} = \frac{1}{2^5(n+2)} \begin{pmatrix} (n+2)^2(5n+4) & 3n^2(n-2) \\ 3(n+2)^2(n+4) & n(n-2)(5n+16) \end{pmatrix} \begin{pmatrix} p_n(0000) \\ p_n(1111) \end{pmatrix} \quad (6)$$

n-4.

Examples:

³Note that a product of four Macdonald functions is needed here: for example

$$\int_0^{Z_1} du u K_0(u)^3 = \frac{3}{2} \sum_{p=0}^{\infty} \frac{1}{(3p+1)^2} - \frac{2}{3} \quad (2)$$

is not simply given in terms of ζ 's.

by direct computation 0 $n < 4$

$$p_0(0000) = \frac{\sim(3)}{2^2}$$

$$i_1(0001) = \frac{\sim(3)}{2^3}$$

$$p_2(0000) = \frac{3 + \sim(3)}{2^4}$$

$$p_2(0011) = \frac{1 + \sim(3)}{2^4} \text{ known [1]}$$

$$i_3(0001) = \frac{3 + \sim(3)}{2^4}$$

$$i_3(0111) = \frac{1}{2^2} \text{ obvious}$$

and $n = 4$ (initial conditions)

$$p_4(0000) = \frac{3^3 + 7\sim(3)}{2^6}$$

$$p_4(1111) = \frac{53 - 3^2\sim(3)}{2^6}$$

then the recurrence (6) gives for $n > 4$

$$p_6(0000) = \frac{37 + 3^2\sim(3)}{2^4}$$

$$p_6(1111) = 3 \frac{67 - (3)(5)\sim(3)}{2^6}$$

$$p_8(0000) = \frac{(5)(19)(269) + (3^2)(7)(97)\sim(3)}{2^{10}}$$

$$p_8(1111) = 3 \frac{(13)(811) - (3^2)(5^2)(11)\sim(3)}{2^{10}}$$

$$p_{10}(0000) = \frac{9304913 + (3^2)(5^3)(11)(179)\sim(3)}{(2^{12})(5)}$$

$$p_{10}(1111) = 3^4 \frac{(3)(11)(4139) - (5^3)(7)(37)\sim(3)}{(2^{12})(5)}$$

$$p_{12}(0000) = 3^4 \frac{(7)(19)(23909) + (5^3)(23)(263)\sim(3)}{(2^{12})(5)}$$

$$p_{12}(1111) = 3^2 \frac{(43)(67)(11519) - (3^2)(5^3)(7^2)(11)(13)\sim(3)}{(2^{12})(5)}$$

If one sets $n = 2k$ and defines $q_k(0) = p_{2k}(0000) = (2k)!$, $q_k(1) = p_{2k}(1111) = (2k)!$ the recurrence (6) becomes

$$\frac{q_{k+1}(0)}{q_{k+1}(1)} = \frac{1}{2^4(2k+1)} \frac{(5k+2)}{3(k+2)} \frac{3k^2(k-1)}{k(k-1)(5k+8)} = \frac{(k+1)^2}{(k+1)^2} \frac{q_k(0)}{q_k(1)} \quad (7)$$

$k \geq 2$. In the asymptotic regime $k \rightarrow \infty$ the recurrence matrix

$$\begin{pmatrix} \frac{1}{2^5} & \frac{5}{3} & \frac{3}{5} \end{pmatrix}$$

diagonalizes as

$$\begin{pmatrix} \frac{1}{2^5} & \frac{5}{3} & \frac{3}{5} \end{pmatrix} = \begin{pmatrix} \frac{1}{2^5} & \frac{5}{3} & \frac{3}{5} \end{pmatrix} \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix} = \begin{pmatrix} \frac{1}{2^4} & \frac{1}{0} & \frac{0}{4} \end{pmatrix}$$

with eigenvalues $\lambda_1 = 4$ and $\lambda_2 = 16$. In this convenient diagonal basis (7) becomes

$$\frac{q_{k+1}(1) + q_{k+1}(0)}{q_{k+1}(1) - q_{k+1}(0)} = \frac{\frac{1}{2^4(2k+1)} \frac{k-2+k(k-1)(k+4)}{4((k+1)+k(k-1))} = \frac{(k-2)+k(k-1)(k+4)}{4(k+1+k(k-1))} \frac{q_k(1) + q_k(0)}{q_k(1) - q_k(0)}$$

$k \geq 2$ with by definition $q_k(1) + q_k(0) > q_k(1) - q_k(0) > 0$.

Examples:

$k = 2$ (initial conditions)

$$q_2(1) + q_2(0) = \frac{13}{(2^8)} \sim (3) \quad (8)$$

$$q_2(1) - q_2(0) = \frac{5}{(2^5)} \sim (3) \quad (9)$$

$k > 2$

$$q_3(1) + q_3(0) = \frac{53}{(2^{10})} \frac{(3)^2 \sim (3)}{(3^2)(5)}$$

$$q_3(1) - q_3(0) = \frac{349}{(2^{10})} \frac{(3)^4 \sim (3)}{(3^2)(5)}$$

$$\begin{aligned}
q_4(1) + q_4(0) &= \frac{3037 (3)^2 (73) \sim (3)}{(2^{16}) (3^2) (5) (7)} \\
q_4(1) - q_4(0) &= \frac{1787 (3)^2 (47) \sim (3)}{(2^{12}) (3^2) (5) (7)} \\
q_5(1) + q_5(0) &= \frac{(439) (2003) (3^2) (5^3) (181) \sim (3)}{(2^{19}) (3^4) (5^3) (7)} \\
q_5(1) - q_5(0) &= \frac{(7) (73) (1993) (3^2) (5^4) (43) \sim (3)}{(2^{18}) (3^4) (5^2) (7)} \\
q_6(1) + q_6(0) &= \frac{(2283583) (3^2) (5^3) (479) \sim (3)}{(2^{21}) (3^3) (5^3) (7) (11)} \\
q_6(1) - q_6(0) &= \frac{(127) (1901) (3^3) (5^3) (17) \sim (3)}{(2^{14}) (3^3) (5^3) (7) (11)} \\
q_7(1) + q_7(0) &= \frac{(53) (1708543) (3) (5^3) (7^3) (167) \sim (3)}{(2^{21}) (3^2) (5^3) (7^3) (11) (13)} \\
q_7(1) - q_7(0) &= \frac{(13) (61485173) (3) (5^3) (7^4) (211) \sim (3)}{(2^{23}) (3^2) (5^3) (7^2) (11) (13)}
\end{aligned}$$

Both $q_k(1) + q_k(0)$ and $q_k(1) - q_k(0)$ decrease (faster than $(1=16)^{k-2}$) as $(1=16)^{k-2}$ (when k is finite) when $k \rightarrow 1$, i.e. their asymptotic behavior is governed by the smallest of the eigenvalues. This is due to the initial conditions tuning (8) and (9), which defines a $\sim(3)$ dependant initial orientation in the 2 dimensional vector space leading to this particular asymptotic behavior. This situation is analogous to the one encountered for establishing the irrationality of (3) (see [3] for a review of various demonstrations of the irrationality of (3)): in that case a second order recurrence relation (or equivalently a family of integrals) with (3) dependant initial conditions such that an asymptotic convergence governed by the smallest of eigenvalues is achieved. It is however manifest in the present case that both $q_k(1) + q_k(0)$ and $q_k(1) - q_k(0)$ have denominators which increase too fast with k to obtain a convergence allowing for an alternative proof of the irrationality of (3) . This is still the case for any k -independant initial conditions rotation in the plane. If one allows for more general k -dependant combinations, note that if one redefines $q_k(01) = p_{2k}(0011) = (2k)!$ then

$$q_k(01) = q_k(1) + q_k(0) - (q_k(1) - q_k(0)) = k$$

if one redefines $j_k(0111) = i_{2k+1}(0111) = (2k+1)!$ then

$$j_k(0111) = \frac{k-1}{2(2k+1)} q_k(1)$$

and finally if one redefines $j_k(0001) = i_{2k+1}(0001) = (2k+1)!$ then

$$j_k(0001) = \frac{k+1}{2(2k+1)} q_k(0)$$

Particular k -dependant linear combinations of $q_k(1) + q_k(0)$ and $q_k(1) - q_k(0)$ follow as

$$j_k(0111) + j_k(0001) = \frac{k(q_k(1) + q_k(0)) - (q_k(1) - q_k(0))}{2(2k+1)} q_k(1) + q_k(0)$$

$$j_k(0111) - j_k(0001) = \frac{k(q_k(1) - q_k(0)) - (q_k(1) + q_k(0))}{2(2k+1)} q_k(1) - q_k(0)$$

with by definition $j_k(0111) - j_k(0001) > 0$. However both $j_k(0111) - j_k(0001)$ and $j_k(0111) + j_k(0001)$ have again denominators which increase too fast with k .

3 Further Generalization: (5); (7);

In view of obtaining (5), a natural generalization of (2) which would still involve Macdonald functions would be, in the simplest case

$$\int_{a \neq b \neq c \neq d \neq e \neq f} da db dc dd de df \frac{1}{abde + bdef + cdefa + defab + efabc + fabcd} e^{(a+b+c+d+e+f)}$$

nothing but

$$2^6 \int_0^1 \frac{u du}{2} K_0(u)^6$$

Numerics indicates however that this integral is not a linear combination with rational coefficients of $1, (5), (3), (2), (3), \dots$. Introduce rather for $n \geq 2$ the $2n$ integration variables a_i and b_i , with $0 \leq i \leq n$, and consider

$$\int_0^1 \prod_{i=1}^n da_i db_i (1 - (a_i + b_i)) \frac{P(a_1, b_1; \dots; a_n, b_n)}{a_1 b_1 (a_2 + b_2) \dots (a_n + b_n) + \varphi}$$

where cp means cyclic permutation over $1;2;\dots;n$ and $P(a_1;b_1;\dots;a_n;b_n)$ is any given monomial of the variables $a_i;b_i$. The case $n = 2$ corresponds to the variables $a;b;c;d$ used up to now. This is a family of integrals with root

$$I_n = \int_0^1 \int_{i=1}^n Y^n da_i db_i \prod_{i=1}^n (1 - (a_i + b_i)) \frac{1}{a_1 b_1 (a_2 + b_2) \dots (a_n + b_n) + \text{cp}}$$

Change variables

$$u_i = a_i + b_i \quad v_i = \frac{a_i - b_i}{a_i + b_i}$$

to get

$$I_n = 4 \int_0^1 \int_{i=1}^n du_i \prod_{i=1}^n (1 - u_i) \int_0^1 \int_{i=1}^n dv_i \frac{1}{(1 - v_i^2)u_1 + \dots + (1 - v_n^2)u_n}$$

In order to integrate over $u_1 \dots u_n$, follow the procedure used above: Laplace transform the measure and exponentiate the denominator

$$I_n = \frac{4}{(n-1)!} \int_0^1 \int_{i=1}^n dv_i \int_0^1 dt \int_{i=1}^n du_i \exp \left(- \sum_{i=1}^n u_i (1 + t(1 - v_i^2)) \right)$$

Integration over u_i and v_i leads to

$$I_n = \frac{4}{2^n (n-1)!} \int_0^1 dt \frac{1}{((1+t)t)^n} \int_0^1 \frac{1}{\left(\frac{1+t}{t} + 1 \right)^{n-1}} \frac{1}{\left(\frac{1+t}{t} - 1 \right)^{n-1}} dt$$

which can be rewritten⁴ as

$$I_n = \frac{2^n}{(n-1)!} \int_0^1 \frac{dy}{y} \frac{1}{1 - y^2} \left(\log \frac{1}{y} \right)^{n-1}$$

Using

$$\int_0^1 dy y^p \left(\log \frac{1}{y} \right)^n = \frac{(n+1)}{(p+1)^{n+1}}$$

leads to

$$I_n = 2^n \frac{(n+1)}{(n-1)!} \sum_{p=0}^{n-1} \frac{(p+1)(p+2)\dots(p+n-2)}{(n+2p-1)^{n+1}} \quad (10)$$

⁴or as

$$I_n = \frac{2}{(n-1)!} \int_0^1 dy y \left(\frac{y}{1-y^2} \right)^{n-1}$$

Examples:

$$I_2 = 7 \quad (3)$$

$$I_3 = 3 \quad (3)$$

$$I_4 = 3(7 \quad (3) \quad \frac{31}{2^2} \quad (5))$$

$$I_5 = \frac{5}{3} (\quad (3) \quad (5))$$

$$I_6 = \frac{5}{2^3} (7 \quad (3) \quad \frac{(5)(31)}{2} \quad (5) + \frac{(3^2)(127)}{2^4} \quad (7))$$

Only (odd) does appear in (10) as can be seen by distinguishing between n odd and n even

$$I_n = \frac{4}{(n-1)^2} \sum_{p=0}^{\lfloor \frac{n-1}{2} \rfloor} \frac{\sum_{q=1}^Q ((n+2p-1)^2 - (2q-1)^2)}{(n+2p-1)^{n+1}} \quad n \text{ even}$$

$$I_n = \frac{4}{(n-1)^2} \sum_{p=0}^{\lfloor \frac{n-3}{2} \rfloor} \frac{\sum_{q=1}^Q ((n+2p-1)^2 - (2q)^2)}{(n+2p-1)^n} \quad n \text{ odd}$$

Acknowledgments: One of us (S.O.) would like to thank J. Desbois, C. Jacquemin, J. Myrheim, C. Schmit and P. Zinn-Justin for useful conversations.

[1] J. Desbois, C. Furtlehner and S. Ouvry, Nucl. Phys. B 453 [FS] (1995) 759

[2] see Gradshteyn and Ryzhik, Table of Integrals, Series and Products, Fifth Edition, Academic Press, 8.432.1. and 8.432.6.

[3] for a review see S. Fischler, Seminaire Bourbaki, Novembre 2002, n°910