

# A lower bound to the spectral threshold in curved tubes

P. Exner<sup>1,2</sup>, P. Freitas<sup>3</sup> and D. Krejčířík<sup>1,3</sup>

<sup>1</sup> Department of Theoretical Physics, Nuclear Physics Institute,  
Academy of Sciences, 250 68 Řež near Prague, Czech Republic

<sup>2</sup> Doppler Institute, Czech Technical University,  
Břehova 7, 115 19 Prague, Czech Republic

<sup>3</sup> Departamento de Matemática, Instituto Superior Técnico,  
Av. Rovisco Pais, 1049-001 Lisboa, Portugal

E-mail: exner@ujf.cas.cz, pfreitas@math.ist.utl.pt, dkrej@math.ist.utl.pt

28 April 2004

## Abstract

We consider the Laplacian in curved tubes of arbitrary cross-section rotating together with the Frenet frame along curves in Euclidean spaces of arbitrary dimension, subject to Dirichlet boundary conditions on the cylindrical surface and Neumann conditions at the ends of the tube. We prove that the spectral threshold of the Laplacian is estimated from below by the lowest eigenvalue of the Dirichlet Laplacian in a torus determined by the geometry of the tube.

## 1 Introduction

Problems linking the shape of a region to the spectrum of the associated Laplacian, subject to various boundary conditions, have been considered for more than a century. While classical motivations came from theories of elasticity, acoustics, electromagnetism, etc, in the quantum-mechanical context a strong fresh impetus is mostly due to the recent technological progress in semiconductor physics.

More specially, the Dirichlet Laplacian in finite plane strips or space tubes of constant cross-section is widely used as a mathematical model for the Hamiltonian of a quantum particle in mesoscopic structures called quantum waveguides [DE95, LCM99, Hur00]. The existence of geometrically induced bound states in curved asymptotically straight waveguides is probably the most interesting theoretical result for these systems [ES89, GJ92, RB95, DE95, KK, CDFK]. Indeed, these bound states, which are known to perturb the particle transport, are of pure quantum origin because there are no classical closed

trajectories in the tubes in question, apart from a zero measure set of initial conditions in the phase space. Mathematically, one deals with the discrete spectrum of the Dirichlet Laplacian, which is a non-trivial property for unbounded regions. The principal objective of this paper is to establish a lower bound to the ground-state energies of curved quantum waveguides.

We proceed in greater generality by considering  $d$ -dimensional tubes, unbounded or bounded, with any  $d \geq 2$  and arbitrary cross-section rotating along a reference curve together with the Frenet frame. At the same time, we do not restrict ourselves to asymptotically straight tubes, i.e., if the tube is unbounded, the estimated spectral threshold of the Laplacian may not be a discrete eigenvalue, but rather the threshold of the essential spectrum; this happens, for instance, if the tube is periodically curved.

To state the main result of the paper, let us introduce some notation. Given a bounded or unbounded open interval  $I \subset \mathbb{R}$ , let  $\gamma : I \rightarrow \mathbb{R}^d$  be a unit-speed curve with curvatures  $\kappa_i : I \rightarrow \mathbb{R}$ ,  $i = 1, \dots, d-1$ , w.r.t. an appropriate smooth Frenet frame  $\{e_1, \dots, e_d\}$ , cf. the assumption H1 below. Given a bounded open connected set  $\Omega \subset \mathbb{R}^{d-1}$  with the centre of mass at the origin, we define the tube  $\Omega_\gamma$  by rotating  $\Omega$  along the curve together with the Frenet frame, i.e.,

$$\gamma(s) = L(I, \gamma); \quad L(s; u_2, \dots, u_d) = \gamma(s) + e(s)u; \quad (1)$$

(the repeated indices convention is adopted throughout the paper, the Latin and Greek indices run through  $1, 2, \dots, d$  and  $2, \dots, d$ , respectively). We make the assumption H2 below (cf Remark 3) in order to ensure that  $L : I \rightarrow \mathbb{R}^d$  is a diffeomorphism. Our object of interest is the non-negative Laplacian

$$\Delta_{\Omega_\gamma} \text{ on } L^2(\Omega_\gamma); \quad (2)$$

subject to Dirichlet boundary conditions on the cylindrical part of the boundary  $L(I \times \partial\Omega)$  and, if  $\partial I$  is not empty, Neumann boundary conditions on the remaining boundary  $L(\partial I \times \Omega)$ . Our main result reads as follows.

**Theorem 1.** Suppose the assumptions H1 and H2 are satisfied. Then

$$\inf_{\Omega} \lambda_1(\Omega_\gamma) \geq \min_{\Omega} \lambda_0(\sup_{\Omega} \kappa_1); \quad \lambda_0(\inf_{\Omega} \kappa_1)g; \quad (3)$$

where  $\lambda_0(\kappa) \geq c > 0$  denotes the spectral threshold of  $\Delta_\Omega$  in the tube of cross-section  $\Omega$  built either over a circle of curvature  $\kappa \neq 0$  or over a straight line if  $\kappa = 0$ ;  $c$  is a constant depending only on  $\Omega$  and  $d$ .

The lower bound of Theorem 1 holds, of course, for other boundary conditions imposed on  $L(I \times \partial\Omega)$ , cf. Section 5.

Note that  $\lambda_0(\kappa)$  is the lowest eigenvalue of the Dirichlet Laplacian in a torus of cross-section  $\Omega$  if  $\kappa \neq 0$  or the threshold of the essential spectrum of the Dirichlet Laplacian in an infinite straight tube of cross-section  $\Omega$  (which is the lowest eigenvalue  $\lambda_0$  of the Dirichlet Laplacian in  $\Omega$ ) if  $\kappa = 0$ , cf. Section 4. Thus the claim of Theorem 1 can be expressed illustratively as follows: take

an "osculation torus" at each point of  $\Gamma$  (i.e. the torus with the identical cross-section built over the osculation circle to  $\Gamma$  at the point), then the bound (3) corresponds to the smallest of this tori spectral thresholds. The uniform lower bound given by the geometric constant  $c$  is a consequence of the Faber-Krahn inequality, cf Proposition 4.

We stress again that while the spectrum of (2) is purely discrete whenever  $I$  is bounded,  $(\lambda_1)$  has in general both discrete and essential parts in the unbounded case. For instance, if  $I = \mathbb{R}$ ,  $\Omega = B_a$  (ball of radius  $a > 0$ ),  $\lambda_1 \notin \mathbb{R}$  but  $\lambda_1(s) \neq 0$  as  $|s| \rightarrow 1$ , it is known from [CDFK] that  $\sigma_{\text{ess}}(\Delta) = [0; 1)$  and there are always discrete eigenvalues in  $(0; 0)$ .

While bounds on the eigenvalues for the Laplacian on bounded subsets of  $\mathbb{R}^d$  have been studied by many authors (see [Hen03] for an overview), to the best of our knowledge there is only one previous result on the lower bound to the spectral threshold of the Laplacian in unbounded tubes. Using the Payne-Polya-Weinberger conjecture [PPW 55, PPW 56] proved then in [AB91] (see also [AB92]), M. S. Ashbaugh and the first author derived in [AE90] a lower bound in the situation when  $I = \mathbb{R}$ ,  $d = 2, 3$ , the cross-section was circular and the discrete spectrum of  $\Delta$  was not empty but finite. As we discuss at the end of Section 5, our Theorem 1 provides a better bound and applies to tubes with an infinite number, or without any, discrete eigenvalues, too. On the other hand, the approach of [AE90] applies to more general forms of  $\Omega$  than the regular tubes considered here. Let us also mention that one can use the results of [EW 01] to derive a Lieb-Thirring-type inequality for  $\Delta$ .

The heuristic idea behind the proof of Theorem 1 is as follows. For a moment, let us assume that  $\lambda_1$  is piece-wise constant and all  $\lambda_1 = 0$ , so that  $I$  is a closure of the union of  $L$  (possibly  $L = 1$ ) open subintervals  $I_i$ ,  $i = 1, \dots, L$ , and each  $I_i = (I_i)$  is a circular or straight segment. We have  $\lambda_1 = \lambda_1(I_i)$ ; where each  $\lambda_1$  is the Laplacian on  $L^2(I_i)$  with Dirichlet boundary conditions on  $L(I_i \cap \partial \Omega)$  and the Neumann ones on  $L((I_i) \cap \partial \Omega)$ . Note that  $\inf(\lambda_1)$  does not depend on the length of  $I_i$  because the first (generalised) eigenfunction of the Dirichlet Laplacian in a torus or an infinite straight tube is invariant w.r.t. to rotations or translations, respectively. Consequently,  $\inf(\lambda_1) = \lambda_1(I_1)$ , where  $I_1$  denotes the first curvature of  $\Omega$ . The spectral threshold of  $\Delta$  is thus estimated from below by  $\min \lambda_1(I_1)$  and an analysis of the properties of the first eigenvalue in the torus (Section 4) shows that this minimum is equal to  $\min \lambda_1(\max \lambda_1(I_1); 0 \leq \lambda_1(I_1) \leq g)$  (note that  $\lambda_1(I_1)$  may not be even for a general cross-section!). An important consequence of (geometric) Lemma 1 below is that this lower bound is not affected by higher curvatures  $\lambda_2, \dots$ . Then the general result of Theorem 1 follows by the above procedure at once if one considers the Laplacian through its quadratic form (because the supplementary Neumann conditions do not appear explicitly in the form domain).

The organisation of the paper is as follows. The tube  $\Omega$  and the corresponding Laplacian  $\Delta$  are properly defined in the preliminary Section 2. In Section 3, we prove the geometric Lemma 1 and an intermediate lower bound, Theorem 2, as its direct consequence. Theorem 1 then immediately follows from

Theorem 2 and results in Section 4, which is devoted to a detailed analysis of spectral properties of  $\mathcal{H}_\gamma$  in the case where the reference curve  $\gamma$  is a circular segment. Finally, in Section 5, we summarise the results obtained, discuss possible extensions and refer to some open problems. We conclude the paper by comparing our result with the lower bound found in [AE90] for a special case of infinite tubes in two and three dimensions.

## 2 Preliminaries

### 2.1 The reference curve

Given an open interval  $I \subset \mathbb{R}$  and an integer  $d \geq 2$ , let  $\gamma : I \rightarrow \mathbb{R}^d$  be a unit-speed  $C^{d-1}$ -smooth curve satisfying

**H1**  $\gamma$  possesses a positively oriented  $C^1$ -smooth Frenet frame  $\{e_1, \dots, e_d\}$  with the properties that  $e_1 = \dot{\gamma}$  and

$$\text{span}\{e_2, \dots, e_d\} \subset T_{\gamma(s)} \text{ for all } s \in I; \quad e_i(s) \text{ lies in the span of } e_1(s), \dots, e_{i-1}(s);$$

**Remark 1.** We refer to [Kli78, Sec. 1.2] for the notion of Frenet frames. A sufficient condition to ensure the existence of the Frenet frame of  $\gamma$  is to require that for all  $s \in I$ , the vectors  $\dot{\gamma}(s), \ddot{\gamma}(s), \dots, \gamma^{(d-1)}(s)$  are linearly independent, cf [Kli78, Prop. 1.2.2]. This is always satisfied if  $d = 2$ . However, we prefer not to assume a priori this non-degeneracy condition for  $d \geq 3$  because then one excludes the curves such that  $\gamma|_{I_1}$  lies in a lower-dimensional subspace of  $\mathbb{R}^d$  for some open  $I_1 \subset I$ . Further comments on the assumption **H1** will be given in the closing section.

We have the Serret-Frenet formulae, cf [Kli78, Sec. 1.3],

$$\dot{e}_i = K_{ij} e_j \quad (4)$$

where  $K = (K_{ij})$  is the skew-symmetric  $d \times d$  matrix defined by

$$K = \begin{pmatrix} 0 & & & & 0 \\ & \kappa_1 & & & \\ & & \ddots & & \\ & & & \ddots & \\ 0 & & & & 0 \end{pmatrix} : \quad (5)$$

Here  $\kappa_i$  is called the  $i^{\text{th}}$  curvature of  $\gamma$  which is, under our assumptions, a continuous function of the arc-length parameter  $s \in I$ .

## 2.2 Tubes

Let  $\Omega$  be an arbitrary bounded open connected set in  $\mathbb{R}^{d-1}$ . Without loss of generality, we assume that  $\Omega$  is translated so that its centre of mass is at the origin. Put  $\Omega_0 \equiv \Omega$  and  $u \equiv (u_2; \dots; u_d) \in \Omega_0$ . We define the tube built over  $\Omega$  as the image of the mapping  $L : \Omega_0 \times \mathbb{R}^d$  defined in (1), i.e.  $\Omega \equiv L(\Omega_0)$ . Assuming that

$$L : \Omega_0 \times \mathbb{R}^d \rightarrow \mathbb{R}^d : (s; u) \mapsto L(s; u)g \text{ is a } C^1\text{-diffeomorphism ;} \quad (6)$$

we can identify  $\Omega$  with the Riemannian manifold  $(\Omega_0; G)$ , where  $G = (G_{ij})$  is the metric tensor induced by the immersion  $L$ , i.e.  $G_{ij} \equiv L_{,i} \cdot L_{,j}$ . (Here and in the sequel, the dot denotes the scalar product in  $\mathbb{R}^d$  and the comma with an index means the partial derivative w.r.t.  $x_i$ ,  $x_{,i} \equiv \partial L / \partial x_i$ .) Using (4), we find

$$G = \begin{pmatrix} 0 & h_1 & h_2 & h_3 & \dots & h_{d-1} & h_d \\ h_1 & 1 & 0 & \dots & 0 & 0 & 0 \\ h_2 & 0 & 1 & \dots & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ h_{d-1} & 0 & 0 & \dots & 1 & 0 & 0 \\ h_d & 0 & 0 & \dots & 0 & 0 & 1 \end{pmatrix}; \quad \begin{aligned} h_1 &\equiv h^2 + h \cdot h ; \\ h(s; u) &\equiv 1 - \frac{1}{2} (s) u_2 ; \\ h(s; u) &\equiv K(s) u : \end{aligned} \quad (7)$$

Furthermore,  $\det G = h^2$  which defines through  $d\text{vol} \equiv h(s; u) ds du$  the volume element of  $\Omega$ ; here and in the sequel  $du = du_2 \dots du_d$  denotes the  $(d-1)$ -dimensional Lebesgue measure in  $\Omega$ .

It can be checked by induction that the inverse  $G^{-1} = (G^{ij})$  of the metric tensor (7) satisfies

$$G^{-1} = \frac{1}{h^2} \begin{pmatrix} 0 & 1 & h_2 & h_3 & h_4 & \dots & h_d \\ 1 & h^2 + h_2^2 & h_2 h_3 & h_2 h_4 & \dots & h_2 h_d \\ h_2 & h_2 h_3 & h^2 + h_3^2 & h_3 h_4 & \dots & h_3 h_d \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ h_{d-1} & h_{d-1} h_2 & h_{d-1} h_3 & h_{d-1} h_4 & \dots & h_{d-1} h_d \\ h_d & h_d h_2 & h_d h_3 & h_d h_4 & \dots & h^2 + h_d^2 \end{pmatrix} : \quad (8)$$

**Remark 2** (Low-dimensional examples). When  $d = 2$ , the cross-section  $\Omega$  is an interval, the curve has only one curvature  $\kappa = \kappa_1$  and  $G$  is diagonal with

$$h(s; u) = 1 - \frac{1}{2} \kappa(s) u :$$

When  $d = 3$ , one finds

$$G(s; u) = \begin{pmatrix} 0 & (1 - u_2)^2 + \frac{1}{2} \kappa^2 u^2 & u_3 & u_2 \\ (1 - u_2)^2 + \frac{1}{2} \kappa^2 u^2 & u_3 & 1 & 0 \\ u_3 & u_2 & 0 & 1 \end{pmatrix} ;$$

where  $\kappa = \kappa_1$  and  $\tau = \tau_2$  denote the curvature and torsion of  $\gamma$ , respectively.

Remark 3 (On the assumption (6)). Let  $\|u\| = \sqrt{\int \overline{u} u}$  and define

$$a = \sup_{\|u\|=1} \|u\|.$$

By virtue of the inverse function theorem,  $L$  is a local  $C^1$ -diffeomorphism provided  $h$  does not vanish on  $\Gamma_0$ . It becomes a global diffeomorphism if it is required to be injective in addition. Hence, (6) holds true provided

- H2i (i)  $\|u\| \leq L^{-1}(I)$  and  $\|u\|_{k_1} < 1$ ,  
(ii)  $\Gamma_0$  does not overlap itself,

which we shall assume henceforth. Let us point out two facts. First, if  $\Gamma_0$  were a compact embedded curve, then the condition (ii) could always be achieved for a sufficiently small  $\epsilon$ . Second, we do not need to assume the condition (ii) if we consider  $(\Gamma_0; G)$  as an abstract Riemannian manifold where only the curve is embedded in  $\mathbb{R}^d$ .

For further purposes, we introduce

$$\|u\| = \int_{\Gamma_0} \sqrt{G}^{-1/2} (u_2, u_3, \dots, u_d)^2 \sqrt{G};$$

i.e. the mirror image of  $\|u\|$  w.r.t. the hyperplane  $\int_{\Gamma_0} u_2 = 0$ .

## 2.3 The Laplacian

Introducing the unitary transformation  $\mathcal{T} : L^2(\Gamma_0; G) \rightarrow L^2(\Gamma_0; \text{dvol})$ ; we may identify the Hilbert space  $L^2(\Gamma_0; G)$  with  $H = L^2(\Gamma_0; \text{dvol})$  and the Laplacian (2) with the self-adjoint operator  $H$  associated with the quadratic form  $Q$  on  $H$  defined by

$$Q[u] = \int_{\Gamma_0} \overline{u}_i(s; u) G^{ij}(s; u) u_{,j}(s; u) h(s; u) ds; \quad (9)$$

$$\text{Dom } Q = \{u \in W^{1,2}(\Gamma_0; \text{dvol}) \mid u_{,j}(s; u) = 0 \text{ for a.e. } (s; u) \in \Gamma_0\}.$$

Here  $(x)$  for  $x \in \Gamma_0$  means the corresponding trace of the function  $u$  on the boundary.

We have

$$H = \int_{\Gamma_0} \frac{1}{2} \partial_i \mathcal{T}^{-1} G^{ij} \mathcal{T}^{-1} \partial_j;$$

which is a general expression for the Laplace-Beltrami operator in a manifold equipped with a metric  $G$ . However, we stress that the equality must be understood in the form sense if  $\mathcal{T}^{-1}$  are not differentiable (which is the case we are particularly concerned to deal with in this paper).

### 3 An intermediate lower bound

In this section, we derive an intermediate lower bound to the spectral threshold of  $\mathcal{H}_1$  which is crucial for the proof of Theorem 1.

It is worth to notice that one has the decomposition

$$G^{-1} = \text{diag}(h^{-2}; 1; \dots; 1) + h^{-2} T; \quad (10)$$

where the matrix  $T$  depends on the higher curvatures  $\kappa$ , but not on  $\gamma_1$ , in such a way that  $T = 0$  if  $\kappa = 0$ . Hence, if the reference curve is planar (i.e.  $\kappa = 0$ ) then the norm of a covector  $\gamma \in T_{(s,u)}^* \mathcal{H}_0$  w.r.t. the metric  $G$  is clearly estimated from below by the norm of its projection to  $T_u^* \mathcal{H}_0$  w.r.t. the Euclidean norm, i.e.  $|\gamma|_{G^{-1}} \geq |\gamma|_{T_u^* \mathcal{H}_0}$ . An important observation is that this property is not influenced by the presence of higher curvatures:

Lemma 1. One has

$$G^{-1} \geq \text{diag}(0; 1; \dots; 1)$$

in the matrix-inequality sense.

Proof. In view of (8) and (10), one has  $G^{-1} \geq \text{diag}(0; 1; \dots; 1) = h^{-2} A$  where  $A = \text{diag}(1; 0; \dots; 0) + T$  is positive definite since

$$A_{ij} = \delta_{ij} + 2\gamma_i \gamma_j + (h^{-2})^2 = (\gamma_i + h^{-2})^2 \geq 0$$

for any  $\gamma \in \mathbb{R}^d$ . □

Lemma 1 has the following crucial corollary.

Theorem 2. Suppose the assumptions H1i and H2i are satisfied. Then

$$\inf_{s \in \mathcal{I}} \lambda_1(s) \geq \inf_{s \in \mathcal{I}} \lambda_1(s);$$

where

$$\lambda_1(s) := \inf_{\gamma \in W_0^{1,2}(\mathcal{H}_0)} \frac{\int_{\mathcal{H}_0} |\gamma|^2 ds du}{\int_{\mathcal{H}_0} \gamma^2 ds du}; \quad (11)$$

Proof. The definition of the form (9), Lemma 1 and (11) yield

$$\begin{aligned} Q[\gamma] &= \int_{\mathcal{H}_0} |\gamma|^2 ds du - \frac{1}{2} \int_{\mathcal{H}_0} \gamma^2 ds du \\ &= \int_{\mathcal{H}_0} |\gamma|^2 ds du - \frac{1}{2} \int_{\mathcal{H}_0} \gamma^2 ds du \\ &= \int_{\mathcal{H}_0} |\gamma|^2 ds du - \frac{1}{2} \int_{\mathcal{H}_0} \gamma^2 ds du \\ &= \inf_{s \in \mathcal{I}} \lambda_1(s) \int_{\mathcal{H}_0} |\gamma|^2 ds du \\ &= \inf_{s \in \mathcal{I}} \lambda_1(s) \|\gamma\|_H^2 \end{aligned}$$

for any  $\gamma \in \text{Dom } Q$ . □

## 4 Toroidal segments

In this section, we give a geometrical meaning to the quantity (11) and examine its properties, which then yield Theorem 1 as a consequence of Theorem 2. In particular, the monotonicity properties of Proposition 1 below establish the bound (3) of Theorem 1, while the uniform lower bound follows from Proposition 4 below.

Consider now the situation when  $I$  is bounded,  $\gamma = \gamma_1$  is constant and all  $\gamma_j = 0$ , i.e.  $\Omega$  is either a circular segment of length  $|\mathbb{I}|$  and radius  $l = |\mathbb{J}|$  if  $\gamma \neq 0$  or a straight line of length  $|\mathbb{J}|$  if  $\gamma = 0$ . The assumption  $H2$  holds true provided

$$a|\mathbb{J}| < 1 \quad \text{and} \quad |\mathbb{J}|^2 = |\mathbb{I}|. \quad (12)$$

If  $\gamma = |\mathbb{J}|$ , then  $\Omega$  is a circle with one point removed and  $\Omega$  is a torus of cross-section  $|\mathbb{I}|$  about it (more precisely, depending on the sign of  $\gamma$ ,  $\Omega$  can be identified either with  $(\mathbb{C} \setminus \{0\}) \times \mathbb{I}$  or  $(\mathbb{C} \setminus \{0\}) \times \mathbb{I}$ ), where  $\mathbb{C}$  stands for the one-dimensional sphere of radius  $l = |\mathbb{J}|$ .

Let  $H$  denote the operator associated with (9) in this constant case. The spectrum of  $H$  consists of discrete eigenvalues which we denote by

$$0(\gamma; |\mathbb{I}|) < \lambda_1(\gamma; |\mathbb{I}|) < \lambda_2(\gamma; |\mathbb{I}|) < \dots;$$

where the first one is positive. Since  $K = 0$  and  $\gamma_1$  is constant, the metric (7) is diagonal and independent of the "angular" variables. Consequently, the coefficients of  $H$  do not depend on  $s$  either and the Laplacian can be decomposed w.r.t. the angular momentum subspaces represented by the eigenfunctions of  $\Delta_N^I$ , i.e. the Neumann Laplacian on  $L^2(I)$ .

Lemma 2. Let  $\varphi_n, n \in \mathbb{N}$ , denote the normalised eigenfunction corresponding to the  $(n+1)^{\text{th}}$  eigenvalue  $E_n = (\gamma|\mathbb{J}|)^2 n^2$  of  $\Delta_N^I$ . Then  $H$  is unitarily equivalent to the direct sum  $\bigoplus_{n \in \mathbb{N}} H_n$ ; where each  $H_n$  acts on  $f_n g \in L^2(\Omega; (1-u_2) du)$  and it is defined in the form sense by

$$H_n := \frac{E_n}{(1-u_2)^2} - \frac{1}{1-u_2} \partial_{u_2} (1-u_2) \partial_{u_2}; \quad \text{Dom}(H_n)^{\frac{1}{2}} := f_n g \in W_0^{1,2}(\Omega);$$

Furthermore, each  $H_n$  is unitarily equivalent to the operator  $\hat{H}_n$  on  $f_n g \in L^2(\Omega)$  defined in the form sense by

$$\hat{H}_n := \Delta_D^I + V_n; \quad \text{Dom}(\hat{H}_n)^{\frac{1}{2}} := f_n g \in W_0^{1,2}(\Omega);$$

where

$$V_n(u_2) := \frac{E_n}{(1-u_2)^2} \quad (13)$$

and  $\Delta_D^I$  denotes the Dirichlet Laplacian on  $L^2(I)$ .



Proof. Since  $\gamma$  is constant,  $h(s; u)$  is independent of  $s$  and we have the following natural isomorphism

$$\begin{aligned} H & \cong L^2(I) \otimes L^2(\Omega; (1 - u_2) du); \\ \text{Dom } Q & \cong \text{Dom} \left( \frac{I}{N} \right)^{\frac{1}{2}} \otimes W_0^{1,2}(\Omega; (1 - u_2) du); \end{aligned}$$

Since the family  $\{f_n\}_{n \in \mathbb{N}}$  forms a complete orthonormal basis in  $L^2(I)$ , the Hilbert space  $H$  admits a direct sum decomposition  $H = \bigoplus_{n \in \mathbb{N}} H_n$ ; where  $H_n = \{f_n \otimes g \in L^2(\Omega; (1 - u_2) du) : g \in W_0^{1,2}(\Omega; (1 - u_2) du)\}$ . Noticing that the spaces  $W_0^{1,2}(\Omega; (1 - u_2) du)$  and  $W_0^{1,2}(\Omega)$  can be identified as sets, we arrive at the first claim of the Lemma because  $Q[f] = \sum_{n \in \mathbb{N}} \langle f, H_n \rangle^2$  for any  $f \in \text{Dom}(H_n)^{\frac{1}{2}}$ . The second claim follows by means of the transformation  $\mathcal{T}(1 - u_2)^{\frac{1}{2}}$ ; which is unitary from  $H_n$  to  $f_n \otimes L^2(\Omega)$  and leaves invariant  $\text{Dom}(H_n)^{\frac{1}{2}}$ .  $\square$

Let us recall that the spectrum of  $\frac{I}{D}$  consists of discrete eigenvalues which we denote by

$$0 < \lambda_1 < \lambda_2 < \dots; \quad \lambda_n \rightarrow \infty \text{ as } n \rightarrow \infty;$$

where the lowest eigenvalue  $\lambda_1$  is positive.

Lemma 2 is useful in order to investigate the spectrum of  $H$ . Here we employ it just to establish some properties of the first eigenvalue. Since the spectrum of a direct sum of self-adjoint operators is given by the sum of the individual spectra, cf [RS72, Corol. of Thm. V III.33],  $\lambda_1(\gamma; \mathbb{J})$  is just the first eigenvalue of  $\hat{H}_0$  (and  $H_0$ ).

The first observation is that  $\lambda_1(\gamma; \mathbb{J})$  does not depend on  $|\mathbb{J}|$  because  $E_0 = 0$ . This fact is easy to understand because  $\lambda_1(\gamma; \mathbb{J})$ , with  $\gamma \neq 0$ , is nothing else than the first eigenvalue of the Dirichlet Laplacian in a torus of cross section  $|\mathbb{J}|$  and it is known that the corresponding eigenfunction is invariant w.r.t. the rotations around the point of symmetry  $(0, 0; \mathbb{J})$  is the spectral threshold of an infinite straight tube of cross-section  $|\mathbb{J}|$  which is equal to  $\lambda_1(\gamma; \mathbb{J})$ . In fact, as a direct consequence of a variational formula for the lowest eigenvalue of  $H_0$ , we get the identity

$$\lambda_1(\gamma; \mathbb{J}) = \lambda_1(\gamma); \quad (14)$$

where the latter is given by (11).

Henceforth, we consider  $\mathcal{T} \lambda_1(\gamma)$  as a function on  $(1=a; 1=a)$  and examine its properties by means of the second part of Lemma 2 (an alternative, equivalent, approach is to make the change of trial function  $\mathcal{T}(1 - u_2)^{\frac{1}{2}}$  directly in (11), which makes the denominator of the Rayleigh quotient independent of  $\gamma$ , while the potential  $V_0$  appears in the numerator).

The following result together with Theorem 2 establishes the lower bound (3) of Theorem 1.

**Proposition 1 (Monotonicity).** The function  $\mathcal{T} \lambda_1(\gamma)$  is

- (i) continuous on  $(1=a; 1=a)$ ;

(ii) increasing on  $(-1/a; 0]$ ;

(iii) decreasing on  $[0; 1/a)$ .

Proof. ad (i). This is immediate from the minimax principle applied to  $\hat{H}_0$ .  
ad (ii) and (iii). Calculating

$$\frac{\partial V_0}{\partial u_2}(u_2) = \frac{1}{2(1-u_2)^3};$$

we see that the potential (13) as a function of  $u_2$  is increasing for  $u_2 < 0$  and decreasing for  $u_2 > 0$ . The claim then follows easily by the minimax principle.  $\square$

The following result follows from the fact that the operator  $\hat{H}_0$  is invariant w.r.t. the simultaneous change  $u_1 \rightarrow -u_1$  and  $u_2 \rightarrow -u_2$ .

Proposition 2 (Symmetry). If  $\lambda = \lambda_0$ , then the function  $\lambda_0(\cdot)$  is even on  $(-1/a; 1/a)$ .

We note that  $\lambda_0$ , as an eigenvalue of the Dirichlet Laplacian, has the asymptotics  $\lambda_0 = O(a^{-2})$  as  $a \rightarrow 0$ . Since one is dealing with Dirichlet boundary conditions on  $I \times \mathbb{R}^d$ , one expects the same behaviour from  $\lambda_0(\cdot)$ . We derive the following asymptotics.

Proposition 3 (Thin-width asymptotics). One has

$$\lambda_0(\cdot) = \lambda_0 - \frac{1}{4}a^2 + O(a^4) \quad \text{as } a \rightarrow 0:$$

Proof. Since  $V_0(u_2) = \frac{1}{4}u_2^2 + O(u_2^4)$ ; the result immediately follows by the minimax principle.  $\square$

Finally, applying the Faber-Krahn inequality to  $\lambda_0(\cdot)$  with help of Proposition 1, one obtains the uniform lower bound of Theorem 1.

Proposition 4 (Uniform bound). One has

$$\lambda_0(-1/a; 1/a) \geq c := \frac{\mathcal{P}^{d-1}(\frac{1}{2})}{d \mathcal{P}^{d-1}(\frac{1}{2})} \frac{1}{j_{(d-2)/2,1}^2};$$

where  $j_{(d-2)/2,1}$  denotes the first zero of the Bessel function  $J_{(d-2)/2}$ .

## 5 Conclusions

The main goal of this paper was to derive a lower bound to the spectral threshold of the Laplacian (2) in curved tubes (1). Our Theorem 1 states that this bound is given by  $\lambda_0(\cdot)$ , i.e. the lowest eigenvalue of the Dirichlet Laplacian in a torus of curvature  $\kappa$ , with  $\kappa$  being determined uniquely by the first curvature of the reference curve and the tube cross-section. It follows from Section 4 that

$\lambda_0(\cdot)$  is a decreasing function (cf Proposition 1), i.e. bending diminishes

the lower bound (see also Proposition 3). Another interesting observation is that the lower bound does not depend on higher curvatures of the reference curve (technically, this is a consequence of Lemma 1), i.e. twisting does not diminish the lower bound.

We note that Proposition 2 yields  $\inf_{\mathcal{C}} \lambda_0(k_1, k_1)$  provided  $\ell = \ell_1$ , and Proposition 3 implies asymptotics of the lower bound for thin tubes.

It follows immediately from the minimax principle that the lower bound of Theorem 1 also applies to other boundary conditions imposed on  $L((\partial I) \cup \{0\})$ , e.g., Dirichlet, Robin, periodic, etc.

Adapting the approach of Section 4 to the case of Dirichlet boundary conditions imposed everywhere on  $\partial$ , one reveals interesting isoperimetric inequalities for the first eigenvalue, denoted here by  $\lambda_0^D(\ell; j, j)$ , of the Laplacian in a toroidal segment of curvature  $\ell$ , length  $j$  and cross-section  $j$ . In particular,  $\lambda_0^D(\ell; j, j)$  attains its minimum for  $\ell = 2 = j^2$ , i.e. when  $\ell$  is the whole torus with a supplementary Dirichlet condition imposed on a transverse cross-section  $j$  (cf the beginning of Section 4). This minimum is equal to the first eigenvalue  $\lambda_0$  of the Dirichlet Laplacian in  $j$  and therefore it depends neither on  $j$ , nor on the rotations of  $\ell$ . At the same time, it can be shown that  $\lambda_0^D(\ell; j, j)$  is decreasing on the interval  $[4a^2 = j^2; 2 = j^2]$ . Furthermore, if  $d = 2$ , one can modify the proof of Theorem 2 in [Lau98] and show that the maximum is attained for  $\ell = 0$ , i.e. when  $\ell$  is a rectangle. An open problem is to prove (or disprove) the monotonicity on  $[0; 4a^2 = j^2]$ .

Let us also mention that the lower bound of Theorem 1 is optimal in the sense that the equality is achieved for a tube geometry (a torus or a straight tube). However, the question about an optimal lower bound in an unbounded curved tube is more difficult and remains open.

The hypothesis H2i was discussed in Remark 3. As mentioned in Remark 1, our hypothesis H1i allows us to consider some curves which do not possess a distinguished Frenet frame. However, there still exist curves for which the hypothesis H1i fails; see [Spi79, Chap. 1, p. 34] for an example of such a  $C^1$ -smooth but not analytic curve in  $\mathbb{R}^3$ . Without going into details, let us only mention that the hypothesis H1i is not necessary for the lower bound (3) to hold. For instance, using a Neumann bracketing argument, it suffices to assume that the hypothesis H1i is satisfied "piece-wise"; this may happen if there are isolated points when some of the curvatures vanish.

Let us conclude this paper by comparing the result of Theorem 1 with the lower bound established in [AE90] in the situation when  $I = \mathbb{R}$ ,  $d = 2; 3$ , the cross-section was circular and the discrete spectrum of  $\Delta$  was not empty but finite. The results of [AE90] read as

$$\inf_{\mathcal{C}} \lambda_0 = \begin{cases} 3^{1/N} (j_{0,1} = j_{1,1})^2 & \text{if } d = 2; \\ \min_{j=2;1} \lambda_{j,1}^2 & \text{if } d = 3; N = 1; \end{cases}$$

where  $N$  is the number of discrete eigenvalues (counting multiplicity). Our

uniform lower bound given by Proposition 4 can be written as

$$\inf_{(j_0, j_1) \in \mathbb{N}^2} \lambda_{j_0, j_1}^2 \geq \begin{cases} 0.5860 & \text{if } d = 2; \\ (2 - (3 - \epsilon))^{2/3} \lambda_{j_1=2, j_0=1}^2 \geq 0.6072 & \text{if } d = 3; \end{cases}$$

which is evidently better and applies to tubes with an infinite number, or without any, discrete eigenvalues, too; we also emphasise that we have compared the results of [AE 90] with a crude bound of Proposition 4, a better bound to  $\inf_{(j_0, j_1) \in \mathbb{N}^2} \lambda_{j_0, j_1}^2$  is contained in (3) of our Theorem 1.

## Acknowledgements

The authors thank the referee for helpful suggestions which have improved the presentation. This work was partially supported by FCT/POCTI/FEDER, Portugal, and GAASCR grant TAA 1048101.

## References

- [AB 91] M. S. Ashbaugh and R. D. Benguria, Proof of the Payne-Polya-W einberger conjecture, *Bull. Amer. Math. Soc. (N.S.)* 25 (1991), no. 1, 19{29.
- [AB 92] ———, A second proof of the Payne-Polya-W einberger conjecture, *Comm. Math. Phys.* 147 (1992), no. 1, 181{190.
- [AE 90] M. S. Ashbaugh and P. Exner, Lower bounds to bound state energies in bent tubes, *Phys. Lett. A* 150 (1990), no. 3,4, 183{186.
- [CDFK] B. Chenaud, P. Duclos, P. Freitas, and D. Krejčíř, Geometrically induced discrete spectrum in curved tubes, submitted; preprint available.
- [DE 95] P. Duclos and P. Exner, Curvature-induced bound states in quantum waveguides in two and three dimensions, *Rev. Math. Phys.* 7 (1995), 73{102.
- [ES 89] P. Exner and P. Seba, Bound states in curved quantum waveguides, *J. Math. Phys.* 30 (1989), 2574{2580.
- [EW 01] P. Exner and T. Weidl, Lieb-Thirring inequalities on trapped modes in quantum wires, *Proceedings of the XIIIth International Congress on Mathematical Physics (London 2000)*, Int. Press, Boston, MA, 2001, pp. 437{443.
- [GJ 92] J. Goldstone and R. L. Jaffe, Bound states in twisting tubes, *Phys. Rev. B* 45 (1992), 14100{14107.

- [Hen03] A. Henrot, Minimization problems for eigenvalues of the Laplacian, *Journal of Evolution Equations* 3 (2003), 443{461.
- [Hur00] N. E. Hurt, Mathematical physics of quantum wires and devices, Kluwer, Dordrecht, 2000.
- [KK ] D. Krejčíř and J. Kríž, On the spectrum of curved quantum waveguides, submitted; preprint on [mp-arc 03-264] or [math-ph/0306008].
- [Kli78] W. Klingenberg, A course in differential geometry, Springer-Verlag, New York, 1978.
- [Lau98] R. S. Laugesen, Eigenvalues of the Laplacian on inhomogeneous membranes, *Amer. J. Math.* 120 (1998), 305{344.
- [LCM 99] J. T. Londergan, J. P. Carini, and D. P. Murdoch, Binding and scattering in two-dimensional systems, *LNP*, vol. m 60, Springer, Berlin, 1999.
- [PPW 55] L. E. Payne, G. Polya, and H. F. Weinberger, Sur le quotient de deux fréquences propres consécutives, *C. R. Acad. Sci. Paris* 241 (1955), 917{919.
- [PPW 56] ———, On the ratio of consecutive eigenvalues, *J. Math. Phys.* 35 (1956), 289{298.
- [RB 95] W. Renger and W. Bulla, Existence of bound states in quantum waveguides under weak conditions, *Lett. Math. Phys.* 35 (1995), 1{12.
- [RS72] M. Reed and B. Simon, Methods of modern mathematical physics, I. Functional analysis, Academic Press, New York, 1972.
- [Spi79] M. Spivak, A comprehensive introduction to differential geometry, vol. II, Publish or Perish, Houston, Texas, 1979.