

On Conformal d'Alembert-Like Equations¹

E . C apelas de O liveira
 Departam ento de M atem atica Aplicada
 Im ecc C . P . 6065 U nicam p
 13083-859 Cam pinas (SP) B razil
 E-m ail: capelas@ i n e.unicam p.br
 and
 R . da R ocha
 Instituto de F ísica\Gleb W ataghin"
 IFGW C . P . 6165 U nicam p
 13083-970 Cam pinas (SP) B razil
 E-m ail: roldao@ i .unicam p.br

Abstract

U sing conform al coordinates associated w ith projective conform al relativity we obtain a conform al Klein-Gordon partial differential equation. As a particular case we present and discuss a conform al radial d'Alembert-like equation. As a by-product we show that this radial equation can be identified with a one-dimensional Schrödinger-like equation in which the potential is exactly the second Pöschl-Teller potential.

Introduction

After one of the most important of Einstein's² papers [1] concerning Special Relativity was published, several alternative theories were proposed. Among them, some different interpretations and particular generalizations have been presented. In this paper we are interested in one such theory, namely the theory of hyperspherical universes, proposed by Alcidiacocono [2] several years ago and, more specifically, in the so-called conformal case.

¹To celebrate one century of Special Theory of Relativity.

²The year 1905 is known as the Annus Mirabilis, because in it were published three of the most important papers of the century; Complete Papers on Light Quanta, Brownian Motion, and Special Theory of Relativity.

When we write Maxwell equations in six dimensions, with six projective coordinates (we have, in these coordinates, a Pythagorean metric) a natural problem arises, namely, to provide a physical version of the formalism, i.e. to ascribe a physical meaning to the coordinates. For this theory, there are two possible different physical interpretations: a bitemporal interpretation and a biprojective interpretation. In the first case (bitemporal) we introduce a new universal constant c^0 and the coordinate $x_5 = ic^0 t^0$ where t^0 is interpreted as a second time; we thus obtain in cosmic scale the so-called multitemporal relativity, proposed by Kalitzen [3]. The set of Maxwell equations obtained in this theory generalizes the equations of the unitary theory of electromagnetism and gravitation, as proposed by Corben [4].

On the other hand (our second, biprojective case) we can interpret the extra coordinate, x_5 , as a second projective coordinate. We then obtain the so-called conformal projective relativity, proposed by Arcidiacono [5], which extends in cosmic scale the theory proposed by Ingraham [6], but with a different physical interpretation. In this theory we have another universal constant, r_0 , which can be taken as $r=r_0 = N$, where r is the radius of the hypersphere and N is the cosmological number appearing in the Eddington-Dirac theory [7].

Here we consider only the second alternative, i.e., the biprojective interpretation. With this aim we introduce a projective space P_5 tangent to the hypersphere S^4 . We then introduce six projective coordinates \bar{x}_a , with $a = 0;1;...;5$ and normalized as

$$\bar{x}^2 + \bar{x}_0^2 - \bar{x}_5^2 = r^2;$$

where $\bar{x}^2 = x_i x^i$, $i = 1;...;4$ and r is the radius of the hypersphere. These coordinates allow us to construct the conformal projective relativity, using a six-dimensional tensor formalism.

This paper is organized as follows: in Section 1 we present a review of the so-called theory of hyperspherical universes, proposed by Arcidiacono, considering only the six-dimensional case, in which conformal projective relativity appears as a particular case. The choice of convenient coordinates and the link between the derivatives in these two formulations (a geometric version, six-dimensional, and a physical version, the five-dimensional conformal version) are also presented. After this review, we discuss in Section 2 a Klein-Gordon partial differential equation written in conformal coordinates. In Section 3, we show that a conformal 'radial' d'Alembert-like equation, can be led into a Schrodinger differential equation in which the associated potential is exactly a second Pöschl-Teller potential.

1 Hyperspherical Universes

In 1952 Fantappiè proposed the so-called theory of the universes [8]. This theory is based on group theory and on the hypothesis that the universe is endowed with unique physical laws, valid for all observers. As particular cases, Arcidiacono [2] studied a limitation of that theory, i.e., he considered hyperspherical universes with $3; 4; \dots; n$ dimensions where motions are given by $n(n+1)/2$ -parameter rotation group in spaces with $4; 5; \dots; (n+1)$ dimensions, respectively. Those models of hyperspherical $S^3; S^4; \dots; S^n$ universes, can be interpreted as successive physical improvements, because any one of them (after S^4) contains its precedents and is contained in its successors.

After 1955 Arcidiacono studied the case $n = 4$, special projective relativity, based on the de Sitter hyperspherical universe with a group (the so-called Fantappiè-de Sitter group) of ten parameters. This theory is an improvement (in a unique way) of Einstein's special relativity theory and provides a new group-theoretical version of the big-bang cosmology. As a by-product of special projective relativity one can recover several results, for example, Kinematic Relativity, proposed by Milne [9]; Stationary Cosmology, proposed by Bondi-Gold [10] and Plasma Cosmology, proposed by Alfven [11].

Moreover, if we consider a universe S^4 as globally hyperspherical but endowed with a locally variable curvature, we obtain the so-called general projective relativity which was proposed and studied by Arcidiacono after 1964. This theory allows us to recover several results as particular cases, for example, the unitary theories proposed by Weyl [12], Straneo [13], Kaluza-Klein [14, 15], Veblen [17] and Jordan-Thiry [18] and some generalizations of the gravitational field, as those proposed by Brans-Dicke [19], Rosen [20] and Sciama [21].

In this paper we are interested only in the case $n = 5$, i.e., conformal projective relativity based on the hyperspherical universe S^4 and its associated rotation group, with fifteen parameters, which contains the accelerated motions. We remember that, whereas for $n = 4$ we have a unitary theory (a magnetohydrodynamic field), for $n = 5$ we have another unitary theory, i.e., the magnetohydrodynamics and Newton's gravitation. We also present the relations between Cartesian, projective and conformal coordinates and the link involving derivatives in the six- and five-dimensional formulations.

1.1 Conformal Coordinates

We use the notation x_i , ($i = 1;2;3;4$) and x_5 for conformal coordinates and \bar{x}_a , ($a = 0;1;2;3;4;5$) for projective coordinates. The relations between these coordinates are

$$x_i = r_0 \frac{\bar{x}_i}{\bar{x}_0 + \bar{x}_5} \quad \text{and} \quad x_5 = r_0 \frac{r}{\bar{x}_0 + \bar{x}_5};$$

which satisfy the condition

$$x_5^2 - x^2 = r_0^2 \frac{\bar{x}_0 - \bar{x}_5}{\bar{x}_0 + \bar{x}_5};$$

where $x^2 = x_i x^i$, and r_0 and r are constants. After these considerations, the transformations of the so-called conformal projective group are obtained using the quadratic form in projective coordinates

$$\bar{x}^2 + \bar{x}_0^2 - \bar{x}_5^2 = r^2;$$

decomposing the elements of the six-dimensional rotation group (with fifteen parameters) in fifteen simple rotations $(\bar{x}_a; \bar{x}_b)$ [22].

1.2 Connection Between Derivatives

Our main objective is to write down a differential equation, more precisely a Klein-Gordon-like equation, associated with conformal coordinates. We first obtain the relation between the six projective derivatives $\bar{\partial}_a = \partial/\partial \bar{x}_a$ and the five-dimensional derivatives $\partial_i = \partial/\partial x_i$ and $\partial_5 = \partial/\partial x_5$. We can then write the differential equations in the projective formalism, with six dimensions, in physical, i.e., conformal coordinates, with five dimensions.³

Taking $\phi = \phi(x_i; x_5)$, a scalar field, and using the chain rule we can write

$$\begin{aligned} \partial_i &= (\bar{\partial}_i \bar{x}_k) \bar{\partial}_k + (\bar{\partial}_i \bar{x}_5) \bar{\partial}_5 + (\bar{\partial}_i \bar{x}_0) \bar{\partial}_0 \\ \partial_5 &= (\bar{\partial}_5 \bar{x}_k) \bar{\partial}_k + (\bar{\partial}_5 \bar{x}_5) \bar{\partial}_5 + (\bar{\partial}_5 \bar{x}_0) \bar{\partial}_0 \end{aligned}$$

with $\bar{\partial} = \bar{\partial}(\bar{x}_i; \bar{x}_5; \bar{x}_0)$ and $i; k = 1;2;3;4$:

From now on we take $r = 1 = r_0$. We consider $\bar{\phi}(\bar{x}_a)$ a homogeneous function with degree N in all six projective coordinates \bar{x}_a . Using Euler's theorem associated with homogeneous function, we get

$$\bar{x}_i \bar{\partial}_i + \bar{x}_5 \bar{\partial}_5 + \bar{x}_0 \bar{\partial}_0 = N$$

³As we already know, in five dimensions we must impose a condition on space in order to account for the fact that we are aware of only four dimensions. We have the same situation here, i.e., we must impose an additional condition.

where $\bar{\partial}_a = \partial/\partial \bar{x}_a$ and N is the degree of homogeneity of the function.

Then, the link between the derivatives can be written as follows [23]

$$\begin{aligned}\bar{\partial}_0^2 &= N \frac{A^+}{x_5} + B^+ \partial_5 - x_5 x_i \partial_i \\ \bar{\partial}_5^2 &= N \frac{A}{x_5} - B^+ \partial_5 - x_5 x_i \partial_i \\ \bar{\partial}_i^2 &= N \frac{x_i}{x_5} + x_i \partial_5 + x_5 \partial_i\end{aligned}$$

where we have introduced a convenient notation

$$2A = 1 - x^2 - x_5^2 \quad \text{and} \quad 2B = 1 - x^2 - x_5^2.$$

We observe that for $\bar{x}_5 = 0$ and considering $\bar{\partial}_5^2 = 0$ we obtain

$$\begin{aligned}\bar{\partial}_i^2 &= A \partial_i + \frac{N}{A} x_i \\ \bar{\partial}_0^2 &= A x_i \partial_i + \frac{N}{A}\end{aligned}$$

where $A^2 = 1 - x^2$. These expressions are the same expressions obtained in special projective relativity [25, 26] and provide the link between the projective derivatives and the four derivatives in Cartesian coordinates, i.e., the relation between five-dimensional (de Sitter) universe and four-dimensional (Minkowski) universe.

2 Conformal Klein-Gordon Equation

In this section we use the previous results to calculate the so-called generalized Klein-Gordon differential equation

$$\frac{\partial^2}{\partial \bar{x}_a^2} + m^2 = 0$$

where m^2 is a constant and $a = 0; 1; \dots; 5$. Introducing projective coordinates (in this case we have a Pythagorean metric) we obtain⁴

$$\frac{\partial^2 U}{\partial \bar{x}_i^2} + \frac{\partial^2 U}{\partial \bar{x}_0^2} - \frac{\partial^2 U}{\partial \bar{x}_5^2} + m^2 U = 0$$

⁴ Hereafter we consider $m = m_0 c \sim$ where m_0 , c and \sim have the usual meanings.

where $i = 1; 2; 3; 4$ and $U = U(\bar{x}_i; \bar{x}_0; \bar{x}_5)$.

Using the relations between projective and conformal coordinates and the link (involving the derivatives) in the two formulations we can write

$$x_5^2 \frac{\partial^2}{\partial x_5^2} + 3x_5 \frac{\partial}{\partial x_5} + N(N+5) + m^2 u(x_i; x_5) = 0$$

where N and m^2 are constants, \square is the D'Alembertian operator given by

$$\square = \frac{1}{c^2} \frac{\partial^2}{\partial t^2}$$

and Δ is the Laplacian operator. This partial differential equation is the so-called Klein-Gordon differential equation written in conformal coordinates or a conformal Klein-Gordon equation.

The case $m^2 = 0$ transforms this equation in the so-called generalized d'Alembert differential equation. Another way to obtain this differential equation is to consider the conformal metric in cartesian coordinates, which furnishes the so-called Beltrami metric [2] where the d'Alembert equation appears naturally. This equation can also be obtained by means of the second order Casimir invariant operator⁵ associated with the conformal group.

To solve the conformal Klein-Gordon equation, we first introduce the spherical coordinates $(r; \theta; \phi)$ and get

$$\frac{\partial^2 u}{\partial r^2} + \frac{2}{r} \frac{\partial u}{\partial r} + \frac{1}{r^2} L u - \frac{1}{c^2} \frac{\partial^2 u}{\partial t^2} - \frac{\partial^2 u}{\partial x_5^2} + \frac{3}{x_5} \frac{\partial u}{\partial x_5} + \frac{u}{x_5^2} = 0;$$

where we introduced $x_4 = ict$ and defined the operator⁶

$$L = \frac{\partial^2}{\partial \theta^2} + \cot \theta \frac{\partial}{\partial \theta} + \frac{1}{\sin^2 \theta} \frac{\partial^2}{\partial \phi^2}$$

involving only the angular part. In this partial differential equation we have $u = u(r; \theta; \phi; t; x_5)$ with $\square = N(N+5) + m^2$.

Using the method of separation of variables we can eliminate the temporal and angular parts, writing

$$u = u(r; \theta; \phi; t; x_5) = A e^{inct} Y_{lm}(\theta; \phi) f(r; x_5);$$

⁵ Invariant operators associated with dynamic groups furnish mass formulas, energy spectra and, in general, characterize specific properties of physical systems.

⁶ Here r is a coordinate and should not be confused with the radius of the hypersphere. Besides, it is always possible to define a Wick-rotation [24] of the time coordinate, i.e., $ct \rightarrow ict$.

where A is an arbitrary constant, $n > 0$, $\ell = 0; 1; \dots$ and $m = 0; 1; \dots$ with $Y_{\ell m}(\theta; \phi)$ and $Y_m(\ell; \phi)$ are the spherical harmonics, we get the following partial differential equation

$$\frac{\partial^2 f}{\partial r^2} + \frac{2}{r} \frac{\partial f}{\partial r} - \frac{\partial^2 f}{\partial x_5^2} + \frac{3}{x_5} \frac{\partial f}{\partial x_5} + \frac{f}{x_5^2} + n^2 - \frac{\ell(\ell+1)}{r^2} f = 0$$

with $f = f(r; x_5)$. If we impose a regular solution at the origin ($r \neq 0$), the solution of this partial differential equation can be obtained in terms of a product of two Bessel functions [22].

3 Ad'A lem bert-Like Equation

In this section we present and discuss a partial differential equation which can be identified to a d'A lem bert-like equation, which we call a conformal 'radial' d'A lem bert equation. We firstly introduce a convenient new set of coordinates, then we use separation of variables and obtain two ordinary differential equations. One of them can be identified as an ordinary differential equation whose solution is a generalization of Newton's law of gravitation; the other one is identified with an ordinary differential equation similar to a one-dimensional Schrodinger differential equation with a potential equal to the second P oschl-Teller potential.

We introduce the following change of independent variables

$$\begin{aligned} r &= \cosh \xi; \\ x_5 &= \sinh \xi; \end{aligned}$$

with $\xi > 0$ and $\xi = 0$, in the separated Klein-Gordon equation, obtained in the previous section, and after another separation of variables we can write a pair of ordinary differential equations, namely,

$$2 \frac{d^2 U}{d \xi^2} - p(p+1)U = 0;$$

where $U = U(\xi)$ and

$$\frac{d^2 V}{d \eta^2} + (2 \tanh \eta - 3 \coth \eta) \frac{dV}{d \eta} + \frac{\ell(\ell+1)}{\cosh^2 \eta} - \frac{p(p+1)}{\sinh^2 \eta} V = 0$$

where $V = V(\eta)$ and p is a separation constant.

We first discuss the differential equation in the variable ξ . Its general solution is given by

$$U(\xi) = C_1 \xi^{-p} + C_2 \xi^{p+1}$$

where C_1 and C_2 are arbitrary constants.

If we consider the case $p = 1$, introducing the notation $C_1 = gM$ with g and M having the usual meanings, we get

$$U(r) = \frac{gM}{r} + C_2 r^2$$

i.e., a gravitational potential which can be interpreted as a sum of a Kepler-like potential and a harmonic oscillator potential, giving rise to the gravitational force

$$f(r) = -r U' = \frac{gM}{r^2} + 2C_2 (r^2 - r_5^2)^{1/2};$$

with a singularity at $r = r_5$. We note that for $C_2 = 0$ we obtain an expression analogous to Newton's law of gravitation.

Secondly, the equation in the variable ψ . To solve this ordinary differential equation we first introduce the change of dependent variable

$$V(\psi) = \sinh^2 \frac{\psi}{2} \tanh^2 \frac{\psi}{2} F(\psi)$$

and obtain

$$\frac{d^2}{d\psi^2} F(\psi) + \frac{(\lambda - 1)}{\sinh^2 \frac{\psi}{2}} F(\psi) + \frac{(\lambda + 1)}{\cosh^2 \frac{\psi}{2}} F(\psi) + \left(p + \frac{1}{2}\right)^2 F(\psi) = 0; \quad (1)$$

where the parameter λ is given by a root of the following algebraic equation $(\lambda - 1) = N^2 + 5N + 15 = 4 + m^2$.

The differential equation above can be identified with a Schrodinger-like differential equation in which the associated potential is given by

$$V_{\lambda}(\psi) = \frac{(\lambda - 1)}{\sinh^2 \frac{\psi}{2}} - \frac{(\lambda + 1)}{\cosh^2 \frac{\psi}{2}};$$

which is exactly the second Pöschl-Teller potential with energy E given by $E_p = -(p + 1/2)^2 < 0$. The solution of this ordinary differential equation is well known and can be expressed in terms of the hypergeometric function. An algebraic treatment can be found in [27, 28].

We note that the first Pöschl-Teller potential is connected with the study of a Dirac particle on central backgrounds associated with an anti-de Sitter oscillator, i.e., the transformed radial wave functions satisfy the second-order Schrodinger differential equation whose potential is exactly the first Pöschl-Teller potential [29].

Finally, a particular case of Eq.(1), i.e., the case $\lambda = 0$, is related to the anti-de Sitter static frame as shown recently by da Rocha and Capelas de Oliveira [30].

Concluding Remarks

In this paper we discussed the calculation of a conformal d'Alembert-like equation. We used the methodology of projective relativity to obtain a conformal Klein-Gordon differential equation and, after the separation of variables, we got another partial differential equation in only two independent variables, the so-called conformal d'Alembert differential equation. Another separation of variables led to an ordinary differential equation which generalizes Newton's law of gravitation. Finally, we showed that the remaining differential equation, a 'radial' differential equation, is transformed into a one-dimensional Schrodinger differential equation with an associated potential that can be identified exactly with the second Pöschl-Teller potential.

From supersymmetric quantum mechanics with periodic potentials, it can be seen that the most general periodic potentials which can be analytically solved involve Jacobi's elliptic functions, which in various limits become Pöschl-Teller potentials arising in the context of Kaluza-Klein spectrum [14]. Kaluza-Klein modes of the graviton have been widely investigated [31, 32, 33, 34], since the original formulation of Randall and Sundrum necessarily has a continuum of Kaluza-Klein modes without any mass gap, arising from a periodic system of 3-branes. The methods and equations developed here can shed some new light in the calculation of mass gaps from a distribution of D-branes [34] in the context of five-dimensional supergravity in a forthcoming paper.

A natural continuation of this calculation is to prove that all 'radial' problems associated with an equation resulting from a problem involving a light cone can be led into a Schrodinger-like differential equation in which the potential is exactly the Pöschl-Teller potential [35].

Acknowledgment

One of us (RR) is grateful to CAPES for financial support. We are also grateful to Dr. Geraldo A. Gosti for interesting and useful discussions and Dr. J. Emilio Maiorino for several suggestions about the paper.

References

- [1] A .E instein, Zur E lektrodynam ik B ewegter K orper, *Annalen D er P hysik*, 17, 891, (1905).
- [2] G .A rcidiacono, La Teoria D egli U niversi, Volum e II, D i R enzo Editore, Rom a, (2000).
- [3] N .K alitzen, M ultitemporal Theory of Relativity, *Bulgarian A cadem y of Sciences, So a*, (1975).
- [4] H .C .C orben, A C lassical Theory of E lectrom agnetism and G ravitation, *Phys.Rev.* 69, 225-234, (1946).
- [5] G . A rcidiacono, G li U niversi I persferici, il G ruppo C onform e e il C am po G ravitazionale di Newton, *Collectanea M athem atica*, 36, 119-135, (1985).
- [6] R . L . Ingraham , C onform al G eom etry and E lem entary P articles, *N . C im ento*, 12, 825-851, (1954).
- [7] A . S . Eddington, *Space, T im e and G ravitation*, C am bridge University, C am bridge, (1920).
- [8] L . Fantappiè, *Opere Scelte*, Unione M atem atica Italiana, Bologna, (1973).
- [9] E . A . M ilne, *Relativity, G ravitation and W orld Structure*, C larendon P ress, O xford, (1935); *K inem atic Relativity*, C larendon P ress, O xford, (1948).
- [10] H . Bondi and T . G old, The Steady-State Theory of the Expanding U ni-verse, *M on.Nat.Roy.A stron.Soc.*, 108, 252-270, (1948).
- [11] H . A lfven, On H ierarchical C osm ology A strophysics, *A strophys. Space Sci.*, 89, 313-324, (1983) and M odel of the P lasm a U niverse, *IEEE Trans.P lasm a Sci.*, 14, 629-638, (1986).
- [12] H . W eyl, *Raum-Zeit-M aterie*, Berlin, (1918). *Space-T im e-M atter*, Dover, New York, (1952); *G ravitation und E lektrizitat*, *Sitzungs-berichte P reussische A kadem ie W issenschaften P hys.M ath.*, K 1, 465-480, (1918); *E ine neue E rweiterung der Relativitatstheorie*, *Ann. der P hysik*, 59, 101-133, (1919).

- [13] P. Straneo, *Teorie Unitarie Della Gravitazione e Dell'Elettricità*, N. Cimento, 8, 125-145, (1931).
- [14] T. Kaluza, *Zum Unitätsproblem in der Physik*, Sitzungsberichte Preussische Akademie Wissenschaften Phys. Math., K 1, 966-972, (1921). (Communicated to Einstein in 1919.) Also found in English translation in ref.[16] pages 53-58.
- [15] O. Klein, *The Atomicity of Electricity as a Quantum Theory Law*, Nature, 118, 516 (1926); *Zur Fünfdimensionalen Darstellung der Relativitätstheorie*, Z. Phys., 46, 188-208, (1928). Also found in English translation in ref.[16] pages 59-68.
- [16] Lochlainn O'Raiífeartaigh, Editor. *The Dawning of Gauge Theory*, Princeton University Press, Princeton, N.J., (1997). Edited and with introductory essays by Lochlainn O'Raiífeartaigh.
- [17] O. Veblen, *Projektive Relativitätstheorie*, Springer, Berlin, (1933).
- [18] P. Jordan, *Erweiterung der Projektiven Relativitätstheorie*, Ann. Phys. (Leipzig), 1, 219-228, (1947) and Y. Thiry, *Etude Mathématique des Equations d'une Théorie Unitaire a Quinze Variables de Champ*, J. Math. Pures et Appl., 30, 275-396, (1951).
- [19] C. Brans and R. H. Dicke, *Mach's Principle and a Relativistic Theory of Gravitation*, Phys. Rev. D, 124, 925-935, (1961).
- [20] N. Rosen, *A Bimetric Theory of Gravitation*, Gen. Rel. Grav., 4, 435-447, (1973); *Bimetric General Relativity and Cosmology*, Gen. Rel. Grav., 12, 493-510, (1980).
- [21] D. W. Sciama, *On a Non-Symmetric Theory of Pure Gravitational Field*, Proc. Camb. Philos. Soc., 54, 72-80, (1958).
- [22] E. Capelas de Oliveira, *Sobre Transformações Conformes*, (In Portuguese), Private communication, (2005).
- [23] G. Arcidiacono and E. Capelas de Oliveira, *Conformal Relativity and d'Alembert Equation*, in *New Frontiers in Relativities*, Edited by Tepper L. Gill, Hadronic Press, 297-302, (1996).
- [24] J. J. Sakurai, *Modern Quantum Mechanics*, Addison-Wesley, Boston, (1985).

- [25] E. Capelas de Oliveira and E. A. Notte Cuello, A New Construction of the Casimir Operator for the Poincaré-de Sitter Group, *H. Journal*, 18, 181-189, (1995).
- [26] D. Gomes and E. Capelas de Oliveira, The Second Order Klein-Gordon Field Equation, *Int. J. Math. Math. Sci.* 69, 3775-3781, (2004) and references therein.
- [27] A. O. Barut, A. Inomata and R. Wilson, Algebraic Treatment of Second Pöschl-Teller, Morse-Rosen and Eckart Equations *J. Phys. A: Math. Gen.*, 20, 4083-4096, (1987).
- [28] A. O. Barut, A. Inomata and R. Wilson, A New Realisation of Dynamical Groups and Factorisation Method, *J. Phys. A: Math. Gen.*, 20, 4075-4083, (1987).
- [29] I. I. Cotaescu, The Dirac Particle on Central Backgrounds and the Anti-de Sitter Oscillator, *Int. J. Mod. Phys.*, 13A, 2923-2935, (1998).
- [30] R. da Rocha and E. Capelas de Oliveira, The Casimir Operator of $SO(1;2)$ and the Pöschl-Teller Potential: An AdS Approach, *Rev. Mex. Fis.*, To Appear (2005).
- [31] L. Randall and R. Sundrum, A large mass hierarchy from a small extra dimension, *Phys. Rev. Lett.* 83, 3370-3373, (1999) [[hep-ph/9905221](#)]; An alternative to compactification, *Phys. Rev. Lett.* 83, 4690-4693, (1999) [[hep-th/9906064](#)].
- [32] H. Hatanaka, M. Sakamoto, M. Tachibana and K. Takenaga, M any-brane extension of the Randall-Sundrum solution, *Prog. Theor. Phys.* 102, 1213-1218, (1999) [[hep-th/9909076](#)].
- [33] A. Brandhuber and K. Sfetsos, Non-standard compactifications with mass gaps and Newton's law, *J. High Energy Phys.* 10, 13-32, (1999) [[hep-th/9908116](#)].
- [34] S. Nam, Mass gap in Kaluza-Klein spectrum in a network of brane worlds, *J. High Energy Phys.* 04, 2-8, (2000).
- [35] E. Capelas de Oliveira, A Radial-Like Equation and the Light Cone, to be submitted for publication (2005).