

A sym ptotic corrections to the eigenvalue density of the G U E and L U E

T . M . Garoni

Institute for M athem atics and its Applications, University of M innesota,
400 Lind Hall, 207 Church Street S.E., M inneapolis, M N 55455-0436, USA

P . J. Forrester

D epartm ent of M athem atics and Statistics, University of M elbourne, Parkville, V ictoria 3010, Australia^y

N . E . Frankel

School of Physics, University of M elbourne, Parkville, V ictoria 3010, Australia^z

(D ated: January 4, 2022)

W e obtain correction terms to the large N asym ptotic expansions of the eigenvalue density for the Gaussian unitary and Laguerre unitary ensembles of random $N \times N$ matrices, both in the bulk of the spectrum and near the spectral edge. This is achieved by using the well known orthogonal polynomial expression for the kernel to construct a double contour integral representation for the density, to which we apply the saddle point method. The main correction to the bulk density is oscillatory in N and depends on the distribution function of the limiting density, while the corrections to the Airy kernel at the soft edge are again expressed in terms of the Airy function and its first derivative. We demonstrate numerically that these expansions are very accurate. A matching is exhibited between the asym ptotic expansion of the bulk density, expanded about the edge, and the asym ptotic expansion of the edge density, expanded into the bulk.

PACS num bers: 02.50.Cw, 05.90.+m, 02.30.Gp

I. INTRODUCTION

We consider in this paper two classical ensembles of random matrices, the Gaussian unitary ensemble (GUE), and the Laguerre unitary ensemble (LUE). These ensembles can be characterized by their joint eigenvalue probability density functions

$$P_N(x_1; \dots; x_N) / \prod_{l=1}^N l_N(x_l) \prod_{1 \leq j < k \leq N} (x_k - x_j)^2; \quad x_l^2; \quad (1)$$

with

$$l_N(x) = \begin{cases} \exp(-2Nx^2); & \text{GUE;} \\ x \exp(-4Nx); & \text{LUE;} \end{cases} \quad (2)$$

and

$$= \begin{cases} R; & \text{GUE;} \\ (0; 1); & \text{LUE;} \end{cases} \quad (3)$$

The GUE consists of $N \times N$ Hermitian matrices with independent normally distributed entries on and above the diagonal. It is the cornerstone of random matrix theory^{1,2,3}. The LUE has fundamental applications in mathematical statistics and quantum field theory since it includes Wishart matrices and the Chiral GUE as special cases (the latter after a straightforward change of variables); see e.g. Ref.¹.

We are interested in the large N behavior of the marginal eigenvalue probability density $N_N(x)$, which we hereafter refer to simply as "the density", and which is defined by

$$N_N(x) := \int_{N-1}^Z P_N(x; x_2; \dots; x_N) dx_2 \dots dx_N : \quad (4)$$

The function $N_N(x)$ can be interpreted as the number density of eigenvalues near the point x . We also remark that for the GUE, $N_N(x)$ is equal to the number density of a hamonically trapped system of either non-interacting fermions or impenetrable bosons⁴. There is a similar interpretation for the LUE in terms of a Calogero-Sutherland

type model⁵. For recent advances in asymptotic questions related to these interpretations, complementary to the present study, see Refs.^{6,7,8}

As background to the present study we note that aspects of the large N form of $N(x)$ first arose in studies of field theories related to Hermitian matrix models⁹. There, for the GUE the large N asymptotic expansion of the moments

$$m_N(p) = \int_{\mathbb{R}} x^p N(x) dx \quad (p = 1; 2; \dots)$$

was sort. By a graphical expansion of the matrix integral, involving cataloging the corresponding maps according to their genus, it was predicted that for certain coefficients $a_{2j}(p)$,

$$m_N^{\text{GUE}}(p) = \sum_{j=0}^{p/2} \frac{a_{2j}(p)}{N^{2j}}; \quad (p = 2; 4; \dots) \quad (5)$$

(the odd moments of course vanish). Analogous considerations in the case of the LUE¹⁰ show that

$$m_N^{\text{LUE}}(p) = \sum_{j=0}^{p/2} \frac{a_j(p; \cdot)}{N^j} \quad (6)$$

for certain coefficients $a_j(p; \cdot)$. Observe in particular that (5) contains only even inverse powers in N , while (6) contains both even and odd inverse powers in N .

The graphical methods allow $a_0(p)$ in (5) and $a_0(p; \cdot)$ in (6) to be computed in terms of binomial coefficients for all $p = 0; 1; \dots$. This knowledge in turn can be used (see e.g.¹) to prove that in the limit $N \rightarrow \infty$ and with x fixed

$$N(x) = \lim_{N \rightarrow \infty} N(x) = \begin{cases} \frac{2}{\pi} \sqrt{1-x^2} & ; x \in [-1; 1]; \text{ GUE}; \\ \frac{2}{\pi} \frac{1}{x} \sqrt{1-x^2} & ; x \in (0; 1]; \text{ LUE}; \\ 0; & \text{otherwise}; \end{cases} \quad (7)$$

The first functional form in (7) is referred to as the Wigner semi-circle law, while the second is sometimes named after Marcenko-Pastur. See e.g. Refs.^{1,2,3}.

The expansions (5) and (6) provide a motivation to undertake a study of the asymptotic form of (4). In the case of the GUE such a result has been given by Kalish and Braak¹¹. It states that for $|x| < 1$ and x fixed

$$N(x) = (x) \frac{2 \cos[2N P(x)]}{3 - 2(x)} \frac{1}{N} + O\left(\frac{1}{N^2}\right); \quad (8)$$

where

$$P(x) = 1 + \frac{x}{2} (x) - \frac{1}{2} A \cos(x):$$

Thus one sees that unlike the situation with the moments (5), the leading correction term is $O(1/N)$. Of course this term is oscillatory so one might anticipate that after integration it contributes at a higher order. However inspection of (8) reveals that the situation is more complex: the oscillatory term is not integrable at the endpoints of the support $|x| = 1$. Indeed, it is well known that with the boundary of the eigenvalue support taken as the origin, a scaling regime distinct from that of the bulk becomes relevant. Explicitly, with $Ai(x)$ denoting the Airy function, it has been proved that¹²

$$\begin{aligned} \lim_{N \rightarrow \infty} \frac{N^{1/3}}{2} m_N^{\text{GUE}} - 1 + \frac{1}{2N^{2/3}} &= \lim_{N \rightarrow \infty} \frac{(2N)^{1/3}}{2} m_N^{\text{LUE}} - 1 + \frac{1}{(2N)^{2/3}} \\ &= [Ai(\cdot)]^2 - [Ai(\cdot)]^2; \end{aligned} \quad (9)$$

where \cdot is fixed. In view of the breakdown of (8) in the vicinity of the spectrum edge, (referred to as the soft edge, since although it defines the edge of the support of (x) , for any finite N there is a nonzero probability of finding eigenvalues lying beyond it), we are thus led to also investigate the large N asymptotic expansion extending the limit law (9).

At a technical level, the main achievement of this paper is the derivation of the first correction terms to the limit laws (7) and (9). We do this by utilizing the well known orthogonal polynomial expression for $\rho_N(x)$ to obtain a double integral representation which is amenable to the saddle point method. In the bulk, i.e. in the interior of the support of $\rho(x)$, we show that the asymptotic series progresses in powers of $1/N$, and we obtain the explicit form of the $1/N$ correction. We find for the LUE that the coefficient of the $1/N$ term consists of a component which is oscillatory in N as well as a component which is non-oscillatory in N , whereas for the GUE it consists of only an oscillatory component, as shown in (8). For the soft edge we will see that the asymptotic series progresses in powers of $N^{-1/3}$, and we obtain explicit expressions for the coefficients of the $N^{-1/3}$ and $N^{-2/3}$ terms, which again involve Airy functions.

Due to the similarity in the structure of $\rho_N(x)$ for the GUE and LUE, it is convenient to consider both cases simultaneously to avoid unnecessary repetition, and so at each step of our presentation we discuss the GUE and LUE in parallel. In Section II we discuss the double contour integral expression for $\rho_N(x)$ to which we shall apply the saddle point method. Section III contains our discussion of the asymptotics of $\rho_N(x)$ in the bulk while Section IV discusses the soft edge. In Section V we discuss the extent to which our expansions in the bulk match up with those for the soft edge.

II. CONTOUR INTEGRAL EXPRESSION FOR $\rho_N(x)$

For the unitary ensembles there is a well known and very neat expression for $\rho_N(x)$ in terms of orthogonal polynomials, valid for any N and $x \in \mathbb{R}$. If we let $f_j(x)_{j=0}^N$ denote the monic polynomials orthogonal with respect to $\rho_N(x)$ on \mathbb{R} , then

$$\rho_N(x) = \frac{!_N(x)}{N k_N \frac{1}{N} k^2} [\frac{0}{N}(x) \rho_{N-1}(x) - \frac{0}{N-1}(x) \rho_N(x)] : \quad (10)$$

The norm in the denominator of (10) is just the L^2 -norm associated with $\rho_N(x)$ and \mathbb{R} . For a derivation of (10) the reader is referred to Refs.^{1,2,3}. We remark at this point that there is no universally agreed scale by which the GUE and LUE are defined. To match our choice of scale and notation in (2) to that employed in Ref.¹ for instance, we observe that

$$\rho_N(x) = \begin{cases} \frac{1}{2N} P_{N-1/2} \left(\frac{x}{\sqrt{2N}} \right); & \text{GUE;} \\ \frac{1}{4N} P_N(4Nx); & \text{LUE;} \end{cases}$$

where $P_N(x;y)$ (not to be confused with our definition (1) above) is the kernel defined in Chapter 4 of Ref.¹ (the kernel is often also denoted $K_N(x;y)$ in the literature).

To investigate the large N behavior of $\rho_N(x)$ it is obviously advantageous to start with the expression (10) rather than with the $(N-1)$ -fold integral (4). The $\rho_{N+j-1}(x)$ can be expressed in terms of the standard Hermite and Laguerre polynomials found in Szegő's classic book¹³ as follows

$$\rho_{N+j-1}(x) = \begin{cases} \frac{1}{2^{3(N+j-1)/2}} H_{N+j-1} \left(\frac{x}{\sqrt{2(N+j-1)}} \right); & \text{GUE;} \\ \frac{1}{(N+j-1)! (4N)^{N+j-1}} L_{N+j-1}^{(N)}(4Nx); & \text{LUE;} \end{cases} \quad (11)$$

The required asymptotic expansions of the scaled Hermite and Laguerre polynomials appearing in (11) are known to any order both in the bulk and near the soft edge^{14,15}, and such asymptotic expansions of scaled orthogonal polynomials are now generically said to be of Plancherel-Rotach type (Plancherel and Rotach were the first to compute such asymptotics for the Hermite polynomials). It is reasonable to assume that the most straightforward procedure to obtain the desired asymptotic corrections for $\rho_N(x)$ in each region of interest is to simply insert the corresponding asymptotic expansion for $\rho_{N+j-1}(x)$ into (10). While this is certainly legitimate in principle, and does indeed recover the leading term fairly easily, to derive the correction terms it turns out that such a procedure is rather tedious, and provides little if any insight into the resulting expressions. The source of the complication is that the asymptotic expansions for $\rho_{N+j-1}(x)$ contain a large amount of superfluous information which is canceled when the expansions are substituted into (10). To avoid this, we shall pursue a related, but more direct route.

The Plancherel-Rotach asymptotics for the Hermite and Laguerre polynomials were originally derived by first expressing the polynomials in terms of contour integrals, and then applying the saddle point method. By suitably

massaging the standard results in Szego's book¹³ one finds that

$$c_j(N) = \begin{cases} \frac{1}{2^N} \int_{-2}^2 e^{2Nzx} \frac{e^{-Nz^2/2}}{z^{N+j}} dz; & \text{GUE;} \\ (-1)^{N+j-1} c_j(N) \frac{1}{2^N} \int_{-2}^2 e^{-2Nzx} \frac{(z+2)^{N+1}}{z^{N+1}} dz; & \text{LUE;} \end{cases} \quad (12)$$

$$c_j(N) = \frac{(N+j-1)!}{(2N)^{N+j-1}}; \quad (13)$$

In both cases the contour of integration is a closed positively oriented contour which encircles the origin; in the Laguerre case we further demand that it not contain the point $z = -2$.

Instead of applying the saddle point method to (12) and then substituting the expansions into (10), we shall first insert the contour integrals (12) into (10) to obtain a double integral expression for $J_N(x)$, and then perform the saddle point method on this double integral. To highlight the similarity between the GUE and LUE it is convenient in the GUE case to substitute the contour integral for $c_{N+j-1}(x)$ into (10) rather than that for $c_{N+j-1}(x)$; since $H_j(x) = (-1)^j H_j(x)$ this abuse is perfectly harmless. This results in

$$J_N(x) = 2 \frac{c_0(N) c_1(N)}{k_{N-1} k^2} J_N(x) J_N(x); \quad (14)$$

where

$$J_N(x) = \frac{1}{2^N} \int_{-2}^2 \frac{dz_1}{2-i} \int_{-2}^2 \frac{dz_2}{2-i} e^{N S(z_1; x) + N S(z_2; x)} G(z_1; z_2) \quad (15)$$

and

$$S(z; x) = \begin{cases} 2zx - \log(z) - z^2/2; & \text{GUE;} \\ 2zx - \log(z) + \log(1+z)/2; & \text{LUE;} \end{cases} \quad (16)$$

$$G(z_1; z_2) = u(z_1) u(z_2) \begin{cases} 1 - \frac{z_1}{z_2}; \\ \end{cases} \quad (17)$$

$$u(z) = \begin{cases} 1; & \text{GUE;} \\ (1+z)^{-1}; & \text{LUE;} \end{cases} \quad (18)$$

The remainder of this paper will involve a careful asymptotic analysis of the double integral (15).

Before proceeding we note that it is straightforward to show, using standard results in the orthogonal polynomial literature¹³, that

$$k_{N-1} k^2 = \begin{cases} \frac{2^{2N-3/2}}{N!} \frac{N^{N+1/2}}{N!}; & \text{GUE;} \\ \frac{(4N)^{2N+1}}{(N) (N+1)}; & \text{LUE;} \end{cases} \quad (19)$$

and hence the asymptotics of the prefactors in (14) is

$$2 \frac{c_0(N) c_1(N)}{k_{N-1} k^2} = \begin{cases} \frac{1}{2^N} \frac{1}{N^{3/2}} \frac{1}{N!}; & \text{GUE;} \\ \frac{4^{N+1}}{(N+1)}; & \text{LUE;} \end{cases} \quad (20)$$

$$= \begin{cases} 2N e^{-N} \left(1 + \frac{1}{12N} + O\left(\frac{1}{N^2}\right) \right); & \text{GUE;} \\ 4^{N+1} \frac{(1)}{2N} + O\left(\frac{1}{N^2}\right); & \text{LUE;} \end{cases} \quad (21)$$

Saddle points

Before applying the saddle point method to (15) we need to identify and classify the saddle points of (16). The functions $S(z; x)$ in general have two saddle points at $z = z^*$ where

$$z^* = \begin{cases} x - i \sqrt{x^2 - 1}; & \text{GUE;} \\ 1 - i \sqrt{x^2 - 1}; & \text{LUE;} \end{cases} \quad (22)$$

and

$$x = \begin{cases} \frac{1}{2} \sqrt{1 - x^2}; & \text{GUE;} \\ \frac{1}{x} - 1; & \text{LUE;} \end{cases} \quad (23)$$

We note that for both the GUE and LUE we have

$$x = \frac{1}{2} \sqrt{x^2 - 1}; \quad \text{for } |x| > 1; \quad (24)$$

with x as defined in (7).

Since for $|x| > 1$ we have

$$\frac{S^0(z; x)}{2} = \begin{cases} x e^{-i(A \arcsin(x))}; & \text{GUE;} \\ 2x^2 - x e^{-i\pi/2}; & \text{LUE;} \end{cases} \quad (25)$$

the saddle points $z = z^*$ are both simple when $|x| < 1$, i.e. $S^0(z; x) \neq 0$. However, when $x = 1$ the two simple saddle points given in (22) coalesce to $z = 1$ and $S^0(z; 1)$ vanishes, so we obtain one double saddle point in this case. Thus we already see why the regions $|x| < 1$ and $x = 1$ have qualitatively distinct asymptotic behavior. Simple saddle points generically produce Gaussian integrals whereas double saddle points generically produce Airy functions (see e.g. Ref.¹⁶).

III. BULK ASYMPTOTICS FOR THE GUE AND LUE

In the bulk of the spectrum, i.e. for $|x| < 1$, we hold x fixed and investigate the asymptotics of (15) as N becomes large. From (22) we see that there are two distinct simple saddle points of $S(z; x)$, which form a complex conjugate pair in this case. Let's define $S_{\pm} = S(z_{\pm}; x)$. Then since

$$\text{Re}(S_+) = \text{Re}(S_-); \quad (26)$$

both saddle points contribute to the same order and we deform our contour through both of them. Denoting by a contour passing through z^* along a path of steepest descent, the standard arguments of the saddle point method (see for e.g. Ref.¹⁶) yield

$$J_N(x) = \int_{\gamma_+} \frac{dz_1}{2 - i} + \int_{\gamma_-} \frac{dz_1}{2 - i} + \int_{\gamma_+} \frac{dz_2}{2 - i} + \int_{\gamma_-} \frac{dz_2}{2 - i} e^{N S(z_1; x) + N S(z_2; x)} G(z_1; z_2) + O(e^{2N \text{Re}(S_+) - N}); \quad (27)$$

$$= \int_{\gamma_+} \frac{dz_1}{2 - i} + \int_{\gamma_-} \frac{dz_1}{2 - i} + \int_{\gamma_+} \frac{dz_2}{2 - i} + \int_{\gamma_-} \frac{dz_2}{2 - i} e^{N S(z_1; x) + N S(z_2; x)} G(z_1; z_2) + O(e^{2N \text{Re}(S_+) - N}); \quad (28)$$

for suitably small $\epsilon > 0$.

We now need to parameterize the contours. From (25) we see that if we set

$$= \begin{cases} A \arcsin(x) = 2; & \text{GUE;} \\ = 4; & \text{LUE;} \end{cases} \quad (29)$$

then a suitable parameterization of is

$$z = z^* + e^{-i\pi/2} t; \quad t \in [-\epsilon; \epsilon]; \quad (30)$$

for sufficiently small $\epsilon > 0$. In (29) the function $\text{Arcsin} : [-1; 1] \rightarrow [-\pi/2; \pi/2]$ denotes the principle branch of arcsine. With the parameterization (30) the contour γ_+ is traversed in the negative direction, so we need to compensate for this with an explicit minus sign.

By choosing sufficiently small $S(z; x)$ is analytic on γ_+ and so using the parameterization (30) we see that

$$S(z; x) = S(a t^2 - a t'^2; t); \quad z \in \gamma_+; \quad (31)$$

where

$$a = \begin{cases} (x); & \text{GUE}; \\ 2x^2 - (x); & \text{LUE}; \end{cases} \quad (32)$$

and

$$S'_+(t) = \sum_{k=3}^{\infty} \frac{S^{(k)}(z; x)}{S^{(2)}(z; x) k!} e^{i(k-2)t} t^{k-2}; \quad (33)$$

We note that $S'_-(t) = \overline{S'_+(t)}$, where $\overline{}$ denotes complex conjugation.

It is a straightforward exercise to show that $S = S_+$, and that

$$\text{Re}(S_+) = \begin{cases} < \frac{1}{2} + x^2; & \text{GUE}; \\ = 2x - \log(2); & \text{LUE}; \end{cases} \quad (34)$$

$$\text{Im}(S_+) = -P(x); \quad (35)$$

where

$$P(x) = \int_{x_0}^z (t) dt; \quad (36)$$

$$\begin{aligned} & \geq 1 + \frac{x}{2} - (x) - \frac{1}{2} \text{Arcos}(x); & \text{GUE}; \\ & = \geq 1 + x - (x) - \frac{2}{\pi} \text{Arcos}\left(\frac{P(x)}{2}\right); & \text{LUE}; \end{aligned} \quad (37)$$

Here x_0 is the left edge of the support of (x) given in (7), i.e. $x_0 = 0; 1$, for the LUE and GUE respectively. We note that $P(x)$ is the probability distribution function corresponding to (x) . The limiting distribution function $P(x)$ will play a significant role in the bulk asymptotic expansion of $J_N(x)$.

With the results (31) and (35) for $S(z; x)$, and the definitions

$$G(t) = G(z_1; z_2) = \frac{z_1! z_2!}{z_1! z_2!} e^{i t_1} e^{i t_2}; \quad (38)$$

$$E_1^{(B)} = \int_0^z \frac{e^{-a N t_1^2}}{2} dt_1; \quad (39)$$

we can make the change of variables (30) in (28) to obtain

$$\begin{aligned} J_N(x) &= e^{2N \text{Re}(S_+)} E_1^{(B)} E_2^{(B)} \int_{\gamma_+} \int_{\gamma_+} e^{i(1+\frac{1}{2})(N-P(x))} e^{-a N t_1'^2} e^{-a N t_2'^2} G(t) \\ &\quad + O(e^{2N \text{Re}(S_+)} N^{-1}); \end{aligned} \quad (40)$$

A small amount of massaging shows that the $+$ term in (40) is the complex conjugate of the $+$ term, and likewise the $+$ term is conjugate to the $+$ term. Suppose now that we define the function $F_+(\cdot; t)$ by

$$F_+(\cdot; t) = e^{i t_1} e^{i t_2} G(t) \quad (41)$$

then

$$J_N(x) = 2 e^{2N \text{Re}(S_+)} \text{Re} E_1^{(B)} E_2^{(B)} F_+(\cdot; t) + O(e^{2N \text{Re}(S_+)} N^{-1}); \quad (42)$$

We shall discuss the purpose of the parameters ϵ_i shortly. Note that the two terms in (42) are qualitatively distinct { the second term is oscillatory in N while the first is not.

It is convenient to pause for a moment and multiply (42) by the explicit forms for the prefactors required in (14) using (21) and (34) to obtain the corresponding expression for $N(x)$. Since

$$\frac{2c_0(N)c_1(N)}{k_{N-1}k^2}!_N(x)e^{2N\text{Re}(S_+)} = h_0 N^{-1} + \frac{h_1}{N} + O\left(\frac{1}{N^2}\right) \quad (43)$$

with

$$h_0 = \begin{pmatrix} 2; & \text{GUE}; \\ 4x; & \text{LUE}; \end{pmatrix} \quad (44)$$

$$h_1 = \begin{pmatrix} 8 \\ \geq \frac{1}{12}; & \text{GUE}; \\ \geq \frac{(1)}{2}; & \text{LUE}; \end{pmatrix} \quad (45)$$

we have

$$N(x) = h_0 N^{-1} + \frac{h_1}{N} + O\left(\frac{1}{N^2}\right) 2\text{Re} N E_1^{(B)} E_2^{(B)} F_+; (\epsilon; t) \Big|_{\epsilon_i = aN t_1^2} e^{2i(N - P(x))} F_+; (\epsilon; t) \Big|_{\epsilon_i = aN t_1^2} + O(e^{-N}): \quad (46)$$

The introduction of the auxiliary variables $\epsilon_1; \epsilon_2$ in (41) is a common ruse applied in the saddle point method (see e.g. Ref.¹⁶) which we now discuss. Suppose that we construct the Maclaurin expansion in $t_1; t_2$ of $F(\epsilon; t)$ with ϵ considered as a fixed parameter

$$F(\epsilon; t) = \sum_{j=0}^{\infty} \sum_{k=0}^{\infty} \frac{X^j X^k}{k!(j-k)!} \frac{t_1^j t_2^k}{\partial s_1^k \partial s_2^{j-k}} F(\epsilon; s_1; s_2) \Big|_{s_1; s_2=0} + O(t_1^{p_1} t_2^{p_2})_{p_1+p_2=p+1} : \quad (47)$$

If we now set $\epsilon_i = aN t_1^2$ in (47) and perform the integrations required in (46) then we find that each term corresponding to a given value of j in (47) has the same resulting N dependence. This then gives a systematic way of obtaining the corrections out to any given order in N . To see why this occurs, first note that we need only consider the terms in (47) for which both j and k are even since any odd monomials are annihilated by (39), and then further note that with $\epsilon_i = aN t_1^2$ we have

$$N E_1^{(B)} E_2^{(B)} t_1^{2m_1} t_2^{2m_2} \epsilon_1^{-l_1} \epsilon_2^{-l_2} = (-1)^{l_1+l_2} \frac{(m_1+l_1+1)(m_2+l_2+1)}{4^{l_1+l_2} a^{m_1+m_2+1}} N^{-(m_1+m_2)} + O(e^{-a^2 N}); \quad (48)$$

for any $l_1, l_2 \geq 2N$. Hence despite the fact that various powers of ϵ_1 and ϵ_2 arise when $F(\epsilon; t)$ is differentiated, for a given value of j all terms end up with the same N dependence after setting $\epsilon_i = aN t_1^2$ and performing the integrations.

Hence, substituting (47) into (46) one can construct the asymptotic series for $N(x)$ out to any desired order. We shall explicitly construct this series out to order $1=N$ but the generalization to higher orders is obvious. However, as we shall see the resulting asymptotic expansions obtained by keeping only the $1=N$ correction are already extremely good, and it's quite likely that optimal truncation occurs at this order, providing little incentive to construct higher order corrections.

Let us denote by $c_m^{(1)}(x)$ the coefficient of $1=N^m$ in the $1=N$ expansion generated by acting on (47) with $N E_1^{(B)} E_2^{(B)}$. Then since the only terms which contribute are those for which j and k are even we have

$$\frac{c_m^{(1)}(x)}{N^m} = \sum_{k=0}^{\infty} \frac{1}{(2k)!(2[m-k])!} N E_1^{(B)} E_2^{(B)} t_1^{2k} t_2^{2(m-k)} \frac{\partial^{2k}}{\partial s_1^{2k}} \frac{\partial^{2(m-k)}}{\partial s_2^{2(m-k)}} F(\epsilon; s) \Big|_{s_i=0} \Big|_{\epsilon_i = aN t_1^2} : \quad (49)$$

We can re-express $N(x)$ from (46) in terms of the $c_m^{(1)}(x)$ as follows

$$N(x) = 2h_0 \text{Re} c_0^{(1;1)}(x) + 2h_0 \text{Re} e^{2i(N - P(x))} c_0^{(1;1)}(x) + 2h_0 \text{Re} e^{2i(N - P(x))} c_1^{(1;1)}(x) + \frac{h_1}{N} + 2h_0 \text{Re} c_1^{(1;1)}(x) + h_1 \frac{1}{N} + O\left(\frac{1}{N^2}\right) : \quad (50)$$




FIG. 1: Comparison of the asymptotic expansion (51), shown as the dashed line, and the exact result (10), shown as the solid line, for the eigenvalue density of the GUE with $N = 10$.

It is not hard to show that the term $2h_0 \operatorname{Re} c_0^{(1;1)}(x)$ is equal to the limiting density $\rho(x)$ from (7) when $|x| < 1$ as required, and also that $c_0^{(1;1)}(x)$ vanishes identically (this is actually obvious from (17) and (38)). The explicit construction of the remaining $c_m^{(1)}(x)$ required in (50) is straightforward and we finally obtain the following.

Proposition 1. Let $\rho_N(x)$ be as defined in (4), and let x be fixed with $|x| < 1$. Then as $N \rightarrow \infty$ we have the following:
For the GUE

$$\rho_N(x) = \rho(x) \left[\frac{2 \cos[2N P(x)]}{3 \rho^2(x)} \frac{1}{N} + O\left(\frac{1}{N^2}\right) \right]; \quad (51)$$

while for the LUE

$$\begin{aligned} \rho_N(x) &= \rho(x) \left[\frac{\cos(2N P(x)) [1 + x \rho(x) P(x)]}{3 x^2 \rho^2(x)} - \frac{1}{2 x \rho(x)} \frac{1}{N} + O\left(\frac{1}{N^2}\right) \right]; \\ &= \rho(x) \left[\frac{\cos(2N P(x)) - 2 \operatorname{Arccos}\left(\frac{P(x)}{x}\right)}{3 x^2 \rho^2(x)} - \frac{1}{4(1-x)} \frac{1}{N} + O\left(\frac{1}{N^2}\right) \right]; \end{aligned} \quad (52)$$

where $\rho(x)$ is given in (7) and $P(x)$ is the corresponding probability distribution function, given explicitly in (37).

As remarked in the Introduction, the result (51) was obtained previously in¹¹ again by steepest descent, but starting from an integral representation derived by supersymmetric arguments rather than orthogonal polynomials.

It is interesting to note that it is the distribution function $P(x)$ of the limiting density $\rho(x)$ which controls the large N oscillations in the $1/N$ correction to $\rho_N(x)$. Note also that while the non-oscillatory correction vanishes at order $1/N$ for the GUE leaving only an oscillatory correction at this order, the LUE has both an oscillatory and a non-oscillatory component to its $1/N$ correction.

To demonstrate to the reader just how accurate the expansions given by Proposition 1 are we provide in Figures 1 and 2 a numerical comparison of the asymptotic expansions with the exact results computed using the expression (10) in terms of orthogonal polynomials.




FIG. 2: Comparison of the asymptotic expansion (52), shown as the dashed line, and the exact result (10), shown as the solid line, for the eigenvalue density of the LUE with $\beta = 1/2$ and $N = 10$.

IV. SOFT EDGE ASYMPTOTICS FOR THE GUE AND LUE

The appropriate scaling to elucidate the behavior of $p_N(x)$ near the soft edge is to set $x = 1 + \frac{b^{1/3}}{N^{2/3}}$ for fixed b , as appears in (9). Substituting such a scaling into (16) we find

$$N S\left(z; 1+\frac{b^{1/3}}{2 N^{2/3}}\right)=b^{1/3} N^{1/3} z+N S(z); \quad (53)$$

where we've introduced the shorthand $S(z) \equiv S(z; 1)$. Here $b > 0$ is a free parameter that we can fix later as convenient. Defining

$$\tilde{J}_N(\epsilon) \equiv J_N\left(1+\frac{b^{1/3}}{2 N^{2/3}}\right); \quad (54)$$

we see that (53) leads to

$$\tilde{J}_N(\epsilon)=\frac{1}{2 \pi i} \int_{\Gamma} d z_1 \exp \left[N S\left(z_1\right)+b^{1/3} N^{1/3} z_1\right]-\frac{1}{2 \pi i} \int_{\Gamma} d z_2 \exp \left[N S\left(z_2\right)+b^{1/3} N^{1/3} z_2\right]-G\left(z_1 ; z_2\right) ; \quad (55)$$

The reader might be concerned by the slightly unorthodox term in the exponent proportional to $N^{1/3}$, however it is subdominant to the $N S(z)$ term and its presence does not affect any of the usual arguments of the saddle point method; the asymptotic behavior of (55) is determined by $S(z)$. From (22), (23) and (25) we see for both the GUE and LUE that $S(z)$ has one double saddle point, located at $z = 1$. We can deform the contour of integration to a new contour which passes through $z = 1$ along paths of steepest descent. Note that although in (12) the integrals must be positively oriented, since we have two integrals in (55) we are free to orient the integrals in the negative direction since the consequent minus signs cancel. Let us denote by C the contour consisting of the union of two rays of unit length, the first starting at $z = e^{i \pi/3} - 1$ and ending at $z = 1$ and the second starting at $z = 1$ and ending at $z = e^{i \pi/3} - 1$. If we then denote by A any suitable arc such that $C \cup A$ is a simple closed curve enclosing the origin we can write

$$\begin{aligned} \tilde{J}_N(\cdot) = & \int_C^Z + \int_A^Z \frac{dz_1}{2-i} \exp \left(-N S(z_1) - b^{1=3} N^{1=3} z_1 \right) + \int_C^Z + \int_A^Z \frac{dz_2}{2-i} \exp \left(-N S(z_2) - b^{1=3} N^{1=3} z_2 \right) G(z_1; z_2); \\ & (56) \end{aligned}$$

$$\begin{aligned} = & \int_C^Z \frac{dz_1}{2-i} \exp \left(-N S(z_1) - b^{1=3} N^{1=3} z_1 \right) + \int_C^Z \frac{dz_2}{2-i} \exp \left(-N S(z_2) - b^{1=3} N^{1=3} z_2 \right) G(z_1; z_2) + o(e^{2N S(\cdot-1)} N^{-p}); \\ & (57) \end{aligned}$$

The error bound in (57) holds for all $p \geq 2$, and so in what follows we consider p as arbitrarily large. The equality between (57) and (56) can be obtained by noting that we can choose A to consist of two rays lying along the path of steepest descent away from the endpoints of C , which extend as far as we like into the right half plane, together with an arc to close the contour which we can choose to be as far into the right half plane as desired. With such a choice for A one can obtain the required bounds by a straightforward generalization of the usual argument used in the saddle point method. For a careful discussion of the saddle point method suitable for this purpose see for example Section 2.5 of Ref.¹⁶. We note that Ref.¹⁶ refers to the saddle point method as Perron's method.

Now let us change variables in (57) according to $t = z + 1$, so that the vertex of our contour is now at the origin. We shall denote the image of C under this change of variables by B . Further, since $S(z)$ is analytic on C we have

$$S(t-1) = S(\cdot-1) + b \frac{z^3}{3} + b \frac{z^3}{3} \cdot (t); \quad (58)$$

where

$$\cdot (t) := \sum_{k=4}^{\infty} \frac{S^{(k)}(\cdot-1) 3!}{S^{(3)}(\cdot-1) k!} t^{k-3}; \quad (59)$$

and we have now chosen

$$b = \frac{S^{(3)}(\cdot-1)}{(\cdot-1)^2}; \quad (60)$$

$$= \begin{cases} 1; & \text{GUE;} \\ 2; & \text{LUE;} \end{cases} \quad (61)$$

We also note that by setting $x = 1$ in (34) and (35) we have

$$S(\cdot-1) = \begin{cases} < \frac{3}{2} & \text{GUE;} \\ \geq \frac{3}{2} & \text{LUE;} \end{cases} \quad (62)$$

Finally, defining

$$F(\cdot; t) := G(t_1-1; t_2-1) e^{-i \cdot (t_1) + i \cdot (t_2)}; \quad (63)$$

and

$$E_i^{(S)} := \int_B^Z \exp \left(b N \frac{t_i^3}{3} - b^{1=3} N^{1=3} t_i \right) \frac{dt_i}{2-i} \quad (64)$$

we have

$$\tilde{J}_N(\cdot) = e^{2N \operatorname{Re}[S(\cdot-1)] + 2b^{1=3} N^{1=3}} \int_{i=bN}^h E_1^{(S)} E_2^{(S)} F(\cdot; t) j_{i=bN}^{t_i^3=3} + o(N^{-p}); \quad (65)$$

If we now multiply (65) by the prefactors required in (14) using (21) we obtain

$$\frac{(bN)^{1=3}}{2} N^{-1} + \frac{b^{1=3}}{2} \frac{1}{N^{2=3}} = g_0(\cdot) + \frac{g_1(\cdot)}{N^{1=3}} + \frac{g_2(\cdot)}{N^{2=3}} + o\left(\frac{1}{N}\right) N^{4=3} E_1^{(S)} E_2^{(S)} F(\cdot; t) j_{i=bN}^{t_i^3=3} + o(N^{-p}); \quad (66)$$

where $g_m(\cdot)$ is the coefficient of $N^{-m/3}$ in the large N fixed expansion of

$$\begin{aligned} & \frac{8}{e^{2=2N^{1=3}}}; \quad \text{GUE}; \\ & : 2^{2-2=3} \left(1 + \frac{1}{(2N)^{2=3}} \right); \quad \text{LUE}; \end{aligned} \quad (67)$$

In (66) we have presented only terms $O(1/N)$ in the first factor since this will be sufficient for our purposes in what follows. Higher order terms are easily retained if desired.

Our work is now essentially done. One expands $F(\cdot; t)$ around $t = 0$ for fixed λ as in (47) and then sets $\lambda = bN t_1^3 = 3$, analogous to the bulk case. Again, after integration, each value of j in the Maclaurin expansion (47) contributes to the same order in N . To see this explicitly we can use the following lemma.

Lemma 1. Let B be the contour consisting of the union of a ray starting at $e^{i=3}$ and ending at the origin, and a ray starting at the origin and ending at $e^{i=3}$. For any $b > 0$ and $0 < \epsilon < 1=3$ we have for large N that

$$\oint_B z^m \exp(bN \frac{z^3}{3}) b^{1=3} N^{1=3} z \frac{dz}{2\pi i} = (-1)^m b^{(m+1)=3} N^{(m+1)=3} \text{Ai}^{(m)}(\cdot) + O(e^{-bN});$$

where $\text{Ai}^{(m)}(\cdot)$ is the m^{th} derivative of the Airy function $\text{Ai}(\cdot)$.

Proof. This follows from the standard entire contour integral expression for $\text{Ai}(\cdot)$ (see e.g. [17]) by simply changing variables $z \mapsto b^{1=3} N^{1=3} z$, and noting that rays defining the contour B can be extended to infinity at the cost of introducing exponentially subdominant corrections. \square

An immediate consequence of Lemma 1 is that with $\lambda = bN t_1^3 = 3$ we have

$$N^{2=3} E_1^{(S)} E_2^{(S)} t_1^{m_1} t_2^{m_2} \frac{\lambda_1}{1} \frac{\lambda_2}{2} = \frac{(-1)^{\lambda_1 + \lambda_2 + m_1 + m_2}}{3^{\lambda_1 + \lambda_2} b^{(m_1 + m_2 + 2)=3}} \text{Ai}^{(m_1 + 3\lambda_1)}(\cdot) \text{Ai}^{(m_2 + 3\lambda_2)}(\cdot) \frac{1}{N^{1=3}} + O(e^{-bN}); \quad (68)$$

Hence, if we construct the Maclaurin expansion of $F(\cdot; t)$ with λ fixed as in (47), and set $\lambda = bN t_1^3 = 3$ and integrate using (68), we obtain an expansion for

$$N^{4=3} E_1^{(S)} E_2^{(S)} F(\cdot; t) \Big|_{\lambda = bN t_1^3 = 3} \quad (69)$$

in powers of $N^{-1=3}$. Denoting the coefficient of $N^{-m/3}$ in this expansion by $g_m(\cdot)$ we have explicitly that

$$g_m(\cdot) = \sum_{k=0}^{m+2} \frac{1}{k!(m+2-k)!} N^{4=3} E_1^{(S)} E_2^{(S)} t_1^k t_2^{m+2-k} \frac{\partial^k}{\partial s_1^k} \frac{\partial^{m+2-k}}{\partial s_2^{m+2-k}} F(\cdot; s_1; s_2) \Big|_{s_1; s_2 = 0} \Big|_{\lambda = bN t_1^3 = 3}; \quad (70)$$

The reader might be concerned that according to (68) the $k_1 + k_2 = 0$ and $k_1 + k_2 = 1$ terms grow with N ; however it is not hard to show from (70) that the coefficients $c^{(2)}(\cdot)$ and $c^{(1)}(\cdot)$ vanish identically.

We can now express (66) in terms of the coefficients $c_m(\cdot)$ as

$$\begin{aligned} \frac{(bN)^{1=3}}{2} N^{-1} + \frac{b^{1=3}}{2} \frac{1}{N^{2=3}} &= g_0(\cdot) q_0(\cdot) + g_1(\cdot) q_0(\cdot) + g_0(\cdot) q_1(\cdot) \frac{1}{N^{1=3}} \\ &+ g_2(\cdot) q_0(\cdot) + g_1(\cdot) q_1(\cdot) + g_0(\cdot) q_2(\cdot) \frac{1}{N^{2=3}} + O\left(\frac{1}{N}\right); \end{aligned} \quad (71)$$

We have explicitly displayed terms $O(1/N)$ here, but it is straightforward to retain as many terms as desired. The expansion constructed from terms $O(1/N)$ however is extremely accurate, as we demonstrate in Figures 3 and 4, and it appears to be the numerically optimal order at which to truncate the expansions.

The explicit forms for the coefficients $c_m(\cdot)$ can be constructed from (70) and substituted into (71). We can also further simplify the Airy derivatives appearing in (68) using the Airy differential equation $\text{Ai}^{(0)}(\cdot) = -\text{Ai}(\cdot)$ so that only $\text{Ai}(\cdot)$ and its first derivative appear. We finally obtain the following

Proposition 2. Let $N(x)$ be as defined in (4). Then with λ fixed, as $N \rightarrow \infty$ we have the following:




FIG. 3: Comparison of the asymptotic expansion (72), shown as the dashed line, and the exact result (10), shown as the solid line, for the eigenvalue density near the soft edge at $x = 1$, for the GUE with $N = 10$.

For the GUE

$$\begin{aligned} \frac{N^{1/3}}{2} p_N(1 + \frac{\xi}{2N^{2/3}}) &= \text{Ai}^0(\xi)^2 - \text{Ai}(\xi)^2 \\ &+ \frac{1}{20} \xi^2 \text{Ai}(\xi)^2 - 2 \text{Ai}^0(\xi)^2 - 3 \text{Ai}(\xi) \text{Ai}^0(\xi) \frac{1}{N^{2/3}} \\ &+ O\left(\frac{1}{N}\right); \end{aligned} \quad (72)$$

while for the LUE

$$\begin{aligned} \frac{(2N)^{1/3}}{2} p_N(1 + \frac{\xi}{(2N)^{2/3}}) &= \text{Ai}^0(\xi)^2 - \text{Ai}(\xi)^2 + \frac{1}{2^{1/3}} \text{Ai}(\xi)^2 \frac{1}{N^{1/3}} \\ &+ \frac{2^{1/3}}{10} \xi^2 \text{Ai}(\xi)^2 - 2 \text{Ai}^0(\xi)^2 + (2 - 5\xi^2) \text{Ai}(\xi) \text{Ai}^0(\xi) \frac{1}{N^{2/3}} \\ &+ O\left(\frac{1}{N}\right); \end{aligned} \quad (73)$$

Figures 3 and 4 provide a numerical comparison of the asymptotic expansions given in Proposition 2 with the exact results computed using the expression (10) in terms of orthogonal polynomials.

Note that it appears that the GUE converges much faster than the LUE, since while the two curves in (3) are almost indistinguishable at $N = 10$ for the GUE, the asymptotic expansion for the LUE begins to diverge from the exact result in (4) already by $N = 2$, and both curves are rather different from the limiting Airy kernel expression (9). We investigated the effects of retaining more terms in the expansion for the LUE case; keeping terms $O(1/N)$ did not noticeably change the plots, while keeping terms higher than $1/N$ caused significant divergence of the asymptotic expansion from the exact result. The explanation for this is most likely that, as made precise in the next section, the edge expansions match onto the bulk expansions, and these in turn become more divergent near the edge at each order in $1/N$.




FIG. 4: Comparison of the asymptotic expansion (73), shown as the dashed line, and the exact result (10), shown as the solid line, for the eigenvalue density near the soft edge at $x = 1$, for the LUE with $\beta = 1/2$ and $N = 20$. Also shown is the limit as $N \rightarrow 1$ given by the Airy kernel (9), shown as the dotted line lying below the other two curves.

V. MATCHING OF THE BULK AND EDGE EXPANSIONS

In Figures 1{4 plots of the bulk and edge asymptotic expansions have separately been compared against the exact density for $N = 10$. Although the scale of the independent variable is different, we can see from Figures 3 and 4 that the edge asymptotic expansions are accurate approximations to the exact density up to the neighborhood of the first local maximum (relative to the edge $\xi = 0$) at least and thus should be used instead of the bulk asymptotic expansion in this region.

At a quantitative level, it is possible to exhibit a matching between the various asymptotic expansions. Suppose in (51) we set $x = 1 + \beta/(2N)^{2/3}$, and in (52) we set $x = 1 + \beta/(2N)^{2/3}$, and take $\beta < 0$ and fixed. Expanding the right hand sides as an asymptotic series in N gives

$$\begin{aligned} N^{1/3} \rho_N^{\text{GUE}}(1 + \beta/(2N)^{2/3}) &: \frac{2^{-1/3}}{2} \sum_{j=0}^{\infty} \frac{\cos(4j \beta^{3/2})}{2^j j} \\ &+ \frac{j \beta^{3/2}}{4} + \frac{\cos(4j \beta^{3/2})}{8} + \frac{j \beta^{3/2} \sin(4j \beta^{3/2})}{20} + \frac{1}{N^{2/3}} \\ &+ O\left(\frac{1}{N^{4/3}}\right) : \end{aligned} \quad (74)$$

$$\begin{aligned} (2N)^{1/3} \rho_N^{\text{LUE}}(1 + \beta/(2N)^{2/3}) &: \frac{2^{-1/3}}{2} \sum_{j=0}^{\infty} \frac{\cos(4j \beta^{3/2})}{2^j j} \\ &+ \frac{(1 + \sin(4j \beta^{3/2}))}{2^j j} \frac{1}{(2N)^{1/3}} + O\left(\frac{1}{N^{2/3}}\right) \end{aligned} \quad (75)$$

where the symbol $:$ denotes that the asymptotic series have been expanded as specified. An important feature is that this procedure mixes the terms which are at different orders in N in (51) and (52).

Let us now compute the $N \rightarrow 1$ asymptotic expansions of the right hand sides of the first two terms in each of (72) and (73), multiplied by 2. Using the fact that for $x \rightarrow 1$ (see e.g.¹⁷)

$$\text{Ai}(-x) = \frac{1}{\sqrt[3]{x^{1/4}}} \cos \frac{2}{3} x^{3/2} - \frac{5\sqrt[3]{3}}{24x^{1/2}} \cos \frac{2}{3} x^{3/2} + \frac{2}{3} x^{3/2} + O\left(\frac{1}{x^{3/4}}\right)$$

we obtain expansions which reproduce the N -independent terms in (74) and (75), giving furthermore, terms of higher order in $1=j$. In (75) the term proportional to $1=(2N)^{1=3}$ is reproduced, and this too is accompanied by terms of higher order in $1=j$. In (74) the term $j^{j=2}=4 N^{2=3}$ is reproduced, while the other terms proportional to $1=N^{2=3}$ are out by rational factors. The explanation for the missing higher order terms in $1=j$, and incorrect rational factors is most likely due to the fact that terms of all orders in $1=N$ in (51) and (52) contribute to each distinct order in the expansions (74) and (75). Specifically, from the results exhibited above, it would seem that expanding the complete large N asymptotic series for $\frac{GUE}{N}(x)$ and $\frac{LUE}{N}(x)$ as in (74) and (75) would give precisely the large N expansion of (72) and (73), extended to all orders in N .

Electronic address: garoni@ima.umn.edu; URL: <http://www.ima.umn.edu/~garoni>

^y Electronic address: P.Forrester@msunimelb.edu.au; URL: <http://www.ms.unimelb.edu.au/~matpjf/matpjf.html>

^z Electronic address: n.frankel@physics.unimelb.edu.au

¹ P. J. Forrester, Log-gases and Random Matrices, URL <http://www.ms.unimelb.edu.au/~matpjf/matpjf.html>.

² M. L. Mehta, Random Matrices (Academic Press, New York, 1991).

³ P. Deift, Orthogonal Polynomials and Random Matrices: A Riemann-Hilbert Approach, no. 3 in Courant Lecture Notes in Mathematics (Courant Institute of Mathematical Sciences, 1999).

⁴ P. J. Forrester, N. E. Frankel, T. M. Garoni, and N. S. Witte, Commun. Math. Phys. 238, 257 (2003).

⁵ P. J. Forrester and N. E. Frankel, J. Math. Phys. 45, 2003 (2004).

⁶ P. J. Forrester, N. E. Frankel, T. M. Garoni, and N. S. Witte, Phys. Rev. A. 67, 043607 (2003).

⁷ T. M. Garoni, J. Math. Phys. 46, 043516 (2005).

⁸ I. V. Krasovsky, math-ph/0411016.

⁹ E. Brezin, C. Itzykson, G. Parisi, and J. Zuber, Commun. Math. Phys. 59, 35 (1978).

¹⁰ P. Di Francesco, Nucl. Phys. B 648, 461 (2002).

¹¹ F. Kalisch and D. Braak, J. Phys. A. 35, 9957 (2002).

¹² P. J. Forrester, Nuclear Phys. B 402, 709 (1993).

¹³ G. Szego, Orthogonal polynomials (American Mathematical Society, Rhode Island, 1967).

¹⁴ M. Plancherel and W. Rotach, Commentarii Mathematici Helvetici 1, 227 (1929).

¹⁵ E. Moecklin, Commentarii Mathematici Helvetici 7, 24 (1934).

¹⁶ R. Wong, Asymptotic Approximations of Integrals (Academic Press, New York, 1989).

¹⁷ F. W. J. Olver, Asymptotics and special functions (Academic Press, New York, 1974).