

Topology and Phase Transitions II. Theorem on a necessary relation

Roberto Franzosi

*Dipartimento di Fisica dell'Università di Firenze, Via G. Sansone 1,
I-50019 Sesto Fiorentino, and C.N.R.-I.N.F.M., Italy*

Marco Pettini*

*Istituto Nazionale di Astrofisica – Osservatorio Astrofisico di Arcetri,
Largo E. Fermi 5, 50125 Firenze, Italy
and I.N.F.M., Unità di Firenze,
and I.N.F.N., Sezione di Firenze
(Dated: March 28, 2022)*

* Corresponding author. e-mail: pettini@arcetri.astro.it, Phone: +39-055-2752282, Fax: +39-055-220039.

Abstract

In this second paper, we prove a necessity Theorem about the topological origin of phase transitions. We consider physical systems described by smooth microscopic interaction potentials $V_N(q)$, among N degrees of freedom, and the associated family of configuration space submanifolds $\{M_v\}_{v \in \mathbb{R}}$, with $M_v = \{q \in \mathbb{R}^N | V_N(q) \leq v\}$. On the basis of an analytic relationship between a suitably weighed sum of the Morse indexes of the manifolds $\{M_v\}_{v \in \mathbb{R}}$ and thermodynamic entropy, the Theorem states that any possible unbound growth with N of one of the following derivatives of the configurational entropy $S^{(-)}(v) = (1/N) \log \int_{M_v} d^N q$, that is of $|\partial^k S^{(-)}(v)/\partial v^k|$, for $k = 3, 4$, can be entailed only by the weighed sum of Morse indexes. Since the unbound growth with N of one of these derivatives corresponds to the occurrence of a first or of a second order phase transition, and since the variation of the Morse indexes of a manifold is in one-to-one correspondence with a change of its topology, the Main Theorem of the present paper states that a phase transition *necessarily* stems from a topological transition in configuration space. The proof of the Theorem given in the present paper cannot be done without Main Theorem of paper I.

PACS numbers: 05.70.Fh; 05.20.-y; 02.40.-k

Keywords: Statistical Mechanics, Phase Transitions, Topology

1. INTRODUCTION

In Statistical Mechanics, a central task of the mathematical theory of phase transitions has been to prove the loss of differentiability of the pressure function – or of other thermodynamic functions – with respect to temperature, or volume, or an external field. The first rigorous results of this kind are the exact solution of $2d$ Ising model due to Onsager [1], and the Yang-Lee theorem [2] showing that, despite the smoothness of the canonical and grand canonical partition functions respectively, in the $N \rightarrow \infty$ limit also piecewise differentiability of pressure or other thermodynamic functions becomes possible.

Another approach to the problem has considerably grown after the introduction of the concept of a Gibbs measure for infinite systems by Dobrushin, Lanford and Ruelle. In this framework, the phenomenon of phase transition is seen as the consequence of non-uniqueness of a Gibbs measure for a given type of interaction among the particles of a system [3, 4].

Recently, it has been conjectured[5, 6, 7, 8, 9] that the origin of the phase transitions singularities could be attributed to suitable topology changes within the family of equipotential hypersurfaces $\{\Sigma_v = V_N^{-1}(v)\}_{v \in \mathbb{R}}$ of configuration space. These level sets of V_N naturally foliate the support of the statistical measures (canonical or microcanonical) so that the mentioned topology change would induce a change of the measure itself at the transition point. In a few particular cases, the truth of this *topological hypothesis* has been given strong evidence: *i)* through the numerical computation of the Euler characteristic for the $\{\Sigma_v\}_{v \in \mathbb{R}}$ of a two-dimensional lattice φ^4 model [7]; *ii)* through the exact analytic computation of the Euler characteristic of $\{M_v = V_N^{-1}((-\infty, v])\}_{v \in \mathbb{R}}$ submanifolds of configuration space for two different models, the mean-field XY model[10] and the k -trigonometric model [11].

In the present paper we prove a necessity Theorem which implies that for a wide class of potentials (good Morse functions), a first or a second order phase transition can only be the consequence of a topology change of the submanifolds M_v of configuration space, and this appears to be the truly primitive and

deep mathematical origin of the phase transition phenomena, at least for the mentioned class of potentials.

The Theorem is enunciated as follows:

Theorem *Let $V_N(q_1, \dots, q_N) : \mathbb{R}^N \rightarrow \mathbb{R}$, be a smooth, non-singular, finite-range potential. Denote by $M_v := V_N^{-1}((-\infty, v])$, $v \in \mathbb{R}$, the generic submanifold of configuration space bounded by Σ_v . Let $\{q_c^{(i)} \in \mathbb{R}^N\}_{i \in [1, \mathcal{N}(v)]}$ be the set of critical points of the potential, that is s.t. $\nabla V_N(q)|_{q=q_c^{(i)}} = 0$, and $\mathcal{N}(v)$ be the number of critical points up to the potential energy value v . Let $\Gamma(q_c^{(i)}, \varepsilon_0)$ be pseudo-cylindrical neighborhoods of the critical points, and $\mu_i(M_v)$ be the Morse indexes of M_v , then there exist real numbers $A(N, i, \varepsilon_0)$, g_i and real smooth functions $B(N, i, v, \varepsilon_0)$ such that the following equation for the micro-canonical configurational entropy $S_N^{(-)}(v)$ holds*

$$S_N^{(-)}(v) = \frac{1}{N} \log \left[\int_{M_v \setminus \bigcup_{i=1}^{\mathcal{N}(v)} \Gamma(q_c^{(i)}, \varepsilon_0)} d^N q + \sum_{i=0}^N A(N, i, \varepsilon_0) g_i \mu_i(M_{v-\varepsilon_0}) + \sum_{n=1}^{\mathcal{N}_{cp}^{\nu(v)+1}} B(N, i(n), v - v_c^{\nu(v)}, \varepsilon_0) \right],$$

(details and appropriate definitions are given in Section 2), and an unbound growth with N of one of the derivatives $|\partial^k S^{(-)}(v)/\partial v^k|$, for $k = 3, 4$, and thus the occurrence of a first or of a second order phase transition respectively, can be entailed only by the topological term $\sum_{i=0}^N A(N, i, \varepsilon_0) g_i \mu_i(M_{v-\varepsilon_0}) + \sum_{n=1}^{\mathcal{N}_{cp}^{\nu(v)+1}} B(N, i(n), v - v_c^{\nu(v)}, \varepsilon_0)$.

The above given expression for the entropy stems from a decomposition of the volume of a generic submanifold M_v into two parts: the volume of the disjoint union of suitably defined neighborhoods of the critical points of the potential $V_N(q)$, and the volume of its complement. The latter is represented by the first term in square parentheses, and, after Main Theorem of paper I, it cannot entail unbounded growth with N of up to the fourth derivative of the entropy. Only the second and third terms in square parentheses could do it. These two terms, representing the volume of the neighborhoods of critical

points, are of topological meaning.

Thus the proof of the present Theorem *crucially* relies on Main Theorem of paper I which is a first step toward proving the necessity of topology changes of configuration space submanifolds (either the level sets Σ_v of the potential function, or the manifolds M_v bounded by them) for the occurrence of phase transitions [12, 13]. Main Theorem of paper I is based on the assumption of the existence – at any number N of degrees of freedom – of an energy density interval $[\bar{v}_0, \bar{v}_1]$ free of critical values. Under this assumption, in paper I, we proved the uniform convergence in the thermodynamic limit of configurational entropy in the class of three times differentiable functions. However, since in general it is a very hard task to locate all the critical points of a given potential function, it is also very hard to ascertain whether Main Theorem of paper I applies to it or not. To overcome this limitation, in the present paper we prove a new Theorem, based on a less restrictive assumption (allowing the existence of critical points), which we could not prove without the Main Theorem of paper I.

As already remarked above, the necessity Theorem proved in the present paper applies to a very broad class of systems: those described by finite range potentials which are good Morse functions. In fact, checking whether a given potential is a good Morse function or not is not difficult and amounts to control whether the potential is smooth, bounded below and whether its Hessian is non-degenerate, that is free of vanishing eigenvalues; degeneracy typically occurs in presence of continuous symmetries and can be removed by arbitrarily small and standard perturbations [14] which do not alter neither the microscopic dynamics produced by the gradients of the potential nor the thermodynamics.

2. BASIC DEFINITIONS

For a physical system \mathcal{S} of n particles confined in a bounded subset Λ^d of \mathbb{R}^d , $d = 1, 2, 3$, and interacting through a real valued potential function V_N

defined on $(\Lambda^d)^{\times n}$, with $N = nd$, the *configurational microcanonical volume* $\Omega(v, N)$ is defined for any value v of the potential V_N as

$$\Omega(v, N) = \int_{(\Lambda^d)^{\times n}} dq_1 \dots dq_N \delta[V_N(q_1, \dots, q_N) - v] = \int_{\Sigma_v} \frac{d\sigma}{\|\nabla V_N\|}, \quad (1)$$

where $d\sigma$ is a surface element of $\Sigma_v := V_N^{-1}(v)$; in what follows $\Omega(v, N)$ is also called *structure integral*. The norm $\|\nabla V_N\|$ is defined as $\|\nabla V_N\| = [\sum_{i=1}^N (\partial_{q_i} V_N)^2]^{1/2}$.

Now we can define the configurational thermodynamic functions to be used in this paper.

Henceforth, according to the need for explicit reference to the N -dependence of V , we shall use both V and V_N to denote the potential.

Definition 1. *Using the notation $\bar{v} = v/N$ for the value of the potential energy per particle, we introduce the following functions:*

- Configurational microcanonical entropy, relative to Σ_v . *For any $N \in \mathbb{N}$ and $\bar{v} \in \mathbb{R}$,*

$$S_N(\bar{v}) \equiv S_N(\bar{v}; V_N) = \frac{1}{N} \log \Omega(N\bar{v}, N).$$

- Configurational microcanonical entropy, relative to the volume bounded by Σ_v . *For any $N \in \mathbb{N}$ and $\bar{v} \in \mathbb{R}$,*

$$S_N^{(-)}(\bar{v}) \equiv S_N^{(-)}(\bar{v}; V_N) = \frac{1}{N} \log M(N\bar{v}, N)$$

where

$$M(v, N) = \int_{(\Lambda^d)^{\times n}} dq_1 \dots dq_N \Theta[V_N(q_1, \dots, q_N) - v] = \int_0^v d\eta \int_{\Sigma_\eta} \frac{d\sigma}{\|\nabla V_N\|}, \quad (2)$$

with $\Theta[\cdot]$ the Heaviside step function; $M(v, N)$ is the codimension-0 subset of configuration space enclosed by the equipotential hypersurface Σ_v . The representation of $M(v, N)$ given in the r.h.s. stems from the already mentioned co-area formula in [15].

Definition 2 (First and second order phase transitions). *We say that a physical system \mathcal{S} undergoes a phase transition if there exists a thermodynamic*

function which – in the thermodynamic limit ($N \rightarrow \infty$ and $\text{vol}(\Lambda^d)/N = \text{const}$) – is only piecewise analytic. In particular, if the second-order derivative of the entropy $S_\infty^{(-)}(\bar{v})$ is discontinuous at some point \bar{v}_c , then we say that a first-order phase transition occurs. If the third-order derivative of the entropy $S_\infty^{(-)}(\bar{v})$ is discontinuous at some point \bar{v}_c , then we say that a second-order phase transition occurs. These definitions stem from the standard definitions of first and second order phase transitions as due to a discontinuity of the first or second derivatives of the Helmholtz free energy, respectively, and from the existing relationship – through a Legendre transform – between the Helmholtz free energy and the entropy (see Definition 1 in paper I).

Definition 3 (Standard potential, fluid case). We say that an N degrees of freedom potential V_N is a standard potential for a fluid if it is of the form

$$V_N : \quad \mathcal{B}_N \subset \mathbb{R}^N \rightarrow \mathbb{R}$$

$$V_N(q) = \sum_{i \neq j=1}^n \Psi(\|\vec{q}_i - \vec{q}_j\|) + \sum_{i=1}^n U_\Lambda(\vec{q}_i) \quad (3)$$

where \mathcal{B}_N is a compact subset of \mathbb{R}^N , $N = nd$, Ψ is a real valued function of one variable such that additivity holds, and where U_Λ is any smoothed potential barrier to confine the particles in a finite volume Λ , that is

$$U_\Lambda(\vec{q}) = \begin{cases} 0 & \text{if } \vec{q} \in \Lambda' \\ +\infty & \text{if } \vec{q} \in \Lambda^c, \text{ complement in } \mathbb{R}^N \\ \mathcal{C}^\infty & \text{function for } \vec{q} \in \Lambda \setminus \Lambda' \end{cases}$$

where $\Lambda' \subset \Lambda$ and Λ' arbitrarily close to $\Lambda \subset \mathbb{R}^N$, closed and bounded. U_Λ is a confining potential in a limited spatial volume with the additional property that given two limited d -dimensional regions of space, Λ_1 and Λ_2 , having in common a $d - 1$ -dimensional boundary, $U_{\Lambda_1} + U_{\Lambda_2} = U_{\Lambda_1 \cup \Lambda_2}$. By additivity we mean what follows. Consider two systems \mathcal{S}_1 and \mathcal{S}_2 , having $N_1 = n_1 d$ and $N_2 = n_2 d$ degrees of freedom, occupying volumes Λ_1^d and Λ_2^d , having potential energies v_1 and v_2 , for any $(q_1, \dots, q_{N_1}) \in (\Lambda_1^d)^{\times n_1}$ such that $V_{N_1}(q_1, \dots, q_{N_1}) = v_1$, for any $(q_{N_1+1}, \dots, q_{N_1+N_2}) \in (\Lambda_2^d)^{\times n_2}$ such that $V_{N_2}(q_{N_1+1}, \dots, q_{N_1+N_2}) = v_2$, for

$(q_1, \dots, q_{N_1+N_2}) \in (\Lambda_1^d)^{\times n_1} \times (\Lambda_2^d)^{\times n_2}$ let $V_N(q_1, \dots, q_{N_1+N_2}) = v$ be the potential energy v of the compound system $\mathcal{S} = \mathcal{S}_1 + \mathcal{S}_2$ which occupies the volume $\Lambda^d = \Lambda_1^d \cup \Lambda_2^d$ and contains $N = N_1 + N_2$ degrees of freedom. If

$$v(N_1 + N_2, \Lambda_1^d \cup \Lambda_2^d) = v_1(N_1, \Lambda_1^d) + v_2(N_2, \Lambda_2^d) + v'(N_1, N_2, \Lambda_1^d, \Lambda_2^d) \quad (4)$$

where v' stands for the interaction energy between \mathcal{S}_1 and \mathcal{S}_2 , and if $v'/v_1 \rightarrow 0$ and $v'/v_2 \rightarrow 0$ for $N \rightarrow \infty$ then V_N is additive. Moreover, at short distances Ψ must be a repulsive potential so as to prevent the concentration of an arbitrary number of particles within small, finite volumes of any given size.

Definition 4 (Standard potential, lattice case). We say that an N degrees of freedom potential V_N is a standard potential for a lattice if it is of the form

$$V_N : \quad \mathcal{B}_N \subset \mathbb{R}^N \rightarrow \mathbb{R}$$

$$V_N(q) = \sum_{\underline{i}, \underline{j} \in \mathcal{I} \subset \mathbb{N}^d} C_{\underline{i}\underline{j}} \Psi(\|\vec{q}_{\underline{i}} - \vec{q}_{\underline{j}}\|) + \sum_{\underline{i} \in \mathcal{I} \subset \mathbb{N}^d} \Phi(\vec{q}_{\underline{i}}) \quad (5)$$

where \mathcal{B}_N is a compact subset of \mathbb{R}^N . Denoting by a_1, \dots, a_d the lattice spacings, if $\underline{i} \in \mathbb{N}^d$, then $(i_1 a_1, \dots, i_d a_d) \in \Lambda^d$. We denote by m the number of lattice sites in each spatial direction, by $n = m^d$ the total number of lattice sites, by D the number of degrees of freedom on each site. Thus $\vec{q}_{\underline{i}} \in \mathbb{R}^D$ for any \underline{i} . The total number of degrees of freedom is $N = m^d D$. Having two systems made of $N = m^d D$ degrees of freedom, whose site indexes $\underline{i}^{(1)}$ and $\underline{i}^{(2)}$ run over $1 \leq i_1^{(1)}, \dots, i_d^{(1)} \leq m$, and $1 \leq i_1^{(2)}, \dots, i_d^{(2)} \leq m$, after gluing together the two systems through a common $d - 1$ dimensional boundary the new system has indexes \underline{i} running over, for example, $1 \leq i_1 \leq 2m$ and $1 \leq i_2, \dots, i_d \leq m$. If

$$v(N + N, \Lambda_1^d \cup \Lambda_2^d) = v_1(N, \Lambda_1^d) + v_2(N, \Lambda_2^d) + v'(N, N, \Lambda_1^d, \Lambda_2^d) \quad (6)$$

where v' stands for the interaction energy between the two systems and if $v'/v_1 \rightarrow 0$ and $v'/v_2 \rightarrow 0$ for $N \rightarrow \infty$ then V_N is additive.

Definition 5 (Short-range potential). In defining a short-range potential, a distinction has to be made between lattice systems and fluid systems. Given

a standard potential V_N on a lattice, we say that it is a short-range potential if the coefficients $C_{\underline{i}\underline{j}}$ are such that for any $\underline{i}, \underline{j} \in \mathcal{I} \subset \mathbb{N}^d$, $C_{\underline{i}\underline{j}} = 0$ iff $|\underline{i} - \underline{j}| > c$, with c is definitively constant for $N \rightarrow \infty$.

Given a standard potential V_N for a fluid system, we say that it is a short-range potential if there exist $R_0 > 0$ and $\epsilon > 0$ such that for $\|\mathbf{q}\| > R_0$ it is $|\Psi(\|\mathbf{q}\|)| < \|\mathbf{q}\|^{-(d+\epsilon)}$, where $d = 1, 2, 3$ is the spatial dimension.

Definition 6 (Stable potential). We say that a potential V_N is stable [16] if there exists $B \geq 0$ such that

$$V_N(q_1, \dots, q_N) \geq -NB \quad (7)$$

for any $N > 0$ and $(q_1, \dots, q_N) \in (\Lambda^d)^{\times n}$, or for $\vec{q}_{\underline{i}} \in \mathbb{R}^D$, $\underline{i} \in \mathcal{I} \subset \mathbb{N}^d$, $N = m^d D$, for lattices.

Definition 7 (Confining potential). With the above definitions of standard potentials V_N , in the fluid case the potential is said to be confining in the sense that it contains U_Λ which constrains the particles in a finite spatial volume, and in the lattice case the potential V_N contains an on-site potential such that – at finite energy – $\|\vec{q}_{\underline{i}}\|$ is constrained in compact set of values.

Remark 1 (Compactness of equipotential hypersurfaces). From the previous definition it follows that, for a confining potential, the equipotential hypersurfaces Σ_v are compact (because they are closed by definition and bounded in view of particle confinement).

In view of formulating and proving the Main Theorem of the present paper, we have to define some neighborhoods, that we call “pseudo-cylindrical”, of critical points of a potential function V_N . Before defining these pseudo-cylindrical neighborhoods of critical points, let us remember the following basic result in Morse theory.

Theorem. Let f be a smooth real valued function on a compact finite dimensional manifold M . Let $a < b$ and suppose that the set

$$f^{-1}([a, b]) \equiv \mathcal{M} = \{x \in M | a \leq f(x) \leq b\} \quad (8)$$

is compact and contains no critical points of f , that is $\|\nabla f\| \geq C > 0$ with C a constant. Let $y \in (a, b)$. Then there exists a diffeomorphism

$$\sigma : (a, b) \times f^{-1}(y) \rightarrow f^{-1}[(a, b)] \quad \text{by} \quad (v, x) \mapsto \sigma(v, x). \quad (9)$$

Corollary. *The manifolds $f^{-1}(y)$, $a < y < b$, are all diffeomorphic.*

This result is based on the existence of a one-parameter group of diffeomorphism

$$\sigma_v : M \rightarrow M \quad \text{by} \quad x \mapsto \sigma(v, x) \quad (10)$$

associated with the vector field $X = \nabla f(x)/\|\nabla f(x)\|^2$ with $v \mapsto \sigma(v, x)$ a solution of the differential equation on M

$$\frac{d\sigma(v, x)}{dv} = \frac{\nabla f[\sigma(v, x)]}{\|\nabla f[\sigma(v, x)]\|^2}, \quad \sigma(0, x) = x. \quad (11)$$

$\sigma(v, x)$ is defined for all $v \in \mathbb{R}$ and $x \in M$. Details can be found in standard references as [17, 18, 19].

Applied to the configuration space M , if the function f is identified with the potential V_N , then in the absence of critical points of V in the interval (v_0, v_1) the hypersurfaces $\Sigma_v = V_N^{-1}(v)$, $v \in (v_0, v_1)$, are all diffeomorphic.

Definition 8 (Pseudo-cylindrical neighborhoods). *Let Σ_{v_c} be a critical level set of V_N , that is a level set containing at least one critical point of V_N . Around any critical point $q_c^{(i)}$, consider the set of points $\gamma(q_c^{(i)}, \rho; v_c) \subset \Sigma_{v_c}$ at a distance equal to $\rho > 0$ from $q_c^{(i)}$, that is $q \in \gamma(q_c^{(i)}, \rho; v_c) \Rightarrow d(q - q_c^{(i)}) = \rho$, where $d(\cdot, \cdot)$ is the distance measured through the metric induced on Σ_{v_c} by the euclidean metric of the immersion space, and ρ is such that $\rho < \frac{1}{2} \min_{i,j} d(q_c^{(i)} - q_c^{(j)})$, i, j label all the critical points on the given critical level set. Moreover, set the thickness of all the pseudo-cylinders equal to $\varepsilon_0 = \min_{j \in \mathbb{N}} (v_c^{j+1} - v_c^j)$. After Sard Theorem, both ρ and ε_0 are finite because, at finite dimension, there is a finite number of isolated critical points and, consequently, a finite number of critical values. We define a pseudo-cylindrical neighborhood $\Gamma(q_c^{(i)}, \varepsilon_0) \subset M$ of $q_c^{(i)}$ as the open subset of M bounded by the following set of points. By mapping $\gamma(q_c^{(i)}, \rho; v_c)$ from Σ_{v_c} to $\Sigma_{(v_c+\varepsilon_0)}$, and from Σ_{v_c} to $\Sigma_{(v_c-\varepsilon_0)}$, through the flow*

generated by the vector field $X = \nabla V_N(q)/\|\nabla V_N(q)\|^2$, we obtain the walls of $\Gamma(q_c^{(i)}, \varepsilon_0)$, which are transverse to the Σ_v , and then we close the neighborhood with the pieces of $\Sigma_{(v_c+\varepsilon_0)}$ and $\Sigma_{(v_c-\varepsilon_0)}$ bounded by the images $\gamma(q_c^{(i)}, \rho; (v_c+\varepsilon_0))$ and $\gamma(q_c^{(i)}, \rho; (v_c-\varepsilon_0))$ of $\gamma(q_c^{(i)}, \rho; v_c)$ through $\sigma(v, x)$, respectively.

Lemma 1 (Generalization of Corollary 1, paper I). *Let V_N be a standard, smooth, confining, short-range potential bounded from below (Definitions 3–7)*

$$V_N : \mathcal{B}_N \subset \mathbb{R}^N \rightarrow \mathbb{R}$$

with V_N given by Definition 3 (fluid case), or by Definition 4 (lattice case).

Let $\{\Sigma_v\}_{v \in \mathbb{R}}$ be the family of $N-1$ dimensional hypersurfaces $\Sigma_v := V_N^{-1}(v)$, $v \in \mathbb{R}$, of \mathbb{R}^N . Let $\{M_v\}_{v \in \mathbb{R}}$ be the family of N dimensional subsets $M_v := V_N^{-1}((-\infty, v])$, $v \in \mathbb{R}$, of \mathbb{R}^N . Let $\{\overline{M}_v\}_{v \in \mathbb{R}}$ be the family of N dimensional subsets $\overline{M}_v := M_v \setminus \bigcup_{i=1}^{\mathcal{N}(v)} \Gamma(q_c^{(i)}, \varepsilon)$, $v \in \mathbb{R}$, of \mathbb{R}^N , where $\Gamma(q_c^{(i)}, \varepsilon)$ are the pseudo-cylindrical neighborhoods of the critical points $q_c^{(i)}$ of $V_N(q)$ contained in M_v and $\mathcal{N}(v)$ is the number of critical points in M_v . Let $\{\overline{\Sigma}_v\}_{v \in \mathbb{R}}$ be the family of $N-1$ dimensional subsets of \mathbb{R}^N defined as $\overline{\Sigma}_v := \Sigma_v \setminus \bigcup_{i=1}^{\mathcal{N}(v)} [\Gamma(q_c^{(i)}, \varepsilon) \cap \Sigma_v]$. Let $\bar{v}_0 = v_0/N$, $\bar{v}_1 = v_1/N \in \mathbb{R}$, $\bar{v}_0 < \bar{v}_1$ and let $\bar{v}_c = v_c/N$ be the only critical value of V_N in the interval $I_{\bar{v}} = [\bar{v}_0, \bar{v}_1]$, and let $\Gamma^*(q_c^{(i)}, \varepsilon^*)$, with $q_c^{(i)} \in V_N^{-1}(v_c)$ and ε^* such that $\varepsilon^* > \max(v_1 - v_c, v_c - v_0)$. The following two statements hold:

a) for any $\bar{v}, \bar{v}' \in [\bar{v}_0, \bar{v}_1]$ it is

$$\overline{\Sigma}_{N\bar{v}} \text{ is } \mathcal{C}^\infty - \text{diffeomorphic to } \overline{\Sigma}_{N\bar{v}'};$$

b) putting $\overline{M}(v, N) = \text{vol}(\overline{M}_v)$, the quantities $[d\overline{M}(v, N)/dv]/\overline{M}(v, N)$ and $(d^k/dv^k)\{[d\overline{M}(v, N)/dv]/\overline{M}(v, N)\}$, $k = 1, 2, 3$, are uniformly bounded in N in the interval $[\bar{v}_0, \bar{v}_1]$.

Proof. For what concerns point (a), we note that the flow associated with the \mathcal{C}^∞ vector field $X = \nabla V_N(q)/\|\nabla V_N(q)\|^2$ is well defined at any point $q \in \overline{M}_{v_1} \setminus \overline{M}_{v_0}$. Thus the set $\overline{M}_{v_1} \setminus \overline{M}_{v_0}$ is diffeomorphic to the non-critical neck $\partial\overline{M}_{v_0} \times [v_0, v_1]$. Then, after the “non-critical neck theorem” [17], for

any $\bar{v}, \bar{v}' \in I_{\bar{v}} = [\bar{v}_0, \bar{v}_1]$ it is $\bar{\Sigma}_{N\bar{v}} \approx \bar{\Sigma}_{N\bar{v}'}$. Incidentally, this entails also $\bar{M}_{N\bar{v}} \approx \bar{M}_{N\bar{v}'}$ for any $\bar{v}, \bar{v}' \in [\bar{v}_0, \bar{v}_1]$.

Now let us consider point (b). Define $\bar{S}_N(\bar{v}) = \frac{1}{N} \log[\bar{\Omega}(N\bar{v}, N)]$, where $\bar{\Omega}(N\bar{v}, N) = \text{vol}(\bar{\Sigma}_{\bar{v}N})$, having proved the statement (a), we can apply Lemma 4 of paper I which entails that

$$\sup_{N, \bar{v} \in I_{\bar{v}}} |\bar{S}_N(\bar{v})| < \infty \quad \text{and} \quad \sup_{N, \bar{v} \in I_{\bar{v}}} \left| \frac{\partial^k \bar{S}_N}{\partial \bar{v}^k}(\bar{v}) \right| < \infty, \quad k = 1, 2, 3, 4.$$

Whence, after Lemma 3 of paper I, it follows $\bar{S}_{\infty}(\bar{v}) = \lim_{N \rightarrow \infty} \bar{S}_N(\bar{v}) \in \mathcal{C}^3(I_{\bar{v}})$.

The next step is to prove that also $\bar{S}_{\infty}^{(-)}(\bar{v}) \in \mathcal{C}^3(I_{\bar{v}})$, where

$$\bar{S}_{\infty}^{(-)}(\bar{v}) := \lim_{N \rightarrow \infty} \bar{S}_N^{(-)}(\bar{v}) = \lim_{N \rightarrow \infty} \frac{1}{N} \log[\text{vol}(\bar{M}_{\bar{v}N})],$$

because, after Lemmas 3 and 4 of paper I, this entails the truth of statement (b). Let us begin by considering the microcanonical configurational inverse temperature. From its definition $\beta_N(\bar{v}) = \partial S_N^{(-)} / \partial \bar{v}$ one obtains $\beta_N(\bar{v}) = \Omega(N\bar{v}, N) / M(N\bar{v}, N)$. The function $\beta_N(\bar{v})$ is well known to be intensive and well defined also in the thermodynamic limit, at least for extensive potential energy functions. Then we work out a representation of $\beta_N(\bar{v})$ in the form of a microcanonical average of a suitable function. To this purpose we derive $\Omega(N\bar{v}, N)$ with respect to v by means of Federer's derivation formula [15, 20], and then we integrate it. Federer's derivation formula (see Lemma 5 of paper I) states that

$$\frac{d}{dv} \Omega(v, N) = \int_{\Sigma_v} \|\nabla V\| A \left(\frac{1}{\|\nabla V\|} \right) \frac{d\sigma}{\|\nabla V\|} \quad , \quad (12)$$

where A stands for the operator

$$A(\bullet) = \nabla \left(\frac{\nabla V}{\|\nabla V\|} \bullet \right) \frac{1}{\|\nabla V\|} \quad .$$

Then we can write

$$\begin{aligned} \Omega(N\bar{v}, N) &= \int_0^v d\eta \int_{\Sigma_{\eta}} \|\nabla V\| A \left(\frac{1}{\|\nabla V\|} \right) \frac{d\sigma}{\|\nabla V\|} \\ &= \int_{M_v} d\mu \|\nabla V\| A \left(\frac{1}{\|\nabla V\|} \right) \end{aligned} \quad (13)$$

where $d\mu = d^N q$, so that we finally obtain

$$\beta_N(\bar{v}) = \left[\int_{M_v} d\mu \right]^{-1} \int_{M_v} d\mu \|\nabla V\| A\left(\frac{1}{\|\nabla V\|}\right) = \left\langle \|\nabla V\| A\left(\frac{1}{\|\nabla V\|}\right) \right\rangle_{M_v} \quad (14)$$

which holds for $\bar{v} \in [0, \bar{v}_0)$. An important remark is in order. We have used Federer's derivation formula apparently ignoring that it applies in the absence of critical points of the potential function. However, if the potential V is a good Morse function (not a very restrictive condition at all) we know, after Sard theorem [17], that the ensemble of critical values, here of the potential, is a point set. Therefore, any finite interval of values of the potential is the union of a finite number of open intervals where no critical value is present, and correspondingly no critical point on the $\{\Sigma_v\}$ exists. On all these open sets, free of critical points, Federer's derivation formula can be legally applied. Moreover, the results found by applying Federer's formula on each open interval free of critical values of V can be regularly glued together because of the existence of the thermodynamic limit of $\beta_N(\bar{v})$.

Let us now consider $\bar{\Omega}(N\bar{v}, N)$ for $\bar{v} \in [\bar{v}_0, \bar{v}_1]$. As all the hypersurfaces $\bar{\Sigma}_{\bar{v}}$ labeled by $\bar{v} \in [\bar{v}_0, \bar{v}_1]$ are diffeomorphic, we can use Federer's derivation formula to obtain an expression for $\bar{\Omega}(N\bar{v}, N)$ similar to that given in Eq.(13) for $\Omega(N\bar{v}, N)$, that is

$$\begin{aligned} \bar{\Omega}(N\bar{v}, N) &= \int_{v_0}^v d\eta \int_{\bar{\Sigma}_\eta} \|\nabla V\| A\left(\frac{1}{\|\nabla V\|}\right) \frac{d\sigma}{\|\nabla V\|} + \Omega(N\bar{v}_0, N) \\ &= \int_{\bar{M}_v \setminus M_{v_0}} d\mu \|\nabla V\| A\left(\frac{1}{\|\nabla V\|}\right) + \int_{M_{v_0}} d\mu \|\nabla V\| A\left(\frac{1}{\|\nabla V\|}\right) \\ &= \int_{M_v \setminus \Gamma^*} d\mu \|\nabla V\| A\left(\frac{1}{\|\nabla V\|}\right) \end{aligned} \quad (15)$$

where Γ^* stands for the union of all the pseudo-cylindrical neighborhoods of the critical points of V in the interval $[\bar{v}_0, \bar{v}_1]$. Then we consider the restriction $\bar{\beta}_N(\bar{v})$ of the function $\beta_N(\bar{v})$ to the subset $M_v \setminus \Gamma^*$; from

$$\bar{\beta}_N(\bar{v}) = \frac{\bar{\Omega}(N\bar{v}, N)}{\bar{M}(N\bar{v}, N)} \quad (16)$$

we get

$$\begin{aligned}\bar{\beta}_N(\bar{v}) &= \left[\int_{M_v \setminus \Gamma^*} d\mu \right]^{-1} \int_{M_v \setminus \Gamma^*} d\mu \|\nabla V\| A\left(\frac{1}{\|\nabla V\|}\right) \\ &= \left\langle \|\nabla V\| A\left(\frac{1}{\|\nabla V\|}\right) \right\rangle_{M_v \setminus \Gamma^*}.\end{aligned}\quad (17)$$

By comparing Eq.(14) with Eq.(17), we see that also $\bar{\beta}_N(\bar{v})$ has to be intensive up to the $N \rightarrow \infty$ limit, like $\beta_N(\bar{v})$. In fact the excision of the set Γ^* out of M_v , no matter how the measure of Γ^* depends on N , cannot change the intensive character of $\bar{\beta}_N(\bar{v})$. The relationship among $\bar{S}_N(\bar{v})$, $\bar{S}_N^{(-)}(\bar{v})$ and $\bar{\beta}_N(\bar{v})$ is given by the logarithm of both sides of (16)

$$\frac{1}{N} \log \bar{\Omega}(\bar{v}N, N) = \frac{1}{N} \log \bar{M}(\bar{v}N, N) + \frac{1}{N} \log \bar{\beta}_N(\bar{v}).$$

whence, using $\lim_{N \rightarrow \infty} \frac{1}{N} \log \bar{\beta}_N(\bar{v}) = 0$, we obtain $\bar{S}_\infty^{(-)}(\bar{v}) = \bar{S}_\infty(\bar{v})$ and thus $\bar{S}_\infty^{(-)}(\bar{v}) \in \mathcal{C}^3(I_{\bar{v}})$.

Finally, $\bar{S}_\infty^{(-)}(\bar{v}) \in \mathcal{C}^3(I_{\bar{v}})$ entails

$$\sup_{N, \bar{v} \in I_{\bar{v}}} \left| \bar{S}_N^{(-)}(\bar{v}) \right| < \infty \quad \text{and} \quad \sup_{N, \bar{v} \in I_{\bar{v}}} \left| \frac{\partial^k \bar{S}_N^{(-)}(\bar{v})}{\partial \bar{v}^k} \right| < \infty, \quad k = 1, 2, 3, 4.$$

so that, resorting to Lemma 3 of paper I, the truth of statement (b) follows. \square

3. MAIN THEOREM

In this Section we prove the following

Theorem 1 (Entropy and Topology). *Let $V_N(q_1, \dots, q_N) : \mathbb{R}^N \rightarrow \mathbb{R}$, be a smooth, non-singular, finite-range potential. Denote by $M_v := V_N^{-1}((-\infty, v])$, $v \in \mathbb{R}$, the generic submanifold of configuration space bounded by Σ_v .*

Let $\{q_c^{(i)} \in \mathbb{R}^N\}_{i \in [1, \mathcal{N}(v)]}$ be the set of critical points of the potential, that is s.t. $\nabla V_N(q)|_{q=q_c^{(i)}} = 0$, and $\mathcal{N}(v)$ be the number of critical points up to the potential energy value v . Denote by $\bar{v} = v/N$ the potential energy density, and assume that for any given interval $[\bar{v}_0, \bar{v}_1]$ the number of critical values \bar{v}_c^j contained in it is at most a linearly growing function of N . Let $\Gamma(q_c^{(i)}, \varepsilon_0)$ be the pseudo-cylindrical neighborhood of the critical point $q_c^{(i)}$, and $\mu_i(M_v)$ be

the Morse indexes of M_v , then there exist real numbers $A(N, i, \varepsilon_0)$, g_i and real smooth functions $v \mapsto B(N, i, v, \varepsilon_0)$ such that the following equation for the microcanonical configurational entropy $S_N^{(-)}(v)$ holds

$$S_N^{(-)}(v) = \frac{1}{N} \log \left[\int_{M_v \setminus \bigcup_{i=1}^{\mathcal{N}_{cp}^{\nu(v)}} \Gamma(q_c^{(i)}, \varepsilon_0)} d^N q + \sum_{i=0}^N A(N, i, \varepsilon_0) g_i \mu_i(M_{v-\varepsilon_0}) + \sum_{n=1}^{\mathcal{N}_{cp}^{\nu(v)}} B(N, i(n), v - v_c^{\nu(v)}, \varepsilon_0) \right], \quad (18)$$

where $\nu(v) = \max\{j | v_c^j \leq v\}$, and an unbound growth with N of one of the derivatives $|\partial^k S^{(-)}(v)/\partial v^k|$, for $k = 3, 4$, and thus the occurrence of a first or of a second order phase transition respectively, can be entailed only by the topological term $\sum_{i=0}^N A(N, i, \varepsilon_0) g_i \mu_i(M_{v-\varepsilon_0}) + \sum_{n=1}^{\mathcal{N}_{cp}^{\nu(v)}} B(N, i(n), v - v_c^{\nu(v)}, \varepsilon_0)$.

The proof of formula (18) is worked out constructively. This formula relates thermodynamic entropy, defined in the microcanonical configurational ensemble, with quantities of topological meaning (the Morse indexes) of the configuration space submanifolds $M_v = V_N^{-1}((-\infty, v]) = \{q = (q_1, \dots, q_N) \in \mathbb{R}^N | V_N(q) \leq v\}$.

After Morse theory, topology changes of the manifolds M_v can be put in one-to-one correspondence with the existence of critical points of the potential function $V_N(q_1, \dots, q_N)$. A point q_c is a *critical point* if $\nabla V_N(q)|_{q=q_c} = 0$. The potential energy value $v_c = V_N(q_c)$ is said to be a *critical value* for the potential function. Passing a critical value v_c , the manifolds M_v change topology. Within the framework of Morse theory, if the potential V_N is a *good Morse function*, that is a regular function bounded below and with non-degenerate Hessian (that is the Hessian has no vanishing eigenvalue), then topology changes occur through the *attachment of handles* in the neighborhoods of the critical points. Therefore, in order to establish the relationship between entropy and configuration space topology, we have to unfold the contribution given to the volume of M_v by suitably defined neighborhoods of all the critical points contained

in M_v because it is within these neighborhoods that the relevant information about topology is contained.

This result is made possible by the idea of exploiting the existence of the so-called Morse chart in the neighborhood of any nondegenerate critical point of the potential function V_N . In fact, the Morse chart allows to represent the *local* analytic form of the equipotential hypersurfaces in an universal form independent of the potential energy value at the critical point, and only dependent upon the index of the critical point (equal to the number of negative eigenvalues of the Hessian of the potential) and, obviously, upon the dimension N of configuration space. Hence the possibility of a formal computation of the contribution of the neighborhoods of all the critical points to the volume of M_v as a function of v .

Proof. Let us consider the definition of the configurational microcanonical entropy $S_N^{(-)}(v)$, already given in Eq.(2),

$$S_N^{(-)}(v) = \frac{1}{N} \log M(v, N) , \quad (19)$$

with

$$\begin{aligned} M(v, N) &= \int_{V_N(q) \leq v} d^N q = \int_0^v d\eta \int_{(\Lambda^d)^{\times n}} d^N q \delta[V_N(q) - \eta] \\ &= \int_0^v d\eta \int_{\Sigma_\eta} \frac{d\sigma}{\|\nabla V_N\|} , \end{aligned} \quad (20)$$

where we have set equal to zero the minimum of the potential. Let $\{\Sigma_{v_c^j}\}$ be the family of all the critical level sets (in general not manifolds) of the potential, that is the constant potential energy hypersurfaces that contain at least one *critical point* $q_c^{(i)}$, where $\nabla V_N(q)|_{q=q_c^{(i)}} = 0$. For a potential which is a good Morse function, after the Sard Theorem (see Corollary 2 at p.200 of Ref.[17]), at any finite dimension N , and below any finite upper bound of the potential energy, the number of critical points in configuration space and thus also the set of critical values $\{v_c^j\}_{j \in \mathbb{N}}$, are *finite, isolated* and such that $v_c^j < v_c^k$ if $j < k$, so that any energy interval $v_0 \leq v \leq v_1$ is the union of a finite number of open intervals free of critical points.

In order to split the integration on M_v into two parts: the integration on the union of the neighborhoods of all the critical points contained in M_v and the integration on its complement in M_v , we have defined for each critical point $q_c^{(i)}$ its *pseudo-cylindrical* neighborhood $\Gamma(q_c^{(i)}, \varepsilon_0)$; ε_0 is the thickness – in potential energy – of the neighborhood. The assumption that the number of critical values \bar{v}_c^j is at most linearly growing with N entails, together with Sard theorem, that ε_0 is finite.

Let us now split the integration on M_v into the integration on $M_v \cap \bigcup_i \Gamma(q_c^{(i)}, \varepsilon_0)$ and on its complement $M_v \setminus \bigcup_i \Gamma(q_c^{(i)}, \varepsilon_0)$. We have

$$\int_{M_v} d^N q = \int_{M_v \setminus \bigcup_{i=1}^{\mathcal{N}(v+\varepsilon_0)} \Gamma(q_c^{(i)}, \varepsilon_0)} d^N q + \int_{M_v \cap \bigcup_{i=1}^{\mathcal{N}(v+\varepsilon_0)} \Gamma(q_c^{(i)}, \varepsilon_0)} d^N q \quad , \quad (21)$$

where $\mathcal{N}(v)$ is the number of critical points of $V_N(q)$ up to the level v . We can equivalently write

$$\text{vol}(M_v) = \int_{M_v \setminus \bigcup_{i=1}^{\mathcal{N}(v+\varepsilon_0)} \Gamma(q_c^{(i)}, \varepsilon_0)} d^N q + \sum_{j=1}^{\mathcal{N}_{cl}(v+\varepsilon_0)} \sum_{m=1}^{\mathcal{N}_{cp}^j} \int_{M_v \cap \Gamma_j(q_c^{(m)}, \varepsilon_0)} d^N q \quad (22)$$

where $\mathcal{N}_{cl}(v)$ is the number of critical levels $\Sigma_{v_c^j}$ such that $v_c^j < v$, and \mathcal{N}_{cp}^j is the number of critical points on the critical hypersurface $\Sigma_{v_c^j}^j$ and where we have changed the notation of the pseudo-cylindrical neighborhoods to $\Gamma_j(q_c^{(m)}, \varepsilon_0)$ labelling with j the level set to which it belongs and numbering with m the critical points on the j -th level set. Notice that $\mathcal{N}(v) = \sum_{j=1}^{\mathcal{N}_{cl}(v)} \mathcal{N}_{cp}^j$.

Then we use the co-area formula in the r.h.s. of Eq.(20) to rewrite Eq.(22); a distinction is necessary between two cases for $\Sigma_v = \partial M_v$: its label v is closer than ε_0 to a critical level or not; thus we obtain

$$\begin{aligned} \text{vol}(M_v) = & \int_{M_v \setminus \bigcup_{i=1}^{\mathcal{N}(v+\varepsilon_0)} \Gamma(q_c^{(i)}, \varepsilon_0)} d^N q \\ & + \sum_{j=1}^{\mathcal{N}_{cl}(v+\varepsilon_0)} \sum_{m=1}^{\mathcal{N}_{cp}^j} \int_{v_c^j - \varepsilon_0}^{v_c^j + \varepsilon_0} d\eta \int_{\Gamma_j(q_c^{(m)}, \varepsilon_0)} d^N q \delta[V_N(q) - \eta] \end{aligned} \quad (23)$$

when $v > v_c^{\nu(v)} + \varepsilon_0$ and $v < v_c^{\nu(v)+1} - \varepsilon_0$, where $\nu(v)$ is such that $v_c^{\nu(v)} < v <$

$v_c^{\nu(v)+1}$; whereas

$$\begin{aligned}
\text{vol}(M_v) &= \int_{M_v \setminus \bigcup_{i=1}^{\mathcal{N}^{(v+\varepsilon_0)}} \Gamma(q_c^{(i)}, \varepsilon_0)} d^N q \\
&+ \sum_{j=1}^{\mathcal{N}_{cl}(v)-1} \sum_{m=1}^{\mathcal{N}_{cp}^j} \int_{v_c^j - \varepsilon_0}^{v_c^j + \varepsilon_0} d\eta \int_{\Gamma_j(q_c^{(m)}, \varepsilon_0)} d^N q \delta[V_N(q) - \eta] \\
&+ \sum_{m=1}^{\mathcal{N}_{cp}^{\nu(v+\varepsilon_0)}} \int_{v_c^{\nu(v)} - \varepsilon_0}^v d\eta \int_{\Gamma_{\nu(v)}(q_c^{(m)}, \varepsilon_0)} d^N q \delta[V_N(q) - \eta] . \quad (24)
\end{aligned}$$

when $v_c^{\nu(v)+1} - \varepsilon_0 < v$ or $v < v_c^{\nu(v)} + \varepsilon_0$.

Near to any critical point, a second order power series expansion of $V(q)$ reads

$$V_N^{(2)}(q) = V_N(q_c) + \frac{1}{2} \sum_{i,j} \frac{\partial^2 V_N}{\partial q_i \partial q_j} (q^i - q_c^i) (q^j - q_c^j)$$

For sufficiently small ε_0 , the integrals $\int_{\Gamma_j(q_c, \varepsilon_0)} d^N q \delta[V_N(q) - \eta]$ can be replaced with arbitrary precision by $\int_{\Gamma_j(q_c, \varepsilon_0)} d^N q \delta[V_N^{(2)}(q) - \eta]$. Moreover, if $V_N(q)$ is a good Morse function, then a coordinate transformation exists to the so-called *Morse chart* [18] such that

$$\tilde{V}_N^{(2)}(x) = V_N(q_c) - \sum_{l=1}^k x_l^2 + \sum_{l=k+1}^N x_l^2$$

where k is the Morse index of q_c . Using Morse chart we have

$$\int_{\Gamma_j(q_c, \varepsilon_0)} d^N q \delta[V_N^{(2)}(q) - \eta] = \int_{\Gamma_j(q_c, \varepsilon_0)} d^N x |\det J| \delta[\tilde{V}_N^{(2)}(x) - \eta] \quad (25)$$

where J is the Jacobian of the coordinate transformation.

Using Morse coordinates inside the pseudo-cylinder $\Gamma_j(q_c^{(m)}, \varepsilon_0)$ around the critical point $q_c^{(m)}$, we see that each part of an hypersurface $\Sigma_\eta \cap \Gamma_j(q_c^{(m)}, \varepsilon_0)$ is a quadric

$$\xi = \eta - v_c^j = - \sum_{l=1}^{k_m} x_l^2 + \sum_{l=k_m+1}^N x_l^2 = - |X|^2 + |Y|^2 \quad , \quad (26)$$

where the Morse index of $q_c^{(m)}$ is denoted by k_m , so that $|X|^2 = \sum_{l=1}^{k_m} x_l^2$ and

$|Y|^2 = \sum_{l=k_m+1}^N x_l^2$. Thus we rewrite the r.h.s. of Eq.(25) as

$$\begin{aligned}
& |\det J| \int_{\Gamma_j(q_c^{(m)}, \varepsilon_0)} d^N x \delta(-|X|^2 + |Y|^2 - \xi) \\
&= |\det J| \int_{\Gamma_j(q_c^{(m)}, \varepsilon_0)} d\Omega^{k_m-1} d\Omega^{N-k_m-1} d|X| d|Y| |X|^{k_m-1} |Y|^{N-k_m-1} \delta(-|X|^2 + |Y|^2 - \xi)
\end{aligned} \tag{27}$$

where $d\Omega^r$ is the solid angle element in r dimension, whose integration yields the surface C_r of the r -dimensional hypersphere of unit radius. Putting $z = |X|^2$ and integrating on the angular coordinates we get

$$\frac{1}{2} |\det J| C_{N-k_m-1} C_{k_m-1} \int_0^{\alpha(\xi, r)} dz \int_{\sqrt{\xi}}^{\beta(\xi, r)} d|Y| |Y|^{N-k_m-1} z^{(k_m-2)/2} \delta(-z + |Y|^2 - \xi) \tag{28}$$

where $\alpha(\xi, r) = (\sqrt{\xi^2 + 4r^2} - \xi)/2$ and $\beta(\xi, r) = \sqrt{(\sqrt{\xi^2 + 4r^2} + \xi)/2}$. These expressions stem from the definition of $\Gamma_j(q_c, \varepsilon_0)$ whose boundaries have to be orthogonal to the potential level sets described by Eq.(26). These boundaries are given by the equation $|X||Y| = r$. Putting $y = |Y|$, from Eq.(28) when $\xi > 0$ we obtain

$$\frac{1}{2} J_{jm} C_{N-k_m-1} C_{k_m-1} \int_{\sqrt{\xi}}^{\beta(\xi, r)} dy y^{N-k_m-1} (y^2 - \xi)^{(k_m-2)/2} \tag{29}$$

and when $\xi < 0$ we obtain

$$\frac{1}{2} J_{jm} C_{N-k_m-1} C_{k_m-1} \int_0^{\beta(\xi, r)} dy y^{N-k_m-1} (y^2 - \xi)^{(k_m-2)/2} \tag{30}$$

where C_{N-k_m-1} and C_{k_m} are surfaces of hyperspheres of unit radii, that is $C_n = 2\pi^{n/2}/(n/2 - 1)!$ (for n even) and $C_n = 2^{(n+1)/2}\pi^{(n-1)/2}/(n-2)!!$ (for n odd); J_{jm} stands for the numerical absolute value of the determinant of J computed at the critical level v_c^j and at the critical point $q_c^{(m)}$. By defining

$$F_+(\xi, k_m, N) = \int_{\sqrt{\xi}}^{\beta(\xi, r)} dy y^{N-k_m-1} (y^2 - \xi)^{(k_m-2)/2} \tag{31}$$

and

$$F_-(\xi, k_m, N) = \int_0^{\beta(\xi, r)} dy y^{N-k_m-1} (y^2 - \xi)^{(k_m-2)/2} \tag{32}$$

we can now write

$$\begin{aligned} \text{vol}(M_v) &= \int_{M_v \setminus \bigcup_{i=1}^{\mathcal{N}(v+\varepsilon_0)} \Gamma(q_c^{(i)}, \varepsilon_0)} d^N q \\ &+ \sum_{j=1}^{\mathcal{N}_{cl}(v+\varepsilon_0)} \sum_{m=1}^{\mathcal{N}_{cp}^j} \frac{1}{2} C_{N-k_m-1} C_{k_m-1} J_{jm} \int_{-\varepsilon_0}^{\varepsilon_0} d\xi F(\xi, k_m, N). \end{aligned} \quad (33)$$

when $v > v_c^{\nu(v)} + \varepsilon_0$ and $v < v_c^{\nu(v)+1} - \varepsilon_0$, where $\nu(v) = \max\{j | v_c^j \leq v\}$, or

$$\begin{aligned} \text{vol}(M_v) &= \int_{M_v \setminus \bigcup_{i=1}^{\mathcal{N}(v+\varepsilon_0)} \Gamma(q_c^{(i)}, \varepsilon_0)} d^N q \\ &+ \sum_{j=1}^{\mathcal{N}_{cl}(v)-1} \sum_{m=1}^{\mathcal{N}_{cp}^j} \frac{1}{2} C_{N-k_m-1} C_{k_m-1} J_{jm} \int_{-\varepsilon_0}^{\varepsilon_0} d\xi F(\xi, k_m, N) \\ &+ \sum_{m=1}^{\mathcal{N}_{cp}^{\nu(v)+\varepsilon_0}} \frac{1}{2} C_{N-k_m-1} C_{k_m-1} J_{\tilde{j}m} \int_{-\varepsilon_0}^{v-v_c^{\nu(v)}} d\xi F(\xi, k_m, N) \end{aligned} \quad (34)$$

with $\tilde{j} = \mathcal{N}_{cl}(v)$, when $v_c^{\nu(v)+1} - \varepsilon_0 < v$ or $v < v_c^{\nu(v)} + \varepsilon_0$. In Eqs.(33) and (34) we have put $\int_{-\varepsilon_0}^{\varepsilon_0} d\xi F(\xi, k_m, N) = \int_{-\varepsilon_0}^0 d\xi F_-(\xi, k_m, N) + \int_0^{\varepsilon_0} d\xi F_+(\xi, k_m, N)$.

Notice that it is $\mathcal{N}(v) = \sum_{i=0}^N \mu_i(M_v)$, where $\mu_i(M_v)$ are the multiplicities of the critical points of index i (there are at most $N+1$ values for the indexes of critical points at dimension N) below the energy value v .

Therefore, we can rearrange the double summation in Eqs.(33), (34) by expressing it as a double summation on all the possible values of the Morse indexes and on the number of critical points for each value of the Morse index, that is

$$\sum_{i=0}^N \sum_{k=1}^{\mu_i(M_v)} A(N, i, \varepsilon_0) J_{j(i,k)m(i,k)} \quad (35)$$

where, since the integrals in Eqs.(33), (34) are independent of the index j , we have defined a set of positive coefficients $A(N, i, \varepsilon_0)$ as

$$A(N, i, \varepsilon_0) = \frac{1}{2} C_{N-i-1} C_i \int_{-\varepsilon_0}^{\varepsilon_0} d\xi F(\xi, i, N). \quad (36)$$

We remark that the term (35), being a function of the Morse indexes $\mu_i(M_v)$, has topological meaning. In order to make clearer this topological meaning, we rewrite (35) in the equivalent form of a weighed sum of Morse indexes as

follows. From the set of positive numbers $J_{j(i,k)m(i,k)}$ we define

$$g_i = \frac{1}{\mu_i(M_v)} \sum_{k=1}^{\mu_i(M_v)} J_{j(i,k)m(i,k)} \quad (37)$$

and rewrite the second term of the r.h.s. of Eq.(33) as

$$\sum_{i=0}^N A(N, i, \varepsilon_0) g_i \mu_i(M_v). \quad (38)$$

Equation (35) is here recast in a form which makes somewhat clearer its dependence on Morse indexes. Moreover, we introduce also the coefficients

$$B(N, i, v - v_c^{\nu(v)}, \varepsilon_0) = \frac{1}{2} C_{N-i-1} C_i J_{\tilde{j}m(i,k(i))} \int_{-\varepsilon_0}^{v-v_c^{\nu(v)}} d\xi F(\xi, i, N), \quad (39)$$

where $k(i)$ stems from $j(i, k) = \tilde{j}$, such that for $v = v_c^{\nu(v)}$ it is $B(N, i, 0, \varepsilon_0) = 0$, and for $v - v_c^{\nu(v)} = \varepsilon_0$ it is $B(N, i, \varepsilon_0, \varepsilon_0) = A(N, i, \varepsilon_0) g_i$.

For the purposes of the present proof, we are not concerned about the complication of the coefficients $A(N, i, \varepsilon_0)$ and $B(N, i, v - v_c^{\nu(v)}, \varepsilon_0)$ because all what we need, in order to make the link between configuration space topology and thermodynamics, is that the second term in the volume splitting in Eq.(21) can be written in the form (38). In fact, now we can write the entropy per degree of freedom as

$$\begin{aligned} S_N^{(-)}(v) &= \frac{1}{N} \log M(v, N) = \frac{1}{N} \log \int_{M_v} d^N q \\ &= \frac{1}{N} \log \left[\int_{M_v \setminus \bigcup_{i=1}^{\mathcal{N}(v+\varepsilon_0)} \Gamma(q_c^{(i)}, \varepsilon_0)} d^N q + \int_{M_v \cap \bigcup_{i=1}^{\mathcal{N}(v+\varepsilon_0)} \Gamma(q_c^{(i)}, \varepsilon_0)} d^N q \right] \\ &= \frac{1}{N} \log \left[\int_{M_v \setminus \bigcup_{i=1}^{\mathcal{N}(v)} \Gamma(q_c^{(i)}, \varepsilon_0)} d^N q + \sum_{i=0}^N A(N, i, \varepsilon_0) g_i \mu_i(M_v) \right], \end{aligned} \quad (40)$$

when $v < v_c^{\nu(v)+1} - \varepsilon_0$ or $v > v_c^{\nu(v)} + \varepsilon_0$, or

$$\begin{aligned} S_N^{(-)}(v) &= \frac{1}{N} \log \left[\int_{M_v \setminus \bigcup_{i=1}^{\mathcal{N}(v+\varepsilon_0)} \Gamma(q_c^{(i)}, \varepsilon_0)} d^N q + \sum_{i=0}^N A(N, i, \varepsilon_0) g_i \mu_i(M_{v-\varepsilon_0}) \right. \\ &\quad \left. + \sum_{n=1}^{\mathcal{N}_{cp}^{\nu(v+\varepsilon_0)+1}} B(N, i(n), v - v_c^{\nu(v)}, \varepsilon_0) \right], \end{aligned} \quad (41)$$

when $v > v_c^{\nu(v)+1} - \varepsilon_0$ or $v < v_c^{\nu(v)} + \varepsilon_0$.

The equation above links thermodynamic entropy with the Morse indexes of the configuration space submanifolds M_v , that is with their topology. In fact, according to Bott's "critical-neck theorem" [21], any change with v of any index $\mu_i(M_v)$, $i = 0, \dots, N$, which can only be due to the crossing of a critical level, is associated with a topology change of the M_v .

Conversely, any topology change, in the sense of a loss of diffeomorphicity, occurring to the M_v when v is varied, is signaled by one or more changes of the Morse indexes $\mu_i(M_v)$ because, after the "non-critical neck theorem" [17], this has to be the consequence of the crossing of a critical level.

We remark that the two terms $\sum A_i g_i \mu_i$ and $\sum_i B_i$ in Eq.(41) stem from the same term in the volume splitting (21) (the union of neighborhoods of critical points), so that they will both participate in producing the development of singularities proper to a phase transition (for more details about the role of these two terms, see Remark 2).

Let us now show that the coefficients $B(N, i, v - v_c^{\nu(v)}, \varepsilon_0)$ are smooth functions of v at any finite N .

Noting that $d^k B(N, i, v, \varepsilon_0)/dv^k = d^{(k-1)} F(\xi, k_m, N)/d\xi^{(k-1)}$, we focus on the smoothness of $F_{\pm}(\xi, k_m, N)$. Let us consider $F_+(\xi, k_m, N)$. There are two cases: k_m is even so that $(k_m - 2)/2$ is an integer; k_m is odd so that $(k_m - 2)/2 = n + 1/2$ with $n \in \{-1\} \cup \mathbb{N}$. In the first case, by iteratively applying the derivation formula $\frac{d}{d\alpha} \int_{\phi(\alpha)}^{\psi(\alpha)} dx f(x, \alpha) = \frac{d\psi(\alpha)}{d\alpha} f(\psi(\alpha), \alpha) - \frac{d\phi(\alpha)}{d\alpha} f(\phi(\alpha), \alpha) + \int_{\phi(\alpha)}^{\psi(\alpha)} dx \frac{\partial f(x, \alpha)}{\partial \alpha}$ to Eq. (31), one immediately realizes that $F(\xi, k_m, N)$ is smooth.

In the second case, Eq.(31) is rewritten as

$$\mathcal{I}(\xi, n, N) = \int_{\sqrt{\xi}}^{\beta(\xi, r)} dy y^{N-k_m-1} (y^2 - \xi)^{n+\frac{1}{2}} \quad (42)$$

which, after integration by parts, yields the recursion formula

$$\mathcal{I}(\xi, n, N) = \frac{(y^2 - \xi)^{n+\frac{3}{2}} y^{N-1}}{2 + 2n + N} \Big|_{y=\beta(\xi, r)} + \frac{(N-1)\xi}{2 + 2n + N} \mathcal{I}(\xi, n, N-2) . \quad (43)$$

For this recursion formula there are two possible initial conditions. The first initial condition is obtained by direct integration of $\mathcal{I}(\xi, n, N)$ for $N = k_m + 2$ and $n > 0$, and is

$$\mathcal{I}(\xi, n, N = k_m + 2) = \frac{(y^2 - \xi)^{n+\frac{3}{2}}}{3 + 2n} \Big|_{y=\beta(\xi, r)} ; \quad (44)$$

the second initial condition for the recursion (43) is obtained by working out $\mathcal{I}(\xi, n, N)$ for $N = k_m + 1$ and $n > 0$. From the above definition of $\mathcal{I}(\xi, n, N)$ we get

$$\mathcal{I}(\xi, n, N = k_m + 1) = \frac{(y^2 - \xi)^{n+\frac{3}{2}}}{2(n+1)y} \Big|_{y=\beta(\xi, r)} - \frac{\xi}{2(n+1)} \int_{\sqrt{\xi}}^{\beta(\xi, r)} dy \frac{(y^2 - \xi)^{n+\frac{1}{2}}}{y^2} \quad (45)$$

where the integral in the r.h.s. – that we denote by $I(n)$ – can be solved by introducing the double recursion

$$\begin{aligned} I(n) &= \frac{2(y^2 - \xi)^{n+\frac{3}{2}}}{(4n+3)y^3} \Big|_{y=\beta(\xi, r)} - \frac{3\xi}{4n+3} H(n) \\ H(n) &= I(n-1) - \xi H(n-1) \end{aligned} \quad (46)$$

which leads to

$$H(n+1) = \frac{2(y^2 - \xi)^{n+\frac{3}{2}}}{(4n+3)y^3} \Big|_{y=\beta(\xi, r)} - \left(\frac{3\xi}{4n+3} + \xi \right) H(n) \quad (47)$$

together with

$$H(1) = \left(\frac{\xi}{3y^3} - \frac{4}{3y} \right) \sqrt{y^2 - \xi} + \log(y + \sqrt{y^2 - \xi}) \Big|_{y=\beta(\xi, r)} - \log(\sqrt{\xi}). \quad (48)$$

From Eq. (48) and by Eq. (47) the system of double recursion (46) is solved. From this we get the required second initial condition to be inserted in Eq. (43). Now, notice that all the terms that are either powers of $(y^2 - \xi)$ or simply powers of y , are to be computed at $y = \beta(\xi, r) = \sqrt{(\sqrt{\xi^2 + 4r^2} + \xi)/2}$, so that they are never vanishing functions of ξ , as a consequence these terms are infinitely many times differentiable with respect to ξ . The apparent singularity of $H(1)$ for $\xi = 0$ is cured in Eq. (46) where it enters multiplied by ξ . We

remark that the recursion formula given in Eq. (43) holds also for $n = -1, 0$ which correspond to Morse indexes $k_m = 1, 2$ respectively; the initial conditions for the recursion are to be explicitly computed from Eq. (42) by substituting $n = -1$ or $n = 0$. Also for these special values of n , one is left with manifestly smooth functions, provided that $N > 2$, an obviously acceptable "restriction" in our context. This concludes the proof of the smoothness of the coefficients $B(N, i, v - v_c^{\nu(v)}, \varepsilon_0)$.

Let us now come to the proof of the statement of Theorem 1 which says that the source of a phase transition can only be the second term in square brackets in Eq.(18), which is of topological meaning. To this purpose we have to resort to the Main Theorem of paper I and to its Corollary 1.

Let us assume that only one critical value \bar{v}_c exists in a given interval $[\bar{v}_0, \bar{v}_1]$. After Sard theorem [17], at any finite N there is a finite number of isolated critical points on $\Sigma_{N\bar{v}_c}$. For any arbitrarily small $\delta > 0$, the Main Theorem of paper I and its Corollary 1 still apply to the two subintervals $[\bar{v}_0, \bar{v}_c - \delta]$ and $[\bar{v}_c + \delta, \bar{v}_1]$. In order to understand why a breakdown of uniform boundedness in N of $|\partial^k S_N^{(-)} / \partial v^k|$ for $k = 3$ or $k = 4$ can be originated only by the topological term in r.h.s. of Eq.(18), we consider each critical point $q_c^{(i)}$ on $\Sigma_{N\bar{v}_c}$ enclosed in a small pseudo-cylindrical neighborhood $\Gamma(q_c^{(i)}, \varepsilon_0)$ of thickness ε_0 and we take $\varepsilon < \varepsilon_0$ arbitrarily close to ε_0 . From Morse theory, we know that passing a critical value v_c entails that for each critical point of index k_i a k_i -handle H^{N, k_i} is attached to $M_{v < v_c}$ so that the following diffeomorphism holds

$$M_{(v_c + \varepsilon_0)} \approx M_{(v_c - \varepsilon_0)} \bigcup_{\phi_1} H^{N, k_1} \bigcup_{\phi_2} H^{N, k_2} \dots \bigcup_{\phi_n} H^{N, k_n} \quad (49)$$

where a k_i -handle in N dimensions ($0 \leq k_i \leq N$) is the product of two disks, a k_i -dimensional disk D^{k_i} , and another $(N - k_i)$ -dimensional disk $D^{(N - k_i)}$ s.t. $H^{N, k_i} = D^{k_i} \times D^{(N - k_i)}$, and where \bigcup_{ϕ_i} stands for the attachment of H^{N, k_i} to $M_{(v_c - \varepsilon_0)}$ through the embedding $\phi_i : \mathbb{S}^{k_i - 1} \times D^{N - k_i} \rightarrow \partial M_{(v_c - \varepsilon_0)}$ (where \mathbb{S} is an hypersphere; details can be found in [17, 18, 19]).

The excision of the pseudo-cylindrical neighborhoods $\Gamma(q_c^i, \varepsilon_0)$ of all the

critical points $q_c^{(i)} \in \Sigma_{N\bar{v}_c}$ implies that all the manifolds $\overline{M}_v := M_v \setminus \bigcup_{i=1}^{\# \text{ crit.pts.}} \Gamma(q_c^{(i)}, \varepsilon_0)$ with $N\bar{v}_c - \varepsilon_0 < v < N\bar{v}_c + \varepsilon_0$ are free of critical points and, consequently, are diffeomorphic. In fact, for any $v, v' \in \mathbb{R}$ such that $N\bar{v}_c - \varepsilon_0 < v < v' < N\bar{v}_c + \varepsilon_0$, \overline{M}_v is a deformation retraction of $\overline{M}_{v'}$ through the flow associated with the vector field $X = -\nabla V_N / \|\nabla V_N\|^2$ [17, 18].

Now, defining $\overline{M}(v, N) = \text{vol}(\overline{M}_v)$ and $\Gamma(v, N) = \text{vol}[\bigcup_{i=1}^{\# \text{ crit.pts.}} \Gamma(q_c^{(i)}, \varepsilon_0)]$, equation (18) becomes

$$\begin{aligned} S_N^{(-)}(v) &= \frac{1}{N} \log [\overline{M}(v, N) + \Gamma(v, N)] \\ &= \frac{1}{N} \log [\overline{M}(v, N)] + \frac{1}{N} \log \left[1 + \frac{\Gamma(v, N)}{\overline{M}(v, N)} \right]. \end{aligned} \quad (50)$$

By applying the Main Theorem of paper I and its Corollary 1 to the first term in the r.h.s. of the equation above, we know that $\frac{1}{N} |\partial^k \log [\overline{M}(v, N)] / \partial v^k|$ for $k = 1, \dots, 4$, are uniformly bounded in N , and thus no phase transition can be attributed to this term.

Then, let us consider the second term of the r.h.s. of the equation above. By computing its first derivative we obtain

$$\frac{d}{dv} \frac{1}{N} \log \left[1 + \frac{\Gamma(v, N)}{\overline{M}(v, N)} \right] = \frac{1}{N} \frac{\Gamma'}{\overline{M} + \Gamma} - \frac{1}{N} \frac{\Gamma}{\overline{M} + \Gamma} \left(\frac{\overline{M}'}{\overline{M}} \right), \quad (51)$$

where $(\overline{M}'/\overline{M})$ stands for $[d\overline{M}(v, N)/dv]/\overline{M}(v, N)$. After Lemma 1, $(\overline{M}'/\overline{M})$ is uniformly bounded in N and therefore so does the second term in the r.h.s. of Eq.(51). Whence, if $|\partial S_N^{(-)}/\partial v|$ were to grow with N this could not be due to the term $\overline{M}(v, N)$.

Then we compute the second derivative

$$\begin{aligned} \frac{d^2}{dv^2} \frac{1}{N} \log \left[1 + \frac{\Gamma(v, N)}{\overline{M}(v, N)} \right] &= \frac{1}{N} \frac{\Gamma''}{\overline{M} + \Gamma} + \frac{1}{N} \frac{\Gamma \overline{M} (\overline{M}'/\overline{M}) + \Gamma \Gamma'}{(\overline{M} + \Gamma)^2} \left(\frac{\overline{M}'}{\overline{M}} - \frac{\Gamma'}{\Gamma} \right) \\ &\quad - \frac{1}{N} \frac{\Gamma}{\overline{M} + \Gamma} \left(\frac{\overline{M}'}{\overline{M}} \right) + \frac{1}{N} \frac{d}{dv} \left(\frac{\overline{M}'}{\overline{M}} \right). \end{aligned} \quad (52)$$

Again, we can observe that the uniform boundedness with N of both $(\overline{M}'/\overline{M})$ and $(d/dv)(\overline{M}'/\overline{M}) = (\overline{M}''/\overline{M}) - (\overline{M}'/\overline{M})^2$ – after Lemma 1 – entails that if $|\partial^2 S_N^{(-)}/\partial v^2|$ were to grow with N , this could not be due to the term $\overline{M}(v, N)$.

Similarly, after a lengthy but trivial computation of the third and fourth derivatives of the second term in the r.h.s. of Eq.(50), one finds that $\overline{M}(v, N)$ enters the various terms obtained through the ratio $\overline{M}'/\overline{M}$ and through its derivatives $\frac{d^k}{dv^k}[\overline{M}'/\overline{M}]$ with $k = 1, 2, 3$, thus, as a consequence of Lemma 1, the uniform boundedness in N of $[d\overline{M}(v, N)/dv]/\overline{M}(v, N)$ and $\frac{d^k}{dv^k}\{[d\overline{M}(v, N)/dv]/\overline{M}(v, N)\}$ with $k = 1, 2, 3$, implies that if $|\partial^3 S_N^{(-)}/\partial v^3|$ or $|\partial^4 S_N^{(-)}/\partial v^4|$ were to grow with N , this could not be due to the term $\overline{M}(v, N)$.

In conclusion, the first term within square brackets in Eq.(18) cannot be at the origin of a phase transition, nor can it be the third one, which is the sum of smooth functions. Only the second term of the r.h.s. of Eq.(18), which is in one-to-one correspondence with topology changes of the M_v , can originate an unbound growth with N of a derivative $|\partial^k S_N^{(-)}/\partial v^k|$ for some k , thus entailing a phase transition. \square

Remark 2. *A comment about the above considered volume splitting is in order. At any finite N , the volume $M(v, N)$ is a smooth function of v , as is the entropy $S_N^{(-)}(v)$. Since the term $\sum_{i=0}^N A(N, i, \varepsilon_0) g_i \mu_i(M_v)$ entering Eqs.(40) and (41) depends on the integer valued functions $\mu_i(v)$, one could erroneously think that the presence of the functions $\mu_i(v)$ in this term conflicts with the smoothness of volume and entropy. Of course, at finite N , smoothness of volume and entropy is not lost. In fact, the term $\sum_{i=0}^N A(N, i, \varepsilon_0) g_i \mu_i(M_v)$ is constant in any open interval $(v_c^{\nu(v)} + \varepsilon_0, v_c^{\nu(v)+1} - \varepsilon_0)$ and the functions $v \mapsto B(N, i, v)$ smoothly connect in the interval $(v_c^{\nu(v)} - \varepsilon_0, v_c^{\nu(v)} + \varepsilon_0)$ the values that the function $\sum_{i=0}^N A(N, i, \varepsilon_0) g_i \mu_i(M_v)$ takes in the intervals $(v_c^{\nu(v)-1} + \varepsilon_0, v_c^{\nu(v)} - \varepsilon_0)$ and $(v_c^{\nu(v)} + \varepsilon_0, v_c^{\nu(v)+1} - \varepsilon_0)$.*

Loosely speaking, $\sum_i A(N, i, \varepsilon_0) g_i \mu_i(M_v) + \sum_n B(N, i(n), v)$ is shaped as a "staircase" with "rounded corners".

Remark 3. *About the applicability domain of the Main Theorem proved in the present paper, note that V is required to be a finite range potential and a good Morse function. The former is a physical assumption, the latter a mathematical property of V . Finite range potentials are typical in condensed*

matter systems, where even Coulomb interactions are effective only at a finite distance because of the Debye shielding. The mathematical property of being a good Morse function is absolutely generic, in fact it requires that a potential function is bounded from below and that the Hessian of the potential is non-degenerate (i.e. its eigenvalues never vanish). Moreover, given a real-valued function f of class \mathcal{C}^2 defined on an arbitrary open subset X of \mathbb{R}^N , the mapping $x \mapsto f(x) - (a_1x_1 + \dots + a_Nx_N) : X \mapsto \mathbb{R}$ is nondegenerate for almost all $(a_1, \dots, a_N) \in \mathbb{R}^N$ (see Chapter 6 of Ref.[14]). This means that nondegeneracy is generic whereas degeneracy is exceptional. Continuous symmetries are the only physically relevant source of degeneracy, however this kind of degeneracy can be removed by adding a generic term $(a_1x_1 + \dots + a_Nx_N)$ to the potential with an arbitrarily small vector (a_1, \dots, a_N) . This removal of degeneracy is a rephrasing, within the framework of Morse theory, of a standard procedure undertaken in statistical mechanics to explicitly break a continuous symmetry, that is the addition of an external field whose limit to zero is taken after the limit $N \rightarrow \infty$.

Remark 4. Let us briefly compare meaning and strength of the Main Theorems of papers I and II, which we denote by MT-I and MT-II respectively. The proof of MT-I is preliminary to, and independent of, the proof of MT-II. On the other hand, it is important to note that the hypotheses under which MT-II applies are fulfilled by a very broad class of physically meaningful potentials, at variance with the case of MT-I. In fact, MT-I applies in presence of a potential energy density interval of finite length which is definitely (that is asymptotically in N) free of critical points, a rather restrictive assumption which we could hardly relax in the present demonstration scheme. On the contrary, we can approach the problem from a complementary point of view allowing the existence of critical points of the potential – as is assumed in the hypotheses of MT-II – with the only limitation to a linear growth with N of the number of critical values of V in a potential energy density interval of finite length. This assumption is suggested by what happens in lattice systems

where critical levels are separated by a finite minimum energy amount, as is the case of "spin flips" in the 1d-XY model [10], mean-field XY model [10], p-trig model [11], or of elementary configurational changes that, in lattices and fluids, correspond to the appearance of a new critical value of V at an energetic cost which is independent of N .

It would be impossible to prove MT-II without resorting to MT-I, though MT-II is much stronger than MT-I. Actually, the link established by MT-II in Eq.(18), between thermodynamic entropy and topology, tells that only through suitable variations with N of the v -patterns of $\sum_{i=0}^N A(N, i, \varepsilon_0) g_i \mu_i(M_v)$ some of the derivatives $|\partial^k S_N^{(-)} / \partial v^k|$ can cease to be uniformly bounded in N from above, thus giving rise to a phase transition. This is an important hint for a future rigorous investigation of the sufficiency conditions for both MT-I and MT-II. The combination of MT-I and MT-II provides clear evidence of the relevance of topology for the phenomenon of phase transitions.

4. FINAL REMARKS

Let us conclude with a few general comments. Earlier attempts at introducing topological concepts in statistical mechanics concentrated on *macroscopic* low-dimensional parameter spaces. Actually this happened after Thom's remark that the critical point shown by the van der Waals equation corresponds to the Riemann-Hugoniot catastrophe [22]. Hence some applications of the theory of singularities of differentiable maps to the study of phase transitions followed [23]. An elegant formulation of phase transitions as due to a topological change of some abstract manifold of macroscopic variables was obtained by using the Atiyah-Singer index theorem [24, 25] and deserves special attention because it applies to the $2d$ Ising model, whose phase transition is associated with a jump of the Atiyah index of some suitable vector bundle. This shows that also for discrete variables systems, like spin systems, topological concepts can be useful in the study of phase transitions, provided that the relevant manifolds are identified.

The Main Theorem, that we have proved above, makes a new kind of link between the study of phase transitions and differential topology. In fact, in the present work we deal with the high-dimensional *microscopic* configuration space of a physical system. The level sets of the microscopic interaction potential among the particles – or the regions of configuration space bounded by them – are the configuration space submanifolds that necessarily have to change their topology in correspondence with a phase transition point. The topology changes implied here are those described within the framework of Morse theory through *attachment of handles* [18].

We have explicitly investigated these topology changes in some particular models. The results so far obtained, already reported in the literature, are: *i)* numerical results on the lattice φ^4 model [7], *ii)* exact analytical results on the two models considered in Refs.[10, 11], *iii)* analytical results on the coupled rotators model. Although the models in item (*ii*) do not fulfil the condition of short-range interactions assumed by MT-I and MT-II in their present formulations, we already get a coherent scenario illustrating in practice how topology changes in presence and in absence of phase transitions. In fact, we observe that when the \bar{v} -pattern of the Euler characteristic $\chi(\bar{v})$ – which is the probe that we use to detect the variations of topology with \bar{v} – approaches a smooth curve as N increases, then phase transitions are absent; this is the case of $\chi(\Sigma_v)$ for the one-dimensional φ^4 model and of $\chi(M_v)$ for the one-dimensional chain of coupled rotators. At variance, sharp jumps or “cuspy” \bar{v} -patterns of $\chi(\Sigma_v)$ or of $\chi(M_v)$ are associated with phase transitions, in qualitative agreement with what is expected after MT-II; in Ref.[11] it is also shown that first and second order phase transitions are signaled by markedly different \bar{v} -patterns of $\chi(M_v)$.

Notice that in our approach the role of the potential V is twofold: it determines the relevant submanifolds of configuration space and it is a good Morse function on the same space. However, for example, in the case of entropy driven phase transitions occurring in hard sphere gases, the fact that the (sin-

gular) interaction potential cannot play any longer the role of Morse function does not mean that the connection between topology and phase transitions is lost, it rather means that other Morse functions are to be used. Just to give an idea of what a good Morse function could be in this case, let us think of the sum of all the pairwise euclidean distances between the hard spheres of a system: it is real valued, it has a minimum when the density is maximum, that is for close packing, meaning that this function is bounded below. The discussion of non-degeneracy is more involved and here would be out of place, let us simply remark that Morse functions are dense and degeneracy is easily removed when necessary.

The topology of configuration space submanifolds makes also a subtle link between dynamics and thermodynamics because it affects both of them, the former because it can be seen as the geodesic flow of a suitable Riemannian metric endowing configuration space [9], the latter because an analytic (though approximate) relation between thermodynamic entropy and Morse indexes of the critical points of configuration space submanifolds can be worked out [10].

Moreover, there are “exotic” kinds of transitional phenomena in statistical physics, like the glassy transition of amorphous systems to a supercooled liquid regime, or the folding transitions in polymers and proteins, which are qualitatively unified through the so-called *landscape paradigm* [26, 27] which is based on the idea that the relevant physics of these systems can be understood through the study of the properties of the potential energy hypersurfaces and, in particular, of their stationary points, usually called “saddles”. That this landscape paradigm naturally goes toward a link with Morse theory and topology has been hitherto overlooked. However, though at present our Main Theorem only applies to first and second order phase transitions, the topological approach seems to have the potentiality of unifying the mathematical description of very different kinds of phase transitions.

5. ACKNOWLEDGMENTS

The authors wish to thank L. Casetti and S. Schreiber for comments and suggestions. A particularly warm acknowledgment is addressed to G. Vezzosi for his continuous interest in our work and for many helpful discussions and suggestions.

- [1] L. Onsager, Phys. Rev. **65**, (1944) 117.
- [2] C.N. Yang and T.D. Lee, Phys. Rev. **87**, (1952) 404.
- [3] D. Ruelle, *Thermodynamic formalism*, Encyclopaedia of Mathematics and its Applications, (Addison-Wesley, New York, 1978).
- [4] H.O. Georgii, *Gibbs Measures and Phase Transitions*, (Walter de Gruyter, Berlin, 1988).
- [5] L. Caiani, L. Casetti, C. Clementi and M. Pettini, Phys. Rev. Lett. **79**, (1997) 4361.
- [6] R. Franzosi, L. Casetti, L. Spinelli and M. Pettini, Phys. Rev. **E60**, (1999) R5009.
- [7] R. Franzosi, M. Pettini, and L. Spinelli, Phys. Rev. Lett. **84**, (2000) 2774.
- [8] L. Casetti, E.G.D. Cohen and M. Pettini, Phys. Rev. Lett. **82**, (1999) 4160.
- [9] L. Casetti, M. Pettini, and E.G.D. Cohen, Phys. Rep. **337**, (2000) 237-341.
- [10] L. Casetti, M. Pettini, and E.G.D. Cohen, J. Stat. Phys. **111**, (2003) 1091.
- [11] L. Angelani, L. Casetti, M. Pettini, G. Ruocco, and F. Zamponi, Europhys. Lett. **62**, (2003) 775; Phys. Rev. **E71**, (2005) 036152.
- [12] R. Franzosi, M. Pettini, and L. Spinelli, *Topology and Phase Transitions I. Preliminary Results*, archived in: math-ph/0505057.
- [13] R. Franzosi, and M. Pettini, Phys. Rev. Lett. **92**, (2004) 060601.
- [14] M. Morse and S. S. Cairns, *Critical Point Theory in Global Analysis and Differential Topology*, (Academic Press, New York, 1969).
- [15] H. Federer, *Geometric Measure Theory*, (Springer, New York 1969), p. 249.
- [16] D. Ruelle, *Statistical Mechanics. Rigorous results*, (Benjamin, Reading, 1969).
- [17] R.S. Palais and C. Terng, *Critical Point Theory and Submanifold Geometry*, (Springer, New York 1988).
- [18] M.W. Hirsch, *Differential Topology*, (Springer, New York 1976).

- [19] J. Milnor, *Morse Theory*, (Princeton University Press, Princeton, 1973).
- [20] P. Laurence, *ZAMP* **40**, (1989) 258.
- [21] R. Bott and J. Mather, *Topics in Topology and Differential Geometry*, in *Battelle Rencontres*, Eds. C.M. De Witt and J.A. Wheeler, p.460. of the Main Theorem of paper I. The combination of the Main Theorem of paper I and Theorem 1 of the present paper
- [22] R. Thom, in *Statistical Mechanics*, Eds. S.A. Rice, K.F. Freed, and J.C. Light, (University of Chicago Press, 1972), p.93.
- [23] T. Poston and I. Stewart, *Catastrophe Theory and its Applications*, (Pitman Press, London, 1978), and references therein quoted.
- [24] M. Rasetti, *Topological concepts in the theory of phase transitions*, in *Differential Geometric Methods in Mathematical Physics*, Ed. H.D. Döbner (Springer-Verlag, New York, 1979).
- [25] M. Rasetti, *Structural Stability in Statistical Mechanics*, in *Springer Tracts in Math.*, Ed. W. Güttinger (Springer-Verlag, New York, 1979).
- [26] F.H. Stillinger, *Science* **267**, (1995) 1935.
- [27] S. Sastry, P.G. Debenedetti and F.H. Stillinger, *Nature* **393**, (1998) 554.