

Distribution of particles which produces a desired radiation pattern

A.G. Ramm

Mathematics Department, Kansas State University,
Manhattan, KS 66506-2602, USA
ramm@math.ksu.edu,
fax 785-532-0546, tel. 785-532-0580

Abstract

A method is given for calculation of a distribution of small particles, embedded in a medium, so that the resulting medium would have a desired radiation pattern for the plane wave scattering by this medium.

1 Introduction

Let $D_0 \subset \mathbb{R}^3$ be a bounded domain, k_0 be the wave number in D_0 and $k < k_0$ be the wave number in $D'_0 = \mathbb{R}^3 \setminus D_0$. We assume that D_0 is a homogeneous medium, so that k_0 is a constant. This assumption can be weakened: we may assume that $k_0 = k_0(x)$ is a known function. Let D_m be a particle, d_m be its diameter, $1 \leq m \leq M$, M is the number of small particles, $a = \max_{1 \leq m \leq M} \frac{d_m}{2}$ is an estimate for the radius of a small particle. We assume that $k_0 a \ll 1$, and then $ka \ll 1$, i.e., particles are small, that $d \gg a$, where $d = \min_{j \neq m} \text{dist}(D_m, D_j)$ and that the particles are acoustically soft, i.e., $u = 0$ on S_m , the boundary of D_m . Denote $\bigcup_{m=1}^M D_m := U$, $\mathbb{R}^3 \setminus U = V$, ∂U is the boundary of U , C_m is the electrical capacitance of a perfect conductor with the shape of D_m .

The Inverse Problem is:

Can one distribute many small particles in D_0 so that a plane wave $u_0 := e^{ik\alpha \cdot x}$, (where α is a given unit vector, $\alpha \in S^2$, S^2 is a unit sphere), scattered by D_0 , would produce a desired radiation pattern (scattering amplitude) $A(\alpha', \alpha)$?

The acoustic pressure u solves the problem (1)-(3):

$$(\nabla^2 + k^2 - q(x))u = 0 \text{ in } V, \quad q(x) = \begin{cases} k^2 - k_0^2 & \text{in } D_0 \\ 0 & \text{in } D'_0, \end{cases} \quad (1)$$

PACS 03.40.Kf MSC 35J05, 35J10, 70F10, 81U40, 35R30

key words: acoustic scattering, many-body problem, nanotechnology, inverse problems

$$u = 0 \text{ on } \partial U, \quad (2)$$

$$u = e^{ik\alpha \cdot x} + A(\alpha', \alpha) \frac{e^{ikr}}{r} + o\left(\frac{1}{r}\right), \quad r := |x| \rightarrow \infty, \quad \alpha' = \frac{x}{r}, \quad (3)$$

and the coefficient $A(\alpha', \alpha)$ is called the scattering amplitude or radiation pattern.

Let $G = G(x, y)$ solve the problem

$$[\nabla^2 + k^2 - q(x)]G = -\delta(x - y) \text{ in } \mathbb{R}^3, \quad r \left(\frac{\partial G}{\partial r} - ikG \right) = o(1) \quad r \rightarrow \infty. \quad (4)$$

This G exists, is unique, and solves the integral equation

$$G(x, y) = g(x, y) - (k^2 - k_0^2) \int_{D_0} g(x, z) G(z, y) dz, \quad g = g(x, y) := \frac{e^{ik|x-y|}}{4\pi|x-y|}. \quad (5)$$

For brevity, we drop the k -dependence in g , G and other functions below. One may consider $G(x, y)$ known since D_0 and k_0 are known.

Let us look for the solution to (1)–(3) of the form

$$u = U_0(x, \alpha) + \sum_{m=1}^M \int_{S_m} G(x, t) \sigma_m(t) dt, \quad (6)$$

where $\sigma_m(s)$ are unknown functions and $U_0(x, \alpha)$ is the scattering solution corresponding to the potential $q(x)$ in the absence of small bodies, i.e., in the whole space. For arbitrary σ_m the right-hand side of (6) solves equation (1) (because G solves (4)), satisfies the radiation condition (3), and

$$A(\alpha', \alpha) = A_q(\alpha', \alpha) + \frac{1}{4\pi} \sum_{m=1}^M \int_{S_m} U_0(s, -\alpha') \sigma_m(s) ds, \quad (7)$$

where the scattering solution $U_0(s, \alpha)$ can be defined by the formula ([1, p.232]):

$$G(x, s) = \frac{e^{ikr}}{4\pi r} U_0(s, \alpha) + o\left(\frac{1}{r}\right), \quad r := |x| \rightarrow \infty, \quad \alpha = -\frac{x}{r}. \quad (8)$$

Formula (8) was proved in [3, p.46], where it was shown that $U_0(x, \alpha)$ solves the problem

$$[\nabla^2 + k^2 - q(x)]U_0(x, \alpha) = 0 \text{ in } \mathbb{R}^3, \quad U_0 = e^{ik\alpha \cdot x} + A_q(\alpha', \alpha) \frac{e^{ikr}}{r} + o\left(\frac{1}{r}\right), \quad (9)$$

where $r := |x| \rightarrow \infty$ along the direction α' . The function U_0 can also be considered known, and G can be considered as Green's function for the Schrödinger equation (9).

The right-hand side of (6) solves problem (1)–(3) if and only if σ_m are such that boundary condition (2) is satisfied:

$$-U_0(s, \alpha) - \sum_{m \neq j} \int_{S_m} G(s, t) \sigma_m(t) dt = \int_{S_j} G(s, t) \sigma_j(t) dt, \quad s \in S_j, \quad 1 \leq j \leq M. \quad (10)$$

So far the smallness of the particles was not used. If $ka \ll 1$, then equation (10) can be simplified:

$$-U_0(s_j, \alpha) - \sum_{m \neq j} G(s_j, t_m) Q_m = \int_{S_j} g_0(s_j, t) \sigma_j dt, \quad 1 \leq j \leq M, \quad Q_m := \int_{S_m} \sigma_m dt, \quad (11)$$

where $s_j \in S_j$ is any point in S_j , and $g_0 = \frac{1}{4\pi|x-y|}$. In equation (11) we have used two approximations. The first one is

$$G(s, t) \approx g(s, t), \quad s, t \in S_j. \quad (12)$$

This approximation is justified by (5) when $k|x-y| \ll 1$, because the term $g(x, y)$ is the main term on the right-hand side of (5) as $x \rightarrow y$, and $|s_j - t| \ll 1$. The second one is $g \approx g_0$ if $k|s-t| \ll 1$, and the error of this approximation is $O(ka)$, so this approximation is also justified because $ka \ll 1$.

Equation (11) is equation for the charge distribution σ_j on the surface S_j of a perfect conductor charged to the potential $-U_0(s_j, \alpha) - \sum_{m \neq j} G(s_j, t_m) Q_m$. Therefore the total charge $Q_j := \int_{S_j} \sigma_j dt$ on the surface S_j can be calculated by the formula

$$Q_j = C_j \left(-U_0(s_j, \alpha) - \sum_{m \neq j} G(s_j, t_m) Q_m \right), \quad 1 \leq j \leq M, \quad (13)$$

where C_j is the electrical capacitance of the perfect conductor D_j . In [1, p.385] formulas for calculation of C_j with arbitrary desired accuracy are given. Equation (13) is a linear algebraic system for finding unknown Q_j , $1 \leq j \leq M$.

Consider the limiting case $M \rightarrow \infty$ of the distribution of particles in D_0 . Define $C(y)$ as follows:

$$\int_D C(y) dy = \lim_{M \rightarrow \infty} \sum_{D_m \subset D} C_m, \quad (14)$$

where $D \subset D_0$ is an arbitrary subdomain of D_0 . The above definition means that $C(y)$ is the limiting density of the capacitances of the small particles in D . Formula (6) shows that the self-consistent field u_e can be defined as

$$u_e(x, \alpha) = \begin{cases} U_0(x, \alpha) + \sum_{m=1}^M G(x, t_m) Q_m, & \min_{1 \leq m \leq M} |x - t_m| \gg a, \\ U_0(x, \alpha) + \sum_{m \neq j} G(x, t_m) Q_m, & |x - t_j| \sim a. \end{cases} \quad (15)$$

Therefore, defining u_e we neglect the influence of any fixed single small particle on the field. This is justified when $M \rightarrow \infty$. Using (13) and (15) one gets

$$u_e(x, \alpha) = U_0(x, \alpha) - \sum_m G(x, t_m) C_m u_e(t_m, \alpha),$$

and in the limit $M \rightarrow \infty$ one obtains the equation:

$$u_e(x, \alpha) = U_0(x, \alpha) - \int_{D_0} G(x, y) C(y) u_e(y, \alpha) dy, \quad (16)$$

where $C(y)$ is defined in (14). Equation (16) is equivalent to the Schrödinger scattering problem:

$$[\nabla^2 + k^2 - q(x) - C(x)]u_e = 0 \text{ in } \mathbb{R}^3, \quad (17)$$

where $q(x)$ is known, the function u_e has the following asymptotics

$$u_e = e^{ik\alpha \cdot x} + A(\alpha', \alpha) \frac{e^{ikr}}{r} + o\left(\frac{1}{r}\right), \quad r := |x| \rightarrow \infty, \quad \alpha' = \frac{x}{r}, \quad (18)$$

and $A(\alpha', \alpha)$ is the scattering amplitude at a fixed $k > 0$.

Therefore the Inverse Problem, stated above, is reduced to inverse scattering problem of finding the potential $q(x) + C(x)$ from the knowledge of the corresponding fixed-energy scattering amplitude $A(\alpha', \alpha)$.

This problem was solved by the author (see [1, Chapter 5] and references therein). In Section 2 we outline the author's algorithm for solving this inverse scattering problem.

If the potential $q(x) + C(x)$ is found and $q(x)$ is known, then $C(x)$ is found and one knows the density of the particle distribution in D which produces the desired radiation pattern $A(\alpha', \alpha)$. Assuming that the particles are identical, one has $C_m = \mathcal{C}$, where \mathcal{C} is the electrical capacitance of one particle, and $C(x) = N(x)\mathcal{C}$, where $N(x)$ is the density of particles, that is, the number of particles per unit volume around point x . See [2] for the theory of wave scattering by small bodies.

2 Solution to inverse scattering problem

We follow [1, p.264] and take $k = 1$ without loss of generality. Given $A(\alpha', \alpha)$ one finds $A_\ell(\alpha) := \int_{S^2} A(\alpha', \alpha) \overline{Y_\ell(\alpha')} d\alpha'$, where $Y_\ell := Y_{\ell m}$ are the normalized spherical harmonics, so that

$$A(\alpha', \alpha) = \sum_{\ell=0}^{\infty} A_\ell(\alpha) Y_\ell(\alpha'), \quad \sum_{\ell=0}^{\infty} := \sum_{\ell=0}^{\infty} \sum_{m=-\ell}^{\ell}. \quad (19)$$

If the data $A(\alpha', \alpha)$ are exact, then ([1, p.262])

$$\max_{\alpha \in S^2} |A_\ell(\alpha)| \leq O\left(\sqrt{\frac{b_0}{\ell}} \left(\frac{b_0 e}{2\ell}\right)^{\ell+1}\right), \quad |Y_\ell(\theta')| \leq \frac{1}{\sqrt{4\pi}} \frac{e^{\kappa r}}{|j_\ell(r)|} \quad \forall r > 0, \quad \theta \in \mathcal{M}, \quad (20)$$

where $b_0 > 0$ is the radius of the smallest ball containing the domain D_0 , $\mathcal{M} := \{\theta : \theta \in \mathbb{C}^3, \theta \cdot \theta = 1\}$, $\theta \cdot \omega := \sum_{j=1}^3 \theta_j \omega_j$, $\kappa = |Im\theta|$, $j_\ell(r)$ is the spherical Bessel function. Fix an arbitrary $\xi \in \mathbb{R}^3$. One can find (nonuniquely and explicitly) $\theta', \theta \in \mathcal{M}$, such that $\theta' - \theta = \xi$, $\theta \rightarrow \infty$. For example, if $\xi = te_3$, $t = |\xi| > 0$, (which can be assumed without loss of generality), then $\theta' = \frac{t}{2}e_3 + z_1e_1 + z_2e_2$, $\theta = -\frac{t}{2}e_3 + z_1e_1 + z_2e_2$, and the condition

$$\frac{t^2}{4} + z_1^2 + z_2^2 = 1, \quad z_1, z_2 \in \mathbb{C}, \quad (*)$$

implies $\theta, \theta' \in \mathcal{M}$. One may find many $z_1, z_2 \in \mathbb{C}$, such that $(*)$ holds and $|z_1| \rightarrow \infty$. For example, take $z_1 = re^{i\varphi}$, $z_2 = re^{-i\varphi}$, then $r^2 \sin(2\varphi) - r^2 \sin(2\varphi) = 0$, $r^2 [\cos(2\varphi + \cos(2\varphi))] = 1 - \frac{t^2}{4}$, so $r^2 \cos(2\varphi) = \frac{1}{2} - \frac{t^2}{8}$. One can take $r \geq |\frac{1}{2} - \frac{t^2}{8}|^{1/2}$ and find φ such that $\cos(2\varphi) = \left(\frac{1}{2} - \frac{t^2}{8}\right) \frac{1}{r^2}$.

In what follows we always assume

$$\theta' - \theta = \xi, \quad \theta', \theta \in \mathcal{M}, \quad |\theta| \rightarrow \infty. \quad (21)$$

Because of (20), the series

$$A(\theta', \alpha) = \sum_{\ell=0}^{\infty} A_\ell(\alpha) Y_\ell(\theta'), \quad \theta \in \mathcal{M}, \quad (22)$$

converges absolutely and uniformly on compact subsets of $S^2 \times \mathcal{M}$.

Fix positive numbers $b_0 < b_1 < b_2$ such that $D_0 \subset B_{b_0} := \{x : |x| \leq b_0\}$. Note that the scattering solution $u(x, \alpha)$ for the scattering potential $q(x) + C(x)$ can be written explicitly in the region $|x| > b_0$:

$$u(x, \alpha) = e^{i\alpha \cdot x} + \sum_{\ell=0}^{\infty} A_\ell(\alpha) Y_\ell(\alpha') h_\ell(r), \quad r := |x| > b_0, \quad (23)$$

where $k = 1$ and $h_\ell(r)$ is the spherical Hankel function which is normalized by the asymptotics

$$h_\ell(r) \sim \frac{e^{ir}}{r} \quad r \rightarrow \infty. \quad (24)$$

Let $\nu(\alpha) \in L^2(S^2)$. Consider the problem:

$$\mathcal{F}(\nu) = \min, \quad (25)$$

where

$$\mathcal{F}(\nu) := \int_{b_1 \leq |x| \leq b_2} \left| e^{-i\theta \cdot x} \int_{S^2} u(x, \alpha) \nu(\alpha) d\alpha - 1 \right|^2 dx. \quad (26)$$

The function $u(x, \alpha)$ in (26) is defined in (23), and the minimization in (25) is with respect to all $\nu \in L^2(S^2)$. One can prove ([1, p.265]) that

$$\inf \mathcal{F}(\nu) := d(\theta) \leq \frac{const}{|\theta|}, \quad \theta \in \mathcal{M}, \quad |\theta| \gg 1, \quad (27)$$

Let $\nu(\alpha, \theta)$ be an arbitrary approximate solution to (25) in the following sense:

$$\mathcal{F}(\nu(\alpha, \theta)) \leq 2d(\theta). \quad (28)$$

For this ν define

$$\hat{C} := -4\pi \int_{S^2} A(\theta', \alpha) \nu(\alpha, \theta) d\alpha, \quad (29)$$

where $A(\theta', \alpha)$ is defined in (22).

Let $\tilde{C}(\xi) = \int_{D_0} C(x) e^{-i\xi \cdot x} dx$, where $C(x) \in L^2(D_0)$ vanishes in D'_0 . Let (21) hold.

The following theorem is proved by the author in [1, p.266].

Theorem. *Under the above assumptions one has*

$$|\hat{C} - \tilde{C}(\xi)| = O\left(\frac{1}{|\theta|}\right). \quad (30)$$

Conclusion: An algorithm is given for embedding many small particles in a domain D_0 in such a way that the plane wave, scattered by such domain, would have a desired radiation pattern $A(\alpha', \alpha)$.

The algorithm consists of solving the inverse scattering problem, namely, finding the potential $q(x) + C(x)$, vanishing outside D_0 , from the fixed-energy ($k^2 = \text{const} > 0$) scattering amplitude $A(\alpha', \alpha)$. If this potential is found, then $C(x)$ is found, and the small acoustically soft identical particles should be distributed in D_0 with the density $N(x) = \mathcal{C}^{-1}C(x)$, where $\mathcal{C} > 0$ is the electrical capacitance of a perfect conductor which has the shape of the particle.

References

- [1] Ramm, A. G. , Inverse problems, Springer, New York, 2005.
- [2] Ramm, A. G. , Wave scattering by small bodies of arbitrary shapes, World Sci. Publishers, Singapore, 2005.
- [3] Ramm, A. G. , Scattering by obstacles, D.Reidel, Dordrecht, 1986.