

POINCARÉ POLYNOMIALS OF HYPERBOLIC LIE ALGEBRAS OF RANK THREE

M. Gungormez

Dept. Physics, Fac. Science, Istanbul Tech. Univ.

80626, Maslak, Istanbul, Turkey

e-mail: gungorm@itu.edu.tr

Abstract

In view of a previous work, we explicitly give the Poincaré polynomials of 19 Hyperbolic Lie algebras of rank 3. It is seen that every one of these polynomials is expressed as the ratio of Poincaré polynomial of B_3 Lie algebra and a polynomial of finite degree.

I. INTRODUCTION

In a previous work [1], we have shown that Poincare polinomials of Hyperbolic Lie algebras are expressed as the ratio of Poincare polynomial of a properly chosen finite Lie algebra and a polynomial of some finite degree. In view of the fact that there is only a finite number of this type of Lie algebras [2], we now present the case for rank 3 for which we have 19 different ones. It is interesting to note that B_3 Lie algebra plays a central role here.

For a brief summary of our method which is presented in [1] in detail, we note that Poincare polinomials[3] could be defined as being in relation with Weyl groups. Let us consider

$$P(t) \equiv \sum_{n=0}^{\infty} q(n) t^n$$

as Poincare polynomial of a Kac-Moody Lie algebra [4] G_r of rank r where t is some indeterminate. $q(n)$ here is the number of Weyl group elements which are defined as products of n simple Weyl reflections. Let $\mathcal{W}(G_r)$ be the Weyl group, ρ Weyl vector and α_i 's be the simple roots of G_r where $i = 1, 2 \dots r$. As is given in [1], our main objective to obtain $q(n)$ is then simply the statement that

$$\gamma \equiv \rho - \Sigma(\rho) \tag{I.1}$$

is unique for each and every element $\Sigma \in \mathcal{W}(G_r)$. Note that γ is by definition an element of positive root lattice. Beside the set Γ^0 which consists of zero root solely, the set

$$\Gamma^1 \equiv \{\alpha_1, \alpha_2, \dots, \alpha_r\}$$

contains the first $r+1$ solutions of (I.1) and Γ^1 corresponds by definition to simple Weyl reflections. All these are related with the fact that $q(0) = 1$ and also $q(1) = r$ as a result of

$$q(0) = \dim \Gamma^0$$

and

$$q(1) = \dim \Gamma^1$$


where $\dim \Gamma^n$ means the number of elements for any Γ^n which corresponds to Weyl group elements consisting of products of n simple Weyl reflections. One can readily see that all the sets Γ^n of solutions of (I.1) are to be constructed recursively from Γ^1 . Our results will be presented for 19 Hyperbolic Lie algebras of rank 3 in the next section.

II. RANK THREE CLASSIFICATION


Let \mathcal{H}^s be one of the 19 Hyperbolic Lie algebra of rank 3 as given in [2] where $s = 1, 2, \dots, 19$. Our main observation here is

$$P^s(t) \equiv \frac{P_{B_3}(t)}{Q_s(t)} \quad (II.1)$$


for Poincare polinomial $P^s(t)$ of \mathcal{H}^s where $P_{B_3}(t)$ is the Poincare polinomial of B_3 Lie algebra and $Q_s(t)$ is a polinomial of some finite degree of indeterminate t . B_3 here is defined by the diagram




and the polinomials $Q_s(t)$ will be given as in the following:




$$Q_1 = (1+t) (1+t^2) (1-t+t^2) (1-t^2-t^3)$$




$$Q_2 = (1+t) (1-t+t^2) (1+t+t^2) (1-t-t^3)$$




$$Q_3 = (1+t) (1+t^2) (1-t+t^2) (1-t-t^2) (1+t+t^2)$$




$$Q_4 = (1+t)^2 (1+t^2) (1-t+t^2) (1-t-t^2-t^3)$$




$$Q_5 = (1+t)^2 (1+t^2) (1-t+t^2) (1-t-t^2)$$




$$Q_6 = (1+t) (1-t+t^2) (1+t+t^2) (1-t-t^2-t^3)$$



$$Q_7 = (1+t)^2 (1-2t) (1+t^2) (1-t+t^2) (1+t+t^2)$$




$$Q_8 = (1+t) (1+t^2) (1-t-t^3-t^5)$$



$$Q_9 = (1+t) (1+t^2) (1-t-t^2-t^3-t^4-t^5)$$



$$Q_{10} = (1+t) (1+t^2) (1-t+t^2) (1-t^2-t^3)$$



$$Q_{11} = (1+t) (1-t+t^2) (1+t+t^2) (1-t-t^3)$$




$$Q_{12} = (1+t) (1-t+t^2) (1+t+t^2) (1-t-t^3)$$




$$Q_{13} = (1+t) (1+t^2) (1-t-t^3-t^5)$$




$$Q_{14} = (1+t) (1+t^2) (1-t-t^3-t^5)$$




$$Q_{15} = (1+t) (1+t^2) (1-t+t^2) (1-t-t^2) (1+t+t^2)$$




$$Q_{16} = (1+t)^2 (1+t^2) (1-t+t^2) (1-t-t^2-t^3)$$



$$Q_{17} = (1+t)(1-t+t^2)(1+t+t^2)(1-t-t^2-t^3)$$



$$Q_{18} = (1+t)(1+t^2)(1-t+t^2)(1-t-t^2)(1+t+t^2)$$



$$Q_{19} = (1+t)^2(1-2t)(1+t^2)(1-t+t^2)(1+t+t^2)$$

Let us finally note that one could ask, beside (I.1), if there is an alternative form, for instance

$$P^s(t) \equiv \frac{P_{A_3}(t)}{R_s(t)} \quad (II.2)$$

where $P(A_3)$ is Poincare polinomial of A_3 Lie algebra and $R_s(t)$'s are some other set of polinomials of some finite order. We also studied this problem and find that one can find solutions except $s=8,9,13,14$ hence B_3 is unique for this instance.

ACKNOWLEDGEMENT

The author would like thank H.R.Karadayi for his continous support during this work.

REFERENCES

- [1] H.R.Karadayi and M.Gungormez, On Poincare Polinomials of Hyperbolic Lie Algebras(submitted for publication)
- [2] C. Saclioglu, J.Phys. A:Math.Gen(1989) 3753-3769
- [3] J.E.Humphreys, Reflection Groups and Coxeter Groups,
Cambridge University Press, 1990
- [4] V.Kac, Infinite Dimensional Lie Algebras
Cambridge University Press, 1982