

Finite vs affine W -algebras

Alberto De Sole¹ *and* Victor G. Kac²

¹ *Department of Mathematics, Harvard University*

1 Oxford Street, Cambridge, MA 02138, USA

& *Istituto Nazionale di Alta Matematica*

Città Universitaria, 00185 Roma, Italy

E-mail: desole@math.harvard.edu

² *Department of Mathematics, MIT*

77 Massachusetts Avenue, Cambridge, MA 02139, USA

E-mail: kac@math.mit.edu

“I am an old man, and I know that a definition cannot be so complicated.”

I.M. Gelfand (after a talk on vertex algebras in his Rutgers seminar)

Abstract

In Section 1 we review various equivalent definitions of a vertex algebra V . The main novelty here is the definition in terms of an indefinite integral of the λ -bracket. In Section 2 we construct, in the most general framework, the Zhu algebra $Zhu_{\Gamma}V$, an associative algebra which “controls” Γ -twisted representations of the vertex algebra V with a given Hamiltonian operator H . An important special case of this construction is the H -twisted Zhu algebra Zhu_HV . In Section 3 we review the theory of non-linear Lie conformal algebras (respectively non-linear Lie algebras). Their universal enveloping vertex algebras (resp. universal enveloping algebras) form an important class of freely generated vertex algebras (resp. PBW generated associative algebras). We also introduce the H -twisted Zhu non-linear Lie algebra Zhu_HR of a non-linear Lie conformal algebra R and we show that its universal enveloping algebra is isomorphic to the H -twisted Zhu algebra of the universal enveloping vertex algebra of R . After a discussion of the necessary cohomological material in Section 4, we review in Section 5 the construction and basic properties of affine and finite W -algebras, obtained by the method of quantum Hamiltonian reduction. Those are some of the most intensively studied examples of freely generated vertex algebras and PBW generated associative algebras. Applying the machinery developed in Sections 3 and 4, we then show that the H -twisted Zhu algebra of an affine W -algebra is isomorphic to the finite W -algebra, attached to the same data. In Section 6 we define the Zhu algebra of a Poisson vertex algebra, and we discuss quasiclassical limits.

0 Introduction

0.1 What is a vertex algebra?

According to the general principles of quantum field theory, the observables are quantum fields, which are operator valued distributions on the space-time, satisfying certain axioms, called Wightman's axioms. In one sentence, a vertex algebra is a chiral part of a 2-dimensional conformally covariant quantum field theory.

The data of a vertex algebra consist of the space of states V (an arbitrary vector space), the vacuum vector $|0\rangle \in V$, the infinitesimal translation operator $T \in \text{End}V$ and a collection of $\text{End}V$ -valued quantum fields \mathcal{F} , subject to the axioms formulated below (which are “algebraic” consequences of Wightman's axioms).

A quantum field $a(z)$ is a formal $\text{End}V$ -valued distribution, that is an $\text{End}V$ -valued linear function on the space of Laurent polynomials $\mathbb{C}[z, z^{-1}]$ (the space of test functions), which satisfies the condition that $a(z)v$ is a Laurent series in z for every $v \in V$. We can write $a(z)$ in the form

$$a(z) = \sum_{n \in \mathbb{Z}} a_{(n)} z^{-n-1}, \quad \text{where } a_{(n)} \in \text{End}V, \quad (0.1)$$

so that $a_{(n)} = \text{Res}_z z^n a(z)$, where, as usual, Res_z stands for the coefficient of z^{-1} . Then the above condition means that $a_{(n)}v = 0$ for n sufficiently large.

The following are the four axioms of a vertex algebra $(V, |0\rangle, T, \mathcal{F} = \{a^j(z) = \sum_{n \in \mathbb{Z}} a_{(n)}^j z^{-n-1}\}_{j \in J})$:

(vacuum)	$T 0\rangle = 0,$
(translation covariance)	$[T, a^j(z)] = \partial_z a^j(z),$
(locality)	$(z-w)^{N_{jk}}[a^j(z), a^k(w)] = 0$ for some $N_{jk} \in \mathbb{Z}_+,$
(completeness)	vectors $a_{(n_1)}^{j_1} \dots a_{(n_s)}^{j_s} 0\rangle$ span $V.$

This definition allows one to easily construct examples. In Section 1 we give several more equivalent definitions of vertex algebras, which allow us to work with them. First, we show that, if we enlarge \mathcal{F} to the maximal collection $\bar{\mathcal{F}}$ of quantum fields for which the axioms still hold, then the map $\bar{\mathcal{F}} \rightarrow V$, defined by $a(z) \mapsto a_{(-1)}|0\rangle$, is bijective. We thus get the state-field correspondence, defined as the inverse map $V \rightarrow \bar{\mathcal{F}}$

$$V \ni a \mapsto Y(a, z) \in \bar{\mathcal{F}},$$

which leads to the second definition of a vertex algebra. We shall use the customary notation $Y(a, z)$ for the quantum field, corresponding to the state $a \in V$, and the customary name “vertex operators” for these quantum fields.

The state-field correspondence allows one to introduce bilinear products on V for each $n \in \mathbb{Z}$ by letting

$$a_{(n)}b = \text{Res}_z z^n Y(a, z)b.$$

This leads to the third, the original Borchers definition of a vertex algebra as a vector space V with the vacuum vector $|0\rangle$ and bilinear products $a_{(n)}b$ for each $n \in \mathbb{Z}$, satisfying a simple vacuum identity and a quite complicated cubic identity, called the Borchers identity [B]. This identity is somewhat similar to the Jacobi identity; the former is as important for the theory of vertex algebras as the latter is for the theory of Lie algebras.

A part of the structure of a vertex algebra is the λ -bracket, defined by

$$[a_\lambda b] = \text{Res}_z e^{\lambda z} Y(a, z)b = \sum_{n \in \mathbb{Z}_+} \frac{\lambda^n}{n!} a_{(n)}b \quad (a, b \in V),$$

which satisfies axioms similar to the axioms of a Lie algebra. This structure, extensively studied in the past few years, is called a Lie conformal algebra [K]. The only additional operation needed to get a vertex algebra structure is the -1^{st} product $a_{(-1)}b$, usually called the normally ordered product. The vacuum vector is the identity element for this product, which is neither commutative nor associative. However, the normally ordered product is quasicommutative and quasiassociative, meaning that the “quantum corrections” to the commutator and the associator are certain explicit expressions in terms of the λ -bracket. Finally, the λ -bracket and the normally ordered product are related by a “non-commutative Wick formula”, which is an analogue of the Leibniz rule, where the “quantum correction” is expressed in terms of a double λ -bracket. This is the fourth description of a vertex algebra, given in [BK], similar to the definition of a Poisson algebra. On the other hand, this definition is the most convenient one for the theory of quantum integrable systems.

Remarkably, the fourth definition can be interpreted by saying that a vertex algebra is a Lie conformal algebra whose λ -bracket can be “integrated”. This leads to a new, fifth definition of a vertex algebra, which is discussed in Section 1.9.

0.2 Non-linearities

Some of the most important examples of vertex algebras, such as affine, fermionic and Virasoro vertex algebras are the universal enveloping vertex algebras of certain Lie conformal algebras. However, the majority of affine W -algebras, which are central to the present paper, cannot be obtained in this way due to the “non-linearities” in the commutation relations. The first example of this kind is Zamolodchikov’s W_3 algebra [Za]. (The two generating quantum fields of the W_3 algebra correspond to the basic invariants of the Weyl group of sl_3 , hence the name “ W -algebra”.)

In order to take into account the above mentioned “non-linearities”, one needs the notion of a non-linear Lie conformal algebra, introduced in [DK]. This and the parallel, simpler notion of a non-linear Lie algebra, are discussed in Section 3. It follows from [DK, KW1] that all affine W -algebras can be obtained as universal enveloping vertex algebras of certain non-linear Lie conformal algebras. Likewise, all “finite” W -algebras are universal enveloping algebras of certain non-linear Lie algebras.

0.3 Hamiltonian operators

We shall consider most of the time H -graded vertex algebras. A vertex algebra V is called H -graded if one fixes a Hamiltonian operator H , i.e. a diagonalizable operator on V such that

$$[H, Y(a, z)] = z\partial_z Y(a, z) + Y(Ha, z). \quad (0.2)$$

The main source of Hamiltonians is an energy-momentum field, that is a vertex operator of the form $Y(\nu, z) = \sum_{n \in \mathbb{Z}} L_n z^{-n-2}$, with the following three properties:

- (i) the operators L_n form the Virasoro algebra

$$[L_m, L_n] = (m - n)L_{m+n} + \frac{c}{12}(m^3 - m)\delta_{m, -n}, \quad (0.3)$$

where $c \in \mathbb{C}$ is called the central charge,

- (ii) $L_{-1} = T$,

- (iii) L_0 is a diagonalizable operator.

Then $H = L_0$ is a Hamiltonian operator.

In connection to twisted representation theory of a vertex algebra V one also considers an abelian group A of automorphisms of the vertex algebra V , commuting with H . It turns out that A -twisted irreducible positive energy (with respect to H) modules over V are in a canonical one-to-one correspondence with irreducible modules over an associative algebra $Zhu_{H,A}V$, canonically associated to the triple (V, H, A) . In the case when the group A is trivial and all the eigenvalues of H are integers this construction was introduced by Zhu [Z]. Our construction generalizes those of [Z, KWa, DLM], and the proof of the main theorem is closer to [DLM]. Especially important for this paper is the case when the grading defined by the action of A is given by the eigenspaces of $e^{2\pi i H}$. In this case we denote $Zhu_{H,A}V$ by $Zhu_H V$, and we call it the H -twisted Zhu algebra.

In order to define and study the properties of the Zhu algebra, we introduce and study in Section 2 a certain deformation of the vertex operators of V . This deformed vertex algebra satisfies all the axioms of a vertex algebra, except that the translation covariance is replaced by a “gauged” translation covariance with the potential $\hbar H/(1 + \hbar z)$. We show that for this particular choice of the potential the deformed vertex operators satisfy a very simple deformation of the Borcherds identity. This makes the exposition of the theory of Zhu algebras very simple and transparent.

In Section 3.3 we introduce the (non-linear) Zhu Lie algebra $Zhu_H R$ of a non-linear Lie conformal algebra R and we show that the Zhu algebra $Zhu_H V(R)$ of the universal enveloping vertex algebra $V(R)$ is the universal enveloping algebra of $Zhu_H R$. As an application, we show in Section 3.4 that the $N = 1$ extension of the Sugawara construction, given in [KT], is a “chiralization” of

the cubic Dirac operator, defined in [AM, Ko1], which immediately implies all its properties.

In Section 4 we introduce the notion of an almost linear differential of a non-linear Lie algebra (resp. a non-linear Lie conformal algebra) and show that, under certain conditions, the cohomology of the universal enveloping algebra (resp. universal enveloping vertex algebra) equals the universal enveloping algebra (resp. universal enveloping vertex algebra) of the cohomology of the non-linear Lie algebra (resp. Lie conformal algebra).

0.4 What is a W -algebra?

In the remainder of the paper we apply the developed machinery to finite and affine W -algebras. They are universal enveloping algebras of certain non-linear Lie algebras and non-linear Lie conformal algebras respectively, obtained by the method of quantum Hamiltonian reduction.

Let us first explain the classical Hamiltonian reduction in a special situation that shall be used. Let \mathfrak{g} be a simple finite-dimensional Lie algebra with a non-degenerate symmetric invariant bilinear form $(\cdot|\cdot)$, and let $x \in \mathfrak{g}$ be an ad-diagonalizable element with eigenvalues in $\frac{1}{2}\mathbb{Z}$. We have the decomposition into a sum of subalgebras: $\mathfrak{g} = \mathfrak{g}_- \oplus \mathfrak{g}_0 \oplus \mathfrak{g}_+$, where \mathfrak{g}_+ (resp. \mathfrak{g}_-) is the sum of all eigenspaces of $\text{ad } x$ with positive (resp. negative) eigenvalues and \mathfrak{g}_0 is the 0th eigenspace. Let G_+ be the connected Lie group with the Lie algebra \mathfrak{g}_+ . Recall that the algebra $S(\mathfrak{g})$ of polynomial functions on \mathfrak{g}^* carries the classical Poisson algebra structure, defined by $\{a, b\} = [a, b]$ if $a, b \in \mathfrak{g}$, and extended to $S(\mathfrak{g})$ by the Leibniz rule. We shall identify \mathfrak{g}^* with \mathfrak{g} via the bilinear form $(\cdot|\cdot)$. The coadjoint action of G_+ on \mathfrak{g}_+^* can be conveniently thought of as the composition of its adjoint action on $\mathfrak{g}_- \subset \mathfrak{g}$ with the projection π_- of \mathfrak{g} onto \mathfrak{g}_- , the latter being canonically identified with \mathfrak{g}_+^* via the invariant form $(\cdot|\cdot)$. Let $f \in \mathfrak{g}_-$, let $\mathcal{O} = G_+ \cdot f$ be the coadjoint orbit of f in \mathfrak{g}_- , and consider the G_+ -invariant submanifold $\tilde{\mathcal{O}} = \pi_-^{-1}(\mathcal{O})$ of $\mathfrak{g} = \mathfrak{g}^*$. The ideal I consisting of functions from $S(\mathfrak{g})$ which vanish on $\tilde{\mathcal{O}}$ is not a Poisson ideal. Nevertheless, it is easy to check that the algebra $P = (S(\mathfrak{g})/I)^{G_+}$ of G_+ -invariants (which can be thought of as the algebra of functions on $\tilde{\mathcal{O}}/G_+$) carries a well-defined Poisson bracket induced from that on $S(\mathfrak{g})$. The Poisson algebra P is called the Hamiltonian reduction of the (Hamiltonian) action of G_+ on \mathfrak{g}^* . (A more general setup is a Hamiltonian action of a group N on a Poisson manifold X with the moment map $\mu : X \rightarrow \mathfrak{n}^*$. Picking an N -orbit \mathcal{O} in \mathfrak{n}^* , one obtains a Poisson manifold $\mu^{-1}(\mathcal{O})/N$.)

According to the general BRST idea, in order to quantize the Poisson algebra P , one should try to represent it as a homology of a complex (C, d) , which can be quantized.

This can be done, when the pair (x, f) is *good*, meaning that $f \in \mathfrak{g}_-$ is an eigenvector of $\text{ad } x$ with eigenvalue -1 , such that its centralizer in \mathfrak{g} lies in $\mathfrak{g}_- + \mathfrak{g}_0$. The most important examples of such pairs come from the \mathfrak{sl}_2 -triples $\{e, h = 2x, f\}$ in \mathfrak{g} .

One may replace \mathfrak{g}_+ by a slightly smaller subalgebra $\mathfrak{n}_\mathfrak{s} = \mathfrak{s} + \mathfrak{g}_{\geq 1}$, where \mathfrak{s} is an arbitrary subspace of $\mathfrak{g}_{1/2}$ and $\mathfrak{g}_{\geq 1}$ is the sum of all eigenspaces of $\text{ad } x$ with eigenvalues ≥ 1 . It is easy to see [GG] that this does not change the Poisson algebra P .

Let $N_\mathfrak{s}$ be the connected Lie group with the Lie algebra $\mathfrak{n}_\mathfrak{s}$. Then the orbit $N_\mathfrak{s} \cdot f$ of f in $\mathfrak{g}_+^* = \mathfrak{g}_-$ is $\mathcal{O}_\mathfrak{s} = f + [f, \mathfrak{s}]$, hence its preimage in $\mathfrak{g}^* = \mathfrak{g}$ is $\tilde{\mathcal{O}}_\mathfrak{s} = f + [f, \mathfrak{s}] + \mathfrak{g}_0 + \mathfrak{g}_+$. This is defined by the following equations in $\xi \in \mathfrak{g}$:

$$(\xi|a) = (f|a), \quad a \in \mathfrak{m}_\mathfrak{s},$$

where $\mathfrak{m}_\mathfrak{s} = ([f, \mathfrak{s}]^\perp \cap \mathfrak{g}_{1/2}) + \mathfrak{g}_{\geq 1}$, and \perp stands for the orthocomplement in \mathfrak{g} with respect to $(\cdot|\cdot)$. Hence the ideal $I_\mathfrak{s} \subset S(\mathfrak{g})$ of all functions from $S(\mathfrak{g})$ which vanish on $\tilde{\mathcal{O}}_\mathfrak{s}$ is generated by $\{a - (f|a) \mid a \in \mathfrak{m}_\mathfrak{s}\}$. It follows that the coordinate ring $\mathbb{C}[\tilde{\mathcal{O}}_\mathfrak{s}] = S(\mathfrak{g})/I_\mathfrak{s}$ can be represented as the 0th cohomology of the Koszul complex

$$(S(\mathfrak{g}) \otimes \wedge(\mathfrak{m}_\mathfrak{s}), d_K),$$

where the degrees of elements of \mathfrak{g} (resp. $\mathfrak{m}_\mathfrak{s}$) are equal to 0 (resp. -1), and

$$d_K(a) = 0, \quad d_K(\varphi_a) = a - (f|a).$$

Here $a \in \mathfrak{g}$ and φ_a denotes the element of $\mathfrak{m}_\mathfrak{s}$ corresponding to a , under the inclusion $\mathfrak{m}_\mathfrak{s} \subset \mathfrak{g}$.

The algebra P is the algebra of invariants of the group $N_\mathfrak{s}$ acting on the algebra $\mathbb{C}[\tilde{\mathcal{O}}_\mathfrak{s}]$, which is the 0th Lie algebra cohomology of the complex

$$(\mathbb{C}[\tilde{\mathcal{O}}_\mathfrak{s}] \otimes \wedge(\mathfrak{n}_\mathfrak{s}^*), d_L),$$

where d_L is the standard Lie algebra cohomology differential of the $\mathfrak{n}_\mathfrak{s}$ -module $\mathbb{C}[\tilde{\mathcal{O}}_\mathfrak{s}]$. Therefore the algebra P is isomorphic to the 0th cohomology of the complex [KS]

$$(S(\mathfrak{g}) \otimes \wedge(\mathfrak{m}_\mathfrak{s} \oplus \mathfrak{n}_\mathfrak{s}^*), d_K + d_L). \quad (0.4)$$

If \mathfrak{s} is chosen in such a way that $\mathfrak{m}_\mathfrak{s} = \mathfrak{n}_\mathfrak{s}$, it is clear how to quantize this complex. Just replace $S(\mathfrak{g})$ by the universal enveloping algebra $U(\mathfrak{g})$ and replace $\wedge(\mathfrak{n}_\mathfrak{s} \oplus \mathfrak{n}_\mathfrak{s}^*)$ by the Clifford algebra on the space $\mathfrak{n}_\mathfrak{s} \oplus \mathfrak{n}_\mathfrak{s}^*$ with the symmetric bilinear form defined by the pairing between $\mathfrak{n}_\mathfrak{s}$ and $\mathfrak{n}_\mathfrak{s}^*$. One can find such \mathfrak{s} by introducing the following non-degenerate skewsymmetric bilinear form on $\mathfrak{g}_{1/2}$:

$$\langle a, b \rangle = (f|[a, b]), \quad (0.5)$$

and taking \mathfrak{s} to be any maximal isotropic subspace of $\mathfrak{g}_{1/2}$.

Unfortunately in the Lie superalgebra case this may not work (since the bilinear form (0.5) is symmetric on the odd part of $\mathfrak{g}_{1/2}$). The way out is to choose \mathfrak{s} to be a co-isotropic subspace of $\mathfrak{g}_{1/2}$ (the favorite choice is $\mathfrak{s} = \mathfrak{g}_{1/2}$), so that $\mathfrak{s}^\perp \subset \mathfrak{s}$, hence $\mathfrak{m}_\mathfrak{s} \subset \mathfrak{n}_\mathfrak{s}$, and the space $\mathfrak{p} = \mathfrak{s}/\mathfrak{s}^\perp$ carries the induced non-degenerate bilinear form, also denoted by $\langle \cdot, \cdot \rangle$. One then replaces the complex (0.4) by a quasi-isomorphic complex

$$(S(\mathfrak{g}) \otimes \wedge(\mathfrak{n}_\mathfrak{s} \oplus \mathfrak{n}_\mathfrak{s}^*) \otimes S(\mathfrak{p}), d) \quad (0.6)$$

where the degree of the elements of \mathfrak{p} is 0, and d is a suitable modification of the differential $d_K + d_L$. This complex is again easy to quantize by replacing, as before, $S(\mathfrak{g})$ by $U(\mathfrak{g})$, $\wedge(\mathfrak{n}_{\mathfrak{s}} \oplus \mathfrak{n}_{\mathfrak{s}}^*)$ by the corresponding Clifford algebra and $S(\mathfrak{p})$ by the Weil algebra on the space \mathfrak{p} with the bilinear form $\langle \cdot, \cdot \rangle$. Finally, we write for the differential d the same formula as in the classical complex (0.6).

The cohomology of the quantization of the complex (0.6) is an associative algebra, called the finite W -algebra associated to the triple (\mathfrak{g}, x, f) , and denoted by $W^{\text{fin}}(\mathfrak{g}, x, f)$. (In fact, this is the 0th cohomology since the rest vanishes.)

One associates to the same triple a family of vertex algebras $W_k(\mathfrak{g}, x, f)$, called affine W -algebras, depending on a complex parameter k , in a similar fashion, replacing the associative algebras by the corresponding vertex algebras. Namely, $U(\mathfrak{g})$ is replaced by the universal affine vertex algebra $V^k(\mathfrak{g})$, and Clifford and Weil algebras are replaced by the vertex algebras of free superfermions $F(\Pi\mathfrak{n}_{\mathfrak{s}} \oplus \Pi\mathfrak{n}_{\mathfrak{s}}^*)$ and $F(\mathfrak{p})$, where Π stands for the change of parity. The differential d is replaced by $d_{(0)} = \text{Res}_z d(z)$, where the vertex operator $d(z)$ is obtained by writing $d = \text{ad } D$, where D is an element of the quantization of (0.6), and replacing all factors in D by the corresponding vertex operators and the associative products by normally ordered products.

The first main result of the theory of W -algebras states that the associative algebra $W^{\text{fin}}(\mathfrak{g}, x, f)$ (resp. vertex algebra $W_k(\mathfrak{g}, x, f)$) is freely generated in the sense of Poincaré–Birkhoff–Witt by elements attached to a basis of the centralizer of f in \mathfrak{g} (see [KW1]). Thus, according to [DK], all of these W -algebras are universal enveloping algebras (resp. enveloping vertex algebras) of non-linear Lie algebras (resp. non-linear Lie conformal algebras).

0.5 Some background remarks

After the work of Zamolodchikov [Za], a number of papers on affine W -algebras, appeared in physics literature, mainly as “extended conformal algebras”, i.e. vertex algebra extensions of the Virasoro vertex algebra. A review of the subject up to the early 1990s may be found in the collection of a large number of reprints on W -algebras [BS]. The most important work of this period is the work by Feigin and Frenkel [FF1, FF2], where the approach described in the previous subsection was introduced in the case of the principal nilpotent element f . Namely, they quantize the classical Hamiltonian reduction, introduced by Drinfeld and Sokolov [DS]. For example, if $\mathfrak{g} = \mathfrak{sl}_n$, the Drinfeld–Sokolov classical reduction produces the Gelfand–Dikii Poisson algebras, while their Feigin–Frenkel quantization produces the Virasoro vertex algebra for $n = 2$ and Zamolodchikov’s W_3 algebra for $n = 3$. Further important contributions were made in [BT, ST, FKW, FB], and many other works. The framework described in (0.4) was developed in [KRW, KW1, KW2] and applied to representation theory of superconformal algebras: it turned out that in the case of minimal nilpotent orbits of small simple Lie superalgebras these W -algebras are the $N=1,2,3,4$ and many other well-known superconformal algebras, which led to a further progress in their representation theory. Of course, in the quasi-classical limit of $W_k(\mathfrak{g}, x, f)$ one recovers the Drinfeld–Sokolov reduction and its

generalizations. For the most recent important development in representation theory of W -algebras see [A1, A2].

The finite W -algebras were much less studied. Some of the few references in physics literature that we know of are [BT1, DV, RS]. More recently there has been a revival of interest in them in connection to geometry and representation theory of simple finite-dimensional Lie algebras, see [M, P1, P2, GG, BrK]. The earliest work is [Ko], where it was shown that the algebra $W^{\text{fin}}(\mathfrak{g}, x, f)$ for the principal nilpotent f is canonically isomorphic to the center of $U(\mathfrak{g})$.

It is proved in the Appendix, written jointly with A. D'Andrea, C. De Concini and R. Heluani, that the quantum Hamiltonian reduction definition of finite W -algebras adopted in the paper and the definition via the Whittaker models which goes back to [Ko] (see [P1, GG]) are equivalent. This result was independently proved in [A2]. In fact in [A2] the result is stated only in the case of principal nilpotent f , cf. [KS], but the proof there can be adapted to the general case. The connection to the Whittaker models may be given as another reason for the name "W-algebras".

0.6 The main results of the paper

The definition and structure theory of finite and affine W -algebras are very similar, but, as far as we know, the precise connection between them has been unknown. The main new result of the present paper answers this question: the algebra $W^{\text{fin}}(\mathfrak{g}, x, f)$ is the H -twisted Zhu algebra $Zhu_H W_k(\mathfrak{g}, x, f)$, where $H = L_0$ comes from the energy-momentum field of the vertex algebra $W_k(\mathfrak{g}, x, f)$. In the case of the principal nilpotent f this result has been independently obtained by Arakawa [A2] (cf. [FKW]).

In conclusion of the paper we introduce the notion of a Zhu algebra of a Poisson vertex algebra and discuss its connection to the Zhu algebra of a vertex algebra via the quasiclassical limit. As a result, the quantum and classical affine W -algebras on the one hand, and the finite W -algebra and the Poisson algebra of the Slodowy slice on the other hand, are tied in one picture.

Though, for simplicity, we assumed above that \mathfrak{g} is a Lie algebra, we work in the more general situation of a simple finite-dimensional Lie superalgebra \mathfrak{g} with a non-degenerate supersymmetric invariant bilinear form, as in [KRW, KW1, KW2]. This is important due to applications to representation theory of superconformal algebras. Thus, throughout the paper we shall always talk about Lie superalgebras, Lie conformal superalgebras, associative superalgebras, etc., unless otherwise stated. However, we shall often drop the prefix "super".

We wish to thank Bojko Bakalov for many illuminating discussions. In particular it was his suggestion to introduce \hbar in Zhu's definition of the Zhu algebra, which led us to the definition of \hbar -deformed vertex operators. He also helped us with the proof of Theorem 2.30.

The main results of the paper were reported in June 2005 at the summer school in ESI, Vienna.

All vector spaces, algebras and their tensor products are considered over

the field of complex numbers \mathbb{C} , unless otherwise stated. We denote by \mathbb{R} , \mathbb{R}_+ , \mathbb{Z} , \mathbb{Z}_+ and \mathbb{N} the sets of real numbers, non negative real numbers, integers, non-negative integers and positive integers, respectively.

1 Five equivalent definitions of a vertex algebra

1.1 Vector superspaces

Let $V = V_{\bar{0}} \oplus V_{\bar{1}}$ be a vector superspace, i.e. a vector space, endowed with a decomposition in a direct sum of subspaces V_{α} , where $\alpha \in \mathbb{Z}/2\mathbb{Z} = \{\bar{0}, \bar{1}\}$. If $v \in V_{\alpha}$, one writes $p(v) = \alpha$ and calls it the *parity* of v . One has the associated vector superspace decomposition, which makes $\text{End}V$ an associative superalgebra (i.e. a $\mathbb{Z}/2\mathbb{Z}$ -graded associative algebra)

$$\text{End}V = (\text{End}V)_{\bar{0}} \oplus (\text{End}V)_{\bar{1}},$$

where $(\text{End}V)_{\alpha} = \{a \in \text{End}V \mid aV_{\beta} \subset V_{\alpha+\beta}\}$. One defines the bracket on an associative superalgebra A by

$$[a, b] = ab - p(a, b)ba, \quad (1.1)$$

which makes A a Lie superalgebra. Here and further we use the notation

$$p(a, b) = (-1)^{p(a)p(b)}, \quad s(a) = (-1)^{p(a)}, \quad (1.2)$$

and, when a formula involves $p(a)$, we assume that a is homogeneous with respect to the $\mathbb{Z}/2\mathbb{Z}$ -grading, and extend this formula to arbitrary a by linearity.

1.2 The first definition and the Extension Theorem

The first definition of a vertex algebra, given in the beginning of the introduction in the case of a vector space V , extends to the super case as follows. (We shall work in this generality.)

Definition 1.1. A *vertex algebra* is a quadruple $(V, |0\rangle, T, \mathcal{F})$, where $V = V_{\bar{0}} \oplus V_{\bar{1}}$ is a vector superspace, called the *space of states*, $|0\rangle$ is an element of $V_{\bar{0}}$, called the *vacuum vector*, T is an even endomorphism of V , called the *infinitesimal translation operator*, and \mathcal{F} is a collection of $\text{End}V$ -valued quantum fields, i.e.

$$\mathcal{F} = \{a^j(z) = \sum_{n \in \mathbb{Z}} a_{(n)}^j z^{-n-1}\}_{j \in J},$$

where for each $j \in J$ all $a_{(n)}^j \in \text{End}V$ have the same parity, and for every $b \in V$ we have $a_{(n)}^j b = 0$ for $n \gg 0$. The above data satisfy the following axioms:

- (i) vacuum: $T|0\rangle = 0$,
- (ii) translation covariance: $[T, a^j(z)] = \partial_z a^j(z)$,

(iii) locality: $(z-w)^{N_{jk}}[a^j(z), a^k(w)] = 0$ for some $N_{jk} \in \mathbb{Z}_+$,

(iv) completeness: $V = \text{span}_{\mathbb{C}} \{a_{(n_1)}^{j_1} \dots a_{(n_s)}^{j_s} | 0\rangle\}$.

The commutator in the locality axiom is defined by (1.1).

In order to pass to the second definition, we need the following simple

Lemma 1.2. *Let V be a vector superspace and let $\Omega \in V_0, T \in (\text{End}V)_0$, be such that $T\Omega = 0$. Let $a(z) = \sum_{n \in \mathbb{Z}} a_{(n)} z^{-n-1}$ be a translation covariant $\text{End}V$ -valued quantum field, i.e. $[T, a(z)] = \partial_z a(z)$. Then $a_{(n)}\Omega = 0$ for all $n \in \mathbb{Z}_+$.*

Proof. Recall that $a_{(n)}\Omega = 0$ for all $n \geq N$, where N is a sufficiently large non-negative integer. Choose minimal such N ; we have to show that $N = 0$. By assumption we have for all $n \in \mathbb{Z}$:

$$[T, a_{(n)}] = -na_{(n-1)}. \quad (1.3)$$

Applying both sides to Ω for $n = N$, we have: $Ta_{(N)}\Omega = -Na_{(N-1)}\Omega$. Hence, if $N > 0$, we obtain $a_{(N-1)}\Omega = 0$, a contradiction. \square

Let $(V, |0\rangle, T, \mathcal{F})$ be a vertex algebra. Let $\tilde{\mathcal{F}}$ be the set of all $\text{End}V$ -valued quantum fields $a(z)$, which are translation covariant, i.e. $[T, a(z)] = \partial_z a(z)$. By Lemma 1.2, we have

$$a(z)|0\rangle \in V[[z]], \quad \text{if } a(z) \in \tilde{\mathcal{F}}, \quad (1.4)$$

which allows us to define the linear map

$$s : \tilde{\mathcal{F}} \rightarrow V, \quad s(a(z)) = a(z)|0\rangle|_{z=0}. \quad (1.5)$$

We next define the n^{th} product of $\text{End}V$ -valued quantum fields, for $n \in \mathbb{Z}$:

$$a(w)_{(n)}b(w) = \text{Res}_z (i_{z,w}(z-w)^n a(z)b(w) - p(a,b)i_{w,z}(z-w)^n b(w)a(z)), \quad (1.6)$$

where $i_{z,w}$ denotes the power series expansion in the domain $|z| > |w|$.

Lemma 1.3. (a) *The n^{th} product of two $\text{End}V$ -valued quantum fields is an $\text{End}V$ -valued quantum field.*

(b) *If $a(z), b(z)$ are translation covariant $\text{End}V$ -valued quantum fields, so are $\partial_z a(z)$ and $a(z)_{(n)}b(z)$ for every $n \in \mathbb{Z}$.*

(c) *If $(a(z), b(z))$ is a local pair of $\text{End}V$ -valued quantum fields, so is $(\partial_z a(z), b(z))$.*

(d) *If $(a(z), b(z), c(z))$ is a collection of pairwise local $\text{End}V$ -valued quantum fields, then $(a(z)_{(n)}b(z), c(z))$ is a local pair for every $n \in \mathbb{Z}$.*

Proof. (a) is straightforward. (b) follows immediately by the definition (1.6) of n^{th} product of fields. (c) is easy and (d) is known as Dong's Lemma. Its proof can be found in [K, Lemma 3.2]. \square

Lemma 1.4. (a) If $a(z), b(z) \in \tilde{\mathcal{F}}$, $a(z) = \sum_{n \in \mathbb{Z}} a_{(n)} z^{-n-1}$, and $s(a(z)) = a, s(b(z)) = b$, then $s(a(z)_{(n)} b(z)) = a_{(n)} b$ and $s(\partial_z a(z)) = T a$.

(b) If $a(z) \in \tilde{\mathcal{F}}$, then $a(z)|0\rangle = e^{Tz} a$, where $a = s(a(z))$.

Proof. To prove the first part of (a), apply both sides of (1.6) to $|0\rangle$. Since $a(z)|0\rangle \in V[[z]]$ by (1.4) and $i_{w,z}(z-w)^n$ involves only non-negative powers of z as well, the second term of the RHS of (1.6) applied to $|0\rangle$ vanishes. Hence $(a(w)_{(n)} b(w))|0\rangle|_{w=0} = \text{Res}_z z^n a(z) b = a_{(n)} b$. The second part of (a) is immediate by applying the translation covariance to $|0\rangle$ and putting $z = 0$. Finally (b) is obtained by looking at the Taylor expansion of $a(z)|0\rangle$, and using the identities $\partial_z^n a(z)|0\rangle|_{z=0} = T^n a$, given by (a). \square

Denote by \mathcal{F}' the minimal subspace of the space of all $\text{End} V$ -valued quantum fields, containing I_V and \mathcal{F} , which is ∂_z -invariant and closed under all n^{th} products. We can construct \mathcal{F}' as follows. Define by induction an increasing sequence \mathcal{F}_k , $k \geq 0$, of subspaces of the space of $\text{End} V$ -valued quantum fields, by letting $\mathcal{F}_0 = \text{span}_{\mathbb{C}}\{I_V, \mathcal{F}\}$, and $\mathcal{F}_{k+1} = \mathcal{F}_k + (\sum_{i \geq 0} \partial_z^i \mathcal{F}_k) + (\sum_{n \in \mathbb{Z}} \mathcal{F}_{k(n)} \mathcal{F}_k)$. Then $\mathcal{F}' = \bigcup_{k \geq 0} \mathcal{F}_k$. We also let $\bar{\mathcal{F}}$ be the collection of all $\text{End} V$ -valued quantum fields which are translation covariant and local to all elements of \mathcal{F} . By Lemma 1.3 we have the following inclusions:

$$\mathcal{F} \subset \mathcal{F}' \subset \bar{\mathcal{F}} \subset \tilde{\mathcal{F}}. \quad (1.7)$$

Theorem 1.5. Let $(V, |0\rangle, T, \mathcal{F})$ be a vertex algebra. Then

- (i) $(V, |0\rangle, T, \bar{\mathcal{F}})$ is also a vertex algebra.
- (ii) The map $s : \bar{\mathcal{F}} \rightarrow V$, defined by (1.5), is bijective.
- (iii) $\mathcal{F}' = \bar{\mathcal{F}}$.

Proof. Lemma 1.3 guarantees that \mathcal{F}' is a collection of translation covariant pairwise local fields, and that all pairs of quantum fields $(a(z), b(z))$, with $a(z) \in \mathcal{F}'$ and $b(z) \in \bar{\mathcal{F}}$, are local. In particular $(V, |0\rangle, T, \mathcal{F}')$ is a vertex algebra. Hence, by Lemma 1.4(a) and the completeness axiom, the map $s : \mathcal{F}' \rightarrow V$ is surjective. Let now $a(z) \in \bar{\mathcal{F}}$ be such that $s(a(z)) = 0$. In particular $a(z)|0\rangle = 0$ by Lemma 1.4(b). Since $s : \mathcal{F}' \rightarrow V$ is surjective, for each $b \in V$ there exists $b(z) \in \mathcal{F}'$ such that $s(b(z)) = b$. We have, by locality of the pair $(a(z), b(z))$,

$$(z-w)^N a(z) b(w)|0\rangle = p(a, b)(z-w)^N b(w) a(z)|0\rangle = 0.$$

Letting $w = 0$ and canceling z^N , we get $a(z)b = 0$ for every $b \in V$, hence $a(z) = 0$. Thus, the map $s : \bar{\mathcal{F}} \rightarrow V$ is injective. Since $s : \mathcal{F}' \rightarrow V$ is surjective and, by (1.7), $\mathcal{F}' \subset \bar{\mathcal{F}}$, we immediately get that $\mathcal{F}' = \bar{\mathcal{F}}$ and that $s : \bar{\mathcal{F}} \rightarrow V$ is bijective. \square

Remark 1.6. Theorem 1.5 implies the Existence Theorem 4.5 from [K] with the assumption (iii) and part of assumption (ii) removed. Lemma 1.2 was pointed out by Nikolov and Todorov (this property is an axiom in [K]), see also Remark 4.4b in [K].

Lemma 1.4(a), and Theorem 1.5 and its proof imply the following

Corollary 1.7. *Let $(V, |0\rangle, T, \mathcal{F})$ be a vertex algebra. Then for each $a \in V$ there exists a unique $\text{End}V$ -valued quantum field $Y(a, z)$ which is translation covariant and local with all quantum fields from \mathcal{F} , and such that $Y(a, z)|0\rangle|_{z=0} = a$. Furthermore, one has:*

$$Y(|0\rangle, z) = I_V, \quad (1.8)$$

$$Y(Ta, z) = \partial_z Y(a, z), \quad (1.9)$$

$$Y(a_{(n)}b, z) = Y(a, z)_{(n)}Y(b, z). \quad (1.10)$$

1.3 The second definition of a vertex algebra

We arrive at the second, equivalent definition of a vertex algebra, which was given in [K].

Definition 1.8. A *vertex algebra* is a triple $(V, |0\rangle, Y)$, where $V = V_0 \oplus V_1$ is a vector superspace (the space of states), $|0\rangle \in V_0$ is a vector (vacuum vector), and Y is a parity preserving linear map from V to the space of $\text{End}V$ -valued quantum fields: $Y(a) = Y(a, z) = \sum_{n \in \mathbb{Z}} a_{(n)} z^{-n-1}$, called the *state-field correspondence*. The infinitesimal translation operator T is encoded in the state-field correspondence via the formula

$$Ta = a_{(-2)}|0\rangle. \quad (1.11)$$

The axioms are as follows,

- (i) vacuum: $T|0\rangle = 0$, $Y(a, z)|0\rangle|_{z=0} = a$,
- (ii) translation covariance: $[T, Y(a, z)] = \partial_z Y(a, z)$,
- (iii) locality: $(z - w)^N [Y(a, z), Y(b, w)] = 0$ for some $N \in \mathbb{Z}_+$.

Notice that the definition (1.11) of the infinitesimal translation operator follows from (1.9) by applying both sides to $|0\rangle$ and putting $z = 0$. We thus have proved already that the first definition implies the second. The converse is obvious, just define T by (1.11) and let $\mathcal{F} = \{Y(a, z)\}_{a \in V}$. The completeness axiom is automatic since, by the vacuum axiom,

$$a = a_{(-1)}|0\rangle \quad \text{for } a \in V. \quad (1.12)$$

Note also that we automatically have (1.8), (1.9) and (1.10) for all $a, b \in V$.

1.4 Borcherds identity and the third definition of a vertex algebra

Formula (1.10), called the n^{th} product identity, leads to the most important formula of the theory of vertex algebras, called the Borcherds identity. In order to derive it from (1.10), recall the simple decomposition formula, which holds

for any local formal distribution $a(z, w) = \sum_{m, n \in \mathbb{Z}} a_{(m, n)} z^{-m-1} w^{-n-1}$ in z and w , where by locality one means that $(z - w)^N a(z, w) = 0$ for some $N \in \mathbb{Z}_+$ (see [K, Corollary 2.2]):

$$a(z, w) = \sum_{j \in \mathbb{Z}_+} c^j(w) \partial_w^j \delta(z - w) / j!, \quad (1.13)$$

where the sum is finite (actually $j < N$), the quantum fields $c^j(w)$ are given by

$$c^j(w) = \text{Res}_z (z - w)^j a(z, w), \quad (1.14)$$

and $\delta(z - w)$ is the formal delta function, defined by

$$\delta(z - w) = i_{z, w} \frac{1}{z - w} - i_{w, z} \frac{1}{z - w} = z^{-1} \sum_{n \in \mathbb{Z}} \left(\frac{w}{z} \right)^n. \quad (1.15)$$

The Borchers identity reads:

$$\begin{aligned} Y(a, z) Y(b, w) i_{z, w} (z - w)^n - p(a, b) Y(b, w) Y(a, z) i_{w, z} (z - w)^n \\ = \sum_{j \in \mathbb{Z}_+} Y(a_{(n+j)} b, w) \partial_w^j \delta(z - w) / j!, \end{aligned} \quad (1.16)$$

for all $n \in \mathbb{Z}$ and $a, b \in V$. Note that the sum on the right is finite since $a_{(N)} b = 0$ for $N \gg 0$, because $Y(a, z)$ is a quantum field.

In order to prove (1.16), note that the LHS is local, since, multiplied by $(z - w)^N$, it is equal to

$$Y(a, z) Y(b, w) i_{z, w} (z - w)^{n+N} - p(a, b) Y(b, w) Y(a, z) i_{w, z} (z - w)^{n+N}, \quad (1.17)$$

which is $[Y(a, z), Y(b, w)](z - w)^{n+N}$ if $n + N \geq 0$. We thus apply the decomposition formula (1.13) to the LHS. By (1.14), (1.15) and (1.17), we have $c^j(w) = Y(a, w)_{(n+j)} Y(b, w)$. We then apply (1.10) to obtain (1.16).

A special case of (1.16), for $n = 0$, is the commutator formula ($a, b \in V$):

$$[Y(a, z), Y(b, w)] = \sum_{j \in \mathbb{Z}_+} Y(a_{(j)} b, w) \partial_w^j \delta(z - w) / j!. \quad (1.18)$$

We thus arrive at the third definition of a vertex algebra:

Definition 1.9. A *vertex algebra* is a triple $(V, |0\rangle, Y)$, where V and $|0\rangle$ are as in Section 1.3 and Y is an injective state-field correspondence $a \mapsto Y(a, z)$, such that (1.8) and the Borchers identity (1.16) hold.

In order to prove that this definition implies that first one, and for further use, we shall digress now to the important notion of the normally ordered product of $\text{End} V$ -valued quantum fields $a(z) = \sum_{n \in \mathbb{Z}} a_{(n)} z^{-n-1}$ and $b(z)$. Let

$$a(z)_+ = \sum_{n < 0} a_{(n)} z^{-n-1}, \quad a(z)_- = \sum_{n \geq 0} a_{(n)} z^{-n-1},$$

and define the *normally ordered product* by

$$: a(z)b(z) := a(z)_+b(z) + p(a, b)b(z)a(z)_-. \quad (1.19)$$

Using the formal Cauchy formulas

$$\text{Res}_z a(z) i_{z,w} \frac{1}{z-w} = a(w)_+, \quad \text{Res}_z a(z) i_{w,z} \frac{1}{z-w} = -a(w)_-,$$

and the observation that $\partial_z(a(z)_\pm) = (\partial_z a(z))_\pm$, we deduce from (1.6) for $n \in \mathbb{Z}_+$:

$$a(w)_{(-n-1)}b(w) =: (\partial_w^n a(w))b(w) : / n!. \quad (1.20)$$

Returning to the third definition, note that commutator formula (1.18) implies locality of the pair $Y(a, z)$, $Y(b, z)$, since $a_{(j)}b = 0$ for $j \gg 0$ and $(z-w)^{j+1}\partial_w^j \delta(z-w) = 0$. Next, notice that the n^{th} product formula (1.10) follows from the Borchers identity by taking Res_z of both sides. Letting $b = |0\rangle$ in (1.10) and using (1.8) and (1.6), (1.20), we obtain:

$$Y(a_{(n)}|0\rangle, z) = 0, \quad \text{for } n \in \mathbb{Z}_+, \quad (1.21)$$

$$Y(a_{(-n-1)}|0\rangle, z) = \partial_z^n Y(a, z) / n!, \quad \text{for } n \in \mathbb{Z}_+. \quad (1.22)$$

It follows by injectivity of the state-field correspondence that

$$a_{(n)}|0\rangle = \delta_{n,-1}a, \quad \text{for } n \geq -1. \quad (1.23)$$

The latter formula for $n = -1$ implies the completeness axiom of the first definition.

Next, define $T \in \text{End}V$ by (1.11). Then $T|0\rangle = 0$ due to (1.8), which is the vacuum axiom of the first definition. Letting $n = 1$ in (1.22), we obtain (1.9), which implies that $Y(T^n a, z) = \partial_z^n Y(a, z)$. Applying both sides of the latter formula to $|0\rangle$ and letting $z = 0$ (which is possible due to (1.13)), we obtain

$$Y(a, z)|0\rangle = e^{zT}a. \quad (1.24)$$

Applying both sides of the n^{th} product identity to $|0\rangle$ and using (1.24), we obtain:

$$e^{zT}(a_{(n)}b) = (Y(a, z)_{(n)}Y(b, z))|0\rangle.$$

Differentiating both sides by z and letting $z = 0$, we obtain, using (1.9) and (1.23):

$$T(a_{(n)}b) = (Ta)_{(n)}b + a_{(n)}Tb. \quad (1.25)$$

Finally, the equivalent form (1.3) of the translation covariance axiom follows from (1.25) and the formula

$$(Ta)_{(n)} = -na_{(n-1)}, \quad (1.26)$$

which is an equivalent form of (1.9). This completes the proof of the equivalence of the third and the first definitions.

The vertex operators $Y(a, z) = \sum_{n \in \mathbb{Z}} a_{(n)} z^{-n-1}$ allow one to define n^{th} bilinear products on V for each $n \in \mathbb{Z}$ by the obvious formula $a_{(n)}b =$ the operator $a_{(n)}$ applied to vector b . Recall that we have (see (1.8) and (1.23)):

$$|0\rangle_{(n)}a = \delta_{n,-1}a \text{ for } n \in \mathbb{Z}, \quad a_{(n)}|0\rangle = \delta_{n,-1}a \text{ for } n \geq -1. \quad (1.27)$$

Applying both sides of identity (1.16) to $c \in V$ and comparing coefficients of $z^{-m-1}w^{-k-1}$, we obtain the original form of the Borchers identity:

$$\begin{aligned} \sum_{j \in \mathbb{Z}_+} (-1)^j \binom{n}{j} \left(a_{(m+n-j)}(b_{(k+j)}c) - (-1)^n p(a, b) b_{(n+k-j)}(a_{(m+j)}c) \right) \\ = \sum_{j \in \mathbb{Z}_+} \binom{m}{j} (a_{(n+j)}b)_{(m+k-j)}c \end{aligned} \quad (1.28)$$

for all $a, b, c \in V$, $m, n, k \in \mathbb{Z}$.

It is easy to see that the third definition of the vertex algebra is essentially the same as the original Borchers definition as a vector (super)space V with a vacuum vector $|0\rangle$ and \mathbb{C} -bilinear products $a_{(n)}b$ for each $n \in \mathbb{Z}$, such that (1.27) and (1.28) hold.

From this point of view all the standard algebra notions of a homomorphism and isomorphism, subalgebras and ideals, carry over to vertex algebras in the obvious way. Note that the right ideals are automatically T -invariant due to (1.20). Also, right ideals and T -invariant left ideals are automatically two-sided ideals, due to (1.11) and the simple skewsymmetry relation (see [FHL], [K, (4.2.1)]):

$$Y(a, z)b = p(a, b)e^{zT}Y(b, -z)a. \quad (1.29)$$

1.5 Lie conformal algebras and the fourth definition of a vertex algebra

For the fourth definition of a vertex algebra we need the definition of a Lie conformal algebra [K, D'K].

Definition 1.10. A *Lie conformal superalgebra* is a $\mathbb{Z}/2\mathbb{Z}$ -graded $\mathbb{C}[T]$ -module $R = R_{\bar{0}} \oplus R_{\bar{1}}$, endowed with a parity preserving \mathbb{C} -bilinear λ -bracket $R \otimes R \rightarrow \mathbb{C}[\lambda] \otimes R$, denoted by $[a_\lambda b]$, such that the following axioms hold:

$$\begin{aligned} (\text{sesquilinearity}) \quad & [Ta_\lambda b] = -\lambda[a_\lambda b], \quad T[a_\lambda b] = [Ta_\lambda b] + [a_\lambda Tb], \\ (\text{skewsymmetry}) \quad & [b_\lambda a] = -p(a, b)[a_{-\lambda-T}b], \\ (\text{Jacobi identity}) \quad & [a_\lambda[b_\mu c]] - p(a, b)[b_\lambda[a_\mu c]] = [[a_\lambda b]_{\lambda+\mu}c]. \end{aligned}$$

One writes $[a_\lambda b] = \sum_{n \in \mathbb{Z}_+} \frac{\lambda^n}{n!} (a_{(n)}b)$, where the sum is finite, and the bilinear products $a_{(n)}b$ are called the n^{th} products of R . The expression $[a_{-\lambda-T}b]$ in the skewsymmetry relation is interpreted as $\sum_{j \in \mathbb{Z}_+} \frac{(-\lambda-T)^n}{n!} (a_{(n)}b)$. Of course, λ

and μ are (even) indeterminates. Thus, the skewsymmetry relation is equivalent to the following identities ($n \in \mathbb{Z}_+$):

$$b_{(n)}a = p(a, b) \sum_{j \in \mathbb{Z}_+} (-1)^{n+j+1} T^j(a_{(n+j)}b)/j! . \quad (1.30)$$

Similarly, the Jacobi identity is equivalent to the following identities ($m, n \in \mathbb{Z}_+$):

$$a_{(m)}(b_{(n)}c) - p(a, b)b_{(n)}(a_{(m)}c) = \sum_{i \in \mathbb{Z}_+} \binom{m}{i} (a_{(i)}b)_{(m+n-i)}c . \quad (1.31)$$

Define the *formal Fourier transform* by

$$F_z^\lambda a(z) = \text{Res}_z e^{\lambda z} a(z) .$$

It is a linear map from the space of U -valued formal distributions to $U[[\lambda]]$, and it has the following properties, which are immediate to check:

$$F_z^\lambda \partial_z a(z) = -\lambda F_z^\lambda a(z) \quad (1.32)$$

$$F_z^\lambda (e^{zT} a(z)) = F_z^{\lambda+T} a(z) \text{ if } a(z) \in U((z)) , \quad (1.33)$$

$$F_z^\lambda a(-z) = -F_z^{-\lambda} a(z) , \quad (1.34)$$

$$F_z^\lambda \partial_w^n \delta(z-w) = e^{\lambda w} \lambda^n . \quad (1.35)$$

Note that (1.30) for all $n \in \mathbb{Z}$ is an equivalent form of (1.29), and (1.31) for all $m, n \in \mathbb{Z}$ is an equivalent form of (1.18). This is not surprising in view of the following proposition [K].

Proposition 1.11. *Define a λ -bracket on a vertex algebra V by*

$$[a_\lambda b] = F_z^\lambda Y(a, z)b .$$

Then the $\mathbb{C}[T]$ -module V with this λ -bracket is a Lie conformal algebra.

Proof. The sesquilinearity relations follow from (1.26) and (1.25) for $n \in \mathbb{Z}_+$. Applying F_z^λ to both sides of (1.29) and using (1.33) and (1.34), we get the skewsymmetry relation. In order to prove the Jacobi identity, apply F_z^λ to the commutator formula (1.18), applied to c , and use (1.35) to obtain:

$$[a_\lambda Y(b, w)c] = p(a, b)Y(b, w)[a_\lambda c] + e^{\lambda w}Y([a_\lambda b], w)c . \quad (1.36)$$

Applying F_w^μ to both sides of this formula, we get the Jacobi identity. \square

Next, define the *normally ordered product* on V by

$$:ab := a_{(-1)}b \quad (= \text{Res}_z z^{-1} Y(a, z)b)$$

and recall that $a_{(0)} = \text{Res}_z Y(a, z)$. The following proposition is an immediate corollary of identity (1.36).

Proposition 1.12. (a) The operator $a_{(0)}$ is a derivation of all n^{th} products of V , and it commutes with T . Consequently, $\text{Ker } a_{(0)}$ is a subalgebra of the vertex algebra V , and its intersection with $\text{Im } a_{(0)}$ is an ideal of $\text{Ker } a_{(0)}$.

(b) One has the following identity for all $a, b, c \in V$:

$$[a_\lambda : bc :] = [a_\lambda b]c : + p(a, b) : b[a_\lambda c] : + \int_0^\lambda [[a_\lambda b]_\mu c] d\mu. \quad (1.37)$$

Proof. The first (resp. second) part of (a) follows from (1.36) (resp. the second sesquilinearity relation) by letting $\lambda = 0$. (b) is obtained from (1.36) by comparing coefficients of w^0 . \square

Formula (1.37) is called in [K] the *non-commutative Wick formula*. It is extremely useful in calculations of the commutators of normally ordered products of quantum fields.

Skewsymmetry of the λ -bracket and (1.37) imply the right non-commutative Wick formula [BK]:

$$[ab :_\lambda c] = (e^{T\partial_\lambda} a)[b_\lambda c] : + p(a, b) : (e^{T\partial_\lambda} b)[a_\lambda c] : + p(a, b) \int_0^\lambda [b_\mu [a_{\lambda-\mu} c]] d\mu. \quad (1.38)$$

The normally ordered product is not commutative, however, taking the coefficient of z^0 of the skewsymmetry relation (1.29), we obtain the “quasicommutativity”, which can be written in the following form:

$$: ab : - p(a, b) : ba := \int_{-T}^0 [a_\lambda b] d\lambda. \quad (1.39)$$

Likewise, this product is not associative, but, taking the coefficient of z^0 of the -1^{st} product identity, i.e. (1.10) for $n = -1$, we obtain the “quasi-associativity”, which can be written in the following form:

$$:: ab : c : - : a : bc :: = \left(\int_0^T d\lambda a \right) [b_\lambda c] : + p(a, b) : \left(\int_0^T d\lambda b \right) [a_\lambda c] : . \quad (1.40)$$

The integrals in formulas (1.39) and (1.40) are interpreted as follows: expand the λ -bracket and put the powers of λ on the left, under the sign of integral, then take the definite integral by the usual rules inside the parenthesis. For example we have $:(\int_0^T d\lambda a)[b_\lambda c]: = :(\int_0^T \sum_{j \geq 0} \frac{\lambda^j}{j!} d\lambda a)(b_{(j)}c): = \sum_{j \geq 0} : \frac{T^{j+1}}{(j+1)!} a(b_{(j)}c): = \sum_{j \geq 0} a_{(-j-2)}(b_{(j)}c)$.

We thus arrive at the fourth definition of a vertex algebra [BK].

Definition 1.13. A vertex algebra is a quintuple $(V, |0\rangle, T, [\cdot_\lambda \cdot], : :)$, where

- (i) $(V, T, [\cdot_\lambda \cdot])$ is a Lie conformal superalgebra,

- (ii) $(V, |0\rangle, T, ::)$ is a unital differential (i.e. T is a derivation) superalgebra, satisfying (1.39) and (1.40),
- (iii) the λ -bracket $[\cdot, \cdot]_\lambda$ and the product $::$ are related by the non-commutative Wick formula (1.37).

We have shown that the fourth definition follows from the previous ones. A proof of the fact that the fourth definition implies the first one can be found in [BK].

Remark 1.14. It is proved in [BK] that one can replace the quasi-associativity axiom (1.40) with a weaker version of it, namely the condition that the associator $:(:ab:)c:-:a(:bc):$ is symmetric in a and b , or, equivalently,

$$:a(:bc:):-p(a,b):b(:ac:):=:(:ab:)c:-p(a,b):(:ba:)c: . \quad (1.41)$$

1.6 Conformal weight

Let $(V, |0\rangle, T, Y)$ be a vertex algebra. Recall that a diagonalizable operator H on the vertex superspace V is called a *Hamiltonian operator* if equation (0.2) holds; the eigenspace decomposition for H :

$$V = \oplus_{\Delta \in \mathbb{C}} V[\Delta] \quad (1.42)$$

is called the *H-grading* of V . If a is an eigenvector of H , its eigenvalue is called the *conformal weight* of a and is denoted by Δ_a or $\Delta(a)$.

Proposition 1.15. *A decomposition (1.42) is an H-grading of V if and only if either of the following two equivalent properties holds:*

- (i) *If for $a \in V[\Delta]$ we write $Y(a, z) = \sum_{n \in \mathbb{Z}-\Delta} a_n z^{-n-\Delta}$, then*

$$[H, a_n] = -na_n . \quad (1.43)$$

- (ii) *If $a \in V[\Delta_a]$, $b \in V[\Delta_b]$, $n \in \mathbb{Z}$, then*

$$\Delta_{a_{(n)}b} = \Delta_a + \Delta_b - n - 1 . \quad (1.44)$$

Proof. It is immediate to see that (0.2) and (1.43) are equivalent. We have, by definition,

$$a_{(n)} = a_{n-\Delta_a+1} , \quad n \in \mathbb{Z} , \quad a_m = a_{(m+\Delta_a-1)} , \quad m \in \mathbb{Z} - \Delta_a . \quad (1.45)$$

Hence (1.44) is equivalent to

$$\Delta_{a_m b} = \Delta_{a_{(m+\Delta_a-1)}b} = \Delta_b - m , \quad m \in \mathbb{Z} - \Delta_a ,$$

which is in turn equivalent to (1.43), by applying both sides to b . \square

Remark 1.16. Notice that, using (1.11) and (1.12), we get from (1.44) that

$$\Delta_{|0\rangle} = 0, \quad \Delta_{Ta} = \Delta_a + 1. \quad (1.46)$$

Formulas (1.44) and (1.46) show that $e^{2\pi i H}$ is an automorphism of the vertex algebra V . (However, the H -grading is not a vertex algebra grading.)

In the most important case, when the Hamiltonian operator comes from an energy-momentum field $Y(L, z)$, i.e. $H = L_0$ (see Subsection 0.3), its eigenvector $a \in V$ has conformal weight Δ if and only if

$$[L_\lambda a] = (T + \Delta\lambda)a + O(\lambda^2). \quad (1.47)$$

In the case when $[L_\lambda a] = (T + \Delta\lambda)a$, the vector a and the corresponding vertex operator $Y(a, z)$ are called *primary*. This is equivalent to $L_n a = \delta_{n,0}\Delta a$ for all $n \in \mathbb{Z}_+$.

Taking the formal Fourier transform of both sides of (0.2) applied to b , we obtain the definition of a Hamiltonian operator on a Lie conformal algebra R , as a diagonalizable operator H on R such that

$$[Ha \lambda b] + [a \lambda Hb] = (H + \lambda\partial_\lambda + I_R)[a \lambda b]. \quad (1.48)$$

The most important example comes from a conformal derivation L of R , such that $L_{(0)} = T$, and $L_{(1)} = H$ is diagonalizable (cf. (1.47)).

1.7 Universal enveloping vertex algebras of Lie conformal algebras

A fairly general and very important class of vertex algebras is obtained from regular formal distribution Lie algebras as their universal enveloping vertex algebras [K].

Recall that a formal distribution Lie (super)algebra is a pair $(\mathfrak{g}, \mathcal{F})$ where \mathfrak{g} is a Lie (super)algebra and \mathcal{F} is a local family of \mathfrak{g} -valued formal distributions $\{a^j(z) = \sum_{n \in \mathbb{Z}} a_{(n)}^j z^{-n-1}\}_{j \in J}$, whose coefficients $a_{(n)}^j$ span \mathfrak{g} . Recall that locality means that $(z-w)^{N_{ij}}[a^i(z), a^j(w)] = 0$ for some $N_{ij} \in \mathbb{Z}_+$ and all $i, j \in J$. The formal distribution Lie superalgebra $(\mathfrak{g}, \mathcal{F})$ is called regular if there exists a derivation T of the algebra \mathfrak{g} , such that

$$Ta^j(z) = \partial_z a^j(z), \quad j \in J. \quad (1.49)$$

Due to Dong's lemma (see e.g. [K], Lemma 3.2), the minimal family $\bar{\mathcal{F}}$ of \mathfrak{g} -valued formal distributions containing \mathcal{F} , which is ∂_z -invariant and closed under all products (cf. (1.6))

$$a(w)_{(j)}b(w) = \text{Res}_z(z-w)^j[a(z), b(w)], \quad j \in \mathbb{Z}_+,$$

is again local. Hence, by (1.13) and (1.14) we have for each pair $a(z), b(w) \in \bar{\mathcal{F}}$:

$$[a(z), b(w)] = \sum_{j \in \mathbb{Z}_+} (a(w)_{(j)}b(w))\partial_w^j \delta(z-w)/j!, \quad (1.50)$$

where the sum is finite by locality. In terms of coefficients, equation (1.50) can be written as follows:

$$[a_{(m)}, b_{(n)}] = \sum_{j \in \mathbb{Z}_+} \binom{m}{j} (a_{(j)} b)_{(m+n-j)}, \quad (1.51)$$

where $(a_{(j)} b)_{(n)} = \text{Res}_w w^n a(w) b(w)$.

Denote by $\mathfrak{g}_- \subset \mathfrak{g}$ the span of all $a_{(m)}$ with $a(z) \in \bar{\mathcal{F}}$, $m \in \mathbb{Z}_+$. It follows from (1.51) that \mathfrak{g}_- is a subalgebra of the Lie superalgebra \mathfrak{g} (it is called the *annihilation subalgebra*) and, moreover, since (1.49) for $a(z)$ means that $T(a_{(n)}) = -na_{(n-1)}$, \mathfrak{g}_- is a T -invariant subalgebra. Let $V \equiv V(\mathfrak{g}, \mathcal{F}) = U(\mathfrak{g})/U(\mathfrak{g})\mathfrak{g}_-$, and let $|0\rangle$ denote the image of $1 \in U(\mathfrak{g})$ in V . Extending T to a derivation of $U(\mathfrak{g})$, we get the induced endomorphism of V , which we again denote by T . Finally, each \mathfrak{g} -valued formal distribution $a^j(z) \in \mathcal{F}$ induces an $\text{End} V$ -valued distribution, which we again denote by $a^j(z)$, via the representation given by the left multiplication, so that (1.50) still holds. Hence $\mathcal{F} = \{a^j(z)\}_{j \in \mathbb{Z}}$ is again a local system. Moreover, the $a^j(z)$ are quantum fields. Indeed, for each monomial $v = a_{(n_1)}^{j_1} \dots a_{(n_s)}^{j_s} |0\rangle \in V$ one easily checks by induction on s , using (1.51), that $a_{(n)}^j v = 0$ for $n \gg 0$. Thus, all axioms of the first definition of a vertex algebra hold, and we obtain

Theorem 1.17. $(V(\mathfrak{g}, \mathcal{F}), |0\rangle, T, \mathcal{F})$ is a vertex algebra.

The vertex algebra $V(\mathfrak{g}, \mathcal{F})$ is called *the universal enveloping vertex algebra* of the regular formal distribution Lie algebra $(\mathfrak{g}, \mathcal{F})$.

Remark 1.18. Note that $\bar{\mathcal{F}}$ is a T -module. Define a λ -bracket on $\bar{\mathcal{F}}$ by (cf. (1.50)): $[a_\lambda b] = \sum_{j \in \mathbb{Z}_+} \frac{\lambda^j}{j!} (a_{(j)} b)$. It is not difficult to show (see [K]) that we thus define on $\bar{\mathcal{F}}$ a structure of a Lie conformal algebra.

A special case of the above construction is the universal enveloping vertex algebra $V(R)$ of a Lie conformal algebra $(R, T, [\cdot_\lambda \cdot])$. First we construct the *maximal* formal distribution Lie algebra $(\text{Lie} R, R)$, associated to R , by letting [K]:

$$\text{Lie} R = R[t, t^{-1}] / (T + \partial_t) R[t, t^{-1}]$$

with the bracket given by (1.51), where $a_{(n)}$ stands for the image of at^n in $\text{Lie} R$, and the family of $\text{Lie} R$ -valued formal distributions $\{a(z) = \sum_{n \in \mathbb{Z}} a_{(n)} z^{-n-1}\}_{a \in R}$. Then $(\text{Lie} R, R)$ is regular since $T (= -\partial_t)$ on $\text{Lie} R$ satisfies (1.49), and the image of $R[t]$ in $\text{Lie} R$ is the annihilation subalgebra.

Then $V(R) = V(\text{Lie} R, R)$ is called the *universal enveloping vertex algebra* of the Lie conformal algebra R .

It is easy to show that the map $a \mapsto at^{-1}|0\rangle$ defines a Lie conformal algebra embedding: $R \hookrightarrow V(R)$. Moreover, it is easy to see that in $V(R)$:

$$: aB := aB \text{ if } a \in R, \quad B \in V(R), \quad (1.52)$$

where the product on the right is the product of $U(\text{Lie} R)$.

Definition 1.19. A set of elements $\mathcal{B} = \{a_i \mid i \in \mathcal{I}\}$ *strongly generates* a vertex algebra V if the monomials $: a_{j_1} a_{j_2} \dots a_{j_s} :$ span V . Here and further the normally ordered product of $s > 2$ (resp. $s = 0$) elements is understood from right to left (resp. is $|0\rangle$). An ordered set of elements $\mathcal{B} = \{a_i \mid i \in \mathcal{I}\}$ *freely generates* V if the monomials $: a_{j_1} \dots a_{j_s} :$ with $j_r \leq j_{r+1}$, and $j_r < j_{r+1}$ when $p(a_{j_r}) = \bar{1}$, $1 \leq r < s$, form a basis (over \mathbb{C}) of V .

Proposition 1.20. (a) *The inclusion $R \hookrightarrow V(R)$ satisfies the universality property, similar to that of the inclusion of a Lie algebra in its universal enveloping algebra.*

(b) *Any ordered basis (over \mathbb{C}) of R freely generates $V(R)$.*

Proof. It follows from the corresponding properties of the universal enveloping algebras of Lie (super)algebras. For example, in order to prove (b), we use the decomposition

$$\text{Lie} R = R t^{-1} \oplus (\text{Lie} R)_-,$$

as direct sum of subspaces, which follows by the identity $at^{-n-1} = (T^n a/n!)t^{-1}$ in $(\text{Lie} R)_-$. Hence $R(\simeq R t^{-1})$ freely generates $V(R) \simeq U(R t^{-1})$ by the PBW Theorem for ordinary Lie algebras. \square

An equivalent construction of $V(R)$, purely in terms of the Lie conformal algebra R , is as follows [BK, GMS]. Denote by R_L the Lie (super)algebra structure on R , defined by (cf. (1.39)) $[a, b] = \int_{-T}^0 [a_\lambda b] d\lambda$, and let $U(R_L)$ be its universal enveloping (super)algebra. Then $U(R_L)$ carries a unique structure of a vertex algebra with $|0\rangle = 1$, with T , extended from R_L to $U(R_L)$ as a derivation, the λ -bracket defined on R_L as on R , the normally ordered product defined by (cf. (1.52))

$$: aB := aB \text{ if } a \in R_L, \quad B \in U(R_L),$$

and the λ -bracket and normally ordered product extended to $U(R_L)$ by (1.37), (1.38), (1.39), (1.40). The vertex algebra isomorphism $U(R_L) \xrightarrow{\sim} V(R)$ is induced by the map $a \mapsto a_{(-1)}|0\rangle$, $a \in R_L$.

Remark 1.21. It is easy to see that a Hamiltonian operator on a Lie conformal algebra R , defined by (1.48), extends uniquely to a Hamiltonian operator of the vertex algebra $V(R)$.

1.8 Three examples of vertex algebras and energy-momentum fields

The most important examples of regular formal distribution Lie superalgebras and the associated universal enveloping vertex algebras are the following three.

Universal affine vertex algebra

Let $\mathfrak{g} = \mathfrak{g}_0 + \mathfrak{g}_1$ be a simple finite-dimensional Lie superalgebra with a non-degenerate supersymmetric invariant bilinear form $(\cdot | \cdot)$. Recall that “supersymmetric” means that the form $(\cdot | \cdot)$ is symmetric (resp. skewsymmetric) on \mathfrak{g}_0 (resp. \mathfrak{g}_1) and $(\mathfrak{g}_0 | \mathfrak{g}_1) = 0$, and “invariant” means that $([a, b] | c) = (a | [b, c])$.

The *Kac-Moody affinization* is the formal distribution Lie superalgebra $(\hat{\mathfrak{g}}, \mathcal{F})$, where $\hat{\mathfrak{g}} = \mathfrak{g}[t, t^{-1}] \oplus \mathbb{C}K$ is the Lie superalgebra with parity given by $p(t) = p(K) = 0$, and Lie bracket $(a, b \in \mathfrak{g}, m, n \in \mathbb{Z})$

$$[at^m, bt^n] = [a, b]t^{m+n} + m(a|b)\delta_{m,-n}K, \quad [K, \hat{\mathfrak{g}}] = 0, \quad (1.53)$$

and \mathcal{F} is the following collection of $\hat{\mathfrak{g}}$ -valued formal distributions:

$$\mathcal{F} = \left\{ a(z) = \sum_n (at^n)z^{-n-1} \mid a \in \mathfrak{g} \right\} \cup \{K\}.$$

Notice that the collection \mathcal{F} is local since an equivalent form of (1.53) is

$$[a(z), b(w)] = [a, b](w)\delta(z-w) + (a|b)K\partial_w\delta(z-w), \quad [a(z), K] = 0. \quad (1.54)$$

The pair $(\hat{\mathfrak{g}}, \mathcal{F})$ is regular with $T = -\frac{d}{dt}$. Hence we have the universal enveloping vertex algebra $V(\hat{\mathfrak{g}}, \mathcal{F})$.

Given $k \in \mathbb{C}$, the subspace $(K - k)V(\hat{\mathfrak{g}}, \mathcal{F})$ is an ideal of $V(\hat{\mathfrak{g}}, \mathcal{F})$. The vertex algebra

$$V^k(\mathfrak{g}) = V(\hat{\mathfrak{g}}, \mathcal{F}) / (K - k)V(\hat{\mathfrak{g}}, \mathcal{F})$$

is called the *universal affine* vertex algebra of *level* k . Note that the Lie conformal algebra associated to $(\hat{\mathfrak{g}}, \mathcal{F})$ is

$$\text{Cur}\mathfrak{g} = (\mathbb{C}[T] \otimes \mathfrak{g}) \oplus \mathbb{C}K, \quad [a_\lambda b] = [a, b] + \lambda(a|b)K, \quad [a_\lambda K] = 0 = [K_\lambda K],$$

for $a, b \in 1 \otimes \mathfrak{g}$. It is called the current Lie conformal algebra. It is easy to see that

$$V(\text{Cur}\mathfrak{g}) = V(\hat{\mathfrak{g}}, \mathcal{F}). \quad (1.55)$$

Femionic vertex algebra

Let $A = A_0 + A_1$ be a vector superspace with a non-degenerate skew-supersymmetric bilinear form $\langle \cdot, \cdot \rangle$. Recall that “skew-supersymmetric” means supersymmetric if the parity of A is reversed. The *Clifford affinization* is the formal distribution Lie superalgebra (\hat{A}, \mathcal{F}) , where

$$\hat{A} = A[t, t^{-1}] \oplus \mathbb{C}K, \quad \mathcal{F} = \left\{ a(z) = \sum_{n \in \mathbb{Z}} (at^n)z^{-n-1} \mid a \in A \right\} \cup \{K\},$$

and the bracket on \hat{A} is defined by

$$[at^m, bt^n] = \langle a, b \rangle \delta_{m,-n-1}K, \quad [K, \hat{A}] = 0, \quad (1.56)$$

or, equivalently, by

$$[a(z), b(w)] = \langle a, b \rangle \delta(z - w) K, \quad [a(z), K] = 0. \quad (1.57)$$

For the same reason as above, (\hat{A}, \mathcal{F}) is a regular formal distribution Lie superalgebra, hence we obtain the universal enveloping vertex algebra $V(\hat{A}, \mathcal{F})$. The vertex algebra

$$F(A) = V(\hat{A}, \mathcal{F}) / (K - 1)V(\hat{A}, \mathcal{F})$$

is called the *fermionic* vertex algebra, attached to $(A, \langle \cdot, \cdot \rangle)$.

Remark 1.22. It is important to point out that the properties of $V^k(\mathfrak{g})$ depend essentially on k (for example it is simple for generic k , but not simple for a certain infinite set of values of k), whereas all $F^k(A)$ are isomorphic (and simple) if $k \neq 0$, that is why we put $k = 1$.

The *fermionic* Lie conformal algebra, associated to (\hat{A}, \mathcal{F}) , is

$$(\mathbb{C}[T] \otimes A) \oplus \mathbb{C}K, \quad [a_\lambda b] = \langle a, b \rangle K, \quad [a_\lambda K] = 0 = [K_\lambda K],$$

for $a, b \in 1 \otimes A$, and we have an isomorphism, similar to (1.55).

Universal Virasoro vertex algebra

This example is constructed starting with the formal distribution Lie superalgebra $(\text{Vir}, \mathcal{F})$, where $\text{Vir} = (\bigoplus_{n \in \mathbb{Z}} \mathbb{C}L_n) \oplus \mathbb{C}C$ is the Virasoro Lie algebra, with Lie bracket as in (0.3) with c replaced by C , and $\mathcal{F} = \{L(z) = \sum_n L_n z^{-n-2}, C\}$. Locality holds since (0.3) is equivalent to

$$[L(z), L(w)] = \partial_w L(w) \delta(z - w) + 2L(w) \partial_w \delta(z - w) + \frac{C}{12} \partial_w^3 \delta(z - w), \quad (1.58)$$

and the pair $(\text{Vir}, \mathcal{F})$ is regular with $T = L_{-1}$. The vertex algebra $(c \in \mathbb{C})$

$$V^c = V(\text{Vir}, \mathcal{F}) / (C - c)V(\text{Vir}, \mathcal{F})$$

is called the universal Virasoro vertex algebra with central charge c .

Note that the associated Lie conformal algebra, called the Virasoro Lie conformal algebra, is

$$(\mathbb{C}[T] \otimes L) \oplus \mathbb{C}C, \quad [L_\lambda L] = (T + 2\lambda)L + \frac{\lambda^3}{12}C, \quad [L_\lambda C] = 0 = [C_\lambda C], \quad (1.59)$$

and an isomorphism similar to (1.55) holds.

Energy-momentum fields

We conclude this section by constructing energy-momentum fields $Y(L, z)$ for each of the above examples, so that $H = L_0 = L_{(1)}$ is a Hamiltonian operator (cf. Subsection 0.3).

For $V^k(\mathfrak{g})$ it is the well-known Sugawara construction (defined for $k \neq -h^\vee$)

$$L^{\mathfrak{g}} = \frac{1}{2(k + h^\vee)} \sum_{i \in I} : a^i a_i : , \quad (1.60)$$

where $\{a_i\}_{i \in I}$ and $\{a^i\}_{i \in I}$ is a pair of dual bases of \mathfrak{g} , i.e. $(a_i | a^j) = \delta_{ij}$, we identify $a \in \mathfrak{g}$ with $(at^{-1})|0\rangle \in V^k(\mathfrak{g})$, and h^\vee is the dual Coxeter number, i.e. the eigenvalue of the Casimir operator $\sum_i a^i a_i$ on \mathfrak{g} divided by 2. Then $Y(L^{\mathfrak{g}}, z) = \sum_n L_n^{\mathfrak{g}} z^{-n-2}$ is an energy-momentum field with central charge

$$c_k^{\mathfrak{g}} = \frac{k \operatorname{sdim} \mathfrak{g}}{k + h^\vee}, \text{ where } \operatorname{sdim} \mathfrak{g} = \dim \mathfrak{g}_0 - \dim \mathfrak{g}_1. \quad (1.61)$$

Moreover, all $a(z)$, where $a \in \mathfrak{g}$, are primary of conformal weight 1. This is easy to prove using the non-commutative Wick formula (1.37).

Remark 1.23. The vertex algebra $V^k(\mathfrak{g})$ can be constructed as above for any Lie superalgebra \mathfrak{g} and a supersymmetric invariant bilinear form $(\cdot | \cdot)$. Also, the Hamiltonian operator $H = L_0^{\mathfrak{g}}$ can be constructed in this generality for any k by making use of the derivation $-t \frac{d}{dt}$ of $\hat{\mathfrak{g}}$. More generally, if h is a diagonalizable derivation of \mathfrak{g} , then a Hamiltonian operator can be constructed by using the derivation $-t \frac{d}{dt} - \operatorname{ad} h$ of $\hat{\mathfrak{g}}$. In this case the field $a(z)$ has conformal weight $1 - j$ if $[h, a] = ja$, $j \in \mathbb{C}$. Note also that all derivations $-t^{n+1} \frac{d}{dt}$ of $\hat{\mathfrak{g}}$ with $n \geq -1$ extend to the endomorphisms of the space $V^k(\mathfrak{g})$, which form the "annihilation subalgebra" $\sum_{n \geq -1} \mathbb{C} L_n$ of the Virasoro algebra, and one has:

$$[L_n, Y(a, z)] = \sum_{j \geq -1} \binom{n+1}{j+1} Y(L_j a, z) z^{n-j}, \quad n \geq -1.$$

For $F(A)$ we let

$$L^A = \frac{1}{2} \sum_{i \in I} : (T\Phi^i) \Phi_i : , \quad (1.62)$$

where $\{\Phi_i\}_{i \in I}$ and $\{\Phi^i\}_{i \in I}$ is a pair of dual bases of A , i.e. $\langle \Phi_i, \Phi^i \rangle = \delta_{ij}$, and we identify $\Phi \in A$ with $(\Phi t^{-1})|0\rangle$. Then $Y(L^A, z)$ is an energy-momentum field with central charge

$$c^A = -\frac{1}{2} \operatorname{sdim} A. \quad (1.63)$$

All the fields $\Phi(z)$, where $\Phi \in A$, are primary of conformal weight $1/2$.

Next, we consider the following important modification of this construction. Suppose that $A = A_+ \oplus A_-$, where both A_+ and A_- are isotropic subspaces with respect to the bilinear form $\langle \cdot, \cdot \rangle$. Choose a basis $\{\varphi_i\}$ of A_+ and the dual basis $\{\varphi^i\}$ of A_- , so that $\langle \varphi_i, \varphi^j \rangle = \delta_{ij}$. Then we can define a family of energy-momentum fields, parameterized by a collection $\vec{m} = (m_i)$ of complex numbers:

$$L^{A, \vec{m}} = - \sum_i m_i : \varphi^i (T\varphi_i) : + \sum_i (1 - m_i) : (T\varphi^i) \varphi_i : . \quad (1.64)$$

The central charge of $Y(L^{A,\vec{m}}, z)$ is equal

$$c_{\vec{m}}^A = \sum_i s(\varphi_i)(12m_i^2 - 12m_i + 2), \quad (1.65)$$

and all the fields $\varphi^i(z)$ and $\varphi_i(z)$ are primary of conformal weight m_i and $1 - m_i$, respectively.

Note that the vertex algebra $F(A = A_+ \oplus A_-)$ has an important \mathbb{Z} -grading,

$$F(A) = \oplus_{m \in \mathbb{Z}} F_m(A), \quad (1.66)$$

called the *charge decomposition*, which is defined by

$$\text{charge } |0\rangle = 0, \quad \text{charge } \varphi^i(z) = 1, \quad \text{charge } \varphi_i(z) = -1. \quad (1.67)$$

Finally, for the Virasoro vertex algebra V^c the energy-momentum field is, of course, $L(z) = Y(L_{-2}|0\rangle, z)$, with central charge c .

1.9 The indefinite integral definition of a vertex algebra

In this section we introduce a new, fifth definition of vertex algebras. In some sense it just consists in a change of notation in the fourth definition discussed in Section 1.5, but it is much more elegant, as it encodes all axioms, such as formulas (1.37), (1.38), (1.39) and (1.40), in the axioms of a Lie conformal algebra (see Definition 1.10) written in "integral form".

In a nutshell, the fifth definition says that a vertex algebra is a Lie conformal algebra in which the λ -bracket can be "integrated". More precisely we have

Definition 1.24. A vertex algebra is a $\mathbb{Z}/2\mathbb{Z}$ -graded $\mathbb{C}[T]$ -module V , endowed with an even element $|0\rangle \in V_0$ and an *integral λ -bracket*, namely a linear map $V \otimes V \rightarrow \mathbb{C}[\lambda] \otimes V$, denoted by $\int^\lambda dx [a \text{ }_x \text{ } b]$, such that the following axioms hold:

(unity)

$$\int^\lambda dx [|0\rangle \text{ }_x \text{ } a] = \int^\lambda dx [a \text{ }_x \text{ } |0\rangle] = a,$$

(sesquilinearity)

$$\int^\lambda dx [T a \text{ }_x \text{ } b] = - \int^\lambda dx x [a \text{ }_x \text{ } b], \quad \int^\lambda dx [a \text{ }_x \text{ } T b] = \int^\lambda dx (T + x) [a \text{ }_x \text{ } b], \quad (1.68)$$

(skewsymmetry)

$$\int^\lambda dx [b \text{ }_x \text{ } a] = -p(a, b) \int^\lambda dx [a \text{ }_{-x-T} \text{ } b], \quad (1.69)$$

(Jacobi identity)

$$\int^\lambda dx \int^\mu dy \left\{ [a_x [b_y c]] - p(a, b) [b_y [a_x c]] - [[a_x b]_{x+y} c] \right\} = 0 . \quad (1.70)$$

Before showing that Definition 1.24 is equivalent to Definition 1.13, some words of explanation are needed. In particular, not all expressions above are immediately defined in terms of the integral λ -bracket $\int^\lambda dx [a_x b]$, but in some sense they are "canonically" defined, since there is a unique way to make standard "integral manipulations", such as change of variables and change of order of integration, to rewrite them as expressions which make sense, i.e. which are well defined only in terms of the integral λ -bracket. In all manipulations it is convenient to think of $\int^\lambda dx [a_x b]$ as being a definite integral with lower bound $-\infty$. Moreover, not all relations above are in fact axioms: some of them should be considered as defining relations for some new notation. This is best understood if we write the integral λ -bracket as $\int^\lambda dx [a_x b] = I_\lambda(a, b)$, and we try to write all relations above in terms of I_λ only.

The first sesquilinearity relation (1.68) includes a piece of notation,

$$\int^\lambda dx x [a_x b] := -I_\lambda(Ta, b) , \quad (1.71)$$

and an axiom,

$$\frac{d}{d\lambda} I_\lambda(Ta, b) = -\lambda \frac{d}{d\lambda} I_\lambda(a, b) . \quad (1.72)$$

The latter follows by the "fundamental Theorem of calculus", which is encoded in the integral notation. Using (1.71), the second sesquilinearity relation (1.68) is equivalent to

$$TI_\lambda(a, b) = I_\lambda(Ta, b) + I_\lambda(a, Tb) . \quad (1.73)$$

Notice that here we are implicitly assuming that the action of T commutes with taking integrals. The right hand side of the skewsymmetry relation (1.69) is defined by a change of variable $y = -x - T$:

$$\int^\lambda dx [a_{-x-T} b] = - \int^{-\lambda-T} dy [a_y b] = -I_{-\lambda-T}(a, b) .$$

Hence, in terms of I_λ axiom (1.69) is equivalent to

$$I_\lambda(b, a) = p(a, b) I_{-\lambda-T}(a, b) . \quad (1.74)$$

We are left to consider the Jacobi identity. The first two terms are quite clear. For example, the first term should be understood as

$$\int^\lambda dx [a_x \left(\int^\mu dy [b_y c] \right)] = I_\lambda(a, I_\mu(b, c)) ,$$

and similarly the second term, obtained by exchanging λ with μ and a with b . Less clear is the last term in (1.70). For this, we first need to perform a change of variable $z = x + y$, which gives

$$- \int_{-\infty}^{\lambda} dx \int_{\mu+x}^{\mu+x} dz [[a \ x \ b]_z \ c] . \quad (1.75)$$

Since we want to perform *first* the integral in x (corresponding to the inner bracket) and *then* the integral in z , we now need to exchange the order of integration. For this we have to be careful (as we usually are, when exchanging order of integration in multiple integrals), and, as explained above, we think of $\int^{\lambda} dx$ as $\int_{-\infty}^{\lambda} dx$. With this in mind, (1.75) is an integral on the following region of the (x, z) -plane: $\{(x, z) \mid x \leq \lambda, z \leq \mu + x\}$, which can be equivalently written as $\{(x, z) \mid z \leq \lambda + \mu, z - \mu \leq x \leq \lambda\}$. Hence, after exchanging the order of integration, (1.75) becomes

$$- \int_{\lambda+\mu}^{\lambda+\mu} dz \left(\int_{z-\mu}^{\lambda} dx [a \ x \ b] \right) [z \ c] .$$

In the inner integral we finally replace z by $-T$, thanks to (1.71), and now it is easy to rewrite the above expression in terms of the function I_{λ} :

$$-I_{\lambda+\mu}(I_{\lambda}(a, b) - I_{-\mu-T}(a, b), c) .$$

Thus, the Jacobi identity is equivalent to the following equation:

$$I_{\lambda}(a, I_{\mu}(b, c)) - p(a, b)I_{\mu}(b, I_{\lambda}(a, c)) = I_{\lambda+\mu}(I_{\lambda}(a, b) - I_{-\mu-T}(a, b), c) . \quad (1.76)$$

To go from Definition 1.24 to Definition 1.13 we put

$$: ab : = \int_0^0 dx [a \ x \ b] = I_0(a, b) , \quad [a \ \lambda \ b] = \frac{d}{d\lambda} \int^{\lambda} dx [a \ x \ b] = \frac{d}{d\lambda} I_{\lambda}(a, b) , \quad (1.77)$$

and conversely, to go from Definition 1.13 to Definition 1.24 we let

$$\int^{\lambda} dx [a \ x \ b] = : ab : + \int_0^{\lambda} dx [a \ x \ b] .$$

We then only need to check that the above axioms, written in terms of the normally ordered product and the λ -bracket defined by (1.77), correspond to the axioms of vertex algebra as stated in Definition 1.13.

It is immediate to check that the sesquilinearity relations (1.72) and (1.73) are equivalent both to the sesquilinearity of the λ -bracket $[a \ \lambda \ b]$, and to the fact that T is a derivation of the normally ordered product $: ab :$. Similarly, the skewsymmetry axiom (1.74) gives both the skewsymmetry of the λ -bracket $[a \ \lambda \ b]$ and the skewsymmetry relation (1.39) for the normally ordered product $: ab :$. We are left to consider the Jacobi identity (1.76). It gives three different

conditions. If we look at the constant term in λ and μ , namely we put $\lambda = \mu = 0$ in (1.76), we get

$$: a(: bc :) : -p(a, b) : b(: ac :) : = : \left(\int_{-T}^0 d\lambda [a \underset{\lambda}{} b] \right) c : ,$$

which, in view of (1.39), is equivalent to the "weak" quasi-associativity (1.41). If instead we put $\mu = 0$ in (1.76) and we take the derivative of both sides with respect to λ , we get the left Wick formula (1.37). Notice that (1.76) is unchanged if we exchange λ with μ and a with b . Hence we do not need to look separately at the constant term in λ . Finally, if we take the derivatives with respect to both λ and μ of both sides of (1.70) (equivalent to (1.76), we get the Jacobi identity for the λ -bracket $[a \underset{\lambda}{} b]$.

The above arguments, together with Remark 1.14, clearly prove that Definition 1.13 and Definition 1.24 are indeed equivalent.

2 The Zhu algebra and representation theory

2.1 \hbar -deformation of vertex operators

Let V be a vertex algebra with the vacuum vector $|0\rangle$, the infinitesimal translation operator T and the vertex operators $Y(a, z) = \sum_{n \in \mathbb{Z}} a_{(n)} z^{-n-1}$ ($a \in V$). Fix a Hamiltonian operator H on V (see Subsection 1.6), and let (1.42) be the eigenspace decomposition for H .

Define the \hbar -deformation of the vertex operator $Y(a, z)$ as follows:

$$Y_{\hbar}(a, z) = (1 + \hbar z)^{\Delta_a} Y(a, z) = \sum_{n \in \mathbb{Z}} a_{(n, \hbar)} z^{-n-1}. \quad (2.1)$$

Here and further, as before, when a formula involves Δ_a it is assumed that a is an eigenvector of H with eigenvalue Δ_a , and the formula is extended to arbitrary a by linearity. We have, obviously, the following formula for \hbar -deformed n^{th} products:

$$a_{(n, \hbar)} b = \sum_{j \in \mathbb{Z}_+} \binom{\Delta_a}{j} \hbar^j a_{(n+j)} b. \quad (2.2)$$

We view \hbar as an indeterminate, and we expand $(1 + \hbar z)^{\Delta_a} = \sum_{j \in \mathbb{Z}_+} \binom{\Delta_a}{j} \hbar^j z^j$, so that $Y_{\hbar}(a, z) \in (\text{End} V)[[\hbar]][[z, z^{-1}]]$, i.e. $a_{(n, \hbar)} \in (\text{End} V)[[\hbar]]$ is a formal power series in \hbar (with coefficients in $\text{End} V$). However, $a_{(n, \hbar)} b$ is a polynomial in \hbar (with coefficients in V), so that we can specialize \hbar to any value in \mathbb{C} . Note also that the space $(\text{End} V)[[\hbar]][[z, z^{-1}]]$ is also a module over the algebra $\mathbb{C}[[\hbar z]]$, which allows us to expand binomials $(1 + \hbar z)^{\Delta_a}$ into formal power series.

The 0^{th} product is often viewed as a bracket on V . We introduce the following \hbar -deformation of this bracket (which is different from $a_{(0, \hbar)} b$):

$$[a, b]_{\hbar} = \sum_{j \in \mathbb{Z}_+} (-\hbar)^j a_{(j, \hbar)} b. \quad (2.3)$$

Note that, by (2.1),

$$[a, b]_{\hbar} = \text{Res}_z (1 + \hbar z)^{-1} Y_{\hbar}(a, z) b = \text{Res}_z (1 + \hbar z)^{\Delta_a - 1} Y(a, z) b ,$$

hence we obtain

$$[a, b]_{\hbar} = \sum_{j \in \mathbb{Z}_+} \binom{\Delta_a - 1}{j} \hbar^j a_{(j)} b . \quad (2.4)$$

The following is a simple nice relation between the $(-1, \hbar)$ -product (2.2) and the \hbar -bracket (2.4), which can be checked directly:

$$a_{(-1, \hbar)} b = : ab : + \int_0^{\hbar} dx [H(a), b]_x . \quad (2.5)$$

Lemma 2.1. (a) *The following sesquilinearity properties of the \hbar -deformed vertex operators hold*

$$\begin{aligned} Y_{\hbar}(Ta, z) &= (1 + \hbar z)[T, Y_{\hbar}(a, z)] \\ &= ((1 + \hbar z)\partial_z - \hbar\Delta_a)Y_{\hbar}(a, z) . \end{aligned} \quad (2.6)$$

(b) *The (n, \hbar) -product satisfies the following \hbar -deformation of the sesquilinearity equations for the usual n^{th} product:*

$$((T + \hbar(n + 1 + H))a)_{(n, \hbar)} b = -na_{(n-1, \hbar)} b , \quad (2.7)$$

$$T(a_{(n, \hbar)} b) = \sum_{k \in \mathbb{Z}_+} (-\hbar)^k (Ta)_{(n+k, \hbar)} b + a_{(n, \hbar)} (Tb) . \quad (2.8)$$

In particular,

$$a_{(-2, \hbar)} b = ((T + \hbar H)a)_{(-1, \hbar)} b . \quad (2.9)$$

(c) *The \hbar -bracket satisfies the relations:*

$$[(T + \hbar H)a, b]_{\hbar} = 0 , \quad (2.10)$$

$$(T + \hbar H)[a, b]_{\hbar} = [a, (T + \hbar H)b]_{\hbar} . \quad (2.11)$$

(d) *The following identity holds for $k \in \mathbb{Z}_+$,*

$$a_{(-k-1, \hbar)} b = \sum_{j=0}^k \binom{\Delta_a}{k-j} \hbar^{k-j} (T^{(j)}a)_{(-1, \hbar)} b , \quad (2.12)$$

or, equivalently, in terms of \hbar -deformed vertex operators:

$$Y_{\hbar}(a, z)_+ = (1 + \hbar z)^{\Delta_a} (e^{zT}a)_{(-1, \hbar)} .$$

Proof. Equations (2.6) follow by the definition (2.1) of the \hbar -deformed vertex operator and the translation covariance axiom for vertex algebras. By looking at the coefficients of z^{-n-1} in the second identity in (2.6) we get (2.7), while if we first divide both sides of the first identity in (2.6) by $1 + \hbar z$ and then look at the coefficients of z^{-n-1} we get (2.8). Equations (2.10) and (2.11) are easily checked using the formula (2.4) for the \hbar -bracket. We are left to prove part (d). From (2.7) we have

$$a_{(-k-1, \hbar)} b = \frac{1}{k} (Ta)_{(-k, \hbar)} b + \hbar \frac{\Delta_a - k + 1}{k} a_{(-k, \hbar)} b.$$

Equation (2.12) can be easily proved by induction using the above relation. \square

Proposition 2.2. *One has the following \hbar -deformed n^{th} product identity for \hbar -deformed vertex operators:*

$$(1 + \hbar w)^{n+1} Y_{\hbar}(a_{(n, \hbar)} b, w) = Y_{\hbar}(a, w)_{(n)} Y_{\hbar}(b, w).$$

Proof. By (2.2) and (1.44), the LHS of the \hbar -deformed n^{th} product identity is equal to

$$\sum_{j \in \mathbb{Z}_+} \binom{\Delta_a}{j} \hbar^j (1 + \hbar w)^{\Delta_a + \Delta_b - j} Y(a_{(n+j)} b, w).$$

By the n^{th} product identity for usual vertex operators, $Y(a_{(n+j)} b, w) = \text{Res}_z i_{z, w} (z - w)^{n+j} Y(a, z) Y(b, w) - p(a, b) \text{Res}_z i_{w, z} (z - w)^{n+j} Y(b, w) Y(a, z)$. Hence the LHS of the \hbar -deformed n^{th} product identity is equal to

$$\begin{aligned} \text{Res}_z \sum_{j \in \mathbb{Z}_+} \binom{\Delta_a}{j} \hbar^j (z - w)^j (1 + \hbar w)^{\Delta_a + \Delta_b - j} \left\{ i_{z, w} (z - w)^n Y(a, z) Y(b, w) \right. \\ \left. - p(a, b) i_{w, z} (z - w)^n Y(b, w) Y(a, z) \right\} \end{aligned}$$

But $\sum_{j \in \mathbb{Z}_+} \binom{\Delta_a}{j} \hbar^j (z - w)^j (1 + \hbar w)^{\Delta_a + \Delta_b - j} = (1 + \hbar z)^{\Delta_a} (1 + \hbar w)^{\Delta_b}$, by the binomial formula, which gives the RHS of the \hbar -deformed n^{th} product identity. \square

Theorem 2.3. *One has the following identity for the \hbar -deformed vertex operators (\hbar -deformed Borchers identity) for arbitrary $n \in \mathbb{Z}$:*

$$\begin{aligned} i_{z, w} (z - w)^n Y_{\hbar}(a, z) Y_{\hbar}(b, w) - p(a, b) i_{w, z} (z - w)^n Y_{\hbar}(b, w) Y_{\hbar}(a, z) \\ = \sum_{j \in \mathbb{Z}_+} (1 + \hbar w)^{n+j+1} Y_{\hbar}(a_{(n+j, \hbar)} b, w) \partial_w^j \delta(z - w) / j!. \end{aligned}$$

Proof. It is the same as the proof of the usual Borchers identity, given in Section 1.4, by making use of the \hbar -deformed n^{th} product identity, instead of the n^{th} product identity. \square

Comparing the coefficients of z^{-m-1} of both sides of the \hbar -deformed Borchers identity applied to $c \in V$, we obtain:

$$\begin{aligned} \sum_{j \in \mathbb{Z}_+} (-1)^j \binom{n}{j} & \left(a_{(m+n-j, \hbar)} (Y_{\hbar}(b, w)c) w^j - p(a, b) (-1)^n Y_{\hbar}(b, w) (a_{(m+j, \hbar)} c) w^{n-j} \right) \\ &= \sum_{j \in \mathbb{Z}_+} (1 + \hbar w)^{n+j+1} \binom{m}{j} Y_{\hbar}(a_{(n+j, \hbar)} b, w) c w^{m-j} . \end{aligned} \quad (2.13)$$

Note that, remarkably, the \hbar -deformation does not change the -1^{st} product identity:

$$: Y_{\hbar}(a, w) Y_{\hbar}(b, w) : = Y_{\hbar}(a_{(-1, \hbar)} b, w) .$$

The \hbar -deformed commutator formula

$$[Y_{\hbar}(a, z), Y_{\hbar}(b, w)] = \sum_{j \in \mathbb{Z}_+} (1 + \hbar w)^{j+1} Y_{\hbar}(a_{(j, \hbar)} b, w) \partial_w^j \delta(z - w) / j! \quad (2.14)$$

is a special case of the \hbar -deformed Borchers identity for $n = 0$.

We want to rewrite the Borchers identity (2.13) in terms of the (n, \hbar) -products. We can do it in two ways: either we compare the coefficients of w^{-k-1} in both sides of (2.13), or, for the form it will be used, we first multiply both sides of (2.13) by $(1 + \hbar w)^{-n-1}$ and then compare the coefficients of w^{-k-1} . We then get

$$\begin{aligned} \sum_{i, j \in \mathbb{Z}_+} \binom{n}{j} \binom{-n-1}{i} (-1)^j \hbar^i & \left(a_{(m+n-j, \hbar)} (b_{(k+j+i, \hbar)} c) \right. \\ & \left. - (-1)^n p(a, b) b_{(k+n-j+i, \hbar)} (a_{(m+j, \hbar)} c) \right) \\ &= \sum_{i, j \in \mathbb{Z}_+} \binom{m}{j} \binom{j}{i} \hbar^i (a_{(n+j, \hbar)} b)_{(k+m-j+i, \hbar)} c . \end{aligned} \quad (2.15)$$

We now state some results about \hbar -deformed vertex operators which will be useful in the following sections.

Lemma 2.4. (a) One has for $n, k \in \mathbb{Z}_+$:

$$\begin{aligned} (a_{(n, \hbar)} b)_{(k, \hbar)} c &= \sum_{i, j \in \mathbb{Z}_+} \binom{n}{j} \binom{-n-1}{i} (-1)^j \hbar^i \left(a_{(n-j, \hbar)} (b_{(k+j+i, \hbar)} c) \right. \\ & \left. - (-1)^n p(a, b) b_{(k+n-j+i, \hbar)} (a_{(j, \hbar)} c) \right) . \end{aligned} \quad (2.16)$$

(b) The $(-1, \hbar)$ -product satisfies the following "quasi-associativity" relation

$$\begin{aligned} & (a_{(-1, \hbar)} b)_{(-1, \hbar)} c - a_{(-1, \hbar)} (b_{(-1, \hbar)} c) \\ &= \sum_{j \in \mathbb{Z}_+} \left(a_{(-j-2, \hbar)} (b_{(j, \hbar)} c) + p(a, b) b_{(-j-2, \hbar)} (a_{(j, \hbar)} c) \right) . \end{aligned} \quad (2.17)$$

(c) The following identity holds:

$$\begin{aligned}
& (a_{(-2, \hbar)} b)_{(-1, \hbar)} c = a_{(-2, \hbar)} (b_{(-1, \hbar)} c) \\
& + \sum_{j \in \mathbb{Z}_+} \left((Ta)_{(-j-2, \hbar)} (b_{(j, \hbar)} c) + p(a, b) b_{(-j-2, \hbar)} ((Ta)_{(j, \hbar)} c) \right) \\
& + \hbar \Delta_a \sum_{j \in \mathbb{Z}_+} \left(a_{(-j-2, \hbar)} (b_{(j, \hbar)} c) + p(a, b) b_{(-j-2, \hbar)} (a_{(j, \hbar)} c) \right).
\end{aligned} \tag{2.18}$$

Proof. Equation (2.16) is a special case of (2.15), with $m = 0$. Equation (2.17) is obtained by putting $n = k = -1$ in (2.16). Finally, (2.18) follows immediately by using (2.9) in the LHS and applying (2.17). \square

Lemma 2.5. The following identity holds (where $T^{(n)}$ stands for $T^n/n!$):

$$\begin{aligned}
& a_{(-1, \hbar)} b - p(a, b) b_{(-1, \hbar)} a - \hbar [a, b]_{\hbar} \\
& = \sum_{n \in \mathbb{Z}_+} \sum_{l=0}^n \binom{\Delta_b}{n-l} \hbar^{n-l} (-1)^n \left(T^{(l+1)}(a_{(n)} b) - \hbar^{l+1} \binom{-\Delta_{a_{(n)} b}}{l+1} a_{(n)} b \right).
\end{aligned} \tag{2.19}$$

Proof. Recall the skewsymmetry relation in a vertex algebra (see (1.29)):

$$p(a, b) b_{(n)} a = \sum_{i \in \mathbb{Z}_+} (-1)^{n+i+1} T^{(i)}(a_{(n+i)} b).$$

Using this, we get, by the definition of the $(-1, \hbar)$ -product,

$$\begin{aligned}
p(a, b) b_{(-1, \hbar)} a & = \sum_{k, i \in \mathbb{Z}_+} \binom{\Delta_b}{k} \hbar^k (-1)^{k+i} T^{(i)}(a_{(k+i-1)} b) \\
& = \sum_{n \in \mathbb{Z}_+} \sum_{l=0}^n \binom{\Delta_b}{n-l} \hbar^{n-l} (-1)^{n+1} T^{(l+1)}(a_{(n)} b) \\
& \quad + \sum_{k \in \mathbb{Z}_+} \binom{\Delta_b}{k} (-\hbar)^k a_{(k-1)} b.
\end{aligned} \tag{2.20}$$

The last identity is obtained by separating the summand with $i = 0$ from the rest, and, in the latter, by performing the change of variables $l = i - 1$, $n = k + i - 1 = k + l$. We can add and subtract the appropriate quantity to rewrite the first term in the RHS of (2.20) as

$$\begin{aligned}
& \sum_{n \in \mathbb{Z}_+} \sum_{l=0}^n \binom{\Delta_b}{n-l} \hbar^{n-l} (-1)^{n+1} \left(T^{(l+1)}(a_{(n)} b) - \hbar^{l+1} \binom{-\Delta_{a_{(n)} b}}{l+1} a_{(n)} b \right) \\
& + \sum_{n \in \mathbb{Z}_+} \sum_{l=0}^n \binom{\Delta_b}{n-l} \binom{-\Delta_{a_{(n)} b}}{l+1} (-\hbar)^{n+1} a_{(n)} b.
\end{aligned} \tag{2.21}$$

We now perform the change of variables $k = n + 1$, $i = l + 1$ in the last term of (2.21) to rewrite it as

$$\sum_{k \in \mathbb{Z}_+} \sum_{i=1}^k \binom{\Delta_b}{k-i} \binom{-\Delta_{a_{(k-1)}b}}{i} (-\hbar)^k a_{(k-1)}b ,$$

which, combined to the last term in the RHS of (2.20), gives

$$\sum_{k \in \mathbb{Z}_+} \binom{\Delta_a - 1}{k} \hbar^k a_{(k-1)}b .$$

For this we used the obvious combinatorial identity ($n \in \mathbb{Z}_+$):

$$\binom{x+y}{n} = \sum_{j=0}^n \binom{x}{j} \binom{y}{n-j} \quad (2.22)$$

with $x = -\Delta_a - \Delta_b + k$ and $y = \Delta_b$. The above discussion gives

$$\begin{aligned} p(a, b)b_{(-1, \hbar)}a &= \sum_{n \in \mathbb{Z}_+} \sum_{l=0}^n \binom{\Delta_b}{n-l} \hbar^{n-l} (-1)^{n+1} \left(T^{(l+1)}(a_{(n)}b) \right. \\ &\quad \left. - \hbar^{l+1} \binom{-\Delta_{a_{(n)}b}}{l+1} a_{(n)}b \right) + \sum_{k \in \mathbb{Z}_+} \binom{\Delta_a - 1}{k} \hbar^k a_{(k-1)}b . \end{aligned}$$

Equation (2.19) now follows by the definitions of the $(-1, \hbar)$ -product and \hbar -bracket, and by the relation $\binom{\Delta_a}{k} - \binom{\Delta_a - 1}{k} - \binom{\Delta_a - 1}{k-1} = 0$. \square

Lemma 2.6. *Taking the (n, \hbar) -bracket is a derivation of the \hbar -product:*

$$[a, b_{(n, \hbar)}c]_{\hbar} = ([a, b]_{\hbar})_{(n, \hbar)}c + p(a, b)b_{(n, \hbar)}([a, c]_{\hbar}) , \quad (2.23)$$

or, equivalently, in terms of the \hbar -deformed vertex operators:

$$[a, Y_{\hbar}(b, w)c]_{\hbar} = Y_{\hbar}([a, b]_{\hbar}, w)c + p(a, b)Y_{\hbar}(b, w)[a, c]_{\hbar} .$$

Proof. Take identity (2.13) for $n = 0$, multiply both sides by $(-\hbar)^m$ and sum over $m \in \mathbb{Z}_+$. Then the LHS of the obtained identity is

$$[a, Y_{\hbar}(b, w)c]_{\hbar} - p(a, b)Y_{\hbar}(b, w)[a, c]_{\hbar} ,$$

while the RHS becomes $Y_{\hbar}([a, b]_{\hbar}, w)c$, after changing the order of summation by m and j , and using the expansion in $\mathbb{C}[[x]]$ (for $x = -\hbar w$): $\frac{x^j}{(1-x)^{j+1}} = \sum_{m=0}^{\infty} \binom{m}{j} x^m$. \square

Let $J_{\hbar} \subset V[\hbar]$ be the $\mathbb{C}[\hbar]$ -span of all the elements $a_{(-2, \hbar)}b$ ($a, b \in V$). By (1.27) and (2.9) with $b = |0\rangle$, we have $(T + \hbar H)a \in J_{\hbar}$ for all $a \in V$. Hence, more generally, we obtain by induction on k , using (1.46):

$$T^{(k)}a \equiv \hbar^k \binom{-\Delta_a}{k} a \mod J_{\hbar}. \quad (2.24)$$

Theorem 2.7. (a) J_{\hbar} contains all the elements $a_{(-j, \hbar)}b$, where $j \geq 2$ and $a, b \in V$.

(b) $(T + \hbar H)V \subset J_{\hbar}$.

(c) J_{\hbar} is a two-sided ideal with respect to the product $a_{(-1, \hbar)}b$.

(d) J_{\hbar} is a two-sided ideal with respect to the bracket $\hbar[a, b]_{\hbar}$.

(e) $|0\rangle$ is the identity element for the product $a_{(-1, \hbar)}b$.

(f) $a_{(-1, \hbar)}b - p(a, b)b_{(-1, \hbar)}a - \hbar[a, b]_{\hbar} \in J_{\hbar}$.

(g) $(a_{(-1, \hbar)}b)_{(-1, \hbar)}c - a_{(-1, \hbar)}(b_{(-1, \hbar)}c) \in J_{\hbar}$.

Proof. Claim (a) follows from (2.7); (b) is a special case of (2.24) for $k = 1$; (e) follows from (2.2) and the properties (1.27) of the vacuum; (f) follows from Lemma 2.5 and (2.24); (g) follows from equation (2.17) and (a). Equation (2.18), together with (a), implies $(a_{(-2, \hbar)}b)_{(-1, \hbar)}c \in J_{\hbar}$, hence J_{\hbar} is a right ideal with respect to the $(-1, \hbar)$ -product. Moreover, equation (2.23) with $n = -2$ implies that $[a, b_{(-2, \hbar)}c]_{\hbar} \in J_{\hbar}$, hence J_{\hbar} is a left ideal with respect to the \hbar -bracket. Therefore, due to (f), J_{\hbar} is also a left ideal with respect to the $(-1, \hbar)$ -product, proving (c). Now (d) follows from (c) and (f). \square

By Theorem 2.7(c), (e) and (g), we have a unital associative superalgebra over $\mathbb{C}[\hbar]$:

$$Zhu_{\hbar, H}V := V[\hbar]/J_{\hbar},$$

with the product induced by the $(-1, \hbar)$ -product on $V[\hbar]$, and the image of $|0\rangle$ is the identity element. Furthermore, if we denote by $J_{\hbar=1}$ the \mathbb{C} -span of all elements $\{a_{(-2, \hbar=1)}b \mid a, b \in V\}$, then $V/J_{\hbar=1}$ is a unital associative algebra over \mathbb{C} (cf. [Z]), which will play an important role in the second part of the paper.

Definition 2.8. The H -twisted Zhu algebra of a vertex algebra V with a Hamiltonian H is

$$Zhu_HV := V/J_{\hbar=1}.$$

Remark 2.9. The \hbar -deformation of a vertex algebra is a special case of the following notion of a gauged vertex algebra. In addition to the usual data $(V, |0\rangle, T, Y(a, z))$ of a vertex algebra one is also given an $\text{End}V$ -valued power series $A(z) = \sum_{n=0}^{\infty} A_n z^n$, such that $A(z)|0\rangle = 0$. The vacuum and locality axioms are the same as that for a vertex algebra, but the translation covariance axiom is changed:

$$[T, Y(a, z)] = \partial_z Y(a, z) - Y(A(z)a, z).$$

(For convergence one needs that for each $k \in \mathbb{Z}$, $(A_j a)_{(k+j)}b = 0$ as $j \gg 0$.) The \hbar -deformed vertex algebra corresponds to the choice (cf. (2.6)):

$$A(z) = \hbar H / (1 + \hbar z).$$

Note that vertex operators of a gauged vertex algebra can be obtained from the vertex operators $X(a, z)$ of an ordinary vertex algebra by letting $Y(a, z) = X(B(z)a, z)$, where $B(z)$ is the unique formal power series solution of the equation $B'(z) = A(z)B(z)$, $B(0) = I$. One then has $B(z)|0\rangle = |0\rangle$.

2.2 (\hbar, Γ) -deformation of vertex operators and $Zhu_{\hbar, \Gamma}V$

Let Γ be an additive subgroup of \mathbb{R} , containing \mathbb{Z} . For $\gamma \in \mathbb{R}$ we shall denote the coset $\gamma + \mathbb{Z}$ by $\bar{\gamma}$.

Definition 2.10. A vertex algebra V with a Hamiltonian operator H is said to be Γ/\mathbb{Z} -graded if we have a decomposition

$$V = \bigoplus_{\bar{\gamma} \in \Gamma/\mathbb{Z}} \bar{\gamma}V \quad (2.25)$$

which is a vertex algebra grading, compatible with the Hamiltonian operator H , i.e.

$$\bar{\alpha}V_{(n)}\bar{\beta}V \subset \bar{\alpha} + \bar{\beta}V, \quad H(\bar{\gamma}V) \subset \bar{\gamma}V.$$

In particular, $|0\rangle \in \bar{0}V$. We shall assume for simplicity, that the eigenvalues of H are real.

Given a vector $a \in V$ of conformal weight Δ_a and degree $\bar{\gamma}_a$ (i.e. $a \in \bar{\gamma}_aV$), denote by ϵ_a the maximal non-positive real number in the coset $\bar{\gamma}_a - \bar{\Delta}_a$. This number has the following properties:

$$\epsilon_{|0\rangle} = 0, \quad \epsilon_{Ta} = \epsilon_a, \quad \epsilon_{a_{(n)}b} = \epsilon_a + \epsilon_b + \chi(a, b), \quad (2.26)$$

where $\chi(a, b) = 1$ or 0 , depending on whether $\epsilon_a + \epsilon_b \leq -1$ or not. For example, $\Delta_{|0\rangle} = \epsilon_{|0\rangle} = 0$, so that $\chi(a, |0\rangle) = \chi(|0\rangle, a) = 0$.

Let $\gamma_a = \Delta_a + \epsilon_a$. Then (1.46), (1.44) and (2.26) imply

$$\gamma_{|0\rangle} = 0, \quad \gamma_{Ta} = \gamma_a + 1, \quad \gamma_{a_{(n)}b} = \gamma_a + \gamma_b + \chi(a, b) - n - 1. \quad (2.27)$$

We denote by H' the operator on V , defined by $H'(a) = \gamma_a a$.

Example 2.11. If $\Gamma = \mathbb{Z}$, the decomposition (2.25) is trivial: $V = \bar{0}V$. In this case, for $a \in V$, γ_a is the largest integer less than or equal to Δ_a and $\epsilon_a = 0$ if and only if Δ_a is an integer.

Example 2.12. A case of special interest for the present paper is when the Γ/\mathbb{Z} -grading (2.25) is induced by H , namely

$$\bar{\gamma}V = \bigoplus_{\Delta \in \bar{\gamma}} V[\Delta], \quad (2.28)$$

where Γ is the additive subgroup of \mathbb{R} , generated by 1 and the eigenvalues of H . In this case $\epsilon_a = 0$, $\gamma_a = \Delta_a$ for all $a \in V$, and $\chi(a, b) = 0$ for all $a, b \in V$.

The (\hbar, Γ) -deformation of a vertex operator $Y(a, z)$ is defined as follows:

$$Y_{\hbar, \Gamma}(a, z) = (1 + \hbar z)^{\gamma_a} Y(a, z) = \sum_{n \in \mathbb{Z}} a_{(n, \hbar, \Gamma)} z^{n-1}.$$

We have:

$$a_{(n, \hbar, \Gamma)} b = \sum_{j \in \mathbb{Z}_+} \binom{\gamma_a}{j} \hbar^j a_{(n+j)} b. \quad (2.29)$$

The same proof as that of (2.7) gives

$$-n a_{(n-1, \hbar, \Gamma)} b = ((T + \hbar H')a)_{(n, \hbar, \Gamma)} b + \hbar(n+1) a_{(n, \hbar, \Gamma)} b. \quad (2.30)$$

Define the (\hbar, Γ) -deformed bracket in the same way as (2.3):

$$[a, b]_{\hbar, \Gamma} = \sum_{j \in \mathbb{Z}_+} (-\hbar)^j a_{(j, \hbar, \Gamma)} b,$$

and in the same way as (2.4), we show that

$$[a, b]_{\hbar, \Gamma} = \sum_{j \in \mathbb{Z}_+} \binom{\gamma_a - 1}{j} \hbar^j a_{(j)} b. \quad (2.31)$$

The same proof as that of Proposition 2.2, using (2.27) in place of (1.44), gives the (\hbar, Γ) -deformed n^{th} -product identity:

$$(1 + \hbar w)^{n+1-\chi(a, b)} Y_{\hbar, \Gamma}(a_{(n, \hbar, \Gamma)} b, w) = Y_{\hbar, \Gamma}(a, w)_{(n)} Y_{\hbar, \Gamma}(b, w). \quad (2.32)$$

The same proof as that of Theorem 2.3 gives now the (\hbar, Γ) -deformed Borchers identity:

$$\begin{aligned} & i_{z, w}(z - w)^n Y_{\hbar, \Gamma}(a, z) Y_{\hbar, \Gamma}(b, w) - p(a, b) i_{w, z}(z - w)^n Y_{\hbar, \Gamma}(b, w) Y_{\hbar, \Gamma}(a, z) \\ &= \sum_{j \in \mathbb{Z}_+} (1 + \hbar w)^{n+j+1-\chi(a, b)} Y_{\hbar, \Gamma}(a_{(n+j, \hbar, \Gamma)} b, w) \partial_w^j \delta(z - w) / j!. \end{aligned} \quad (2.33)$$

Denote by V_Γ (resp. $V_{\hbar, \Gamma}$) the \mathbb{C} -span (resp. $\mathbb{C}[\hbar]$ -span in $V[\hbar]$) of all elements $a \in V$ such that $\epsilon_a = 0$, and denote by $J_{\hbar, \Gamma}$ the $\mathbb{C}[\hbar]$ -span of all the elements $a_{(-2+\chi(a, b), \hbar, \Gamma)} b$, where $a, b \in V$ and $\epsilon_a + \epsilon_b \in \mathbb{Z}$. By (2.26), $V_{\hbar, \Gamma}$ is closed under the $(-1, \hbar, \Gamma)$ -product, which coincides on this subspace with the $(-1, \hbar)$ -product, and $J_{\hbar, \Gamma} \subset V_{\hbar, \Gamma}$.

For example, if $\Gamma = \mathbb{Z}$, then we have $V_\mathbb{Z} = \{a \in V \mid \Delta_a \in \mathbb{Z}\}$. On the other hand, if we consider the Γ/\mathbb{Z} -grading of V induced by H , as in Example 2.12, then $V_\Gamma = V$ and $J_{\hbar, \Gamma} = J_\hbar$. In this case the (n, \hbar, Γ) -product coincides with the (n, \hbar) -product (2.2).

The same proof as that of Theorem 2.7, gives the following

Theorem 2.13. (a) $J_{\hbar, \Gamma}$ contains all the elements $a_{(-j+\chi(a, b), \hbar, \Gamma)} b$, where $j \geq 2$ and $a, b \in V$, $\epsilon_a + \epsilon_b \in \mathbb{Z}$.

- (b) $(T + \hbar H)V_{\hbar, \Gamma} \subset J_{\hbar, \Gamma}$.
- (c) $J_{\hbar, \Gamma}$ is a two-sided ideal of $V_{\hbar, \Gamma}$ with respect to the $(-1, \hbar, \Gamma)$ -product.
- (d) $J_{\hbar, \Gamma}$ is a two-sided ideal of $V_{\hbar, \Gamma}$ with respect to the bracket $\hbar[a, b]_{\hbar, \Gamma}$.
- (e) $|0\rangle$ is the identity element for the $(-1, \hbar, \Gamma)$ -product.
- (f) $a_{(-1, \hbar, \Gamma)}b - p(a, b)b_{(-1, \hbar, \Gamma)}a \equiv (1 - \chi(a, b))\hbar[a, b]_{\hbar, \Gamma} \pmod{J_{\hbar, \Gamma}}$, $a, b \in V$.
- (g) The $(-1, \hbar, \Gamma)$ -product is associative on $V_{\hbar, \Gamma} \pmod{J_{\hbar, \Gamma}}$.

Define the following algebra over $\mathbb{C}[\hbar]$:

$$Zhu_{\hbar, \Gamma}V = V_{\hbar, \Gamma}/J_{\hbar, \Gamma}$$

with the algebra structure induced by the $(-1, \hbar, \Gamma)$ -product on $V_{\hbar, \Gamma}$. By Theorem 2.13, this product is well defined, and it is associative. Moreover, by the same theorem, the bracket $\hbar[a, b]_{\hbar, \Gamma}$ induces a bracket on $Zhu_{\hbar, \Gamma}V$, satisfying the relation

$$a_{(-1, \hbar, \Gamma)}b - p(a, b)b_{(-1, \hbar, \Gamma)}a = \hbar[a, b]_{\hbar, \Gamma} . \quad (2.34)$$

Furthermore, the image of $|0\rangle$ in $Zhu_{\hbar, \Gamma}V$, which we denote by 1, is the identity element for the associative product of $Zhu_{\hbar, \Gamma}V$. Also, by Theorem 2.13(b), we have:

$$T + \hbar H = 0 \text{ on } Zhu_{\hbar, \Gamma}V . \quad (2.35)$$

We can specialize $\hbar = c \in \mathbb{C}$, i.e. we let

$$Zhu_{\hbar=c, \Gamma}V = Zhu_{\hbar, \Gamma}V / (\hbar - c)Zhu_{\hbar, \Gamma}V = V_{\Gamma}/J_{\hbar=c, \Gamma} .$$

Note that the algebras $Zhu_{\hbar=c, \Gamma}V$ (over \mathbb{C}) are isomorphic for all $c \in \mathbb{C} \setminus \{0\}$. This can be seen by rescaling $\sigma_c(a) = c^{\Delta_a}a$ and the obvious formula

$$\sigma_c(a_{(n, \hbar=c, \Gamma)}b) = c^{-n-1}\sigma_c(a)_{(n, \hbar=1, \Gamma)}\sigma_c(b) , \quad (2.36)$$

so that σ_c induces an isomorphism $\sigma_c : Zhu_{\hbar=c, \Gamma}V \xrightarrow{\sim} Zhu_{\hbar=1, \Gamma}V$.

We define the (H, Γ) -twisted Zhu algebra by putting $\hbar = 1$:

$$Zhu_{\Gamma}V := Zhu_{\hbar=1, \Gamma}V .$$

(Of course, $Zhu_{\Gamma}V$ depends on H as well, as does $Zhu_{\hbar, \Gamma}$, but we omit H to simplify notation.) Special cases of $Zhu_{\Gamma}V$ appeared in [Z, KWa, DLM].

Example 2.14. The (non- Γ -twisted) Zhu algebra (considered in [Z, KWa] in the most important cases) is defined by putting $\hbar = 1$ and taking $\Gamma = \mathbb{Z}$, as in Example 2.11, i.e.

$$Zhu_{\mathbb{Z}}V = V_{\mathbb{Z}}/J_{\hbar=1, \mathbb{Z}} .$$

Example 2.15. Take the Γ/\mathbb{Z} -grading of V induced by H , as in Example 2.12. Then $V_\Gamma = V$ and the (n, \hbar, Γ) -product on V coincides with the (n, \hbar) -product (2.2). Let

$$a *_n b = a_{(n, \hbar=1)} b, \quad [a_* b] = [a, b]_{\hbar=1} \quad (2.37)$$

in order to simplify notation. Then we obtain the H -twisted Zhu algebra $Zhu_H V = V/J_{\hbar=1}$ introduced in Definition 2.8. Recall that its product is induced by $a *_n b$, and $J_{\hbar=1} = V *_n V$ is the span of all elements $a *_n b$, with $a, b \in V$. Note that, in view of (2.9), $J_{\hbar=1}$ is the two-sided ideal of the algebra $(V, *_n)$, generated by $(T + H)V$.

Later we shall need the following relation, which is obtained from (2.12) by specializing $\hbar = 1$ and using identity (2.22) with $y = -x = \Delta_a$:

$$a *_n b = \sum_{j=1}^n \binom{\Delta_a}{k-j} \left((T^{(j)} a) *_n b - \binom{-\Delta_a}{j} a *_n b \right). \quad (2.38)$$

Remark 2.16. A vertex algebra V with a Hamiltonian operator H is called a *chiralization* of the unital associative algebra $Zhu_H V$. An interesting open question is which unital associative algebras U admit a chiralization. For example, the affine vertex algebra $V^k(\mathfrak{g})$ is a chiralization of $U(\mathfrak{g})$ for any Lie superalgebra \mathfrak{g} . Theorem 5.10 of the present paper says that the affine W -algebra $W_k(\mathfrak{g}, x)$ is a chiralization of the finite W -algebra $W^{\text{fin}}(\mathfrak{g}, x)$. Note that any commutative U admits a chiralization $(V = U, |0\rangle = 1, Y(a, z) = L_a)$ with $H = 0$. Another very interesting example of a chiralization is the so called Kac–Todorov construction [KT] of the Neveu–Schwarz vertex algebra, which is a chiralization of the non-commutative Weil complex [AM], the corresponding coset construction of [KT] being the chiralization of the cubic Dirac operator of [Ko1]. We shall discuss this in more detail in Section 3.4.

The eigenspace decomposition of V with respect to H induces an increasing filtration of V with respect to the product $a_{(-1, \hbar=c, \Gamma)} b$ ($p \in \mathbb{R}$):

$$F^p V = \oplus_{\Delta \leq p} V[\Delta], \quad (F^p V)_{(-1, \hbar=c, \Gamma)} (F^q V) \subset F^{p+q} V, \quad (2.39)$$

which descends to a filtration of the algebra $Zhu_{\hbar=c, \Gamma} V$ by subspaces

$$F^p Zhu_{\hbar=c, \Gamma} V = F^p V_\Gamma / (F^p V_\Gamma \cap J_{\hbar=c, \Gamma}). \quad (2.40)$$

Proposition 2.17. (a) (cf. [Z]) The algebra $Zhu_{\hbar=0, \Gamma} V = V_\Gamma / V_\Gamma \cap (V_{(-2+\chi)} V)$ has a Poisson superalgebra structure, with commutative associative product induced by the -1^{st} product of V_Γ , and Lie bracket induced by the 0^{th} product of V_Γ . Here $V_{(-2+\chi)} V$ denotes the \mathbb{C} -span of all elements $a_{(-2+\chi(a, b))} b$ with $a, b \in V$. Moreover the action of H on V_Γ induces an algebra grading on $Zhu_{\hbar=0, \Gamma} V$.

(b) For every $c \in \mathbb{C}$, the algebra $gr Zhu_{\hbar, \Gamma} V$, associated graded to filtration (2.40), is a Poisson superalgebra, with associative commutative product

$$gr^{\Delta_1} Zhu_{\hbar=c, \Gamma} V \times gr^{\Delta_2} Zhu_{\hbar=c, \Gamma} V \longrightarrow gr^{\Delta_1 + \Delta_2} Zhu_{\hbar=c, \Gamma} V,$$

induced by the $(-1, \hbar = c, \Gamma)$ -product on V , and with Lie bracket

$$\text{gr}^{\Delta_1} Zhu_{\hbar=c, \Gamma} V \times \text{gr}^{\Delta_2} Zhu_{\hbar=c, \Gamma} V \longrightarrow \text{gr}^{\Delta_1 + \Delta_2 - 1} Zhu_{\hbar=c, \Gamma} V ,$$

induced by the $(\hbar = c, \Gamma)$ -deformed bracket (2.31) on V .

(c) There is a canonical surjective homomorphism of graded Poisson algebras

$$Zhu_{\hbar=0, \Gamma} V \twoheadrightarrow \text{gr} Zhu_{\hbar=c, \Gamma} V . \quad (2.41)$$

In particular, $\dim Zhu_{\hbar=c, \Gamma} V \leq \dim Zhu_{\hbar=0, \Gamma} V$.

Proof. Notice that, for $\hbar = 0$, the $(-1, \hbar, \Gamma)$ -product of V_Γ is equal to the -1^{st} product, and the (\hbar, Γ) -bracket on V_Γ is equal to the 0^{th} product, and it is easy to see from the fourth definition of a vertex algebra that they induce a Poisson algebra structure on $V_\Gamma / V_\Gamma \cap (V_{(-2+\chi)} V)$. To conclude the proof of (a) we just notice that $V_\Gamma \cap (V_{(-2+\chi)} V)$ is invariant under the action of H , hence the quotient space $V_\Gamma / V_\Gamma \cap (V_{(-2+\chi)} V)$ has an induced action of H . More explicitly, the eigenspace of H on $V_\Gamma \cap (V_{(-2+\chi)} V)$ with eigenvalue Δ is, by (1.44),

$$\begin{aligned} V_\Gamma \cap (V_{(-2+\chi)} V) &= \sum_{\substack{\Delta_1 + \Delta_2 + 1 = \Delta \\ \gamma_1, \gamma_2 | \epsilon_1 + \epsilon_2 = 0}} V^{\gamma_1} [\Delta_1]_{(-2)} V^{\gamma_2} [\Delta_2] \\ &+ \sum_{\substack{\Delta_1 + \Delta_2 = \Delta \\ \gamma_1, \gamma_2 | \epsilon_1 + \epsilon_2 = -1}} V^{\gamma_1} [\Delta_1]_{(-1)} V^{\gamma_2} [\Delta_2] . \end{aligned} \quad (2.42)$$

To prove (b), just notice that, by Theorem 2.13(f), $ab - ba \equiv c[a, b]_{\hbar=c, \Gamma} \pmod{J_{\hbar=c, \Gamma}}$, for $a, b \in V_\Gamma$, and that, by definition of the $(\hbar = c, \Gamma)$ -bracket, we have

$$[F^p V, F^q V]_{\hbar=c, \Gamma} \subset F^{p+q-1} V .$$

We are left to prove (c). By definition we have, for $c \in \mathbb{C}$,

$$\begin{aligned} \text{gr}^\Delta Zhu_{\hbar=c, \Gamma} V &= \frac{F^\Delta V_\Gamma / (F^\Delta V_\Gamma \cap J_{\hbar=c, \Gamma})}{F^{\Delta-} V_\Gamma / (F^{\Delta-} V_\Gamma \cap J_{\hbar=c, \Gamma})} \\ &\simeq \frac{F^\Delta V_\Gamma}{(F^{\Delta-} V_\Gamma) + (F^\Delta V_\Gamma \cap J_{\hbar=c, \Gamma})} , \end{aligned}$$

and in particular, for $\hbar = 0$,

$$\text{gr}^\Delta Zhu_{\hbar=0, \Gamma} V \simeq \frac{F^\Delta V_\Gamma}{(F^{\Delta-} V_\Gamma) + (F^\Delta V_\Gamma \cap (V_{(-2+\chi)} V))} ,$$

where $F^{\Delta-} V_\Gamma = \bigcup_{\Delta' < \Delta} F^{\Delta'} V_\Gamma$. On the other hand, by the definition of the (n, \hbar, Γ) -products and by (2.42), we have

$$(F^\Delta V_\Gamma) \cap (V_{(-2+\chi)} V) \subset (F^{\Delta-} V_\Gamma) + (F^\Delta V_\Gamma \cap J_{\hbar=c, \Gamma}) ,$$

which induces a quotient map $\text{gr}^\Delta Zhu_{\hbar=0, \Gamma} V \twoheadrightarrow \text{gr}^\Delta Zhu_{\hbar=c, \Gamma} V$. Statement (c) now follows by the fact that, by (a), $\text{gr} Zhu_{\hbar=0, \Gamma} V \simeq Zhu_{\hbar=0, \Gamma} V$. \square

Example 2.18. We will consider in Section 3.3 (see Proposition 3.30) a large class of vertex algebras (i.e. pregraded, freely generated by a free $\mathbb{C}[T]$ -module) for which the map (2.41) is actually an isomorphism. We give here an example of a vertex algebra V , with a Hamiltonian operator H , for which the linear map (2.41) fails to be an isomorphism. Let $V = \bigoplus_{n \in \mathbb{Z}_+} V[n]$ be a unital, associative, commutative, graded algebra, with $V[0] = \mathbb{C}1$. Then V is a vertex algebra with $|0\rangle = 1$, $T = 0$ and $a_{(n)}b = \delta_{n,-1}ab$. It has a Hamiltonian operator, acting as $nI_{V[n]}$ on the subspace $V[n]$. In this case $V_{(-2)}V = 0$, so that $Zhu_{\hbar=0,H}V = V/V_{(-2)}V \simeq V$. On the other hand, since $a_{(-2,\hbar)}|0\rangle = \hbar H(a)$, we have $J_{\hbar=1} = \bigoplus_{n \geq 1} V[n]$. Hence $Zhu_{\hbar=1,H}V = V/J_{\hbar=1} \simeq \mathbb{C}1$. Thus the map (2.41) has non-zero kernel, unless $V = V[0]$.

2.3 Γ -twisted positive energy modules and representations of the algebra $Zhu_{\Gamma}V$

Definition 2.19. A Γ -twisted module over a Γ/\mathbb{Z} -graded vertex algebra V is a vector superspace M and a parity preserving linear map from V to the space of $\text{End}M$ -valued Γ -twisted quantum fields $a \mapsto Y^M(a, z) = \sum_{m \in \bar{\gamma}_a} a_{(m)}^M z^{-m-1}$ (i.e. $a_{(m)}^M \in \text{End}M$ and $a_{(m)}^M v = 0$ for each $v \in M$ and $m \gg 0$), such that the following properties hold:

$$|0\rangle_{(n)}^M = \delta_{n,-1}I_M, \quad (\text{M1})$$

$$\begin{aligned} & \sum_{j \in \mathbb{Z}_+} \binom{m}{j} (a_{(n+j)}b)_{(m+k-j)}^M v \\ &= \sum_{j \in \mathbb{Z}_+} (-1)^j \binom{n}{j} (a_{(m+n-j)}^M b_{(k+j)}^M - p(a, b)(-1)^n b_{(k+n-j)}^M a_{(m+j)}^M) v, \end{aligned} \quad (\text{M2})$$

where $a \in \bar{\gamma}_a V$, $m \in \bar{\gamma}_a$, $n \in \mathbb{Z}$, $k \in \bar{\gamma}_b$.

Formula (M2) is called the Γ -twisted Borcherds identity. In the case when $\Gamma = \mathbb{Z}$, i.e. the grading (2.1) is trivial, M is called a (non-twisted) V -module. A special case of this is $M = V$ with $Y^M(a, z) = Y(a, z)$, when (M2) turns into the Borcherds identity (1.28). Note that an equivalent form of (M2) is

$$\begin{aligned} & i_{z,w}(z-w)^n Y^M(a, z) Y^M(b, w) - p(a, b) i_{w,z}(z-w)^n Y^M(b, w) Y^M(a, z) \\ &= \sum_{j \in \mathbb{Z}_+} Y^M(a_{(n+j)}b, w) \partial_w^j \delta_{\bar{\gamma}_a}(z-w)/j!, \end{aligned} \quad (2.43)$$

where $\delta_{\bar{\gamma}}(z-w) = z^{-1} \sum_{n \in \bar{\gamma}} (\frac{w}{z})^n$ for a \mathbb{Z} -coset $\bar{\gamma}$.

Remark 2.20. For Γ -twisted quantum fields $a(z) = \sum_{m \in \bar{\gamma}_a} a_{(m)} z^{-m-1}$, the usual notion (1.6) of n^{th} product of fields is not quite right, since, for example, it is identically zero if $\gamma_a \notin \mathbb{Z}$. So we introduce the Γ -deformed n^{th} product of Γ -twisted quantum fields ($n \in \mathbb{Z}$):

$$a(w)_{(n,\Gamma)} b(w) = \text{Res}_z (z/w)^{\gamma_a} \left(a(z) b(w) i_{z,w} - p(a, b) b(w) a(z) i_{w,z} \right) (z-w)^n. \quad (2.44)$$

Notice that the above formula depends on the choice of the representative γ_a in the coset $\bar{\gamma}_a = \gamma_a + \mathbb{Z}$. Moreover (2.44) reduces to (1.6) if $\gamma_a = 0$. If we multiply both sides of (2.43) by $(z/w)^{\gamma_a}$ and take residues in z , we get

$$Y^M(a_{(n, \hbar=\frac{1}{w}, \Gamma)} b, w) = Y^M(a, w)_{(n, \Gamma)} Y^M(b, w). \quad (2.45)$$

To get (2.45) we used the definition (2.29) of (\hbar, Γ) -deformed n^{th} products in V , and the definition (2.44) of Γ -deformed n^{th} product of Γ -twisted $\text{End} M$ -valued quantum fields. Equation (2.45) is a generalization of the n^{th} product identity (1.10) to arbitrary Γ -twisted V -modules M , and it reduces to it for $M = V$, the adjoint representation (or, more generally, for a non-twisted V -module M). It has the following interpretation: the space of Γ -twisted $\text{End} M$ -valued quantum fields $\mathcal{F}^M = \{a(x) = Y^M(a, x) \mid a \in V\}$ together with the Γ -deformed n^{th} product (2.44), form an " $(\hbar = \frac{1}{x}, \Gamma)$ -deformed vertex algebra", namely the (\hbar, Γ) -deformed Borcherd's identity (2.33) holds for $\hbar = \frac{1}{x}$ and for any two elements $a(x), b(x) \in \mathcal{F}^M$. This is the first indication that (\hbar, Γ) -deformed vertex operators are related to Γ -twisted representations of the vertex algebra V (as it will become clear from Theorem 2.30).

From now on we fix a Hamiltonian operator H on V with the eigenspace decomposition (1.42). As before, we shall assume that the eigenvalues of H are real. If $a \in V[\Delta_a]$, one often writes (cf. Proposition 1.15) $Y^M(a, z) = \sum_{n \in \bar{\epsilon}_a} a_n^M z^{-n-\Delta_a}$, so that

$$a_{(n)}^M = a_{n-\Delta_a+1}^M, n \in \bar{\gamma}_a, \quad a_n^M = a_{(n+\Delta_a-1)}^M, n \in \bar{\epsilon}_a. \quad (2.46)$$

After making the change of indices (2.46), (M1) becomes $|0\rangle_n = \delta_{n,0}$, and the twisted Borcherd's identity (M2) becomes \mathbb{R} -graded:

$$\begin{aligned} & \sum_{j \in \mathbb{Z}_+} \binom{m + \Delta_a - 1}{j} (a_{(n+j)} b)_{m+k}^M \\ &= \sum_{j \in \mathbb{Z}_+} (-1)^j \binom{n}{j} (a_{m+n-j}^M b_{k+j-n}^M - p(a, b) (-1)^n b_{k-j}^M a_{m+j}^M), \end{aligned} \quad (2.47)$$

where $m \in \bar{\epsilon}_a, k \in \bar{\epsilon}_b, n \in \mathbb{Z}$. Letting in this identity $b = |0\rangle, n = -2$ and $k = 0$, we obtain:

$$(Ta)_m^M = -(m + \Delta_a) a_m^M. \quad (2.48)$$

Letting $n = 0$ in (2.47), we obtain the commutator formula:

$$[a_m^M, b_k^M] = \sum_{j \in \mathbb{Z}_+} \binom{m + \Delta_a - 1}{j} (a_{(j)} b)_{m+k}^M. \quad (2.49)$$

Definition 2.21. The Γ -twisted V -module M is called a *positive energy* (Γ -twisted) V -module if M has an \mathbb{R} -grading $M = \oplus_{j \geq 0} M_j$ such that

$$a_n^M M_j \subset M_{j-n}. \quad (2.50)$$

The subspace M_0 is called the *minimal energy subspace*. Then, of course,

$$a_n^M M_0 = 0 \text{ for } n > 0 \text{ and } a_0^M M_0 \subset M_0. \quad (2.51)$$

Letting $m = \epsilon_a + 1$ and $k = \epsilon_b - 1 + \chi(a, b)$ in (2.47), we obtain:

$$\begin{aligned} \sum_{j \in \mathbb{Z}_+} \binom{\gamma_a}{j} (a_{(n+j)} b)_{\epsilon_a + \epsilon_b + \chi(a, b)}^M v &= \sum_{j \in \mathbb{Z}_+} (-1)^j \binom{n}{j} \\ &\times (a_{n-j+1+\epsilon_a}^M b_{j-n-1+\chi(a, b)+\epsilon_b}^M - p(a, b) (-1)^n b_{\epsilon_b + \chi(a, b) - j - 1}^M a_{j+1+\epsilon_a}^M) v. \end{aligned} \quad (2.52)$$

Recall from Section 2.2 the definition of the (H, Γ) -twisted Zhu algebra $Zhu_\Gamma V$. We are going to establish a correspondence between positive energy Γ -twisted V -modules and $Zhu_\Gamma V$ -modules, generalizing the construction of [Z, KWa, DLM]. For this we need a few lemmas.

Lemma 2.22. *Let M be a Γ -twisted positive energy V -module. Then the map*

$$a \mapsto a_0^M|_{M_0}, \quad a \in V_\Gamma, \quad (2.53)$$

defines a linear map $\varphi : V_\Gamma \mapsto \text{End} M_0$ which is zero on $J_{\hbar=1}$ and induces a homomorphism $Zhu_\Gamma V \rightarrow \text{End} M_0$.

Proof. Let $a, b \in V$ be such that $\epsilon_a + \epsilon_b \in \mathbb{Z}$, i.e. $\epsilon_a + \epsilon_b = 0$ or -1 , hence $\epsilon_a + \epsilon_b + \chi(a, b) = 0$. If $n \leq -1$, and $v \in M_0$, then (2.52) becomes, due to (2.29) and (2.51),

$$(a_{(n, \hbar=1, \Gamma)} b)_0^M v = \sum_{j \in \mathbb{Z}_+} \binom{\gamma_a}{j} (a_{(n+j)} b)_0^M v = a_{n+1+\epsilon_a}^M b_{-n-1+\chi(a, b)+\epsilon_b}^M v.$$

The RHS of this equality is zero if $n \leq -2$, or if $n = -1$ and $\chi(a, b) = 1$. Thus, $\varphi(J_{\hbar=1}) = 0$. Next, if $a, b \in V_\Gamma$, i.e. $\epsilon_a = \epsilon_b = 0$, and $n = -1$, this formula becomes, due to (2.51):

$$(a_{(-1, \hbar=1, \Gamma)} b)_0^M v = a_0^M b_0^M v, \quad v \in M_0.$$

The lemma is proved. \square

The above lemma defines a functor from the category $\text{Mod}_V^{\Gamma, +}$ of Γ -twisted positive energy V -modules, to the category $\text{Mod}_{Zhu_\Gamma V}$ of modules over $Zhu_\Gamma V$:

$$\text{Mod}_V^{\Gamma, +} \rightarrow \text{Mod}_{Zhu_\Gamma V}, \quad M \mapsto M_0.$$

We next want to define a functor in the opposite direction. For this we will need the following "Uniqueness Lemma".

Lemma 2.23. *Let M be a vector superspace and consider a family $\mathcal{F} = \{a^j(z) = \sum_{n \in \gamma_j + \mathbb{Z}} a_{(n)}^j z^{-n-1}\}_{j \in J}$ of Γ -twisted $\text{End} M$ -valued quantum fields, i.e. $a_{(n)}^j \in \text{End} M$ and for each $j \in J$, $v \in M$, $a_{(n)}^j v = 0$ for $n \gg 0$. Assume that all pairs $\{a^i(z), a^j(z)\}$ are local. Let M_0 be a generating subspace of M , i.e. the span of all vectors $a_{(n_1)}^{j_1} \dots a_{(n_s)}^{j_s} M_0$ is M . Let $b(z)$ be another Γ -twisted $\text{End} M$ -valued field such that all pairs $(b(z), a^j(z))$ are local and $b(z)M_0 = 0$. Then $b(z) = 0$.*

Proof. Since, by assumption, M_0 generates the whole space M , we just have to prove that

$$b(z)a^{j_1}(w_1)\dots a^{j_s}(w_s)v = 0, \quad (2.54)$$

for any $j_1, \dots, j_s \in J$ and $v \in M_0$. On the other hand, by locality of the pairs $(b(z), a^{j_k}(z))$, we have for some $N_i \in \mathbb{Z}_+$,

$$\prod_{i=1}^s (z - w_i)^{N_i} b(z)a^{j_1}(w_1)\dots a^{j_s}(w_s)v = 0, \quad (2.55)$$

since we can move $b(z)$ to the right, and $b(z)v = 0$ by assumption. But the LHS of the above expression is a Laurent series in w_s , i.e. the exponents in w_s that occur are bounded below. Hence we can cancel $(z - w_s)^{N_s}$ in (2.55). Multiplying after that both sides of (2.55) by $(w_{s-1} - w_s)^N$ for some $N \in \mathbb{Z}_+$, we can permute the factors $a^{j_{s-1}}(w_{s-1})$ and $a^{j_s}(w_s)$ to get

$$(w_{s-1} - w_s)^N \prod_{i=1}^{s-1} (z - w_i)^{N_i} b(z)a^{j_1}(w_1)\dots a^{j_{s-2}}(w_{s-2})a^{j_s}(w_s)a^{j_{s-1}}(w_{s-1})v = 0.$$

Since the LHS is a Laurent series in w_{s-1} , we can cancel the product $(w_{s-1} - w_s)^N (z - w_{s-1})^{N_{s-1}}$. After finitely many such steps we obtain (2.54) with the $a^{j_i}(w_i)$'s in reversed order. \square

Recall that any vertex algebra V is, in particular, a Lie conformal superalgebra, and to the latter one canonically associates the (maximal formal distribution) Lie superalgebra $L(V)$ [K]. Given a Hamiltonian operator H on V and a Γ/\mathbb{Z} -grading (2.25), one likewise associates to V an \mathbb{R} -graded Lie superalgebra $L_{H,\Gamma}(V)$ as follows.

Definition 2.24. The (H, Γ) -twisted formal distribution Lie superalgebra $L_{H,\Gamma}(V)$ is, as a vector superspace, the quotient space of the vector superspace with the basis $\{a_n \mid a \in {}^{\bar{\gamma}}V[\Delta], n \in \bar{\epsilon}_a\}$, by the span of the vectors $(\lambda a + \mu b)_n - \lambda a_n - \mu b_n$ and (cf. (2.48)):

$$(Ta)_n + (n + \Delta_a)a_n \quad (2.56)$$

where $\lambda, \mu \in \mathbb{C}$, $a, b \in V$, $n \in \bar{\epsilon}_a$. The Lie bracket on $L_{H,\Gamma}(V)$ is defined by (cf. (2.49))

$$[a_m, b_k] = \sum_{j \in \mathbb{Z}_+} \binom{m + \Delta_a - 1}{j} (a_{(j)}b)_{m+k}, \quad (2.57)$$

where $m \in \bar{\epsilon}_a, k \in \bar{\epsilon}_b$. It is easy to see that (2.57) defines a structure of a Lie superalgebra on $L_{H,\Gamma}(V)$. Moreover $L_{H,\Gamma}(V)$ is \mathbb{R} -graded by $\deg a_m = m \in \mathbb{R}$:

$$L_{H,\Gamma}(V) = \bigoplus_{j \in \mathbb{R}} L_j. \quad (2.58)$$

The corresponding family \mathcal{F} of Γ -twisted formal distributions consists of $(a \in V)$:

$$Y_{\Gamma}(a, z) = \sum_{m \in \bar{\epsilon}_a} a_m z^{-m - \Delta_a} \in L_{H,\Gamma}(V)[[z^{\pm 1}]]z^{\gamma_a}. \quad (2.59)$$

Let U be the universal enveloping superalgebra of $L_{H,\Gamma}(V)$, and let $U = \oplus_{j \in \mathbb{R}} U_j$ be the grading induced by (2.58). Consider the following fundamental system of neighborhoods of 0 in $U : \{UF_N\}_{N \geq 0}$, where $F_N = \oplus_{j \geq N} L_j$. Denote by U^{com} the completion of the algebra U in this topology and by \bar{U}_j the closure of U_j in U^{com} . Then $\bar{U} := \oplus_{j \in \mathbb{R}} \bar{U}_j$ is a subalgebra of the algebra U^{com} .

Given $a, b \in V$ (homogeneous in H - and Γ/\mathbb{Z} -grading), $m \in \bar{\epsilon}_a$, $k \in \bar{\epsilon}_b$ and $n \in \mathbb{Z}$, introduce the following element of \bar{U}_{m+k} (cf. (2.47)):

$$\begin{aligned} \text{BI}(a, b; m, k, n) &= \sum_{j \in \mathbb{Z}_+} \binom{m + \Delta_a - 1}{j} (a_{(n+j)} b)_{m+k} \\ &- \sum_{j \in \mathbb{Z}_+} (-1)^j \binom{n}{j} (a_{m+n-j} b_{k+j-n} - p(a, b) (-1)^n b_{k-j} a_{m+j}). \end{aligned} \quad (2.60)$$

Let \mathcal{A} denote the factor algebra of the associative superalgebra \bar{U} by the 2-sided ideal generated by all elements $\text{BI}(a, b; m, k, n)$, and the elements $|0\rangle_n - \delta_{n,0} 1$ ($n \in \mathbb{Z}$).

Remark 2.25. Since relations (2.47) imply the relations (2.48) and (2.49), it follows from the definition, that a Γ -twisted V -module M is the same as a module over the associative superalgebra \mathcal{A} , such that $a_n v = 0$ for any given $a \in V$ and $v \in M$ for $n \gg 0$.

Lemma 2.26. (a) Given $a, b, c \in V$, $m \in \bar{\epsilon}_a, k \in \bar{\epsilon}_b, s \in \bar{\epsilon}_c, n \in \mathbb{Z}$, we have

$$\begin{aligned} [c_s, \text{BI}(a, b; m, k, n)] &= \sum_{j \in \mathbb{Z}_+} \binom{s + \Delta_c - 1}{j} (\text{BI}(c_{(j)} a, b; m + s, k, n) \\ &+ p(a, c) \text{BI}(a, c_{(j)} b, m, k + s, n)). \end{aligned} \quad (2.61)$$

(b) Let B denote the (graded) subspace of the algebra \bar{U} , spanned by all the elements $\text{BI}(a, b; m, k, n)$, with $a, b \in V, m \in \bar{\epsilon}_a, k \in \bar{\epsilon}_b, n \in \mathbb{Z}$. Then $[c_s, B] \subset B$, for any $c \in V$ and $s \in \bar{\epsilon}_c$.

Proof. We denote by $\text{LBI}(a, b; m, k, n)$ and $\text{RBI}(a, b; m, k, n)$ respectively the first and the second sum in the RHS of (2.60). By the definition (2.57) of the Lie bracket in $L_{H,\Gamma}(V)$, we have

$$\begin{aligned} [c_s, \text{LBI}(a, b; m, k, n)] &= \sum_{j \in \mathbb{Z}_+} \binom{m + \Delta_a - 1}{j} [c_s, (a_{(n+j)} b)_{m+k}] \\ &= \sum_{i, j \in \mathbb{Z}_+} \binom{m + \Delta_a - 1}{j} \binom{s + \Delta_c - 1}{i} (c_{(i)} (a_{(n+j)} b))_{m+k+s}. \end{aligned} \quad (2.62)$$

By the commutator formula for the vertex algebra V we have

$$c_{(i)} (a_{(n+j)} b) = \sum_{l \in \mathbb{Z}_+} \binom{i}{l} (c_{(l)} a)_{(n+i+j-l)} b + p(a, c) a_{(n+j)} (c_{(i)} b). \quad (2.63)$$

Now we use the following combinatorial identity, which is a generalization of (2.22) ($l, r \in \mathbb{Z}_+$):

$$\sum_{j=0}^r \binom{x}{j} \binom{y}{r+l-j} \binom{r+l-j}{l} = \binom{y}{l} \binom{x+y-l}{r} . \quad (2.64)$$

By (2.63), and (2.64) with $x = m + \Delta_a - 1$, $y = s + \Delta_c - 1$, and using the change of the summation variable $r = i + j - l$, we can rewrite the RHS of (2.62) as

$$\begin{aligned} & \sum_{l, r \in \mathbb{Z}_+} \binom{s + \Delta_c - 1}{l} \binom{m + s + \Delta_a + \Delta_c - l - 2}{r} ((c_{(l)} a)_{(n+r)} b)_{m+k+s} \\ & + \sum_{i, j \in \mathbb{Z}_+} \binom{m + \Delta_a - 1}{j} \binom{s + \Delta_c - 1}{i} p(a, c) (a_{(n+j)} (c_{(i)} b))_{m+k+s} . \end{aligned}$$

By the definition of $LBI(a, b; m, k, n)$ we thus get

$$\begin{aligned} [c_s, LBI(a, b; m, k, n)] &= \sum_{j \in \mathbb{Z}_+} \binom{s + \Delta_c - 1}{j} (LBI(c_{(j)} a, b; m + s, k, n) \\ &+ p(a, c) LBI(a, c_{(j)} b, m, k + s, n)) . \end{aligned}$$

An easier computation shows that the same formula holds with RBI instead of LBI, thus proving (2.61). The second part of the lemma is now obvious. \square

For $n \in \mathbb{Z}$ we denote by $BI(a, b; z, w; n)$ the generating series of all elements (2.60), namely (see (2.59))

$$\begin{aligned} BI(a, b; z, w; n) &= \sum_{m \in \bar{\epsilon}_a, k \in \bar{\epsilon}_b} z^{-m - \Delta_a} w^{-k - \Delta_b + n} BI(a, b; m, k, n) \\ &= \sum_{j \in \mathbb{Z}_+} Y_\Gamma(a_{(n+j)} b, w) \partial_w^j \delta_{\bar{\gamma}_a}(z - w) / j! \\ &\quad - \left(Y_\Gamma(a, z) Y_\Gamma(b, w) i_{z, w} - p(a, b) Y_\Gamma(b, w) Y_\Gamma(a, z) i_{w, z} \right) (z - w)^n . \end{aligned} \quad (2.65)$$

The second equality follows by a straightforward computation. Notice that, since (2.49) is equivalent to (2.43) with $n = 0$, the Γ -twisted fields $Y_\Gamma(a, z)$, $a \in V$, are pairwise local formal distributions, i.e. one has $(z - w)^N [Y_\Gamma(a, z), Y_\Gamma(b, w)] = 0$ for an integer $N \gg 0$. By comparing the coefficients of $z^{-m - \Delta_a}$ in both sides of equation (2.65) we get, by a straightforward computation,

$$\begin{aligned} \sum_{k \in \bar{\epsilon}_b} w^{-k - \Delta_b + n} BI(a, b; m, k, n) &= \text{Res}_{z=w} Y_\Gamma(Y(a, z - w) b, w) i_{w, z-w} F(z, w) \\ &\quad - \text{Res}_z \left(Y_\Gamma(a, z) Y_\Gamma(b, w) i_{z, w} - p(a, b) Y_\Gamma(b, w) Y_\Gamma(a, z) i_{w, z} \right) F(z, w) , \end{aligned} \quad (2.66)$$

where $F(z, w) = z^{m + \Delta_a - 1} (z - w)^n$, and as before, $i_{z, w}$ denotes expansion in the domain $|z| > |w|$. This formula is analogous to yet another form of Borchers

identity, see [FHL], [K, Sec4.8]. Note that if we replace $F(z, w)$ by $w^l F(z, w)$, the RHS of (2.66) changes by an overall factor w^l . Hence the subspace B of \bar{U} is spanned by the coefficients of all powers of w in the RHS of (2.66), where $F(z, w)$ is $z^{\gamma_a} w^r$, $r \in \mathbb{R}$, times an arbitrary rational function in z and w with poles at $z = 0, w = 0, z = w$.

For every $a \in V$, homogeneous with respect to H - and Γ/\mathbb{Z} -grading, fix an element $m_a \in \bar{\epsilon}_a$ (later we will choose $m_a = \epsilon_a + 1$). We define the *normally ordered product* of Γ -twisted quantum fields as follows (see 2.44):

$$\begin{aligned} : Y_\Gamma(a, z) Y_\Gamma(b, z) : &= \text{Res}_z (z/w)^{m_a + \Delta_a - 1} \left(Y_\Gamma(a, z) Y_\Gamma(b, w) i_{z, w} \right. \\ &\quad \left. - p(a, b) Y_\Gamma(b, w) Y_\Gamma(a, z) i_{w, z} \right) (z - w)^{-1}. \end{aligned} \quad (2.67)$$

By computing the residue in z , we can rewrite the above definition in the more familiar form

$$: Y_\Gamma(a, z) Y_\Gamma(b, z) : = Y_\Gamma(a, z)_+ Y_\Gamma(b, z) + p(a, b) Y_\Gamma(b, z) Y_\Gamma(a, z)_-,$$

where

$$Y_\Gamma(a, z)_+ = \sum_{m \in \bar{\epsilon}_a, m < m_a} a_m z^{-m - \Delta_a}, \quad Y_\Gamma(a, z)_- = \sum_{m \in \bar{\epsilon}_a, m \geq m_a} a_m z^{-m - \Delta_a}.$$

(Notice that the above definition depends on the choice of $m_a \in \bar{\epsilon}_a$.) If we put $n = -1$ and $m = m_a$ in (2.66), we then get, after computing the residue in $z - w$,

$$\begin{aligned} \text{BI}_1(a, b; w) &:= \sum_{k \in \bar{\epsilon}_b} w^{-k - \Delta_b - m_a - \Delta_a} \text{BI}(a, b; m_a, k, -1) \\ &= \sum_{j \in \mathbb{Z}_+} \binom{m_a + \Delta_a - 1}{j} Y_\Gamma(a_{(j-1)} b, w) w^{-j} - : Y_\Gamma(a, w) Y_\Gamma(b, w) :. \end{aligned} \quad (2.68)$$

Lemma 2.27. *The subspace $B \subset \bar{U}$ is spanned by all elements of the form $\text{BI}(a, b; m_a, k, -1)$, with $a, b \in V$ and $k \in \bar{\epsilon}_b$.*

Proof. By the above observations B is spanned by the coefficients of all powers of w in the RHS of (2.66), with $F(z, w) = z^\alpha w^\beta (z - w)^\gamma$, $\alpha \in \bar{\epsilon}_a, \beta, \gamma \in \mathbb{Z}$. Notice that every such function $F(z, w)$ can be written, by partial fraction decomposition, as finite linear combination over $\mathbb{C}[w^{\pm 1}]$ of monomials z^p , with $p \in \bar{\gamma}_a$, and $z^{m_a + \Delta_a - 1} (z - w)^q$, with integer $q \leq -1$. But the RHS of (2.66) with $F(z, w) = z^p$ is identically zero in \bar{U} . This follows by the commutator formula (2.57) for elements of $L_{H, \Gamma}(V) \subset \bar{U}$. Moreover it is easy to show using (2.56) that the RHS of (2.66) with $F(z, w) = z^{m_a + \Delta_a - 1} (z - w)^{-n-1}$, $n \geq 0$, is a finite linear combination over $\mathbb{C}[w^{\pm 1}]$ of expressions as in the RHS of (2.66) with a replaced by $a, Ta, \dots, T^n a$, and $F(z, w) = z^{m_a + \Delta_a - 1} (z - w)^{-1}$. It immediately follows by the above observations that the subspace $B \subset \bar{U}$ is spanned by the coefficients of all powers of w in the RHS of (2.66) with $F(z, w) = z^{m_a + \Delta_a - 1} (z - w)^{-1}$. The claim follows immediately. \square

Consider the subalgebras L_0 , $L_{\geq} = \oplus_{j \geq 0} L_j$, $L_{>} = \oplus_{j > 0} L_j$ and $L_{\leq} = \oplus_{j \leq 0} L_j$ of the Lie superalgebra $L_{H,\Gamma}(V)$. Note that,

$$L_0 = V_{\Gamma}/(T + H)V_{\Gamma},$$

with the Lie bracket (cf. (2.57))

$$[a_0, b_0] = \sum_{j \in \mathbb{Z}_+} \binom{\Delta_a - 1}{j} (a_{(j)} b)_0.$$

Due to Theorem 2.13(c) and (f), and formula (2.4), we have a canonical surjective Lie superalgebra homomorphism

$$\psi : L_0 \rightarrow Zhu_{\Gamma}V, \quad (2.69)$$

induced by the map $a_0 \mapsto a$. Here the associative superalgebra $Zhu_{\Gamma}V$ is viewed as a Lie superalgebra with the usual Lie superbracket.

Now let M_0 be a module over the (associative) superalgebra $Zhu_{\Gamma}V$. Then, in particular, it is a module over $Zhu_{\Gamma}V$ viewed as a Lie superalgebra, hence, using the homomorphism ψ defined by (2.69), M_0 is an L_0 -module. Extend it to a L_{\geq} -module by letting $L_j M_0 = 0$ for $j > 0$ and consider the induced module $\tilde{V}(M_0) := \text{Ind}_{L_{\geq 0}}^L M_0$. The reversed grading of L ($j \mapsto -j$) induces on $\tilde{V}(M_0)$ an \mathbb{R}_+ -grading, which makes it a "positive energy" $L_{H,\Gamma}(V)$ -module, in analogy with Definition 2.21:

$$\tilde{V}(M_0) = \oplus_{j \geq 0} \tilde{M}_j, \quad \tilde{M}_0 = M_0, \quad L_i M_j \subset M_{j-i}.$$

Denote by $V(M_0)$ the quotient of the $L_{H,\Gamma}(V)$ -module $\tilde{V}(M_0)$ by the maximal graded submodule intersecting M_0 trivially. It is again a positive energy $L_{H,\Gamma}(V)$ -module:

$$V(M_0) = \oplus_{j \geq 0} M_j.$$

Of course, $\tilde{V}(M_0)$ and $V(M_0)$ are U -modules and, by the "positive energy" condition, they are also \bar{U} -modules.

Lemma 2.28. *Let M_0 be a module over the associative algebra $Zhu_{\Gamma}V$. Then $V(M_0)$ is naturally a Γ -twisted positive energy module over the vertex algebra V .*

Proof. By the definition of a Γ -twisted positive energy module over V , it suffices to prove that

$$B V(M_0) = 0, \quad (2.70)$$

where $B \subset \bar{U}$ is the subspace spanned by the elements $\text{BI}(a, b; m, k, n)$ in (2.60). For that consider the left submodule $M' = \bar{U} B M_0$ of the \bar{U} -module $\tilde{V}(M_0)$. It is obviously a graded submodule: $M' = \oplus_{j \geq 0} M'_j$. We want to show that M' intersects M_0 trivially, so that it maps to zero in $V(M_0)$. Due to the PBW theorem, M' is the span of elements of the form

$$a_{-m_1}^1 \dots a_{-m_s}^s b_{n_1}^1 \dots b_{n_t}^t B(M_0),$$

where $a_{-m_i}^i \in L_{\leq}$, i.e. $m_i \geq 0$, and $b_{n_i}^i \in L_{>}$, i.e. $n_i > 0$. Since by Lemma 2.26, $[b_i, B] \subset B$, and since $b_n M_0 = 0$ for $n > 0$, we conclude that M' is the span of elements of the form

$$a_{-m_1}^1 \dots a_{-m_s}^s B(M_0), \text{ where } a_{-m_i}^i \in L_{\leq}. \quad (2.71)$$

Obviously $B = \bigoplus_{j \in \mathbb{R}} B_j$ is a graded subspace of U and $B_j M_0 = 0$ for $j > 0$. Hence the zero degree component of M' is

$$M' \cap M_0 = U(L_0) B_0(M_0).$$

We need to show that this is zero, namely that $B_0(M_0) = 0$. By Lemma 2.27, B_0 is spanned by the elements $\text{BI}(a, b; \epsilon_a + 1, -\epsilon_a - 1, -1)$, with $a, b \in V$ such that $\epsilon_a + \epsilon_b \in \mathbb{Z}$. Hence $B_0(M_0)$ is spanned by (cf. (2.29))

$$\begin{aligned} \text{BI}(a, b; \epsilon_a + 1, -\epsilon_a - 1, -1)(M_0) &= ((a_{(-1, \hbar=1, \Gamma)} b)_0 \\ &\quad - \sum_{j \in \mathbb{Z}_+} (a_{\epsilon_a - j} b_{-\epsilon_a + j} + p(a, b) b_{-\epsilon_a - 1 - j} a_{\epsilon_a + j + 1})) M_0. \end{aligned} \quad (2.72)$$

But if $\epsilon_a \neq 0$, then $a_{(-1, \hbar=1, \Gamma)} b \in J_{\hbar=1}$, hence $(a_{(-1, \hbar=1, \Gamma)} b)_0 M_0 = 0$, since M_0 is by assumption a $Zhu_{\Gamma} V$ -module. Moreover, if $\epsilon_a \neq 0$ all summands of the sum in the RHS of (2.72) annihilate M_0 since $-\epsilon_a > 0$. Finally, if $\epsilon_a = 0$ then $a, b \in V_{\Gamma}$ and the RHS of (2.72) becomes $((a_{(-1, \hbar=1, \Gamma)} b)_0 - a_0 b_0) M_0$, which is again zero since, by assumption, M_0 is a $Zhu_{\Gamma} V$ -module.

So far we proved that M' is a submodule of $\tilde{V}(M_0)$ intersecting M_0 trivially, hence in $V(M_0)$ we have $BM_0 = 0$, or equivalently $\text{BI}_1(a, b; z) M_0 = 0$, where $\text{BI}_1(a, b; z)$ was defined in (2.68). We remarked above that all $\text{End} V(M_0)$ -valued fields $Y_{\Gamma}(a, z) = \sum_{n \in \bar{\epsilon}_a} a_n z^{-n - \Delta_a}$ are pairwise local. Moreover the same argument used to prove Dong's Lemma (see e.g. [K, Lemma 3.2]), can be used to show that the Γ -twisted fields $Y_{\Gamma}(a, z)$ are local with the normally ordered products $: Y_{\Gamma}(b, z) Y_{\Gamma}(c, z) :$ defined in (2.67), and hence with the formal distributions $\text{BI}_1(a, b; z)$ in (2.68). Then by Lemma 2.23, $\text{BI}_1(a, b; z) V(M_0) = 0$. But by Lemma 2.27, this implies (2.70). \square

Lemma 2.22 defines a "restriction" functor $M \mapsto M_0$ from the category $\text{Mod}_V^{\Gamma, +}$ of Γ -twisted positive energy V -modules to the category $\text{Mod}_{Zhu_{\Gamma} V}$ of modules over $Zhu_{\Gamma} V$. Lemma 2.28 defines an "induction" functor in the opposite direction, from the category $\text{Mod}_{Zhu_{\Gamma} V}$ to the category $\text{Mod}_V^{\Gamma, +}$, given by $M_0 \mapsto V(M_0)$. We next want to understand in what sense the above functors are inverse of each other, thus establishing an equivalence of categories.

Definition 2.29. We shall call an \mathbb{R}_+ -graded module $M = \bigoplus_{j \geq 0} M_j$ over an \mathbb{R} -graded Lie superalgebra L or over a vertex algebra V , an *almost irreducible module* if M_0 generates the module M and M contains no graded submodules intersecting M_0 trivially. For example, the $L_{H, \Gamma}(V)$ -module $V(M_0)$ is almost irreducible.

It is clear that the restriction functor $M \mapsto M_0$ from the category of almost irreducible $L_{H,\Gamma}(V)$ -modules to the category of L_0 -modules and the induction functor $M_0 \mapsto V(M_0)$ are inverse to each other. Moreover, these functors establish a bijective correspondence between irreducibles of these categories, since clearly an $L_{H,\Gamma}(V)$ -module $V(M_0)$ is irreducible if and only if the L_0 -module M_0 is irreducible.

We can now state the main theorem of this section, which is an immediate consequence of the above results and Definition 2.29.

Theorem 2.30. *The restriction functor $M \mapsto M_0$ is a functor from the category of positive energy Γ -twisted V -modules to the category of $Zhu_\Gamma V$ -modules, which is inverse to the induction functor $M_0 \mapsto V(M_0)$, on the full subcategory of almost irreducible Γ -twisted V -modules. In particular, these functors establish a bijective correspondence between irreducible positive energy Γ -twisted V -modules and irreducible $Zhu_\Gamma V$ -modules.*

3 Non-linear Lie algebras and non-linear Lie conformal algebras

We review here the notions of non-linear Lie superalgebras and non-linear Lie conformal superalgebras, introduced in [DK].

In this section, Γ_+ is a discrete, additively closed subset of \mathbb{R}_+ containing 0. We let $\Gamma'_+ = \Gamma_+ \setminus \{0\}$, and for $\zeta \in \Gamma'_+$ we denote by ζ_- the largest element of Γ_+ strictly smaller than ζ .

3.1 Non-linear Lie superalgebras

Let \mathfrak{g} be a vector superspace, and let $\mathcal{T}(\mathfrak{g})$ denote the tensor superalgebra over \mathfrak{g} . We assume that \mathfrak{g} is endowed with a Γ'_+ -grading

$$\mathfrak{g} = \bigoplus_{\zeta \in \Gamma'_+} \mathfrak{g}_\zeta ,$$

and we extend it to a grading $\mathcal{T}(\mathfrak{g}) = \bigoplus_{\zeta \in \Gamma_+} \mathcal{T}(\mathfrak{g})[\zeta]$ in the obvious way, namely, writing $\zeta(A) = \zeta$ when $A \in \mathcal{T}(\mathfrak{g})[\zeta]$, we require that $\zeta(1) = 0$ and $\zeta(A \otimes B) = \zeta(A) + \zeta(B)$. We have the induced increasing Γ_+ -filtration by subspaces

$$\mathcal{T}_\zeta(\mathfrak{g}) = \bigoplus_{\zeta' \leq \zeta} \mathcal{T}(\mathfrak{g})[\zeta'] .$$

If \mathfrak{g} is endowed with a linear map

$$[\ , \] : \mathfrak{g} \otimes \mathfrak{g} \longrightarrow \mathcal{T}(\mathfrak{g}) ,$$

we extend it to $\mathcal{T}(\mathfrak{g})$ by the Leibniz rule:

$$\begin{aligned} [A, B \otimes C] &= [A, B] \otimes C + p(A, B)B \otimes [A, C] , \\ [A \otimes B, C] &= A \otimes [B, C] + p(B, C)[A, C] \otimes B . \end{aligned}$$

Let, for $\zeta \in \Gamma_+$,

$$\mathcal{M}_\zeta(\mathfrak{g}) = \text{span} \left\{ \begin{array}{c} A \otimes (b \otimes c - p(b, c)c \otimes b) \\ -[b, c] \otimes D \end{array} \mid \begin{array}{c} b, c \in \mathfrak{g}, A, D \in \mathcal{T}(\mathfrak{g}), \\ \zeta(A \otimes b \otimes c \otimes D) \leq \zeta \end{array} \right\},$$

and let $\mathcal{M}(\mathfrak{g}) = \cup_\zeta \mathcal{M}_\zeta(\mathfrak{g})$. Note that $\mathcal{M}(\mathfrak{g})$ is the two sided ideal of the tensor algebra $\mathcal{T}(\mathfrak{g})$ generated by elements $a \otimes b - p(a, b)b \otimes a - [a, b]$, where $a, b \in \mathfrak{g}$.

Definition 3.1. A *non-linear Lie superalgebra* \mathfrak{g} is a Γ'_+ -graded vector superspace with a parity preserving linear map $[\cdot, \cdot] : \mathfrak{g} \otimes \mathfrak{g} \rightarrow \mathcal{T}(\mathfrak{g})$ satisfying the following properties ($a, b, c \in \mathfrak{g}$):

1. grading condition: $[\mathfrak{g}_{\zeta_1}, \mathfrak{g}_{\zeta_2}] \subset \mathcal{T}_{(\zeta_1 + \zeta_2)_-}(\mathfrak{g})$,
2. skewsymmetry: $[a, b] = -p(a, b)[b, a]$,
3. Jacobi identity:

$$[a, [b, c]] - p(a, b)[b, [a, c]] - [[a, b], c] \in \mathcal{M}_{(\zeta(a) + \zeta(b) + \zeta(c))_-}(\mathfrak{g}).$$

The associative superalgebra $U(\mathfrak{g}) = \mathcal{T}(\mathfrak{g})/\mathcal{M}(\mathfrak{g})$ is called the universal enveloping algebra of the non-linear Lie superalgebra \mathfrak{g} .

Definition 3.2. Let U be an associative superalgebra and let $\mathcal{B} = \{E_i\}_{i \in \mathcal{I}}$ be an ordered set of elements of U , compatible with the parity. We shall say that \mathcal{B} is a set of *PBW generators* of U (and that U is *PBW generated* by \mathcal{B}) if the collection of *ordered monomials*

$$S = \left\{ E_{i_1} \dots E_{i_s} \mid i_i \leq \dots \leq i_s \text{ and } i_k < i_{k+1} \text{ if } p(i_k) = \bar{1} \right\}.$$

is a basis of U . If \mathfrak{g} is a Γ'_+ -graded subspace of U and any basis of \mathfrak{g} , compatible with parity and Γ'_+ -grading, PBW generates U , we shall say that U is *PBW generated by* \mathfrak{g} .

It is not difficult to show by the usual method (see e.g. [J]) that the PBW theorem holds for the universal enveloping superalgebra $U(\mathfrak{g})$ of a non-linear Lie superalgebra \mathfrak{g} :

Theorem 3.3. *Let \mathfrak{g} be a non-linear Lie superalgebra. Then $U(\mathfrak{g})$ is PBW generated by \mathfrak{g} .*

In order to state a converse theorem, we need the following

Definition 3.4. Let U be an associative superalgebra, generated by a Γ'_+ -graded subspace $\mathfrak{g} = \bigoplus_{\zeta \in \Gamma'_+} \mathfrak{g}_\zeta$. Define an increasing filtration of U induced by the Γ'_+ -grading of \mathfrak{g} , i.e. the filtration by the subspaces

$$U_\zeta = \sum_{\zeta_1 + \dots + \zeta_s \leq \zeta} \mathfrak{g}_{\zeta_1} \dots \mathfrak{g}_{\zeta_s}. \quad (3.1)$$

We shall say that U is *pregraded by* Γ_+ if

$$[\mathfrak{g}_{\zeta_1}, \mathfrak{g}_{\zeta_2}]^U \subset U_{(\zeta_1 + \zeta_2)_-}, \quad (3.2)$$

where $[\cdot, \cdot]^U$ denotes the commutator in the associative superalgebra U .

Let U be an associative superalgebra and let $\mathfrak{g} = \bigoplus_{\zeta \in \Gamma'_+} \mathfrak{g}_\zeta$ be a subspace of U . We shall assume that U is PBW generated by an ordered basis $\mathcal{B} = \{E_i \mid i \in \mathcal{I}\}$ of \mathfrak{g} compatible with parity and Γ'_+ -grading. Moreover we assume that U is pregraded by Γ_+ . We have $\mathcal{I} = \bigsqcup_\zeta \mathcal{I}_\zeta$, where $\zeta(E_i) = \zeta$ for $i \in \mathcal{I}_\zeta$. It is convenient to introduce the following notation for $\vec{j} = (j_1, \dots, j_s) \in \mathcal{I}^s$ with $j_k \in \mathcal{I}_{\zeta_k}$:

$$\begin{aligned} \zeta(\vec{j}) &= \zeta_1 + \dots + \zeta_s, \\ \#(\vec{j}) &= \# \left\{ (p, q) \mid \begin{array}{l} 1 \leq p < q \leq s \text{ and either } j_p > j_q, \\ \text{or } j_p = j_q \text{ and } p(E_{j_p}) = \bar{1} \end{array} \right\}. \end{aligned} \quad (3.3)$$

Let

$$\begin{aligned} S_\zeta &= \left\{ E_{i_1} \dots E_{i_s} \mid \#(\vec{i}) = 0, \zeta(\vec{i}) \leq \zeta \right\}, \quad S = \bigcup_{\zeta \in \Gamma_+} S_\zeta; \\ \tilde{S}_\zeta &= \left\{ E_{i_1} \otimes \dots \otimes E_{i_s} \mid \#(\vec{i}) = 0, \zeta(\vec{i}) \leq \zeta \right\}, \quad \tilde{S} = \bigcup_{\zeta \in \Gamma_+} \tilde{S}_\zeta. \end{aligned}$$

If $A = E_{j_1} \otimes \dots \otimes E_{j_s}$, we will also let sometimes $\zeta(A) = \zeta(\vec{j})$ and $\#(A) = \#(\vec{j})$.

Let π be the natural quotient map

$$\pi : \mathcal{T}(\mathfrak{g}) \twoheadrightarrow U,$$

given by

$$\pi(a_1 \otimes \dots \otimes a_n) = a_1 \dots a_n, \quad \text{for } a_1, \dots, a_n \in \mathfrak{g},$$

and let ρ be the embedding

$$\rho : U \hookrightarrow \mathcal{T}(\mathfrak{g}),$$

given by $\rho(E_{i_1} \dots E_{i_s}) = E_{i_1} \otimes \dots \otimes E_{i_s}$ for every element from S . Note that $\pi \circ \rho = I_U$. The surjective map π induces the Γ_+ -filtration on U , given by the subspaces (3.1), namely $U_\zeta = \pi(\mathcal{T}_\zeta(\mathfrak{g}))$. On the other hand, since by assumption U is PBW generated by \mathcal{B} , the injective map ρ defines a Γ_+ -grading $U = \bigoplus_\zeta U[\zeta]$ (as a vector space, not as an algebra), by the condition $\rho(U[\zeta]) \subset \mathcal{T}(\mathfrak{g})[\zeta]$, i.e.

$$U[\zeta] = \text{span}_{\mathbb{C}}(S_\zeta \setminus S_{\zeta_-}). \quad (3.4)$$

Moreover, thanks to the embedding $\rho : U \rightarrow \mathcal{T}(\mathfrak{g})$, we can define a linear map

$$[\cdot, \cdot] : \mathfrak{g} \otimes \mathfrak{g} \longrightarrow \mathcal{T}(\mathfrak{g}),$$

given by

$$[a, b] = \rho(ab - p(a, b)ba), \quad (3.5)$$

and we extend it to a linear map $[\cdot, \cdot] : \mathcal{T}(\mathfrak{g}) \otimes \mathcal{T}(\mathfrak{g}) \rightarrow \mathcal{T}(\mathfrak{g})$, by the Leibniz rule, as explained above.

Lemma 3.5. (a) The Γ_+ -filtration (3.1) is induced by the Γ_+ -grading (3.4):

$$U_\zeta = \bigoplus_{\zeta' \leq \zeta} U[\zeta'] .$$

(b) We have

$$\mathcal{T}_\zeta(\mathfrak{g}) = \text{span}_{\mathbb{C}} \tilde{S}_\zeta + \mathcal{M}_\zeta(\mathfrak{g}) .$$

Proof. It is clear by definition that $\bigoplus_{\zeta' \leq \zeta} U[\zeta'] \subset U_\zeta$. In order to prove the reverse inclusion, consider an arbitrary (not necessarily ordered) monomial $A = E_{j_1} \dots E_{j_s}$, and assume that $\zeta(\vec{j}) = \zeta$ (recall notation (3.3)). We want to prove, by induction on $(\zeta(\vec{j}), \#(\vec{j}))$, that $A \in \bigoplus_{\zeta' \leq \zeta} U[\zeta']$. If $\#(\vec{j}) = 0$, we have $A \in U[\zeta]$, so there is nothing to prove. Let $\#(\vec{j}) \geq 1$ and let k be such that either $j_k > j_{k+1}$, or $j_k = j_{k+1}$ and $p(E_{j_k}) = \bar{1}$. In the first case we have

$$\begin{aligned} A &= p(E_{j_k}, E_{j_{k+1}}) E_{j_1} \dots E_{j_{k+1}} E_{j_k} \dots E_{j_s} \\ &+ E_{j_1} \dots [E_{j_k}, E_{j_{k+1}}] \dots E_{j_s} = B + C . \end{aligned}$$

Obviously $\zeta(B) = \zeta(\vec{j})$ and $\#(B) = \#(\vec{j}) - 1$, so by the inductive assumption $B \in \bigoplus_{\zeta' \leq \zeta} U[\zeta']$. Moreover, by the grading condition (3.2), we have $C \in U_{\zeta_-}$, hence, by the inductive assumption, $C \in \bigoplus_{\zeta' \leq \zeta} U[\zeta']$. Suppose now $j_k = j_{k+1}$ and $p(E_{j_k}) = \bar{1}$. In this case we have

$$A = \frac{1}{2} E_{j_1} \dots [E_{j_k}, E_{j_{k+1}}] \dots E_{j_s} ,$$

so that, by (3.2), $A \in U_{\zeta_-}$, and, by inductive assumption, $A \in \bigoplus_{\zeta' \leq \zeta} U[\zeta']$. This proves part (a). In order to prove part (b) we need to show that, given an arbitrary (not necessarily ordered) monomial $A = E_{j_1} \otimes \dots \otimes E_{j_n}$, with $\zeta(\vec{j}) = \zeta$, we can decompose it as

$$A = P + M , \tag{3.6}$$

where $P \in \text{span}_{\mathbb{C}} \tilde{S}_\zeta$ and $M \in \mathcal{M}_\zeta(\mathfrak{g})$. The argument is very similar to that used in part (a), by induction on the pair $(\zeta(\vec{j}), \#(\vec{j}))$. For example, if $\#(\vec{j}) \geq 1$ and k is such that $j_k > j_{k+1}$, we have, by definition of $\mathcal{M}_\zeta(\mathfrak{g})$,

$$\begin{aligned} A &\equiv p(E_{j_k}, E_{j_{k+1}}) E_{j_1} \otimes \dots \otimes E_{j_{k+1}} \otimes E_{j_k} \otimes \dots \otimes E_{j_s} \\ &+ E_{j_1} \otimes \dots \otimes [E_{j_k}, E_{j_{k+1}}] \otimes \dots \otimes E_{j_s} \quad \text{mod } \mathcal{M}_\zeta(\mathfrak{g}) . \end{aligned} \tag{3.7}$$

By the grading condition (3.2) and part (a), we have $[E_{j_k}, E_{j_{k+1}}]^U \in U_{(\zeta_k + \zeta_{k+1})_-} = \bigoplus_{\zeta' < \zeta_k + \zeta_{k+1}} U[\zeta']$. This implies, by the definition (3.4) of the grading of U , that $[E_{j_k}, E_{j_{k+1}}] = \rho([E_{j_k}, E_{j_{k+1}}]^U) \in \bigoplus_{\zeta' < \zeta_k + \zeta_{k+1}} \mathcal{T}(\mathfrak{g})[\zeta'] = \mathcal{T}_{(\zeta_k + \zeta_{k+1})_-}(\mathfrak{g})$. Therefore the second term in the RHS of (3.7) belongs to $\mathcal{T}_{\zeta_-}(\mathfrak{g})$. We thus use induction to conclude that both terms in the RHS of (3.7) decompose as in (3.6). \square

Theorem 3.6. *Let U be an associative algebra, let \mathfrak{g} be a Γ'_+ -graded subspace of U , and let \mathcal{B} be a basis of \mathfrak{g} compatible with parity and Γ'_+ -grading. Suppose that U is PBW generated by \mathcal{B} , and that it is pregraded by Γ_+ . Then the linear map $[\cdot, \cdot] : \mathfrak{g} \otimes \mathfrak{g} \rightarrow \mathcal{T}(\mathfrak{g})$, defined by (3.5), defines a structure of a non-linear Lie superalgebra on \mathfrak{g} , such that U is canonically isomorphic to the universal enveloping algebra $U(\mathfrak{g})$.*

Proof. First, let us list some simple facts about U and $\mathcal{T}(\mathfrak{g})$:

- (a) $[U_{\zeta_1}, U_{\zeta_2}]^U \subset U_{(\zeta_1 + \zeta_2)_-}$, where, as before, $[\cdot, \cdot]^U$ denotes the commutator in U .
- (b) $[\mathcal{T}_{\zeta_1}(\mathfrak{g}), \mathcal{T}_{\zeta_2}(\mathfrak{g})] \subset \mathcal{T}_{(\zeta_1 + \zeta_2)_-}(\mathfrak{g})$, where here $[\cdot, \cdot]$ denotes the "non-linear" bracket of \mathfrak{g} , given by (3.5), extended to $\mathcal{T}(\mathfrak{g})$ by the Leibniz rule.
- (c) $\pi([A, B]) = [\pi(A), \pi(B)]^U$, for $A, B \in \mathcal{T}(\mathfrak{g})$.

By (3.2) we have $[\mathfrak{g}_{\zeta_1}, \mathfrak{g}_{\zeta_2}]^U \subset U_{(\zeta_1 + \zeta_2)_-}$. Hence (a) follows by applying the Leibniz rule to the commutator of arbitrary monomials and by the definition (3.1) of the filtration U_{ζ} . By Lemma 3.5(a) we also have

$$[\mathfrak{g}_{\zeta_1}, \mathfrak{g}_{\zeta_2}] \subset \rho(U_{(\zeta_1 + \zeta_2)_-}) = \rho\left(\bigoplus_{\zeta' < \zeta_1 + \zeta_2} U[\zeta']\right) \subset \bigoplus_{\zeta' < \zeta_1 + \zeta_2} \mathcal{T}(\mathfrak{g})[\zeta'] = \mathcal{T}_{(\zeta_1 + \zeta_2)_-}(\mathfrak{g}).$$

(b) then follows since, by definition, $[\cdot, \cdot]$ is defined on $\mathcal{T}(\mathfrak{g})$ by the Leibniz rule. (c) is obviously true when $A, B \in \mathfrak{g}$, since $\pi \circ \rho = \mathbb{1}|_U$. Hence (c) holds for arbitrary monomials $A, B \in \mathcal{T}(\mathfrak{g})$, since both the left and the right hand side satisfy the Leibniz rule.

We can now prove the theorem. We need to show that $[\cdot, \cdot]$ satisfies the axioms of a non-linear Lie superalgebra (see Definition 3.1). The grading condition is a special case of (b). Skewsymmetry is obvious. It remains to prove the Jacobi identity. We denote, for $a, b, c \in \mathfrak{g}$

$$J(a, b, c) = [a, [b, c]] - p(a, b)[b, [a, c]] - [[a, b], c] \in \mathcal{T}(\mathfrak{g}),$$

and we need to show that $J(a, b, c) \in \mathcal{M}_{\zeta_-}(\mathfrak{g})$, where $\zeta = \zeta(a) + \zeta(b) + \zeta(c)$. By (b) we have $J(a, b, c) \in \mathcal{T}_{\zeta_-}(\mathfrak{g})$. Hence by Lemma 3.5(b), we can decompose $J = P + M$, with $P \in \text{span}_{\mathbb{C}} \tilde{S}_{\zeta_-}$ and $M \in \mathcal{M}_{\zeta_-}(\mathfrak{g})$. By (c) we have $\pi(J(a, b, c)) = [a, [b, c]^U]^U - p(a, b)[b, [a, c]^U]^U - [[a, b]^U, c]^U = 0$, since obviously the Jacobi identity holds on U . On the other hand $\pi(M) = 0$, by definition of $\mathcal{M}_{\zeta}(\mathfrak{g})$. We thus have $\pi(P) = 0$. Since, by assumption, U is PBW generated by \mathcal{B} , we have $\pi : \text{span}_{\mathbb{C}} \tilde{S} \xrightarrow{\sim} U$. Hence $P = 0$, and $J(a, b, c) = M \in \mathcal{M}_{\zeta_-}(\mathfrak{g})$, as required. \square

3.2 Non-linear Lie conformal superalgebras

In this section, following [DK], we recall the definitions and results in the (more difficult) "conformal" case, similar to those in the non-linear Lie algebra case,

discussed in the previous section. In particular we show that the category of non-linear Lie conformal algebras is equivalent to that of pregraded freely generated vertex algebras.

Let R be a $\mathbb{C}[T]$ -module together with a Γ'_+ -grading by $\mathbb{C}[T]$ -submodules:

$$R = \bigoplus_{\zeta \in \Gamma'_+} R_\zeta .$$

We extend the action of T to be a derivation of the tensor algebra $\mathcal{T}(R)$ of R , viewed as a vector superspace, and we extend the Γ_+ -grading to $\mathcal{T}(R) = \bigoplus_{\zeta \in \Gamma_+} \mathcal{T}(R)[\zeta]$ in the obvious way. We also let $\mathcal{T}_\zeta(R) = \bigoplus_{\zeta' \leq \zeta} \mathcal{T}(R)[\zeta']$ be the induced increasing filtration.

We say that R is a *non-linear conformal algebra* if it is endowed with a linear map

$$[\lambda] : R \otimes R \longrightarrow \mathbb{C}[\lambda] \otimes \mathcal{T}(R) ,$$

called the λ -*bracket*, satisfying the following conditions:

1. grading condition: $[R_{\zeta_1} \lambda R_{\zeta_2}] \subset \mathbb{C}[\lambda] \otimes \mathcal{T}_{(\zeta_1 + \zeta_2)_-}(R)$,
2. sesquilinearity: $[Ta \lambda b] = -\lambda[a \lambda b]$, $[a \lambda Tb] = (\lambda + T)[a \lambda b]$.

It is not hard to show, by induction on $\zeta \in \Gamma_+$, that the λ -bracket can be extended to the whole tensor algebra $\mathcal{T}(R)$, by using the axioms in the fourth definition of a vertex algebra (see Section 1.5):

Lemma 3.7. *[DK] If R is a non-linear conformal algebra, there is a unique way to define linear maps*

$$\mathcal{T}(R) \otimes \mathcal{T}(R) \longrightarrow \mathcal{T}(R) , \quad A \otimes B \mapsto :AB: ,$$

$$\mathcal{T}(R) \otimes \mathcal{T}(R) \longrightarrow \mathbb{C}[\lambda] \otimes \mathcal{T}(R) , \quad A \otimes B \mapsto [A \lambda B] ,$$

such that $[a \lambda b]$, with $a, b \in R$, coincides with the λ -bracket on R , $:aB := a \otimes B$ if $a \in R$ and $B \in \mathcal{T}(R)$, and the following relations hold, with $a, b \in R$, $A, B, C \in \mathcal{T}(R)$:

$$\begin{aligned} :1A: &= :A1: = A , \quad [1 \lambda A] = [A \lambda 1] = 0 , \\ :(a \otimes B)C: &= :a(:BC:): + : \left(\int_0^T d\lambda a \right) [B \lambda C] : \\ &+ p(a, B) : \left(\int_0^T d\lambda B \right) [a \lambda C] : , \\ [a \lambda (b \otimes C)] &= :[a \lambda b]C: + p(a, b) : b[a \lambda C] : \\ &+ \int_0^\lambda d\mu [[a \lambda b]_\mu C] , \\ [(a \otimes B) \lambda C] &= : \left(e^{T\partial_\lambda} a \right) [B \lambda C] : + p(a, B) : \left(e^{T\partial_\lambda} B \right) [a \lambda C] : \\ &+ p(a, B) \int_0^\lambda d\mu [B_\mu [a \lambda_{-\mu} C]] . \end{aligned}$$

The above linear maps satisfy the following grading conditions

$$: \mathcal{T}_{\zeta_1}(R) \mathcal{T}_{\zeta_2}(R) : \subset \mathcal{T}_{\zeta_1 + \zeta_2}(R), \quad [\mathcal{T}_{\zeta_1}(R) \lambda \mathcal{T}_{\zeta_2}(R)] \subset \mathbb{C}[\lambda] \mathcal{T}_{(\zeta_1 + \zeta_2)_-}(R).$$

For $\zeta \in \Gamma_+$ we let $\mathcal{M}_\zeta(R)$ be the subspace of $\mathcal{T}(R)$ spanned by the elements

$$A \otimes \left((b \otimes c \otimes D - p(b, c) c \otimes b \otimes D - : \left(\int_{-T}^0 d\lambda [b \lambda c] \right) D : \right), \quad (3.8)$$

with $b, c \in R$, $A, D \in \mathcal{T}(R)$, such that $A \otimes b \otimes c \otimes D \in \mathcal{T}_\zeta(R)$. We also let $\mathcal{M}(R) = \bigcup_{\zeta \in \Gamma_+} \mathcal{M}_\zeta(R)$. This is the analogue of the space $\mathcal{M}(\mathfrak{g})$ defined in the previous section, but now $\mathcal{M}(R)$ is only a left ideal of $\mathcal{T}(R)$, not a two-sided ideal. In particular the quotient space $\mathcal{T}(R)/\mathcal{M}(R)$ is not an associative algebra.

Definition 3.8. A *non-linear Lie conformal algebra* is a non-linear conformal algebra (defined above) satisfying the following additional conditions:

3. skewsymmetry: $[a \lambda b] = -p(a, b)[b \lambda - a]$ for $a, b \in R$,
4. Jacobi identity:

$$[a \lambda [b \mu c]] - p(a, b)[b \mu [a \lambda c]] - [[a \lambda b] \lambda + \mu c] \in \mathbb{C}[\lambda, \mu] \otimes \mathcal{M}_\zeta(R),$$

where $a, b, c \in R$ and $\zeta = (\zeta(a) + \zeta(b) + \zeta(c))_-$.

Example 3.9. Let R be a Lie conformal algebra with λ -bracket $[\cdot \lambda \cdot]$ and let $\hat{R} = R \oplus \mathbb{C}C$ be its central extension, where C is a central element, $TC = 0$ and the λ -bracket of $a, b \in R \subset \hat{R}$ is $[a \lambda b]^\wedge = [a \lambda b] + \varphi(a, b)C$, with $\varphi(a, b) \in \mathbb{C}$. Then we can construct a non-linear Lie conformal algebra structure on R by

$$[a \lambda b]^\wedge = [a \lambda b] + \varphi(a, b),$$

and the trivial grading $R = R_1$. Similarly, a central extension of a Lie algebra \mathfrak{g} can be viewed as a non-linear Lie algebra structure on \mathfrak{g} .

The following theorem, is a "conformal" analogue of the PBW Theorem 3.3 for non-linear Lie algebras. Its proof is much more difficult (see [DK] for details).

Theorem 3.10. [DK] *Let R be a non-linear Lie conformal algebra. Then there is a canonical structure of a vertex algebra on the space $V(R) = \mathcal{T}(R)/\mathcal{M}(R)$, called the universal enveloping vertex algebra of R , such that the vacuum vector is $|0\rangle = 1 \bmod \mathcal{M}(R)$, and such that the infinitesimal translation operator, the normally ordered product and the λ -bracket on $V(R)$ are induced by T , $:$ and $[\lambda]$ respectively. Moreover the canonical map $\pi : \mathcal{T}(R) \rightarrow V(R)$ induces a $\mathbb{C}[T]$ -module embedding $R \subset V(R)$ and $V(R)$ is freely generated by R , namely, any ordered basis \mathcal{B} of R , compatible with parity and Γ'_+ -grading, freely generates $V(R)$ (in the sense of Definition 1.19).*

We next want to state the converse of Theorem 3.10, i.e. the "conformal analogue" of Theorem 3.6 of the previous section.

Definition 3.11. Let V be a vertex algebra, strongly generated by a free $\mathbb{C}[T]$ -submodule $R = \mathbb{C}[T] \otimes \mathfrak{g}$, where $\mathfrak{g} = \bigoplus_{\zeta \in \Gamma'_+} \mathfrak{g}_\zeta$. Define an increasing Γ_+ -filtration by subspaces V_ζ of V , induced by the Γ'_+ -grading of $R = \bigoplus_{\zeta \in \Gamma'_+} (\mathbb{C}[T] \otimes \mathfrak{g}_\zeta)$, where we let $\deg T = 0$. Namely

$$V_\zeta = \sum_{\substack{k_1, \dots, k_s \in \mathbb{Z}_+ \\ \zeta_1 + \dots + \zeta_s \leq \zeta}} : (T^{k_1} \mathfrak{g}_{\zeta_1}) \dots (T^{k_s} \mathfrak{g}_{\zeta_s}) : . \quad (3.9)$$

We shall say that the vertex algebra V is *pregraded* by Γ_+ if the λ -bracket on V satisfies the following *grading condition*

$$[\mathfrak{g}_{\zeta_1} \lambda \mathfrak{g}_{\zeta_2}] \subset \mathbb{C}[\lambda] \otimes V_{(\zeta_1 + \zeta_2)_-} . \quad (3.10)$$

Example 3.12. Suppose the vertex algebra V is freely generated by $\mathbb{C}[T] \otimes \mathfrak{g}$ and it has a Hamiltonian operator H with eigenvalues in Γ_+ , such that $H\mathfrak{g} \subset \mathfrak{g}$ and $\mathfrak{g} \cap V[0] = 0$. Then the corresponding eigenspace decomposition $\mathfrak{g} = \bigoplus_{\Delta \in \text{Spec } H} \mathfrak{g}[\Delta]$ makes V a *pregraded* vertex algebra. Notice that in this case the Γ_+ -filtration V_Δ defined in (3.9) is not induced by the H -eigenspace decomposition of V . Indeed the filtered space V_Δ is preserved by T , while we know that T increases the conformal weight by 1. On the other hand we clearly have

$$\bigoplus_{\Delta' \leq \Delta} V[\Delta'] \subset V_\Delta .$$

From this we get

$$[\mathfrak{g}[\Delta_1] \lambda \mathfrak{g}[\Delta_2]] \subset \mathbb{C}[\lambda] \otimes \left(\bigoplus_{\Delta \leq \Delta_1 + \Delta_2 - 1} V[\Delta] \right) \subset \mathbb{C}[\lambda] \otimes V_{\Delta_1 + \Delta_2 - 1} ,$$

which proves, as we wanted, that V is pregraded by Γ_+ .

It is not hard to show, by induction, that the grading condition (3.10) can be extended to the whole vertex algebra V , namely we have

Lemma 3.13. *[DK] If V is pregraded by Γ_+ , then we have, for $\zeta_1, \zeta_2 \in \Gamma_+$,*

$$: V_{\zeta_1} V_{\zeta_2} : \subset V_{\zeta_1 + \zeta_2} , \quad [V_{\zeta_1} \lambda V_{\zeta_2}] \subset \mathbb{C}[\lambda] \otimes V_{(\zeta_1 + \zeta_2)_-} .$$

The proof of the following result is similar, though somewhat more complicated, than the proof of Theorem 3.6 in the previous section, and it can be found in [DK].

Theorem 3.14. *[DK] Let V be a vertex algebra. Let $R = \bigoplus_{\zeta \in \Gamma'_+} (\mathbb{C}[T] \otimes \mathfrak{g}_\zeta)$ be a Γ'_+ -graded free $\mathbb{C}[T]$ -submodule. Assume that there exists a basis of R , compatible with parity and Γ'_+ -grading of R , which freely generates V . Moreover, assume that V is pregraded by Γ_+ . Then R admits a structure of a non-linear Lie conformal algebra*

$$[\lambda] : R \otimes R \longrightarrow \mathbb{C}[\lambda] \otimes \mathcal{T}(R) ,$$

compatible with the λ -bracket of V , in the sense that $\pi([a \lambda b]^R) = [a \lambda b]^V$ for $a, b \in R$. We are writing upper indices R and V to distinguish the λ -bracket of V from the non-linear λ -bracket of R , and, as before, π denotes the quotient map $\mathcal{T}(R) \twoheadrightarrow V(R)$. Moreover V is canonically isomorphic to the universal enveloping vertex algebra $V(R)$.

Remark 3.15. A homomorphism of Γ'_+ -graded non-linear Lie conformal algebras R and R' is a $\mathbb{C}[T]$ -module homomorphism $\phi : R \rightarrow R'$ preserving $\mathbb{Z}/2\mathbb{Z}$ and Γ'_+ -gradings, such that the induced map $\mathcal{T}(R) \rightarrow \mathcal{T}(R')$ satisfies the following properties:

$$\begin{aligned} \phi(: AB :) - \phi(A)\phi(B) &: \in \mathcal{M}_{\zeta(A)+\zeta(B)}(R') , \\ \phi([A \lambda B]) - [\phi(A) \lambda \phi(B)] &\in \mathbb{C}[\lambda] \otimes \mathcal{M}_{(\zeta(A)+\zeta(B))_-}(R') . \end{aligned}$$

It is not hard to see, using Theorems 3.10 and 3.14, that the category of Γ'_+ -graded free, as $\mathbb{C}[T]$ -modules, non-linear Lie conformal algebras is equivalent to the category of Γ_+ -pregraded vertex algebras, freely generated by a free $\mathbb{C}[T]$ -module. In particular R and R' are isomorphic non-linear Lie conformal algebras if and only if the corresponding universal enveloping vertex algebras $V(R)$ and $V(R')$ are isomorphic vertex algebras. Of course, similar facts hold in the non-linear Lie algebra case as well.

Remark 3.16. Let R be a non-linear conformal algebra which satisfies the skewsymmetry condition 3 of Definition 3.8. The necessary and sufficient condition for R to be a non-linear Lie conformal algebra, i.e. to satisfy condition 4 of Definition 3.8, is as follows. Choose an ordered basis \mathcal{B} of R , compatible with $\mathbb{Z}/2\mathbb{Z}$ and Γ'_+ -grading. Write the LHS of the Jacobi identity $J(a, b, c; \lambda, \mu)$ of three elements $a, b, c \in \mathcal{B}$, as an element of $\mathcal{T}(R)[\lambda, \mu]$ in terms of linear combinations of products of a^i 's. Using skewsymmetry and quasiassociativity rules, rewrite $J(a, b, c; \lambda, \mu)$ as a sum of an element of $\mathcal{M}(R)$, which we ignore, and a linear combination of ordered monomials. Then all coefficients of this linear combination should be zero. It immediately follows from Definition 3.8 and Theorem 3.14 that this condition is respectively sufficient and necessary. If R is freely generated, as a $\mathbb{C}[T]$ -module, by a set $\tilde{\mathcal{B}}$, we can take $\mathcal{B} = \{T^{(k)}a_i \mid k \in \mathbb{Z}_+, a_i \in \tilde{\mathcal{B}}\}$. Then it suffices to check the above condition for $J(a, b, c; \lambda, \mu)$ for all triples of elements of $\tilde{\mathcal{B}}$. Similar remarks hold in the ordinary algebra case.

3.3 The H -twisted Zhu algebra of a non-linear Lie conformal algebra

Let V be a vertex algebra with a Hamiltonian operator H . Let $R \subset V$ be a free $\mathbb{C}[T]$ -submodule $R = \mathbb{C}[T] \otimes \mathfrak{g}$, where \mathfrak{g} is an H -invariant subspace of V . As usual, we let $V[\Delta]$ be the eigenspace of H with eigenvalue Δ , and we denote $\mathfrak{g}[\Delta] = \mathfrak{g} \cap V[\Delta]$. We also fix a Γ'_+ -grading of \mathfrak{g} by H -invariant subspaces:

$$\mathfrak{g} = \bigoplus_{\zeta \in \Gamma'_+} \mathfrak{g}_\zeta , \quad (3.11)$$

which we extend to a Γ'_+ -grading of R by letting $\deg(T) = 0$. We assume that there exists an ordered basis of R , compatible with parity and Γ'_+ -grading, which freely generates V . Moreover, we assume that V is pregraded by Γ_+ , according to Definition 3.11.

By Theorems 3.10 and 3.14, the above assumptions are equivalent to say that $R = \mathbb{C}[T]\mathfrak{g}$ admits a structure of non-linear Lie conformal algebra, and that V is isomorphic to its universal enveloping vertex algebra $V(R)$.

Thanks to Lemma 3.13, condition (3.10) immediately implies the following stronger grading conditions:

$$: V_{\zeta_1} V_{\zeta_2} : \subset V_{\zeta_1 + \zeta_2}, TV_{\zeta} \subset V_{\zeta}, [V_{\zeta_1} \lambda V_{\zeta_2}] \subset \mathbb{C}[\lambda] \otimes V_{(\zeta_1 + \zeta_2)_-}. \quad (3.12)$$

Recall the definitions of the $*_n$ -products and the $*$ -bracket on V , given in Example 2.15. We immediately get from (3.12) the following further grading conditions, for $k \in \mathbb{Z}_+$:

$$\begin{aligned} V_{\zeta_1} *_k V_{\zeta_2} &\subset V_{\zeta_1 + \zeta_2}, & V_{\zeta_1} * V_{\zeta_2} &\subset V_{(\zeta_1 + \zeta_2)_-}, \\ [V_{\zeta_1} * V_{\zeta_2}] &\subset V_{(\zeta_1 + \zeta_2)_-}. \end{aligned} \quad (3.13)$$

Remark 3.17. If the Hamiltonian operator H has eigenvalues in Γ'_+ , then we can just take $\mathfrak{g}_{\zeta} = \mathfrak{g}[\zeta]$, and the grading condition (3.10) is satisfied. Notice that, in this case, the Γ_+ -filtration V_{ζ} is not induced by the H -grading, as the subspaces V_{ζ} are preserved by T , while T increases the conformal weight by 1. On the other hand $\bigoplus_{\Delta \leq \zeta} V[\Delta] \subset V_{\zeta}$, from which condition (3.10) follows immediately.

We fix an ordered basis $\bar{\mathcal{B}}$ of \mathfrak{g} , compatible with the Γ'_+ -grading (3.11) and the H -grading:

$$\bar{\mathcal{B}} = \{e_{\alpha} \mid \alpha \in \bar{\mathcal{I}}\}.$$

We let $\zeta_{\alpha} = \zeta$ if $e_{\alpha} \in \mathfrak{g}_{\zeta}$, and $\Delta_{\alpha} = \Delta$ if $e_{\alpha} \in \mathfrak{g}[\Delta]$. Moreover we let $\bar{\mathcal{I}}_{\zeta} = \{\alpha \mid \zeta_{\alpha} = \zeta\}$ and $\bar{\mathcal{I}}[\Delta] = \{\alpha \mid \Delta_{\alpha} = \Delta\}$, and we extend $\bar{\mathcal{B}}$ to the following ordered basis of $R = \mathbb{C}[T]\mathfrak{g}$:

$$\mathcal{B} = \left\{ e_i = T^{(k)} e_{\alpha} \mid i = (\alpha, k) \in \bar{\mathcal{I}} \times \mathbb{Z}_+ =: \mathcal{I} \right\}.$$

As before, $T^{(k)} = T^k/k!$. Recalling that $TV_{\zeta} \subset V_{\zeta}$ and $TV[\Delta] \subset V[\Delta + 1]$, we let $\mathcal{I}_{\zeta} = \bar{\mathcal{I}}_{\zeta} \times \mathbb{Z}_+$ and $\mathcal{I}[\Delta] = \bigsqcup_{k \in \mathbb{Z}_+} \bar{\mathcal{I}}[\Delta - k] \times \{k\}$. By Theorem 3.10, V is freely generated by R , hence the collection of ordered monomials

$$S = \{ : e_{i_1} \dots e_{i_s} : \mid \#(\vec{i}) = 0 \}.$$

is a basis of V . Here and further, for $\vec{j} = (j_1, \dots, j_s) \in \mathcal{I}^s$, with $j_l = (\alpha_l, k_l)$, $\alpha_l \in \bar{\mathcal{I}}_{\zeta_l} \cap \bar{\mathcal{I}}[\Delta_l]$, $k_l \in \mathbb{Z}_+$, we denote $\zeta(\vec{j})$ and $\#(\vec{j})$ as in (3.3), and we let

$$\Delta(\vec{j}) = \Delta_1 + k_1 + \dots + \Delta_s + k_s.$$

It is clear by definition that, for $\vec{j} = (j_1, \dots, j_s) \in \mathcal{I}^s$, we have $: e_{j_1} \dots e_{j_s} : \in V_{\zeta(\vec{j})} \cap V[\Delta(\vec{j})]$. We define the sets

$$\begin{aligned} S_\zeta &= \left\{ : e_{i_1} \dots e_{i_s} : \mid \zeta(\vec{i}) \leq \zeta, \#(\vec{i}) = 0 \right\}, \\ S_\zeta^* &= \left\{ e_{i_1} *_-1 \dots *_-1 e_{i_s} \mid \zeta(\vec{i}) \leq \zeta, \#(\vec{i}) = 0 \right\}, \end{aligned}$$

so that the basis S can be written as $S = \bigcup_\zeta S_\zeta$, and similarly we define $S^* = \bigcup_\zeta S_\zeta^*$. In the above formula, the $*_{-1}$ -product of more than two elements is defined, as for the normally ordered product, by taking products from right to left: $e_{i_1} *_-1 \dots *_-1 e_{i_s} = e_{i_1} *_-1 (e_{i_2} *_-1 (\dots *_-1 e_{i_s}))$. Recall the filtration V_ζ introduced above; we also define spaces V_ζ^* in a similar way, with $*_{-1}$ -products instead of normally ordered products. In other words,

$$\begin{aligned} V_\zeta &= \text{span}_{\mathbb{C}} \left\{ : e_{j_1} \dots e_{j_s} : \mid \zeta(\vec{j}) \leq \zeta \right\}, \\ V_\zeta^* &= \text{span}_{\mathbb{C}} \left\{ e_{j_1} *_-1 \dots *_-1 e_{j_s} \mid \zeta(\vec{j}) \leq \zeta \right\}. \end{aligned}$$

Lemma 3.18. (a) For every $\vec{j} = (j_1, \dots, j_s)$, we have

$$e_{j_1} *_-1 \dots *_-1 e_{j_s} - : e_{j_1} \dots e_{j_s} : \in V_{\zeta(\vec{j})_-}.$$

(b) The following spaces are equal:

$$V_\zeta = \text{span}_{\mathbb{C}} S_\zeta = V_\zeta^* = \text{span}_{\mathbb{C}} S_\zeta^*.$$

(c) S^* is a basis of V .

Proof. We prove (a) by induction on $\zeta(\vec{j})$. For $s = 1$ there is nothing to prove, so let $s \geq 2$. By inductive assumption we have

$$e_{j_2} *_-1 \dots *_-1 e_{j_s} = : e_{j_2} \dots e_{j_s} : + A,$$

where $A \in V_{(\zeta(\vec{j}) - \zeta(e_{j_1}))_-}$. Hence, by definition of the $*_{-1}$ -product, we have

$$\begin{aligned} e_{j_1} *_-1 \dots *_-1 e_{j_s} &= : e_{j_1} e_{j_2} \dots e_{j_s} : + e_{j_1} *_-1 A \\ &+ \sum_{n \geq 0} \binom{\Delta(e_{j_1})}{n+1} e_{j_1}^{(n)} (: e_{j_2} \dots e_{j_s} :). \end{aligned}$$

The grading conditions (3.12) guarantee that $e_{j_1}^{(n)} (: e_{j_2} \dots e_{j_s} :) \in V_{\zeta(\vec{j})_-}$, and similarly conditions (3.13) guarantee that $e_{j_1} *_-1 A \in V_{\zeta(\vec{j})_-}$. This proves (a).

We now prove, by induction on ζ , that $V_\zeta^* = V_\zeta$. Let $j = (j_1, \dots, j_s)$ be such that $\zeta(\vec{j}) = \zeta$. By (a) we have

$$e_{j_1} *_-1 \dots *_-1 e_{j_s} - : e_{j_1} \dots e_{j_s} : = A \in V_{\zeta_-}. \quad (3.14)$$

Now we can read equation (3.14) in two ways. On one hand, since $:e_{j_1} \cdots e_{j_s}:$ $\in V_\zeta$, and $A \in V_{\zeta_-} \subset V_\zeta$, we get $e_{j_1} *_-1 \cdots *_-1 e_{j_s} \in V_\zeta$. Hence $V_\zeta^* \subset V_\zeta$. On the other hand, we have $e_{j_1} *_-1 \cdots *_-1 e_{j_s} \in V_\zeta^*$, and, by inductive assumption, $A \in V_{\zeta_-} = V_{\zeta_-}^* \subset V_\zeta^*$, so that $:e_{j_1} \cdots e_{j_s}:$ $\in V_\zeta^*$. Hence $V_\zeta \subset V_\zeta^*$.

To show that $V_\zeta = \text{span}_{\mathbb{C}} S_\zeta$, we use the standard argument. First notice that, obviously, $\text{span}_{\mathbb{C}} S_\zeta \subset V_\zeta$. So we only need to show that, given an arbitrary (non necessarily ordered) monomial $:e_{j_1} \cdots e_{j_s}:$ $\in V_\zeta$, $\zeta = \zeta(\vec{j})$, we can write it as a linear combination of ordered monomials $:e_{i_1} \cdots e_{i_s}:$ $\in S_\zeta$. We will prove the statement by induction on the pair $(\zeta(\vec{j}), \#(\vec{j}))$ ordered lexicographically. If $\#(\vec{j}) = 0$, we have $:e_{j_1} \cdots e_{j_s}:$ $\in S_\zeta$, so there is nothing to prove. Suppose then $\#(\vec{j}) \geq 1$, and let p be such that $j_p > j_{p+1}$, or $j_p = j_{p+1}$ and $p(e_{j_p}) = \bar{1}$. In the first case we have:

$$\begin{aligned} :e_{j_1} \cdots e_{j_s}: &= p(e_{j_p}, e_{j_{p+1}}) :e_{j_1} \cdots e_{j_{p+1}} e_{j_p} \cdots e_{j_s}: \\ &+ :e_{j_1} \cdots \left(\int_{-T}^0 d\lambda [e_{j_p} \lambda e_{j_{p+1}}] \right) \cdots e_{j_s}: = B + C. \end{aligned}$$

Obviously $\zeta(B) = \zeta(\vec{j})$ and $\#(B) = \#(\vec{j}) - 1$, so $B \in \text{span}_{\mathbb{C}} S_\zeta$ by inductive assumption. Moreover, it immediately follows by the grading conditions (3.12) that $C \in V_{\zeta_-}$, hence $C \in \text{span}_{\mathbb{C}} S_\zeta$, again by inductive assumption. Consider now the second case, namely $j_p = j_{p+1}$ and $p(e_{j_p}) = \bar{1}$. We have

$$:e_{j_1} \cdots e_{j_s}: = \frac{1}{2} :e_{j_1} \cdots \left(\int_{-T}^0 d\lambda [e_{j_p} \lambda e_{j_{p+1}}] \right) \cdots e_{j_s}:.$$

As before, the RHS belongs to $\text{span}_{\mathbb{C}} S_\zeta$ thanks to the grading conditions (3.12) and inductive assumption.

To conclude the proof of (b), we are left to show that $V_\zeta^* = \text{span}_{\mathbb{C}} S_\zeta^*$. We could proceed in the same way as above. Instead we will prove, equivalently, that $\text{span}_{\mathbb{C}} S_\zeta^* = \text{span}_{\mathbb{C}} S_\zeta$. The proof is again by induction on ζ . Let $\vec{i} = (i_1, \dots, i_s)$ be such that $\zeta(\vec{i}) = \zeta$ and $\#(\vec{i}) = 0$. By (a) we have

$$e_{i_1} *_-1 \cdots *_-1 e_{i_s} - :e_{i_1} \cdots e_{i_s}: = A \in V_{\zeta_-}. \quad (3.15)$$

By the previous result and inductive assumption we have $A \in V_{\zeta_-} = \text{span}_{\mathbb{C}} S_{\zeta_-} = \text{span}_{\mathbb{C}} S_{\zeta_-}^*$. Hence it follows by (3.15) that both $e_{i_1} *_-1 \cdots *_-1 e_{i_s} \in \text{span}_{\mathbb{C}} S_\zeta$ and $:e_{i_1} \cdots e_{i_s}: \in \text{span}_{\mathbb{C}} S_\zeta^*$. This clearly implies that $\text{span}_{\mathbb{C}} S_\zeta^* = \text{span}_{\mathbb{C}} S_\zeta$.

We are left to prove (c). By (b) we only need to show that elements of S^* are linearly independent. Suppose by contradiction that there is a non-trivial linear dependence,

$$\sum_{\vec{i}} c_{\vec{i}} e_{i_1} *_-1 \cdots *_-1 e_{i_s} = 0. \quad (3.16)$$

Let ζ be the largest value of $\zeta(\vec{i})$ for which $c_{\vec{i}} \neq 0$. Thanks to (a), we can rewrite (3.16) as

$$\sum_{\vec{i} \mid \zeta(\vec{i}) = \zeta} c_{\vec{i}} :e_{i_1} \cdots e_{i_s}: + A = 0, \quad (3.17)$$

where $A \in V_{\zeta_-} = \text{span}_{\mathbb{C}} S_{\zeta_-}$. On the other hand, by assumption, S is a basis of V . Hence (3.17) implies that $c_{\vec{i}} = 0$ for all \vec{i} such that $\zeta(\vec{i}) = \zeta$, which is a contradiction. \square

For $\zeta \in \Gamma_+$, we let

$$\Sigma_{\zeta} = \left\{ \begin{array}{l} (T^{(h_1)} e_{\alpha_1}) *_{-1} \cdots *_{-1} (T^{(h_s)} e_{\alpha_s}) \\ - \binom{-\Delta_1}{h_1} \cdots \binom{-\Delta_s}{h_s} e_{\alpha_1} *_{-1} \cdots *_{-1} e_{\alpha_s} \end{array} \middle| \begin{array}{l} \zeta(\vec{i}) \leq \zeta, \#(\vec{i}) = 0, \sum_i h_i > 0 \\ \text{where } \vec{i} = (i_1, \dots, i_s), \\ i_l = (\alpha_l, h_l), \alpha_l \in \bar{\mathcal{I}}[\Delta_l], h_l \in \mathbb{Z}_+ \end{array} \right\},$$

and

$$W_{\zeta} = \text{span}_{\mathbb{C}} \left\{ \begin{array}{l} (T^{(k_1)} e_{\beta_1}) *_{-1} \cdots *_{-1} (T^{(k_s)} e_{\beta_s}) \\ - \binom{-\Delta_1}{k_1} \cdots \binom{-\Delta_s}{k_s} e_{\beta_1} *_{-1} \cdots *_{-1} e_{\beta_s} \end{array} \middle| \begin{array}{l} \zeta(\vec{j}) \leq \zeta, \text{ where} \\ \vec{j} = (j_1 \dots j_s), k_l \in \mathbb{Z}_+ \\ j_l = (\beta_l, k_l), \beta_l \in \bar{\mathcal{I}}[\Delta_l] \end{array} \right\}.$$

Moreover we let $\Sigma = \bigcup_{\zeta} \Sigma_{\zeta}$ and $W = \bigcup_{\zeta} W_{\zeta}$. Obviously $\text{span}_{\mathbb{C}} \Sigma_{\zeta} \subset W_{\zeta}$. We will prove that the converse is also true.

Lemma 3.19. *If $e = T^{(k_1)} e_{\beta_1}$, $\beta_1 \in \bar{\mathcal{I}}_{\zeta_1} \cap \bar{\mathcal{I}}[\Delta_1]$, $k_1 \in \mathbb{Z}_+$, and $A \in W_{\zeta}$, then*

$$e *_{-1} A \in W_{\zeta_1 + \zeta}.$$

Proof. Let

$$A = (T^{(k_2)} e_{\beta_2}) *_{-1} \cdots *_{-1} (T^{(k_s)} e_{\beta_s}) - \binom{-\Delta_2}{k_2} \cdots \binom{-\Delta_s}{k_s} e_{\beta_2} *_{-1} \cdots *_{-1} e_{\beta_s}.$$

The lemma then immediately follows by the following obvious identity, obtained by adding and subtracting terms:

$$\begin{aligned} e *_{-1} A &= \left((T^{(k_1)} e_{\beta_1}) *_{-1} (T^{(k_2)} e_{\beta_2}) *_{-1} \cdots *_{-1} (T^{(k_s)} e_{\beta_s}) \right. \\ &\quad - \binom{-\Delta_1}{k_1} \binom{-\Delta_2}{k_2} \cdots \binom{-\Delta_s}{k_s} e_{\beta_1} *_{-1} e_{\beta_2} *_{-1} \cdots *_{-1} e_{\beta_s} \Big) \\ &\quad - \binom{-\Delta_2}{k_2} \cdots \binom{-\Delta_s}{k_s} \left(T^{(k_1)} e_{\beta_1} - \binom{-\Delta_1}{k_1} e_{\beta_1} \right) *_{-1} e_{\beta_2} *_{-1} \cdots *_{-1} e_{\beta_s}. \end{aligned}$$

\square

Recall from Section 2 that we denote by $J_{\hbar=1} = V *_{-2} V$ the span of elements $a *_{-2} b$ with $a, b \in V$. We also denote, for $\zeta \in \Gamma_+$,

$$(V *_{-2} V)_{\zeta} = \sum_{\substack{\zeta_1, \zeta_2 \in \Gamma_+ \\ \zeta_1 + \zeta_2 \leq \zeta}} V_{\zeta_1} *_{-2} V_{\zeta_2},$$

so that $V *_{-2} V = \bigcup_{\zeta \in \Gamma_+} (V *_{-2} V)_{\zeta}$. We have the following

Lemma 3.20. (a) For $A \in V_{\zeta_1}$, $B \in V_{\zeta_2}$, we have

$$(T^{(k)}A) *_{-1} B - \binom{\Delta(A)}{k} A *_{-1} B \in V_{\zeta_1} *_{-2} V_{\zeta_2}, \quad (3.18)$$

$$\begin{aligned} & [(T^{(h)}A)_*(T^{(k)}B)] \\ & - \binom{-\Delta(A)}{h} \binom{-\Delta(B)}{k} [A_*B] \in (V *_{-2} V)_{(\zeta_1 + \zeta_2)_-}. \end{aligned} \quad (3.19)$$

(b) For $A \in V_{\zeta_1}$, $B \in V_{\zeta_2}$, $C \in V_{\zeta_3}$, we have

$$A *_{-k-1} B \in V_{\zeta_1} *_{-2} V_{\zeta_2}, \quad \forall k \geq 1, \quad (3.20)$$

$$(A *_{-1} B) *_{-1} C - A *_{-1} (B *_{-1} C) \in (V *_{-2} V)_{(\zeta_1 + \zeta_2 + \zeta_3)_-}, \quad (3.21)$$

$$A *_{-1} B - p(A, B) B *_{-1} A - [A * B] \in (V *_{-2} V)_{(\zeta_1 + \zeta_2)_-}. \quad (3.22)$$

(c) The following relations hold

$$V_{\zeta_1} *_{-1} (V_{\zeta_2} *_{-2} V_{\zeta_3}) \subset (V *_{-2} V)_{\zeta_1 + \zeta_2 + \zeta_3}, \quad (3.23)$$

$$(V_{\zeta_1} *_{-2} V_{\zeta_2}) *_{-1} V_{\zeta_3} \subset (V *_{-2} V)_{\zeta_1 + \zeta_2 + \zeta_3}, \quad (3.24)$$

$$[V_{\zeta_1} * (V_{\zeta_2} *_{-2} V_{\zeta_3})] \subset (V *_{-2} V)_{(\zeta_1 + \zeta_2 + \zeta_3)_-}, \quad (3.25)$$

$$[(V_{\zeta_1} *_{-2} V_{\zeta_2}) * V_{\zeta_3}] \subset (V *_{-2} V)_{(\zeta_1 + \zeta_2 + \zeta_3)_-}. \quad (3.26)$$

Proof. We want to prove (3.18) by induction on k . For $k = 1$ it follows immediately by equation (2.9). For $k \geq 2$ we have

$$\begin{aligned} & (T^{(k)}A) *_{-1} B - \binom{-\Delta(A)}{k} A *_{-1} B \\ &= \frac{1}{k} \left((T^{(k-1)}TA) *_{-1} B - \binom{-\Delta(A)-1}{k-1} (TA) *_{-1} B \right) \\ & \quad + \frac{1}{k} \binom{-\Delta(A)-1}{k-1} A *_{-2} B. \end{aligned}$$

The above identity is obtained by adding and subtracting $\frac{1}{k} \binom{-\Delta(A)-1}{k} (TA) *_{-1} B$, and using (2.9) with $\hbar = 1$. Notice that, by definition, the filtered space V_ζ is invariant by the action of T . Hence, by inductive assumption, both terms on the RHS side belong to $V_{\zeta_1} *_{-2} V_{\zeta_2}$, as wanted. By (2.10) we have

$$[(T^{(h)}A)_*B] = - \binom{-\Delta(A)}{h} [A_*B],$$

so it suffices to prove (3.19) for $h = 0$. We will prove it by induction on k . For $k = 1$ we have, by (2.11),

$$[A_*(TB)] + \Delta(B)[A_*B] = (T + H)[A_*B] = [A * B] *_{-2} |0\rangle,$$

and the RHS belongs to $V_{(\zeta_1+\zeta_2)-} *_{-2} V_0 \subset (V *_{-2} V)_{(\zeta_1+\zeta_2)-}$. For $k \geq 2$ we have, by adding and subtracting terms,

$$\begin{aligned} & [A_*(T^{(k)}B)] - \binom{-\Delta(B)}{k} [A * B] \\ &= \frac{1}{k} \left([A_*(T^{(k-1)}TB)] - \binom{-\Delta(B)-1}{k-1} [A * TB] \right) \\ &+ \frac{1}{k} \binom{-\Delta(B)-1}{k-1} \left([A_*(TB)] + \Delta(B)[A * B] \right). \end{aligned}$$

By inductive assumption, both terms on the RHS belong to $(V *_{-2} V)_{(\zeta_1+\zeta_2)-}$, thus proving (3.19).

We can now use the results of the previous section to deduce the relations in part (b). Equation (3.20) follows immediately by (2.38) and (3.18), equation (3.21) follows by (2.17), (3.13) and (3.20), equation (3.22) follows by (2.19), (3.12) and (3.18) with $B = |0\rangle$.

We are left to prove part (c). Condition (3.24) follows by (2.18), (3.13) and (3.20). Condition (3.25) follows by Lemma 2.6 and (3.13). For (3.23) we have, by adding and subtracting terms,

$$\begin{aligned} A *_{-1} (B *_{-2} C) &= p(A, B)p(A, C)(B *_{-2} C) *_{-1} A \\ &+ [A * (B *_{-2} C)] + \left(A *_{-1} (B *_{-2} C) \right. \\ &\left. - p(A, B)p(A, C)(B *_{-2} C) *_{-1} A - [A * (B *_{-2} C)] \right), \end{aligned}$$

so that (3.23) follows by (3.24), (3.25) and (3.22). Finally, with a similar argument, based on adding and subtracting the appropriate terms to $[(A *_{-2} B)_* C]$, it is not hard to show that (3.26) follows by (3.25) and (3.22). \square

Lemma 3.21. (a) *The following spaces are equal:*

$$\text{span}_{\mathbb{C}} \Sigma_{\zeta} = W_{\zeta} = (V *_{-2} V)_{\zeta}.$$

In particular

$$\text{span}_{\mathbb{C}} \Sigma = W = V *_{-2} V.$$

(b) Σ is a basis of $V *_{-2} V$.

Proof. Let $\vec{j} = (j_1, \dots, j_s)$ be such that $j_l = (\beta_l, k_l)$, $\beta_l \in \bar{\mathcal{I}}_{\zeta_l} \cap \bar{\mathcal{I}}[\Delta_l]$, $k_l \in \mathbb{Z}_+$,

and $\zeta(\vec{j}) = \zeta_1 + \cdots \zeta_s = \zeta$. We then have, by adding and subtracting terms,

$$\begin{aligned} & (T^{(k_1)}e_{\beta_1}) *_{-1} \cdots *_{-1} (T^{(k_s)}e_{\beta_s}) - \binom{-\Delta_1}{k_1} \cdots \binom{-\Delta_s}{k_s} e_{\beta_1} *_{-1} \cdots *_{-1} e_{\beta_s} \\ &= \left(T^{(k_1)}e_{\beta_1} - \binom{-\Delta_1}{k_1} e_{\beta_1} \right) *_{-1} \left((T^{(k_2)}e_{\beta_2}) *_{-1} \cdots *_{-1} (T^{(k_s)}e_{\beta_s}) \right) \\ & \quad + \binom{-\Delta_1}{k_1} e_{\beta_1} *_{-1} \left((T^{(k_2)}e_{\beta_2}) *_{-1} \cdots *_{-1} (T^{(k_s)}e_{\beta_s}) \right. \\ & \quad \left. - \binom{-\Delta_2}{k_2} \cdots \binom{-\Delta_s}{k_s} e_{\beta_2} *_{-1} \cdots *_{-1} e_{\beta_s} \right). \end{aligned}$$

The first term in the RHS belongs to $V_{\zeta_1} *_{-2} V_{\zeta-\zeta_1}$ by (3.18), while the second term in the RHS belongs to $(V *_{-2} V)_{\zeta}$ by induction on s and (3.23). This proves the inclusion $W_{\zeta} \subset (V *_{-2} V)_{\zeta}$.

We next prove the reverse inclusion: $(V *_{-2} V)_{\zeta} \subset W_{\zeta}$. Recall that, by Lemma 3.18(b), V_{ζ} is spanned by elements $e_{j_1} *_{-1} \cdots *_{-1} e_{j_s}$ with $\zeta(\vec{j}) \leq \zeta$. Let $\vec{i} = (i_1, \dots, i_s)$, with $i_l = (\alpha_l, h_l)$, $\alpha_l \in \bar{\mathcal{I}}_{\zeta_l^1} \cap \bar{\mathcal{I}}[\Delta_l^1]$, $h_l \in \mathbb{Z}_+$, and $\vec{j} = (j_1, \dots, j_t)$, with $j_l = (\beta_l, k_l)$, $\beta_l \in \bar{\mathcal{I}}_{\zeta_l^2} \cap \bar{\mathcal{I}}[\Delta_l^2]$, $k_l \in \mathbb{Z}_+$, be such that $\zeta(\vec{i}) = \zeta_1^1 + \cdots + \zeta_s^1 = \zeta_1$, $\zeta(\vec{j}) = \zeta_1^2 + \cdots + \zeta_t^2 = \zeta_2$, and $\zeta_1 + \zeta_2 = \zeta$. We need to prove, by induction on ζ , that

$$(e_{i_1} *_{-1} \cdots *_{-1} e_{i_s}) *_{-2} (e_{j_1} *_{-1} \cdots *_{-1} e_{j_t}) \in W_{\zeta}.$$

We will consider separately the two cases $s = 1$ and $s \geq 2$. For $s = 1$ we have

$$\begin{aligned} & e_{i_1} *_{-2} (e_{j_1} *_{-1} \cdots *_{-1} e_{j_t}) = ((T + \Delta)e_{i_1}) *_{-1} e_{j_1} *_{-1} \cdots *_{-1} e_{j_t} \\ &= (h_1 + 1) \left((T^{(h_1+1)}e_{\alpha_1}) *_{-1} (T^{(k_1)}e_{\beta_1}) *_{-1} \cdots *_{-1} (T^{(k_t)}e_{\beta_t}) \right. \\ & \quad \left. - \binom{-\Delta_1^1}{h_1+1} \binom{-\Delta_1^2}{k_1} \cdots \binom{-\Delta_t^2}{k_t} e_{\alpha_1} *_{-1} e_{\beta_1} *_{-1} \cdots *_{-1} e_{\beta_t} \right) \\ & \quad + (\Delta_1^1 + h_1) \left((T^{(h_1)}e_{\alpha_1}) *_{-1} (T^{(k_1)}e_{\beta_1}) *_{-1} \cdots *_{-1} (T^{(k_t)}e_{\beta_t}) \right. \\ & \quad \left. - \binom{-\Delta_1^1}{h_1} \binom{-\Delta_1^2}{k_1} \cdots \binom{-\Delta_t^2}{k_t} e_{\alpha_1} *_{-1} e_{\beta_1} *_{-1} \cdots *_{-1} e_{\beta_t} \right) \in W_{\zeta}. \end{aligned}$$

The last equality is obtained by adding and subtracting the same term and using the obvious identity $(h_1 + 1) \binom{-\Delta_1^1}{h_1+1} = -(\Delta_1^1 + h_1) \binom{-\Delta_1^1}{h_1}$. For $s \geq 2$ we use equation (2.16) with $a = e_{i_1}$, $b = e_{i_2} *_{-2} \cdots *_{-1} e_{i_s}$, $c = e_{j_1} *_{-1} \cdots *_{-1} e_{j_t}$, $n = -1$

and $k = -2$. We get

$$\begin{aligned}
& (e_{i_1} *_{-1} \cdots *_{-1} e_{i_s}) *_{-2} (e_{j_1} *_{-1} \cdots *_{-1} e_{j_t}) \\
&= e_{i_1} *_{-1} ((e_{i_2} *_{-1} \cdots *_{-1} e_{i_s}) *_{-2} (e_{j_1} *_{-1} \cdots *_{-1} e_{j_t})) \\
&+ e_{i_1} *_{-2} ((e_{i_2} *_{-1} \cdots *_{-1} e_{i_s}) *_{-1} (e_{j_1} *_{-1} \cdots *_{-1} e_{j_t})) \\
&+ \sum_{j \geq 0} \left(e_{i_1} *_{-j-3} ((e_{i_2} *_{-1} \cdots *_{-1} e_{i_s}) *_{-1} (e_{j_1} *_{-1} \cdots *_{-1} e_{j_t})) \right. \\
&\quad \left. \pm (e_{i_2} *_{-1} \cdots *_{-1} e_{i_s}) *_{-j-3} (e_{i_1} *_{-1} (e_{j_1} *_{-1} \cdots *_{-1} e_{j_t})) \right),
\end{aligned}$$

where $\pm = p(e_{i_1} e_{i_2}) \cdots p(e_{i_1}, e_{i_s})$. The first term in the RHS belongs to W_ζ by Lemma 3.19 and by induction. Moreover, by the above result (for $s = 1$), also the second term of the RHS belongs to W_ζ . Finally, by (3.13) and (3.20), the last terms of the RHS belong to $(V *_{-2} V)_{\zeta_-}$, so it belongs to W_ζ by induction on ζ .

To conclude part (a) we are left to prove that $W_\zeta \subset \text{span}_{\mathbb{C}} \Sigma_\zeta$. We will use the usual argument. Let $\vec{j} = (j_1, \dots, j_s)$ be an arbitrary (not necessarily ordered) s -tuple with $j_l = (\beta_l, k_l)$, $\beta_l \in \bar{I}_{\zeta_l}[\Delta_l]$, $k_l \in \mathbb{Z}_+$ and such that $\zeta(\vec{j}) = \zeta$. Consider the corresponding element

$$\begin{aligned}
A &= (T^{(k_1)} e_{\beta_1}) *_{-1} \cdots *_{-1} (T^{(k_s)} e_{\beta_s}) \\
&- \left(\begin{smallmatrix} -\Delta_1 \\ k_1 \end{smallmatrix} \right) \cdots \left(\begin{smallmatrix} -\Delta_s \\ k_s \end{smallmatrix} \right) e_{\beta_1} *_{-1} \cdots *_{-1} e_{\beta_s} \in W_\zeta.
\end{aligned}$$

We need to prove, by induction on the pair $(\zeta(\vec{j}), \#(\vec{j}))$, that we can write A as a linear combination of "ordered" elements from Σ_ζ . If $\#(\vec{j}) = 0$, we have $A \in \Sigma_\zeta$, so there is nothing to prove. Then, suppose that $\#(\vec{j}) \geq 1$, and we consider the case in which there is some $p \in \{1, \dots, s\}$ such that $j_p > j_{p+1}$. The case $j_p = j_{p+1}$, $p(e_{j_p}) = \bar{1}$, can be studied in the same way. By adding and subtracting the appropriate terms, we can rewrite A as the sum of the following ten elements:

$$\begin{aligned}
A_1 &= e_{j_1} *_{-1} \cdots *_{-1} \left(e_{j_p} *_{-1} e_{j_{p+1}} *_{-1} \cdots *_{-1} e_{\beta_s} \right. \\
&\quad \left. - \left(e_{j_p} *_{-1} e_{j_{p+1}} \right) *_{-1} \cdots *_{-1} e_{j_s} \right), \\
A_2 &= - \prod_{l=1}^s \left(\begin{smallmatrix} -\Delta_l \\ k_l \end{smallmatrix} \right) e_{\beta_1} *_{-1} \cdots *_{-1} \left(e_{\beta_p} *_{-1} e_{\beta_{p+1}} *_{-1} \cdots *_{-1} e_{\beta_s} \right. \\
&\quad \left. - \left(e_{\beta_p} *_{-1} e_{\beta_{p+1}} \right) *_{-1} \cdots *_{-1} e_{\beta_s} \right),
\end{aligned}$$

$$\begin{aligned}
A_3 &= e_{j_1} *_{-1} \cdots *_{-1} \left(\left(e_{j_p} *_{-1} e_{j_{p+1}} \right) *_{-1} \cdots *_{-1} e_{j_s} \right. \\
&\quad \mp \left. \left(e_{j_{p+1}} *_{-1} e_{j_p} \right) *_{-1} \cdots *_{-1} e_{j_s} - [e_{j_p} * e_{j_{p+1}}] *_{-1} \cdots *_{-1} e_{j_s} \right), \\
A_4 &= - \prod_{l=1}^s \binom{-\Delta_l}{k_l} e_{\beta_1} *_{-1} \cdots *_{-1} \left(\left(e_{\beta_p} *_{-1} e_{\beta_{p+1}} \right) *_{-1} \cdots *_{-1} e_{\beta_s} \right. \\
&\quad \mp \left. \left(e_{\beta_{p+1}} *_{-1} e_{\beta_p} \right) *_{-1} \cdots *_{-1} e_{\beta_s} - [e_{\beta_p} * e_{\beta_{p+1}}] *_{-1} \cdots *_{-1} e_{\beta_s} \right), \\
A_5 &= \pm e_{j_1} *_{-1} \cdots *_{-1} \left(\left(e_{j_{p+1}} *_{-1} e_{j_p} \right) *_{-1} \cdots *_{-1} e_{j_s} \right. \\
&\quad \left. - e_{j_{p+1}} *_{-1} e_{j_p} *_{-1} \cdots *_{-1} e_{j_s} \right), \\
A_6 &= \mp \prod_{l=1}^s \binom{-\Delta_l}{k_l} e_{\beta_1} *_{-1} \cdots *_{-1} \left(\left(e_{\beta_{p+1}} *_{-1} e_{\beta_p} \right) *_{-1} \cdots *_{-1} e_{\beta_s} \right. \\
&\quad \left. - e_{\beta_{p+1}} *_{-1} e_{\beta_p} *_{-1} \cdots *_{-1} e_{\beta_s} \right), \\
A_7 &= e_{j_1} *_{-1} \cdots *_{-1} \left([e_{j_p} * e_{j_{p+1}}] *_{-1} e_{j_{p+2}} *_{-1} \cdots *_{-1} e_{j_s} \right. \\
&\quad \left. - \binom{-\Delta_p}{k_p} \binom{-\Delta_{p+1}}{k_{p+1}} [e_{\beta_p} * e_{\beta_{p+1}}] *_{-1} e_{j_{p+2}} *_{-1} \cdots *_{-1} e_{j_s} \right), \\
A_8 &= \binom{-\Delta_p}{k_p} \binom{-\Delta_{p+1}}{k_{p+1}} e_{j_1} *_{-1} \cdots *_{-1} e_{j_{p-1}} *_{-1} \\
&\quad \left([e_{\beta_p} * e_{\beta_{p+1}}] *_{-1} (T^{(k_{p+2})} e_{\beta_{p+2}}) *_{-1} \cdots *_{-1} (T^{(k_s)} e_{\beta_s}) \right. \\
&\quad \left. - \prod_{l=p+2}^s \binom{-\Delta_l}{k_l} [e_{\beta_p} * e_{\beta_{p+1}}] *_{-1} e_{\beta_{p+2}} *_{-1} \cdots *_{-1} e_{\beta_s} \right), \\
A_9 &= \prod_{l=p}^s \binom{-\Delta_l}{k_l} (T^{(k_1)} e_{\beta_1}) *_{-1} \cdots *_{-1} (T^{(k_{p-1})} e_{\beta_{p-1}}) *_{-1} \\
&\quad \left([e_{\beta_p} * e_{\beta_{p+1}}] *_{-1} e_{\beta_{p+2}} *_{-1} \cdots *_{-1} e_{\beta_s} \right) \\
&\quad - \prod_{l=1}^s \binom{-\Delta_l}{k_l} e_{\beta_1} *_{-1} \cdots *_{-1} e_{\beta_{p-1}} *_{-1} \\
&\quad \left([e_{\beta_p} * e_{\beta_{p+1}}] *_{-1} e_{\beta_{p+2}} *_{-1} \cdots *_{-1} e_{\beta_s} \right), \\
A_{10} &= \pm (T^{(k_1)} e_{\beta_1}) *_{-1} \cdots (T^{(k_{p+1})} e_{\beta_{p+1}}) *_{-1} (T^{(k_p)} e_{\beta_p}) \cdots *_{-1} (T^{(k_s)} e_{\beta_s}) \\
&\quad \mp \prod_{l=1}^s \binom{-\Delta_l}{k_l} e_{\beta_1} *_{-1} \cdots *_{-1} e_{\beta_{p+1}} *_{-1} e_{\beta_p} *_{-1} \cdots *_{-1} e_{\beta_s}.
\end{aligned}$$

Here $\pm = p(e_{j_p}, e_{j_{p+1}})$. All terms A_1, \dots, A_{10} belong to $\text{span}_{\mathbb{C}} \Sigma_{\zeta}$ by inductive assumption. More precisely, A_1, A_2, A_5 and A_6 belong to $(V *_2 V)_{\zeta_-} = W_{\zeta_-}$, thanks to (3.21) and (3.23), hence by induction they belong to $\text{span}_{\mathbb{C}} \Sigma_{\zeta_-} \subset \text{span}_{\mathbb{C}} T_{\zeta}$. Similarly, A_3 and A_4 belong to $(V *_2 V)_{\zeta_-}$, thanks to (3.22), (3.23) and (3.24). A_7 belongs to $(V *_2 V)_{\zeta_-}$, thanks to (3.19), (3.23) and (3.24). A_8 belongs to $(V *_2 V)_{\zeta_-}$, thanks to the equality $W_{\zeta} = (V *_2 V)_{\zeta}$ and conditions (3.13) and (3.23). A_9 belongs to $(V *_2 V)_{\zeta_-}$, by a simple induction argument based on (3.18). Finally, A_{10} is such that $\zeta((j_1, \dots, j_{p+1}, j_p, \dots, j_s)) = \zeta(\vec{j})$, but $\#((j_1, \dots, j_{p+1}, j_p, \dots, j_s)) = \#(\vec{j}) - 1$, hence by inductive assumption it belongs to $\text{span}_{\mathbb{C}} \Sigma_{\zeta}$. This completes the proof of (a).

To prove (b) we only have to show that elements of Σ are linearly independent. But this follows immediately by the definition of Σ and Lemma 3.18(c). \square

Corollary 3.22. *One has*

$$R \cap (V *_2 V) = (T + H)R.$$

Proof. The inclusion $(T + H)R \subset R \cap (V *_2 V)$ is obvious, thanks to equation (2.9). On the other hand, by Lemma 3.21, the set

$$\left\{ T^{(k)} e_{\alpha} - \binom{-\Delta_{\alpha}}{k} e_{\alpha} \mid \alpha \in \hat{\mathcal{I}}_{\Delta}, k \geq 0 \right\},$$

is a basis for $R \cap (V *_2 V)$ and each such basis element is obviously in $(T + H)R$. \square

Recall from Example 2.15 that the H -twisted Zhu algebra of V is, by definition, the space $Zhu_H V = V/(V *_2 V)$, with the associative product induced by the $*_{-1}$ -product on V . We denote by $Zhu_H R$ the image of $R \subset V$ in $Zhu_H V$ via the natural quotient map $\pi_Z : V \twoheadrightarrow Zhu_H V = V/(V *_2 V)$. Moreover, thanks to Corollary 3.22, we have the following identifications: $Zhu_H R = R/(T + H)R \simeq \mathfrak{g}$. If \bar{e}_{α} is the image of the basis element $e_{\alpha} \in \mathfrak{g}$ in $Zhu_H R$, we let, by abuse of notation,

$$\bar{\mathcal{B}} = \{ \bar{e}_{\alpha} \mid \alpha \in \bar{\mathcal{I}} \}.$$

Lemma 3.23. *The associative algebra $Zhu_H V$ is PBW generated by $\bar{\mathcal{B}}$.*

Proof. It is an immediate corollary of Lemma 3.18(c) and Lemma 3.21(b). \square

Notice that $Zhu_H R$ has a natural Γ'_+ -grading induced by the isomorphism $Zhu_H R \simeq \mathfrak{g} = \bigoplus_{\zeta \in \Gamma'_+} \mathfrak{g}_{\zeta}$.

Lemma 3.24. *The Zhu algebra $Zhu_H V$ is pregraded by Γ_+ .*

Proof. Recall the definition of the Γ_+ -filtration of $Zhu_H V$ induced by the Γ'_+ -grading of $Zhu_H R$. It is given by

$$(Zhu_H V)_\zeta = \text{span}_{\mathbb{C}} \left\{ \bar{e}_{\beta_1} \dots \bar{e}_{\beta_s} \mid \beta_k \in \bar{\mathcal{I}}_{\zeta_k}, \zeta_1 + \dots + \zeta_s \leq \zeta \right\}.$$

In particular, by Lemma 3.18(b), we have $(Zhu_H V)_\zeta = \pi_Z(V_\zeta)$. We then have, by (3.13),

$$\begin{aligned} [(Zhu_H R)_{\zeta_1}, (Zhu_H R)_{\zeta_2}] &= \pi_Z([\mathfrak{g}_{\zeta_1} * \mathfrak{g}_{\zeta_2}]) \\ &\subset \pi_Z(V_{(\zeta_1 + \zeta_2)-}) = (Zhu_H V)_{(\zeta_1 + \zeta_2)-}. \end{aligned}$$

According to Definition 3.4, this implies that V is pregraded by Γ_+ . \square

Lemma 3.23 and Lemma 3.24 guarantee that all the hypotheses of Theorem 3.6 are satisfied. In other words $Zhu_H R$ has a structure of non-linear Lie algebra and $Zhu_H V \simeq U(Zhu_H R)$.

We can summarize all the above results in the following

Theorem 3.25. *Let V be an H -graded vertex algebra and let $R \subset V$ be a free $\mathbb{C}[T]$ -module $R = \mathbb{C}[T] \otimes \mathfrak{g}$. Assume that \mathfrak{g} has a an H -invariant Γ'_+ -grading, $\mathfrak{g} = \bigoplus_{\zeta \in \Gamma'_+} \mathfrak{g}_\zeta$, which we extend to R by letting $\deg(T) = 0$. Assume that there is a basis of R , compatible with parity and Γ'_+ -grading which freely generates V (according to Definition 1.19). Moreover we assume that V is pregraded by Γ_+ , i.e. (3.10) holds. Then the space*

$$Zhu_H R = R/(T + H)R \simeq \mathfrak{g},$$

is naturally endowed with the following structure of a non-linear Lie algebra as follows. Let $\{e_\alpha \mid \alpha \in \bar{\mathcal{I}}\}$ be an ordered basis of \mathfrak{g} compatible with parity, Γ'_+ and H -gradings of \mathfrak{g} . Then the non-linear Lie bracket,

$$[\cdot, \cdot] : Zhu_H R \otimes Zhu_H R \longrightarrow \mathcal{T}(Zhu_H R),$$

is given by

$$[\bar{a}, \bar{b}] = \rho(\pi_Z([a * b])),$$

*where $a, b \in R$, \bar{a}, \bar{b} are the corresponding elements in $Zhu_H R$, π_Z denotes the natural quotient map $\pi_Z : V \twoheadrightarrow Zhu_H V = V/(V *_{-2} V)$ and ρ is the embedding $\rho : Zhu_H V \hookrightarrow \mathcal{T}(Zhu_H R)$, defined by $\rho(\bar{e}_{\alpha_1} \dots \bar{e}_{\alpha_s}) = \bar{e}_{\alpha_1} \otimes \dots \otimes \bar{e}_{\alpha_s}$, for all $\vec{\alpha} = (\alpha_1, \dots, \alpha_s)$ such that $\#(\vec{\alpha}) = 0$. Moreover the universal enveloping associative algebra of $Zhu_H R$ is naturally isomorphic to the H -twisted Zhu algebra of V :*

$$Zhu_H V \simeq U(Zhu_H R).$$

Corollary 3.26. *Let $R = \mathbb{C}[T]\mathfrak{g}$ be a non-linear Lie conformal algebra, freely generated as a $\mathbb{C}[T]$ -module, and suppose that \mathfrak{g} is endowed with a Γ'_+ -grading which induces a Hamiltonian operator H on the universal enveloping vertex*

algebra $V(R)$. Then the space $Zhu_H R = R/(T+H)R \simeq \mathfrak{g}$ is naturally endowed with a structure of a non-linear Lie algebra, so that

$$Zhu_H V(R) \simeq U(Zhu_H R) .$$

The non-linear Lie bracket on $Zhu_H R$ is "compatible" with the $*$ -bracket on R in the sense that the following diagram commutes:

$$\begin{array}{ccc} R \otimes R & \xrightarrow{[\ast]} & V(R) \\ \pi_Z \downarrow & & \downarrow \pi_Z \\ Zhu_H R \otimes Zhu_H R & \xrightarrow{[\cdot]} & Zhu_H V(R) \end{array}$$

Definition 3.27. Under the assumptions of Theorem 3.25 or Corollary 3.26, we will call $Zhu_H R$ the H -twisted Zhu non-linear Lie algebra associated to R .

Remark 3.28. It can be useful for many applications to have an explicit way to compute the map $\pi_Z : V \rightarrow Zhu_H V$. Suppose $V \simeq V(R)$ is the universal enveloping vertex algebra of a non-linear Lie conformal algebra $R = \mathbb{C}[T]\mathfrak{g}$, so that $Zhu_H V \simeq U(Zhu_H R)$. As before, for $a \in \mathfrak{g}$, we denote by \bar{a} the same element thought as an element of $Zhu_H R \simeq \mathfrak{g}$. Recall the defining relation of π_Z , i.e.

$$\pi_Z(A \ast_{-1} B) = \pi_Z(A)\pi_Z(B) .$$

We have the following identity, for $a \in \mathfrak{g}$, $B \in V$,

$$\begin{aligned} \pi_Z((T^{(k)}a)B) &= \binom{-\Delta_a}{k} \bar{a} \pi_Z(B) - \int_0^1 d\hbar \pi_Z([H(T^{(k)}a), B]_{\hbar}) \quad (3.27) \\ &= \binom{-\Delta_a}{k} \left(\bar{a} \pi_Z(B) - (\Delta_a + k) \sum_{j \in \mathbb{Z}_+} \binom{\Delta_a - 1}{j} \frac{1}{k + j + 1} \pi_Z(a_{(j)}B) \right) . \end{aligned}$$

In the proof of the first identity we use (2.5), while the second equality follows from the definition (2.4) of the \hbar -bracket. Equation (3.27) can be used inductively to compute the image via π_Z of an arbitrary element of V .

Corollary 3.29. Let $R = \mathbb{C}[T]\mathfrak{g}$ be a non-linear Lie conformal algebra as in Corollary 3.26. Then $Zhu_H V(R)$ is an associative algebra generated by \mathfrak{g} with defining relations $(a, b \in \mathfrak{g})$

$$ab - p(a, b)ba = \sum_{j \in \mathbb{Z}_+} \binom{\Delta_a - 1}{j} \pi_Z(a_{(j)}b) ,$$

where the RHS should be computed with the help of relations (3.27).

Proposition 3.30. Suppose V is a vertex algebra which satisfies all assumptions of Theorem 3.25, and let Γ be as in Example 2.12. Then, for every $c \in \mathbb{C}$, the map (2.41) is an isomorphism of Poisson algebras: $Zhu_{\hbar=0, H} V \simeq gr Zhu_{\hbar=c, H} V$.

Proof. As in (2.39), we denote by $F^\Delta V$ the increasing filtration of V induced by the conformal weight. Recall that, by definition, $J_{\hbar=1} = V *_{-2} V$. It follows by Lemma 3.21 that

$$F^\Delta \cap J_{\hbar=1} \subset \sum_{\Delta_1 + \Delta_2 + 1 \leq \Delta} V[\Delta_1] *_{-2} V[\Delta_2] \subset F^{\Delta-} V + F^\Delta V \cap (V_{(-2)} V). \quad (3.28)$$

The statement then follows from the proof of Proposition 2.17(c) and the fact that, by (2.36), $Zhu_{\hbar=c, H} V \simeq Zhu_{\hbar=1, H} V$ for arbitrary $c \neq 0$. \square

Remark 3.31. We may consider, more generally, the (H, Γ) -twisted Zhu algebra $Zhu_\Gamma V$ defined in Section 2.2, for an arbitrary Γ -grading $V = \bigoplus_{\bar{\gamma} \in \Gamma/\mathbb{Z}} \bar{\gamma} V$ of V (not necessarily induced by the action of H). Recall that we denoted by V_Γ the \mathbb{C} -span of all elements $a \in V$ such that $\Delta_a \in \bar{\gamma}_a$. It is a vertex subalgebra of V , and we clearly have, by definition, $J_\hbar(V_\Gamma) \subset J_{\hbar, \Gamma}(V)$. Hence we have, in general, a surjective associative algebra homomorphism

$$Zhu_H V_\Gamma = V_\Gamma / J_{\hbar=1}(V_\Gamma) \twoheadrightarrow V_\Gamma / J_{\hbar=1, \Gamma}(V) = Zhu_\Gamma V.$$

In particular $Zhu_\Gamma V$ will not be, in general, a PBW generated algebra, even if $V \simeq V(R)$ for some non-linear Lie conformal algebra R . It does not make sense then to talk about the non-linear Lie algebra $Zhu_\Gamma R$.

3.4 Application: the (non-commutative) Weil algebra and the cubic Dirac operator

Let \mathfrak{g} be a finite-dimensional simple or abelian Lie algebra with a non-degenerate symmetric invariant bilinear form $(\cdot | \cdot)$. Here, for simplicity, we assume \mathfrak{g} is an honest, i.e. purely even, Lie algebra, and we denote by $\bar{\mathfrak{g}}$ the space \mathfrak{g} with reversed parity, i.e. purely odd. Consider the following non-linear Lie conformal superalgebra

$$R = \mathbb{C}[T] \otimes \mathfrak{g} + \mathbb{C}[T] \otimes \bar{\mathfrak{g}},$$

where the λ -brackets are $(a, b \in \mathfrak{g}, \bar{a}, \bar{b} \in \bar{\mathfrak{g}})$:

$$\begin{aligned} [a \lambda b] &= [a, b] + \lambda(k + h^\vee)(a|b), \\ [\bar{a} \lambda \bar{b}] &= (k + h^\vee)(a|b), \\ [a \lambda \bar{b}] &= [\bar{a} \lambda b] = \overline{[a, b]}. \end{aligned} \quad (3.29)$$

Let $\{a^i\}$ be an orthonormal basis of \mathfrak{g} . Introduce the following fields of $V(R)$ (see [KT]):

$$\begin{aligned} G &= \frac{1}{k+h^\vee} \left(\sum_i : a^i \bar{a}^i : + \frac{1}{3(k+h^\vee)} \sum_{i,j} : [a^i, a^j] \bar{a}^i \bar{a}^j : \right), \\ L &= \frac{1}{2(k+h^\vee)} \left(\sum_i : a^i a^i : + \sum_i : (T \bar{a}^i) \bar{a}^i : \right. \\ &\quad \left. + \frac{1}{k+h^\vee} \sum_{i,j} : \bar{a}^i [a^i, a^j] \bar{a}^j : \right). \end{aligned} \quad (3.30)$$

The fields G and L satisfy the λ -brackets of the Neveu-Schwarz Lie conformal superalgebra with central charge

$$c = \frac{k \dim \mathfrak{g}}{k+h^\vee} + \frac{\dim \mathfrak{g}}{2},$$

i.e. L is an energy-momentum field with central charge c , G is an odd, primary with respect to L , field of conformal weight $3/2$, and

$$[G_\lambda G] = 2L + \frac{\lambda^2}{3} c. \quad (3.31)$$

Moreover, the fields $a \in \mathfrak{g}$ (respectively $\bar{a} \in \bar{\mathfrak{g}}$) are primary with respect to L of conformal weight 1 (resp. $1/2$), and $[a_\lambda G] = \lambda \bar{a}$, $[\bar{a}_\lambda G] = a$ for $a \in \mathfrak{g}$, $\bar{a} \in \bar{\mathfrak{g}}$.

Let $H = L_{(1)}$ be the Hamiltonian operator on $V(R)$, let \mathcal{W} be the H -twisted Zhu algebra of $V(R)$ and let $\pi_Z : V(R) \rightarrow \mathcal{W}$ be the quotient map. We put $k+h^\vee = 1$ (which can be achieved by rescaling the form $(\cdot | \cdot)$, provided that $k \neq -h^\vee$). By Corollary 3.26, \mathcal{W} is the universal enveloping superalgebra of the non-linear Lie superalgebra $\mathfrak{g} + \bar{\mathfrak{g}}$ with non-linear Lie brackets, extending that on \mathfrak{g} by $(a, b \in \mathfrak{g}, \bar{a}, \bar{b} \in \bar{\mathfrak{g}})$:

$$[a, \bar{b}] = [\bar{a}, b] = \overline{[a, b]}, \quad [\bar{a}, \bar{b}] = (a|b). \quad (3.32)$$

Let $D = \pi_Z(G)$ and $C = \pi_Z(L)$. Then

$$\begin{aligned} D &= \sum_i a^i \bar{a}^i + \frac{1}{3} \sum_{i,j} \overline{[a^i, a^j]} \bar{a}^i \bar{a}^j, \\ C &= \frac{1}{2} \sum_i a^i a^i + \frac{1}{2} \sum_{i,j} [a^i, a^j] \bar{a}^i \bar{a}^j + \frac{1+2h^\vee}{16} \dim \mathfrak{g}. \end{aligned}$$

Equation (3.31) implies, by looking at images via π_Z ,

$$D^2 = C + \left(\frac{h^\vee}{24} - \frac{1}{16} \right) \dim \mathfrak{g}, \quad (3.33)$$

while from the other λ -bracket relations above we get $[D, a] = 0$, $[D, \bar{a}] = a$, C is central. Furthermore, one has the following Lie algebra embedding $a \mapsto \hat{a}$ of \mathfrak{g} in \mathcal{W} , whose image commutes with $\bar{\mathfrak{g}}$:

$$\hat{a} = a - \frac{1}{2} \sum_i \overline{[a, a^i]} \bar{a}^i .$$

Let $\hat{C} = \sum_i \hat{a}^i \hat{a}^i$ be the corresponding Casimir operator. Then

$$\hat{C} = 2C - \frac{1}{8} \dim \mathfrak{g} , \quad D^2 = \frac{1}{2} \hat{C} + \frac{h^\vee}{24} \dim \mathfrak{g} = \frac{1}{2} (\hat{C} + (\rho|\rho)) ,$$

where ρ is half of the sum of positive roots of \mathfrak{g} . The latter relation was derived in [AM, Kol], where the element D was called the cubic Dirac operator, by a direct computation.

Let \mathfrak{h} be a subalgebra of \mathfrak{g} , such that the restriction of $(\cdot|\cdot)$ to \mathfrak{h} is non-degenerate, and let $\mathfrak{h} = \bigoplus_s \mathfrak{h}_s$ be the decomposition of \mathfrak{h} in a direct sum of simple and abelian components. For each \mathfrak{h}_s constructs the elements G_s and L_s as above, and let $G^{\mathfrak{h}} = \sum_s G_s$, $L^{\mathfrak{h}} = \sum_s L_s$. Then each pair G_s , L_s (resp. $G^{\mathfrak{h}}$, $L^{\mathfrak{h}}$) satisfies the relations of the Neveu-Schwarz algebra with central charge c_s (resp. $c^{\mathfrak{h}} = \sum_s c_s$), and we can form the coset Neveu-Schwarz algebra $G - G^{\mathfrak{h}}$, $L - L^{\mathfrak{h}}$ with central charge $c - c^{\mathfrak{h}}$, which commutes with all fields $a \in \mathfrak{h}$, $\bar{a} \in \bar{\mathfrak{h}}$ (cf. [KT]). Applying π_Z , we obtain the relative cubic Dirac operator $D - D^{\mathfrak{h}}$, introduced in [Kol], and the relation, derived there

$$(D - D^{\mathfrak{h}})^2 = \frac{1}{2} (\hat{C} - \hat{C}^{\mathfrak{h}} + (\rho|\rho) - (\rho_{\mathfrak{h}}|\rho_{\mathfrak{h}})) .$$

Starting from the vertex algebra $V^{QFT} = V(R)$, we can construct all "fundamental objects" in diagram (6.16), as explained in Section 6. The "classical analogue" of V^{QFT} , namely its quasiclassical limit V^{clFT} , is the following Poisson vertex algebra. As commutative associative unital superalgebra, $V^{clFT} = S(R)$, where $R = \mathbb{C}[T] \otimes \mathfrak{g} + \mathbb{C}[T] \otimes \bar{\mathfrak{g}}$ as above, and the λ -bracket $\{\cdot_\lambda \cdot\}$ on $S(R)$ is defined by extending formulas (3.29) to $S(R)$ by sesquilinearity and left and right commutative Wick formulas. Consider the elements G^{cl} and L^{cl} of the Poisson vertex algebra $S(R)$, given by the same formulas (3.30) as G and L respectively, with the normally ordered product signs $:$ removed. Then L^{cl} is a Virasoro element with zero central charge, a , \bar{a} , G^{cl} are primary with respect to L^{cl} with conformal weights $1, 1/2$ and $3/2$ respectively and we have (cf (3.31):

$$\{G^{cl}_\lambda \bar{a}\} = a , \quad \{G^{cl}_\lambda a\} = (T + \lambda)\bar{a} , \quad \{G^{cl}_\lambda G^{cl}\} = 2L^{cl} .$$

The "quantum mechanical" analogue V^{QM} of $V(R)$ is given by the H -twisted Zhu algebra. As explained above, it is $V^{QM} = U(r)$, where $r = \mathfrak{g} + \bar{\mathfrak{g}}$ is the non-linear Lie superalgebra with Lie bracket given in (3.32). This algebra is also referred to as the *non-commutative Weil algebra* associated to \mathfrak{g} , [AM]. The "classical mechanical" analogue V^{clM} of $V(R)$ is the quasiclassical limit of $Zhu_H V$, or, equivalently, the H -twisted Zhu algebra of V^{clFT} , as defined

in Section 6. It is the commutative associative unital superalgebra $S(r) = S(\mathfrak{g}) \otimes \Lambda(\bar{\mathfrak{g}})$ together with the Poisson bracket defined by extending formulas (3.32) to $S(r)$ via the Leibniz rule. This algebra is commonly known as the Weil algebra $W_{\mathfrak{g}}$ associated to \mathfrak{g} , [AM]. It is not hard to check that, in this case, the "quantum" and "classical" Zhu algebras at $\hbar = 0$, namely the bottom left and right corners in diagram (6.16), are both isomorphic to the Weil algebra $W_{\mathfrak{g}} = V^{clM}$. Applying the canonical map $\pi_Z^{cl} : V^{clFT} \rightarrow Zhu_H V^{clFT} = V^{clM} \simeq S(r)$ (see Section 6) to G^{cl} and L^{cl} , we obtain the "classical analogues" of the elements D and C in V^{QM} . As explained in Section 6, Theorem 6.4, the associative product in V^{clM} is induced by the associative product in V^{clFT} , while the Poisson bracket is induced by the bracket (6.3) with $\hbar = 1$. Hence we get, for $k + h^\vee = 1$,

$$\begin{aligned} D^{cl} &= \sum_i a^i \bar{a}^i + \frac{1}{3} \sum_{i,j} \overline{[a^i, a^j]} \bar{a}^i \bar{a}^j, \\ C^{cl} &= \frac{1}{2} \sum_i a^i a^i + \frac{1}{2} \sum_{i,j} \bar{a}^i [a^i, a^j] \bar{a}^j. \end{aligned}$$

With respect to the Poisson bracket, C^{cl} is central, $\{D^{cl}, a\} = 0$, $\{D^{cl}, \bar{a}\} = a$ and $\{D^{cl}, D^{cl}\} = 2C^{cl}$. Notice that $\text{ad } D^{cl}$ coincides with the Weil differential of the Weil algebra $W_{\mathfrak{g}}$ in its Koszul form.

4 Almost linear differentials in non-linear Lie algebras and Lie conformal algebras

Throughout this section we let Γ be a discrete additive subgroup of \mathbb{R} containing \mathbb{Z} , i.e. $\Gamma = \frac{1}{N}\mathbb{Z}$ for some positive integer N , and we let $\epsilon = \frac{1}{N}$. We also let Γ_+ (respectively Γ'_+) be the set of non-negative (resp. positive) elements of Γ .

By a *differential* d on a vector superspace U we mean an odd endomorphism of U such that $d^2 = 0$. Moreover, if U is endowed with one or more products (or non-linear products), like a Lie algebra or a Lie conformal algebra, we assume that d is an odd derivation of these products, and in the Lie conformal algebra case we also assume that d commutes with T . For example, in the non-linear Lie algebra case we require that, for $a, b \in U$, $d[a, b] = [da, b] + s(a)[a, db] \in \mathcal{T}(U)$.

4.1 Review of homological algebra: good filtered complexes

Let (U, d) be a filtered complex, namely U is a \mathbb{Z}_+ -graded vector superspace $U = \bigoplus_{n \geq 0} U^n$, with a compatible decreasing Γ -filtration $\{F^p U, p \in \Gamma\}$, i.e. $F^p U = \bigoplus_{n \geq 0} F^p U^n$, where $F^p U^n = F^p U \cap U^n$, and d is a differential of U of degree 1 preserving the filtration: $d(F^p U^n) \subset F^p U^{n+1}$.

The cohomology $H(U, d)$ of this complex is again a \mathbb{Z}_+ -graded and Γ -filtered

space, with

$$F^p H^n(U, d) = \text{Ker}(d : F^p U^n \rightarrow F^p U^{n+1}) / (\text{Im} d \cap F^p U^n) .$$

Its associated graded $\text{gr } H(U, d)$ is then the Γ -bigraded space with

$$\begin{aligned} \text{gr}^{pq} H(U, d) &= F^p H^{p+q}(U, d) / F^{p+\epsilon} H^{p+q}(U, d) \\ &\simeq \frac{F^p U^{p+q} \cap \text{Ker} d}{(F^{p+\epsilon} U^{p+q} \cap \text{Ker} d) + (F^p U^{p+q} \cap \text{Im} d)} . \end{aligned}$$

The associated graded complex $(\text{gr } U, d^{\text{gr}})$ is the Γ -bigraded space $\text{gr } U = \bigoplus_{p,q \in \Gamma} \text{gr}^{pq} U$, where $\text{gr}^{pq} U = F^p U^{p+q} / F^{p+\epsilon} U^{p+q}$, together with the induced differential d^{gr} . Its cohomology $H(\text{gr } U, d^{\text{gr}})$ is then the Γ -bigraded space given by

$$\begin{aligned} H^{pq}(\text{gr } U, d^{\text{gr}}) &= \frac{\text{Ker}(d^{\text{gr}} : \text{gr}^{pq} U \rightarrow \text{gr}^{p,q+1} U)}{\text{Im}(d^{\text{gr}} : \text{gr}^{p,q-1} U \rightarrow \text{gr}^{p,q} U)} \\ &\simeq \frac{F^p U^{p+q} \cap d^{-1}(F^{p+\epsilon} U^{p+q+1})}{F^{p+\epsilon} U^{p+q} + d(F^p U^{p+q-1})} . \end{aligned}$$

Definition 4.1. The complex (U, d) and the differential d are said to be *good* if $H^{pq}(\text{gr } U, d^{\text{gr}}) = 0$ for all $p, q \in \Gamma$ such that $p + q \neq 0$.

Lemma 4.2. Assume U is locally finite, namely for every $n \geq 0$ we have $F^p U^n = 0$ for $p \gg 0$. Then if (U, d) is a good complex we have

$$\text{gr}^{pq} H(U, d) \simeq H^{pq}(\text{gr } U, d^{\text{gr}}) , \quad \forall p, q \in \Gamma .$$

More precisely, we have the following identities ($p \in \Gamma$),

$$F^p U^n \cap \text{Ker} d = F^p U^n \cap \text{Im} d , \quad \text{for } n \neq 0 , \quad (4.1)$$

$$F^p U^0 \cap d^{-1}(F^{p+\epsilon} U^1) = F^{p+\epsilon} U^0 + F^p U^0 \cap \text{Ker} d . \quad (4.2)$$

Proof. The statement of the lemma is standard in the theory of spectral sequences. For completeness, we will provide an elementary proof of it. By assumption we have

$$F^p U^n \cap d^{-1}(F^{p+\epsilon} U^{n+1}) = F^{p+\epsilon} U^n + d(F^p U^{n-1}) , \quad \forall n \neq 0 .$$

Suppose first $n \neq 0$. In this case we have

$$\begin{aligned} F^p U^n \cap \text{Ker} d &= F^p U^n \cap d^{-1}(F^{p+\epsilon} U^{n+1}) \cap \text{Ker} d \\ &= F^{p+\epsilon} U^n \cap \text{Ker} d + d(F^p U^{n-1}) \\ &\subset F^{p+\epsilon} U^n \cap \text{Ker} d + F^p U^n \cap \text{Im} d . \end{aligned} \quad (4.3)$$

The reverse inclusion is obvious. Hence we have

$$F^p U^n \cap \text{Ker} d = F^{p+\epsilon} U^n \cap \text{Ker} d + F^p U^n \cap \text{Im} d .$$

We can now iterate the above result N times to get

$$F^p U^n \cap \text{Ker} d = F^{p+N\epsilon} U^n \cap \text{Ker} d + F^p U^n \cap \text{Im} d .$$

Equation (4.1) then follows by the locally finiteness assumption. We are left to prove (4.2). It follows by (4.3) with $n = 1$ that

$$\begin{aligned} d(F^p U^0) \cap F^{p+\epsilon} U^1 &= d(F^p U^0) \cap F^{p+\epsilon} U^1 \cap \text{Ker} d \\ &= d(F^{p+\epsilon} U^0) + d(F^p U^0) \cap F^{p+2\epsilon} U^1 . \end{aligned}$$

We can now repeat the same argument for the second term in the RHS to get, by induction,

$$d(F^p U^0) \cap F^{p+\epsilon} U^1 = d(F^{p+\epsilon} U^0) + d(F^p U^0) \cap F^{p+r\epsilon} U^1 , \quad \forall r \geq 1 .$$

By locally finiteness, this implies

$$d(F^p U^0) \cap F^{p+\epsilon} U^1 = d(F^{p+\epsilon} U^0) ,$$

which is equivalent to (4.2). \square

4.2 Lie algebras and Lie conformal algebras with a differential

We now review some basic facts about non-linear Lie algebras and non-linear Lie conformal algebras with a "linear" differential. In particular we show that taking cohomology commutes with taking the universal enveloping algebra. In the next section we will see how to generalize these results to the case of an almost linear differential. First, we recall here the well know Künneth Lemma:

Lemma 4.3 (Künneth Lemma). *Let V_1, V_2 be vector superspace with differentials $d_i : V_i \rightarrow V_i$, $i = 1, 2$, and define the differential d on $V = V_1 \otimes V_2$ by $d(a \otimes b) = d(a) \otimes b + s(a)a \otimes d(b)$. Then there is a canonical vector space isomorphism*

$$H(V, d) \simeq H(V_1, d_1) \otimes H(V_2, d_2) .$$

As usual, we denote by $\mathcal{T}(V)$ and $S(V)$ respectively the tensor algebra and the symmetric algebra of the vector superspace V . We have

Lemma 4.4. *Let V be a vector superspace with a differential d . Extend d to a derivation of $\mathcal{T}(V)$ and $S(V)$. Then*

$$H(\mathcal{T}(V), d) \simeq \mathcal{T}(H(V, d)) , \quad H(S(V), d) \simeq S(H(V, d)) .$$

Proof. The statement immediately follows by the Künneth Lemma and the fact that the action of the symmetric group S_n on $V^{\otimes n}$ commutes with the action of d . \square

We can think at V in the previous lemma as an abelian Lie algebra. The following result is then a generalization of Lemma 4.4 to the case in which the space V has a structure of a non-linear Lie algebra.

Theorem 4.5. *Let \mathfrak{g} be a non-linear Lie superalgebra, together with a differential $d : \mathfrak{g} \rightarrow \mathfrak{g}$ preserving the Γ'_+ -grading, i.e. $d(\mathfrak{g}_\zeta) \subset \mathfrak{g}_\zeta$ for $\zeta \in \Gamma'_+$. Consider the space $H(\mathfrak{g}, d)$ with induced Γ'_+ -grading. The non-linear Lie bracket of \mathfrak{g} induces a non-linear product*

$$H(\mathfrak{g}, d) \otimes H(\mathfrak{g}, d) \rightarrow \mathcal{T}(H(\mathfrak{g}, d)) ,$$

and we assume that this defines a structure of a non-linear Lie superalgebra on $H(\mathfrak{g}, d)$. Then there is a canonical associative algebra isomorphism

$$H(U(\mathfrak{g}), d) \simeq U(H(\mathfrak{g}, d)) .$$

Proof. First, let us describe the non-linear product $H(\mathfrak{g}, d) \otimes H(\mathfrak{g}, d) \xrightarrow{[\cdot, \cdot]} \mathcal{T}(H(\mathfrak{g}, d))$. By Lemma 4.4 we have an isomorphism $H(\mathcal{T}(\mathfrak{g}), d) \simeq \mathcal{T}(H(\mathfrak{g}, d))$. Then $[\cdot, \cdot]$ is defined by the following commutative diagram

$$\begin{array}{ccc} H(\mathfrak{g}, d) \otimes H(\mathfrak{g}, d) & \xrightarrow{[\cdot, \cdot]} & \mathcal{T}(H(\mathfrak{g}, d)) = H(\mathcal{T}(\mathfrak{g}), d) \\ \uparrow & & \uparrow \\ \text{Ker}(d : \mathfrak{g} \rightarrow \mathfrak{g})^{\otimes 2} & \longrightarrow & \text{Ker}(d : \mathcal{T}(\mathfrak{g}) \rightarrow \mathcal{T}(\mathfrak{g})) \end{array}$$

where the map on the bottom line is given by the non-linear Lie bracket on \mathfrak{g} . By assumption this makes $H(\mathfrak{g}, d)$ into a non-linear Lie superalgebra, hence its universal enveloping algebra is well defined.

We are left to show that $U(H(\mathfrak{g}, d)) \simeq H(U(\mathfrak{g}), d)$. We will use the same argument as in [KW1, Lemma 3.2]. The Γ'_+ -grading of \mathfrak{g} induces increasing Γ_+ -filtration $\{U_\zeta(\mathfrak{g}), \zeta \in \Gamma_+\}$ of $U(\mathfrak{g})$, and a Γ_+ -grading of the symmetric algebra $S(\mathfrak{g}) = \bigoplus_{\gamma \in \Gamma_+} S(\mathfrak{g})[\zeta]$. By the grading condition of the non-linear Lie bracket, we have an exact sequence

$$0 \rightarrow U_{\zeta-\epsilon}(\mathfrak{g}) \rightarrow U_\zeta(\mathfrak{g}) \rightarrow S(\mathfrak{g})[\zeta] \rightarrow 0 ,$$

which induces an exact triangle in cohomology,

$$\begin{array}{ccc} H(U_{\zeta-\epsilon}(\mathfrak{g}), d) & \longrightarrow & H(U_\zeta(\mathfrak{g}), d) \\ & \nwarrow \delta & \downarrow \\ & & H(S(\mathfrak{g})[\zeta], d) \end{array}$$

We claim that the connecting map δ is identically zero. Let us recall its definition. Given $w \in H(S(\mathfrak{g})[\zeta], d)$, let \tilde{w} be a representative of w in $S(\mathfrak{g})[\Delta]$, and let $\tilde{\tilde{w}}$ be any representative of \tilde{w} in $U_\zeta(\mathfrak{g})$. Clearly $d\tilde{w} = 0$, hence $d\tilde{\tilde{w}} \in U_{\zeta-\epsilon}(\mathfrak{g})$, and δw is the corresponding class in $H(U_{\zeta-\epsilon}(\mathfrak{g}), d)$. On the other hand, by Lemma 4.4 we can choose \tilde{w} to be a symmetric polynomial in the closed elements of \mathfrak{g} , and we let $\tilde{\tilde{w}}$ be the same polynomial, seen as an element of $U_\zeta(\mathfrak{g})$. But with this choice of $\tilde{\tilde{w}}$ we immediately have $d\tilde{\tilde{w}} = 0$, hence $\delta w = 0$ as well. We thus have exact sequences

$$0 \rightarrow H(U_{\zeta-\epsilon}(\mathfrak{g}), d) \rightarrow H(U_\zeta(\mathfrak{g}), d) \rightarrow H(S(\mathfrak{g})[\zeta], d) \rightarrow 0.$$

On the other hand, by Lemma 4.4, we have $H(S(\mathfrak{g})[\zeta], d) \simeq S(H(\mathfrak{g}, d))[\zeta]$. It then follows by induction that $H(U_\zeta(\mathfrak{g}), d) \simeq U_\zeta(H(\mathfrak{g}, d))$, which completes the proof. \square

Remark 4.6. Notice that, if \mathfrak{g} is a "linear" Lie algebra, then its cohomology $H(\mathfrak{g}, d)$ is automatically a Lie algebra. In the non-linear case, though, it is not clear if $H(\mathfrak{g}, d)$ is necessarily a non Lie algebra.

The above argument can be repeated almost verbatim to prove the analogue statement in the context of Lie conformal algebras. In other words we have the following

Theorem 4.7. *Let R be a non-linear Lie conformal algebra, together with a differential $d : R \rightarrow R$, preserving the Γ'_+ -grading of R . Assume moreover that $H(R, d)$, with the induced λ -bracket, is again a non-linear Lie conformal algebra. Then there is a canonical vertex algebra isomorphism*

$$H(V(R), d) \simeq V(H(R, d)).$$

4.3 Non-linear Lie algebras with a good almost linear differential

Let \mathfrak{g} be a finite-dimensional vector superspace graded as follows,

$$\mathfrak{g} = \bigoplus_{\substack{p, q \in \Gamma, \ p+q \in \mathbb{Z}_+ \\ \Delta \in \Gamma'_+}} \mathfrak{g}^{pq}[\Delta]. \quad (4.4)$$

The induced grading of the tensor algebra is $\mathcal{T}(\mathfrak{g}) = \bigoplus_{p, q \in \Gamma, \Delta \in \Gamma'_+} \mathcal{T}^{pq}(\mathfrak{g})[\Delta]$. We further assume \mathfrak{g} has a structure of non-linear Lie algebra (see Definition 3.1) with respect to the Γ'_+ -grading by subspaces $\mathfrak{g}[\Delta] = \bigoplus_{p, q} \mathfrak{g}^{pq}[\Delta]$, $\Delta \in \Gamma'_+$. In particular the non-linear Lie bracket satisfies the grading condition

$$[\mathfrak{g}[\Delta_1], \mathfrak{g}[\Delta_2]] \subset \bigoplus_{\Delta < \Delta_1 + \Delta_2} \mathcal{T}(\mathfrak{g})[\Delta], \quad (4.5)$$

or, equivalently, $\Delta([a, b]) < \Delta(a) + \Delta(b)$. Moreover we assume that the corresponding \mathbb{Z}_+ -grading of \mathfrak{g} , given by $\mathfrak{g}^n = \bigoplus_{p+q=n, \Delta} \mathfrak{g}^{pq}[\Delta]$, $n \in \mathbb{Z}_+$, is a

non-linear Lie algebra grading, and the decreasing Γ -filtration of \mathfrak{g} , given by $F^p \mathfrak{g} = \bigoplus_{p' \geq p, q, \Delta} \mathfrak{g}^{p'q}[\Delta]$, $p \in \Gamma$, is a non-linear Lie algebra filtration. The above assumptions can be combined by the following condition,

$$[\mathfrak{g}^{p_1 q_1}[\Delta_1], \mathfrak{g}^{p_2 q_2}[\Delta_2]] \subset \bigoplus_{\substack{l \geq 0 \\ \Delta < \Delta_1 + \Delta_2}} \mathcal{T}^{p_1 + p_2 + l, q_1 + q_2 - l}(\mathfrak{g})[\Delta]. \quad (4.6)$$

The associated graded $\text{gr } \mathfrak{g}$ of the filtration $\{F^p \mathfrak{g}, p \in \Gamma\}$ is the space \mathfrak{g} with non-linear Lie bracket $[a, b]^{\text{gr}}$, for $a \in \mathfrak{g}^{p_1 q_1}$, $b \in \mathfrak{g}^{p_2 q_2}$, equal to the component of $[a, b]$ in $\mathcal{T}^{p_1 + p_2, q_1 + q_2}(\mathfrak{g})$. It is easy to check that, by definition, $\text{gr } \mathfrak{g}$ is again a non-linear Lie algebra, bigraded by p, q .

Let $U(\mathfrak{g})$ be the universal enveloping associative algebra of the non-linear Lie algebra \mathfrak{g} . It has the \mathbb{Z}_+ -grading $U(\mathfrak{g}) = \bigoplus_{n \geq 0} U(\mathfrak{g})^n$, where

$$U(\mathfrak{g})^n = \text{span}_{\mathbb{C}} \left\{ a_1 \cdots a_s \mid a_i \in \mathfrak{g}^{n_i}, n_1 + \cdots + n_s = n \right\},$$

the decreasing Γ -filtration by subspaces

$$F^p U(\mathfrak{g}) = \text{span}_{\mathbb{C}} \left\{ a_1 \cdots a_s \mid a_i \in F^{p_i} \mathfrak{g}, p_1 + \cdots + p_s \geq p \right\},$$

and the increasing Γ_+ -filtration by subspaces

$$U_{\Delta}(\mathfrak{g}) = \text{span}_{\mathbb{C}} \left\{ a_1 \cdots a_s \mid a_i \in \mathfrak{g}[\Delta_i], \Delta_1 + \cdots + \Delta_s \leq \Delta \right\}.$$

We let $F^p U(\mathfrak{g})^n = F^p U(\mathfrak{g}) \cap U(\mathfrak{g})^n$ and $F^p U_{\Delta}(\mathfrak{g})^n = F^p U(\mathfrak{g}) \cap U(\mathfrak{g})^n \cap U_{\Delta}(\mathfrak{g})$. By construction, grading and filtrations of $U(\mathfrak{g})$ are compatible. Notice that each filtered space $U_{\Delta}(\mathfrak{g})$ is finite-dimensional, hence locally finite, i.e. for each $\Delta \in \Gamma_+$ and $n \geq 0$ we have $F^p U_{\Delta}(\mathfrak{g})^n = 0$ for $p \gg 0$. Moreover, by the grading condition (4.5), we have that

$$[U_{\Delta_1}(\mathfrak{g}), U_{\Delta_2}(\mathfrak{g})] \subset U_{\Delta}(\mathfrak{g}), \quad \text{for some } \Delta < \Delta_1 + \Delta_2. \quad (4.7)$$

The associated graded algebra $\text{gr } U(\mathfrak{g})$ is a Γ -bigraded associative algebra, with a compatible Γ_+ -filtration. Namely the Δ -filtered subspace of bidegree $p, q \in \Gamma$ is

$$\text{gr}^{pq} U_{\Delta}(\mathfrak{g}) = F^p U_{\Delta}(\mathfrak{g})^{p+q} / F^{p+\epsilon} U_{\Delta}(\mathfrak{g})^{p+q}.$$

Lemma 4.8. *We have a canonical isomorphism $\text{gr } U(\mathfrak{g}) \simeq U(\text{gr } \mathfrak{g})$ which preserves Γ -bigrading and Γ_+ -filtration, i.e.*

$$\text{gr}^{pq} U_{\Delta}(\mathfrak{g}) \simeq U_{\Delta}(\text{gr } \mathfrak{g})^{pq}.$$

In particular the natural embedding $\mathfrak{g} = \text{gr } \mathfrak{g} \subset \text{gr } U(\mathfrak{g})$ is such that $\mathfrak{g} \cap \text{gr}^{pq} U_{\Delta}(\mathfrak{g}) = \bigoplus_{\Delta' \leq \Delta} \mathfrak{g}^{pq}[\Delta']$.

Proof. Notice that $\mathfrak{g}^{pq} \subset F^p U(\mathfrak{g})^{p+q}$ and $\mathfrak{g}^{pq} \cap F^{p+\epsilon} U(\mathfrak{g})^{p+q} = 0$. We thus have a natural embedding $\mathfrak{g}^{pq} \subset \text{gr}^{pq} U(\mathfrak{g}) = F^p U(\mathfrak{g})^{p+q} / F^{p+\epsilon} U(\mathfrak{g})^{p+q}$, which induces an injective map $\text{gr } \mathfrak{g} \subset \text{gr } U(\mathfrak{g})$. By universal property we then have an associative algebra homomorphism $U(\text{gr } \mathfrak{g}) \rightarrow \text{gr } U(\mathfrak{g})$. This is a bijection since, by the PBW Theorem, it maps a basis of $U(\text{gr } \mathfrak{g})$ to a basis of $\text{gr } U(\mathfrak{g})$. The remaining part of the statement follows immediately. \square

Definition 4.9. Let $d : U(\mathfrak{g}) \rightarrow U(\mathfrak{g})$ be a differential of \mathbb{Z}_+ -degree 1, preserving both the Γ and the Γ_+ filtrations:

$$d(U(\mathfrak{g})^n) \subset U(\mathfrak{g})^{n+1}, \quad d(F^p U(\mathfrak{g})) \subset F^p U(\mathfrak{g}), \quad d(U_\Delta(\mathfrak{g})) \subset U_\Delta(\mathfrak{g}). \quad (4.8)$$

Let $d^{\text{gr}} : \text{gr } U(\mathfrak{g}) \rightarrow \text{gr } U(\mathfrak{g})$ be the corresponding associated graded differential. We say that d is an *almost linear differential* of \mathfrak{g} if d^{gr} preserves $\text{gr } \mathfrak{g} = \mathfrak{g}$, it has bidegree $(0, 1)$, and it preserves the Γ'_+ -grading of \mathfrak{g} , namely

$$d^{\text{gr}}(\mathfrak{g}^{pq}[\Delta]) \subset \mathfrak{g}^{p, q+1}[\Delta]. \quad (4.9)$$

In this case $d^{\text{gr}}|_{\mathfrak{g}}$ is a differential of the non-linear Lie algebra $\text{gr } \mathfrak{g}$, and the action of d^{gr} on $\text{gr } U(\mathfrak{g}) \simeq U(\text{gr } \mathfrak{g})$ is obtained starting with $d^{\text{gr}}|_{\mathfrak{g}}$ and applying the Leibniz rule. Moreover we say that d is a *good almost linear differential* of \mathfrak{g} if

$$\text{Ker}(d^{\text{gr}}|_{\mathfrak{g}^{pq}}) = \text{Im}(d^{\text{gr}}|_{\mathfrak{g}^{p, q-1}}), \quad \text{for } p+q \neq 0, \quad (4.10)$$

namely the cohomology $H^{pq}(\mathfrak{g}, d^{\text{gr}})$ is concentrated in degree $p+q=0$.

From now on we let d be a good almost linear differential of \mathfrak{g} .

Remark 4.10. An equivalent way of writing condition (4.9) in terms of the differential d is

$$d(\mathfrak{g}^{pq}[\Delta]) \subset \mathfrak{g}^{p, q+1}[\Delta] \oplus F^{p+\epsilon} U_\Delta(\mathfrak{g})^{p+q+1},$$

and $d^{\text{gr}}|_{\mathfrak{g}}$ corresponds to taking only the first component in this decomposition. Moreover, an equivalent way of writing condition (4.10) is

$$\mathfrak{g}^{pq}[\Delta] \cap d^{-1}(F^{p+\epsilon} U_\Delta(\mathfrak{g})^{p+q+1}) \subset d\mathfrak{g}^{p, q-1}[\Delta] + F^{p+\epsilon} U_\Delta(\mathfrak{g})^{p+q}, \quad \text{if } p+q \neq 0.$$

By assumption (4.9), $\text{gr } \mathfrak{g}$ is a bigraded filtered non-linear Lie algebra with differential d^{gr} . Hence its cohomology $H(\mathfrak{g}, d^{\text{gr}})$ is a space with induced bigrading and filtration,

$$H(\mathfrak{g}, d^{\text{gr}}) = \bigoplus_{\substack{p, q \in \Gamma, \ p+q \in \mathbb{Z}_+ \\ \Delta \in \Gamma'_+}} H^{pq}(\mathfrak{g}, d^{\text{gr}})[\Delta], \quad \text{where}$$

$$H^{pq}(\mathfrak{g}, d^{\text{gr}})[\Delta] = (\mathfrak{g}^{pq}[\Delta] \cap \text{Ker } d^{\text{gr}}) / d^{\text{gr}}(\mathfrak{g}^{p, q-1}[\Delta]).$$

The assumption that d^{gr} is good is equivalent to say that $H^{pq}(\mathfrak{g}, d^{\text{gr}})[\Delta] = 0$ if $p + q \neq 0$. Hence the above decomposition reduces to

$$H(\mathfrak{g}, d^{\text{gr}}) = \bigoplus_{p \in \Gamma, \Delta \in \Gamma'_+} H^{p, -p}(\mathfrak{g}, d^{\text{gr}})[\Delta], \quad \text{where}$$

$$H^{p, -p}(\mathfrak{g}, d^{\text{gr}})[\Delta] = \mathfrak{g}^{p, -p}[\Delta] \cap \text{Ker} d^{\text{gr}} = \mathfrak{g}^{p, -p}[\Delta] \cap d^{-1}(F^{p+\epsilon} U_{\Delta}(\mathfrak{g})^1).$$

In particular we have a natural embedding $H(\mathfrak{g}, d^{\text{gr}}) \subset \text{gr } \mathfrak{g}$. It follows that $H(\mathfrak{g}, d^{\text{gr}})$ has the induced structure of a non-linear Lie algebra. Therefore the complex $(\text{gr } \mathfrak{g}, d^{\text{gr}})$ satisfies all assumptions of Theorem 4.5 and we have a canonical isomorphism

$$H(\text{gr } U(\mathfrak{g}), d^{\text{gr}}) \simeq U(H(\text{gr } \mathfrak{g}, d^{\text{gr}})). \quad (4.11)$$

We now fix $\Delta \in \Gamma_+$ and we consider the complex $(U_{\Delta}(\mathfrak{g}), d)$. We want to show that it satisfies all the assumptions of Lemma 4.2. By the above observations, we only need to check that $(U_{\Delta}(\mathfrak{g}), d)$ is good in the sense of Definition 4.1, namely $H^{pq}(\text{gr } U_{\Delta}(\mathfrak{g}), d^{\text{gr}}) = 0$ for $p + q \neq 0$. For this we need the following

Lemma 4.11. *The differential d^{gr} is such that*

$$d^{\text{gr}}(\text{gr}^{pq} U(\mathfrak{g})) \cap \text{gr}^{p, q+1} U_{\Delta}(\mathfrak{g}) = d^{\text{gr}}(\text{gr}^{pq} U_{\Delta}(\mathfrak{g})).$$

Therefore there is a natural embedding

$$H^{pq}(\text{gr } U_{\Delta}(\mathfrak{g}), d^{\text{gr}}) \subset H^{pq}(\text{gr } U(\mathfrak{g}), d^{\text{gr}}).$$

Proof. By assumption on d we have $d(F^p U_{\Delta}(\mathfrak{g})^n) \subset F^p U_{\Delta}(\mathfrak{g})^{n+1}$. It immediately follows that $d^{\text{gr}}(\text{gr}^{pq} U_{\Delta}(\mathfrak{g})) \subset \text{gr}^{p, q+1} U_{\Delta}(\mathfrak{g})$. Hence we only have to show $d^{\text{gr}}(\text{gr}^{pq} U(\mathfrak{g})) \cap \text{gr}^{p, q+1} U_{\Delta}(\mathfrak{g}) \subset d^{\text{gr}}(\text{gr}^{pq} U_{\Delta}(\mathfrak{g}))$.

Let $A \in U(\text{gr } \mathfrak{g})^{pq} \simeq \text{gr}^{pq} U(\mathfrak{g})$ be such that $d^{\text{gr}}(A) \in U_{\Delta}(\text{gr } \mathfrak{g})^{p, q+1}$. We need to show that there is $A' \in U_{\Delta}(\text{gr } \mathfrak{g})^{pq}$ such that $d^{\text{gr}}(A) = d^{\text{gr}}(A')$. If $A \in U_{\Delta}(\text{gr } \mathfrak{g})^{pq}$, there is nothing to prove. Suppose then $\Delta_1 > \Delta$ is such that $A \in U_{\Delta_1}(\text{gr } \mathfrak{g})^{pq}$ and $A \notin U_{\Delta_1 - \epsilon}(\text{gr } \mathfrak{g})^{pq}$.

By the grading condition (4.5), the non-linear Lie algebra $\text{gr } \mathfrak{g}$ is filtered by Γ_+ , and the associated graded Lie algebra, that we denote $\text{gr}_{\Delta} \mathfrak{g}$, is the space \mathfrak{g} with trivial Lie bracket. Hence $\text{gr}_{\Delta} U(\text{gr } \mathfrak{g}) \simeq U(\text{gr}_{\Delta} \mathfrak{g}) = S(\mathfrak{g})$. We will denote by $X \mapsto \bar{X}$ the natural quotient map $U_{\Delta}(\text{gr } \mathfrak{g})^{pq} \twoheadrightarrow S(\mathfrak{g})[\Delta]^{pq} \simeq U_{\Delta}(\text{gr } \mathfrak{g})^{pq} / U_{\Delta - \epsilon}(\text{gr } \mathfrak{g})^{pq}$. By (4.9) d^{gr} is a differential of bidegree $(0, 1)$ of the space \mathfrak{g} preserving the Γ'_+ -grading. Hence d^{gr} extends, by the Leibniz rule, to a differential of $U(\text{gr}_{\Delta} \mathfrak{g}) = S(\mathfrak{g})$ preserving the Γ_+ -grading. It then follows by Lemma 4.4 that $H(S(\mathfrak{g}), d^{\text{gr}}) = S(H(\mathfrak{g}, d^{\text{gr}}))$ or, equivalently,

$$\begin{aligned} \text{Ker} \left(d^{\text{gr}} : S(\mathfrak{g})[\Delta]^{pq} \rightarrow S(\mathfrak{g})[\Delta]^{p, q+1} \right) &= \text{Im} \left(d^{\text{gr}} : S(\mathfrak{g})[\Delta]^{p, q-1} \rightarrow S(\mathfrak{g})[\Delta]^{pq} \right) \\ &+ S \left(\text{Ker} (d^{\text{gr}} : \mathfrak{g} \rightarrow \mathfrak{g}) \right) \cap S(\mathfrak{g})[\Delta]^{pq}, \quad \forall \Delta \in \Gamma_+. \end{aligned} \quad (4.12)$$

By assumption $A \in U_{\Delta_1}(\text{gr } \mathfrak{g})^{pq}$, $A \notin U_{\Delta_1 - \epsilon}(\text{gr } \mathfrak{g})^{pq}$ and $d^{\text{gr}}(A) \in U_{\Delta_1 - \epsilon}(\text{gr } \mathfrak{g})$. This is equivalent to say that $0 \neq \bar{A} \in S(\mathfrak{g})[\Delta_1]^{pq}$ and $d^{\text{gr}}(\bar{A}) = 0$. In other

words $\bar{A} \in \text{Ker}(d^{\text{gr}} : S(\mathfrak{g})[\Delta_1]^{pq} \rightarrow S(\mathfrak{g})[\Delta_1]^{p,q+1})$. By (4.12) we can then find $\bar{B} \in S(\mathfrak{g})[\Delta_1]^{p,q-1}$ and $\bar{C} \in S(\text{Ker}(d^{\text{gr}} : \mathfrak{g} \rightarrow \mathfrak{g})) \cap S(\mathfrak{g})[\Delta_1]^{pq}$ such that

$$\bar{A} = d^{\text{gr}}(\bar{B}) + \bar{C}.$$

Let then $B \in U_{\Delta_1}(\text{gr } \mathfrak{g})^{p,q-1}$ be any preimage of \bar{B} , and $C \in U_{\Delta_1}(\text{gr } \mathfrak{g})^{pq}$ be a preimage of \bar{C} polynomial in the elements of $\text{Ker}(d^{\text{gr}} : \mathfrak{g} \rightarrow \mathfrak{g})$, so that $d^{\text{gr}}(C) = 0$. We have

$$A_1 = A - d^{\text{gr}}(B) - C \in U_{\Delta_1-\epsilon}(\text{gr } \mathfrak{g})^{p,q}, \quad d^{\text{gr}}(A_1) = d^{\text{gr}}(A).$$

We can then repeat the same argument to conclude, by induction, that there is an element $A' \in U_{\Delta}(\text{gr } \mathfrak{g})^{p,q}$ such that $d^{\text{gr}}(A') = d^{\text{gr}}(A)$, thus completing the proof. The second part of the lemma follows immediately from the first part. \square

For every $p \in \Gamma$, $\Delta \in \Gamma'_+$ we fix a basis $\mathcal{A}_{p\Delta} = \{e_i, i \in \mathcal{I}_{p\Delta}\}$ of the space $H^{p,-p}(\text{gr } \mathfrak{g}, d^{\text{gr}})[\Delta] = \mathfrak{g}^{p,-p}[\Delta] \cap \text{Ker } d^{\text{gr}}$.

We also let $\mathcal{A} = \{e_i, i \in \mathcal{I}\} = \bigsqcup_{p\Delta} \mathcal{A}_{p\Delta}$, which is a basis of $H(\text{gr } \mathfrak{g}, d^{\text{gr}})$, and we fix a total ordering of the index set \mathcal{I} . We assume such basis is compatible with the parity, and we denote $p(i) = p(e_i) \in \mathbb{Z}/2\mathbb{Z}$. As in Section 3.1 we denote

$$\Sigma_{p\Delta} = \left\{ e_{i_1} \dots e_{i_s} \left| \begin{array}{l} i_k \in \mathcal{I}_{p_k \Delta_k}, i_1 \leq \dots \leq i_s \\ i_k < i_{k+1} \text{ if } p(i_k) = \bar{1} \\ \sum_k p_k = p, \sum_k \Delta_k \leq \Delta \end{array} \right. \right\},$$

moreover we let $\Sigma_p = \bigcup_{\Delta} \Sigma_{p\Delta}$, $\Sigma_{\Delta} = \bigsqcup_p \Sigma_{p\Delta}$ and $\Sigma = \bigcup_{p,\Delta} \Sigma_{p\Delta}$.

Lemma 4.12. (a) *The complex $(U_{\Delta}(\mathfrak{g}), d)$ is good in the sense of Definition 4.1.*

(b) *For $p \in \Gamma$, $\Sigma_{p\Delta}$ is a vector space basis of $H^{p,-p}(\text{gr } U_{\Delta}(\mathfrak{g}), d^{\text{gr}})$.*

Proof. Since, by (4.11), the non-linear Lie algebra $H(\text{gr } \mathfrak{g}, d^{\text{gr}})$ is concentrated on bidegrees p, q with $p + q = 0$, so must be its universal enveloping algebra,

$$U(H(\text{gr } \mathfrak{g}, d^{\text{gr}})) = \bigoplus_{p \in \Gamma} U(H(\text{gr } \mathfrak{g}, d^{\text{gr}}))^{p,-p}.$$

By (4.11) we have a natural isomorphism $H(U(\text{gr } \mathfrak{g}), d^{\text{gr}}) \simeq U(H(\text{gr } \mathfrak{g}, d^{\text{gr}}))$. Hence, by looking separately at each bidegree $p, q \in \Gamma$, we get

$$\begin{aligned} H^{pq}(U(\text{gr } \mathfrak{g}), d^{\text{gr}}) &= 0, \quad \text{if } p + q \neq 0, \\ H^{p,-p}(U(\text{gr } \mathfrak{g}), d^{\text{gr}}) &\simeq U(H(\text{gr } \mathfrak{g}, d^{\text{gr}}))^{p,-p}. \end{aligned} \quad (4.13)$$

Thanks to Lemma 4.11, there is a natural embedding $H^{pq}(\text{gr } U_{\Delta}(\mathfrak{g}), d^{\text{gr}}) \subset H^{pq}(\text{gr } U(\mathfrak{g}), d^{\text{gr}})$, so that part (a) follows immediately from the first equation above. By comparing the Γ_+ -filtration in both sides of (4.13) we get

$$H^{p,-p}(\text{gr } U_{\Delta}(\mathfrak{g}), d^{\text{gr}}) = U_{\Delta}(\text{gr } \mathfrak{g})^{p,-p} \cap \text{Ker } d^{\text{gr}} = U_{\Delta}(H(\text{gr } \mathfrak{g}, d^{\text{gr}}))^{p,-p}.$$

The second part of the lemma follows by the above identity and the PBW Theorem for (filtered) non-linear Lie algebras. \square

So far we showed that the complex $(U_\Delta(\mathfrak{g}), d)$ satisfies the assumptions of Lemma 4.2. We thus have, from (4.1) and (4.2),

$$F^p U_\Delta(\mathfrak{g})^n \cap \text{Kerd} = F^p U_\Delta(\mathfrak{g})^n \cap \text{Im} d, \quad \text{for } n \neq 0, \quad (4.14)$$

$$F^p U_\Delta(\mathfrak{g})^0 \cap d^{-1}(F^{p+\epsilon} U_\Delta(\mathfrak{g})^1) = F^{p+\epsilon} U_\Delta(\mathfrak{g})^0 + F^p U_\Delta(\mathfrak{g})^0 \cap \text{Kerd}. \quad (4.15)$$

By assumption $e_i \in \mathfrak{g}^{p,-p}[\Delta] \cap \text{Kerd}^{\text{gr}} \subset F^p U_\Delta(\mathfrak{g})^0 \cap d^{-1}(F^{p+\epsilon} U_\Delta(\mathfrak{g})^1)$, for $i \in \mathcal{I}_{p\Delta}$. Thanks to (4.15) we can then find elements $\epsilon_i \in F^{p+\epsilon} U_\Delta(\mathfrak{g})^0$ and $E_i \in F^p U_\Delta(\mathfrak{g})^0 \cap \text{Kerd}$ such that $E_i = e_i + \epsilon_i$. We will assume E_i (hence ϵ_i) has the same parity as e_i . Recall that $H^0(U(\mathfrak{g}), d) = U(\mathfrak{g})^0 \cap \text{Kerd}$, so that $E_i \in H^0(U(\mathfrak{g}), d)$, $\forall i \in \mathcal{I}$.

We then let $\mathcal{B}_{p\Delta} = \{E_i, i \in \mathcal{I}_{p\Delta}\}$ for every $p \in \Gamma$, $\Delta \in \Gamma'_+$, and $\mathcal{B} = \{E_i, i \in \mathcal{I}\} = \bigsqcup_{p\Delta} \mathcal{B}_{p\Delta}$. Moreover we define $H^{p,-p}(\mathfrak{g}, d)[\Delta]$ and $H(\mathfrak{g}, d)$ as the subspace of $H^0(U(\mathfrak{g}), d)$ with basis $\mathcal{B}_{p\Delta}$ and \mathcal{B} respectively:

$$H(\mathfrak{g}, d) = \bigoplus_{p, \Delta} H^{p,-p}(\mathfrak{g}, d)[\Delta], \quad H^{p,-p}(\mathfrak{g}, d)[\Delta] = \text{span}_{\mathbb{C}} \mathcal{B}_{p\Delta}. \quad (4.16)$$

Finally we introduce, as above, the collection of ordered monomials in the elements E_i ,

$$S_{p\Delta} = \left\{ E_{i_1} \dots E_{i_s} \left| \begin{array}{l} i_k \in \mathcal{I}_{p_k \Delta_k}, \quad i_1 \leq \dots \leq i_s \\ i_k < i_{k+1} \text{ if } p(i_k) = \bar{1} \\ \sum_k p_k = p, \quad \sum_k \Delta_k \leq \Delta \end{array} \right. \right\},$$

and we let $S_p = \bigcup_{\Delta} S_{p\Delta}$, $S_\Delta = \bigsqcup_p S_{p\Delta}$ and $S = \bigcup_{p, \Delta} S_{p\Delta}$.

Lemma 4.13. (a) *The cohomology of the complex $(U(\mathfrak{g}), d)$ is concentrated in degree 0: $H^n(U(\mathfrak{g}), d) = 0$, for $n \neq 0$.*

(b) *The 0th cohomology $H^0(U(\mathfrak{g}), d)$ is PBW generated by $H(\mathfrak{g}, d)$. In particular S is a basis of $H^0(U(\mathfrak{g}), d)$ compatible with the Γ and Γ_+ filtrations of $H^0(U(\mathfrak{g}), d)$.*

Proof. Let $A \in U(\mathfrak{g})^n \cap \text{Kerd}$, $n \neq 0$. For some $p \in \Gamma$ and $\Delta \in \Gamma_+$ we have $A \in F^p U_\Delta(\mathfrak{g})^n$, hence, by (4.14), it follows that $A \in F^p U_\Delta(\mathfrak{g})^n \cap \text{Im} d$, namely $\bar{A} = 0$ in $H^n(U(\mathfrak{g}), d) = 0$. This proves the first part of the lemma.

For the second part, we first show that the ordered monomials $E_{i_1} \dots E_{i_s} \in S_{p\Delta}$ span the space $F^p U_\Delta(\mathfrak{g})^0 \cap \text{Kerd}$. For $A \in F^p U_\Delta(\mathfrak{g})^0 \cap \text{Kerd}$ we let $[A]$ be the corresponding element in $\text{gr}^{p,-p} U_\Delta(\mathfrak{g}) = F^p U_\Delta(\mathfrak{g})^0 / F^{p+\epsilon} U_\Delta(\mathfrak{g})^0$. Notice that $d^{\text{gr}}[A] = [dA] = 0$, so that $[A] \in \text{gr}^{p,-p} U_\Delta(\mathfrak{g}) \cap \text{Kerd}^{\text{gr}} = H^{p,-p}(\text{gr } U_\Delta(\mathfrak{g}), d^{\text{gr}})$. It then follows by the second part of Lemma 4.12 that

$$A = A' + \sum_{\vec{i}} c_{\vec{i}} e_{i_1} \dots e_{i_s},$$

where $A' \in F^{p+\epsilon} U_\Delta(\mathfrak{g})^0$ and the sum in the right hand side is a linear combination of ordered monomials $e_{i_1} \dots e_{i_s} \in \Sigma_{p\Delta}$. Recall that, by assumption, for

$i \in \mathcal{I}_{p\Delta}$, we have $E_i = e_i + \epsilon_i \in F^p U_\Delta(\mathfrak{g})^0 \cap \text{Kerd}$ with $\epsilon_i \in F^{p+\epsilon} U_\Delta(\mathfrak{g})^0$. We thus have

$$A_1 = A - \sum_{\vec{i}} c_{\vec{i}} E_{i_1} \dots E_{i_s} \in F^{p+\epsilon} U_\Delta(\mathfrak{g})^0 \cap \text{Kerd}.$$

After repeating the same argument N times we get, by induction, a sequence of elements

$$A_N = A - \sum_{\vec{i}} c_{\vec{i}} E_{i_1} \dots E_{i_s} \in F^{p+N\epsilon} U_\Delta(\mathfrak{g})^0 \cap \text{Kerd},$$

where now the indices $i_k \in \mathcal{I}_{p_k, \Delta_k}$ are such that $p \leq p_1 + \dots + p_s \leq p + N\epsilon$, $\Delta_1 + \dots + \Delta_s \leq \Delta$. Hence $A_N = 0$ by locally finiteness.

We are left to show that the ordered monomials $E_{i_1} \dots E_{i_s} \in S$ are linearly independent. For this, suppose we have a linear combination which is zero,

$$A = \sum_{\vec{i}} c_{\vec{i}} E_{i_1} \dots E_{i_s} = 0, \quad (4.17)$$

where the sum is over "ordered" indices $\vec{i} = (i_1, \dots, i_s)$, $i_k \in \mathcal{I}_{p_k, \Delta_k}$. We can assume the coefficients $c_{\vec{i}}$ are not all zero, and we let

$$p = \min_{\vec{i} \mid c_{\vec{i}} \neq 0} \{p_1 + \dots + p_s\}.$$

We then get by (4.17)

$$A = \sum_{\vec{i} \mid p_1 + \dots + p_s = p} c_{\vec{i}} e_{i_1} \dots e_{i_s} + R = 0,$$

with $R \in F^{p+\epsilon} U(\mathfrak{g})^0$. We can now look at the image $[A]$ of A in $\text{gr}^{p, -p} U(\mathfrak{g})^0 = F^p U(\mathfrak{g})^0 / F^{p+\epsilon} U(\mathfrak{g})^0$. We have

$$[A] = \sum_{\vec{i} \mid p_1 + \dots + p_s = p} c_{\vec{i}} e_{i_1} \dots e_{i_s} = 0. \quad (4.18)$$

By the second part of Lemma 4.12, the ordered monomials $e_{i_1} \dots e_{i_s}$ are linearly independent in $\text{gr}^{p, -p} U(\mathfrak{g})$. Hence (4.18) implies $c_{\vec{i}} = 0$ for all \vec{i} such that $p_1 + \dots + p_s = p$, which is a contradiction. \square

By Lemma 4.13 the associative algebra $H = H^0(U\mathfrak{g}, d)$ is PBW generated by the space $\mathfrak{h} = H(\mathfrak{g}, d)$. By definition (4.16), \mathfrak{h} is Γ'_+ -graded with $\mathfrak{h}[\Delta] = \text{span}_{\mathbb{C}} S_\Delta$. Moreover, by the second part of Lemma 4.13, the natural Γ_+ -filtration $U_\Delta(\mathfrak{g})^0 \cap \text{Kerd} \subset H$ coincides with the Γ_+ -filtration H_Δ induced by the generating space \mathfrak{h} as in (3.1). By the grading condition (4.7), we have $[\mathfrak{h}[\Delta_1], \mathfrak{h}[\Delta_2]] \subset H_\Delta$, for some $\Delta < \Delta_1 + \Delta_2$, namely H is "pregraded" by Γ_+ , in the sense of Definition 3.4. In conclusion, all the assumptions of Theorem 3.6 hold, so that we have

Theorem 4.14. *The space $H(\mathfrak{g}, d)$ admits a structure of non-linear Lie superalgebra such that its universal enveloping associative algebra is canonically isomorphic to $H(U(\mathfrak{g}), d)$:*

$$H(U(\mathfrak{g}), d) \simeq U(H(\mathfrak{g}, d)).$$

4.4 Non-linear Lie conformal algebras with a good almost linear differential

In this section we want to derive the same results as in the previous section in the context of Lie conformal superalgebras.

Let R be a non-linear Lie conformal superalgebra, freely generated as a $\mathbb{C}[T]$ -module by a finite-dimensional subspace \mathfrak{g} : $R = \mathbb{C}[T] \otimes \mathfrak{g}$. We shall assume that \mathfrak{g} is graded as in (4.4). This grading induces a vector space grading of R , by assuming T to have Γ -bidegree $(0, 0)$ and Γ_+ -degree 1:

$$R = \bigoplus_{\substack{p, q \in \Gamma, \ p+q \in \mathbb{Z}_+ \\ \Delta \in \Gamma'_+}} R^{pq}[\Delta] \ , \quad R^{pq}[\Delta] = \bigoplus_{n \geq 0} T^n \mathfrak{g}^{pq}[\Delta - n] \ , \quad (4.19)$$

and of the tensor algebra $\mathcal{T}(R)$. We will assume that R , as a (non-linear) Lie conformal algebra, is \mathbb{Z}_+ -graded by $\mathbb{C}[T]$ -submodules $R^n = \bigoplus_{\substack{p, q, \Delta \\ p+q=n}} R^{pq}[\Delta]$, $n \in \mathbb{Z}_+$, it is Γ -filtered by $F^p R = \bigoplus_{p' \geq p, q, \Delta} R^{p', q}[\Delta]$, $p \in \Gamma$, and moreover we assume that the Γ'_+ -grading by subspaces $R[\Delta] = \bigoplus_{p, q} R^{pq}[\Delta]$, $\Delta \in \Gamma'_+$, is a conformal weight of R . All these conditions mean the following relation

$$R^{p_1, q_1}[\Delta_1]_{(n)} R^{p_2, q_2}[\Delta_2] \subset \bigoplus_{l \geq 0} \mathcal{T}^{p_1+p_2+l, q_1+q_2-l}(R)[\Delta_1 + \Delta_2 - n - 1] \ , \quad n \in \mathbb{Z}_+ \ . \quad (4.20)$$

The associated graded non-linear Lie conformal algebra $\text{gr } R$ is defined as the $\mathbb{C}[T]$ -module R with λ -bracket $[a \lambda b]^{\text{gr}}$, for $a \in R^{p_1, q_1}$, $b \in R^{p_2, q_2}$, equal to the component of $[a \lambda b]$ in $R^{p_1+q_1, p_2+q_2}[\lambda]$. It is not hard to check that $\text{gr } R$ is a Γ -bigraded non-linear Lie conformal algebra, with a conformal weight Δ .

Let $V(R)$ be the universal enveloping vertex algebra of R . Recall that on the vertex algebra $V(R) = \mathcal{T}(R)/\mathcal{M}(R)$ the λ -bracket $[a \lambda b]$ of elements $a, b \in R$ is induced by the non-linear λ -bracket in R , and the normally ordered product $: aB :$ with $a \in R$, $B \in V(R)$ is induced by the associative product in $\mathcal{T}(R)$. These two conditions, together with the axioms of vertex algebra, uniquely define a vertex algebra structure on $V(R)$. Since all axioms of vertex algebra are "homogeneous" with respect to the gradings and the filtration of R , it is not hard to check that the vertex algebra $V(R)$ has induced \mathbb{Z}_+ -grading $V(R) = \bigoplus_{n \geq 0} V(R)^n$, Γ -filtration $F^p V(R) \subset V(R)$, and Γ_+ -grading $V(R) = \bigoplus_{\Delta \in \Gamma'_+} V(R)[\Delta]$, and all are additive with respect to the normally ordered product. Explicitly:

$$\begin{aligned} V(R)^n &= \text{span}_{\mathbb{C}} \left\{ : a_{i_1} \cdots a_{i_s} : \mid a_{i_k} \in R^{p_k, q_k}[\Delta_k], \sum_k (p_k + q_k) = n \right\} \ , \\ F^p V(R) &= \text{span}_{\mathbb{C}} \left\{ : a_{i_1} \cdots a_{i_s} : \mid a_{i_k} \in R^{p_k, q_k}[\Delta_k], \sum_k p_k \geq p \right\} \ , \\ V(R)[\Delta] &= \text{span}_{\mathbb{C}} \left\{ : a_{i_1} \cdots a_{i_s} : \mid a_{i_k} \in R^{p_k, q_k}[\Delta_k], \sum_k \Delta_k = \Delta \right\} \ . \end{aligned}$$

The vertex algebra structure is compatible with the above gradings and filtrations, in the sense that, for each $n \in \mathbb{Z}$, we have:

$$F^{p_1}V(R)^{n_1}[\Delta_1]_{(n)}F^{p_2}V(R)^{n_2}[\Delta_2] \subset F^{p_1+p_2}V(R)^{n_1+n_2}[\Delta_1 + \Delta_2 - n - 1] . \quad (4.21)$$

Notice that, since $\Delta(T) = 1$, each space $V(R)^n[\Delta]$ is finite dimensional, hence the corresponding filtration $F^pV(R)^n[\Delta]$ is locally finite. We have the associated graded vertex algebra, defined by

$$\text{gr } V(R) = \bigoplus_{p,q,\Delta} \text{gr}^{pq}V(R)[\Delta] , \quad \text{gr}^{pq}V(R)[\Delta] = F^pV(R)^n[\Delta]/F^{p+\epsilon}V(R)^n[\Delta] ,$$

with induced vertex algebra structure, which is a Γ -bigraded vertex algebra and it has a Γ_+ -valued conformal weight.

The analogue of Lemma 4.8 holds also in this context, namely we have

Lemma 4.15. *There is a canonical isomorphism $\text{gr } V(R) \simeq V(\text{gr } R)$ which preserves all gradings, namely such that*

$$\text{gr}^{pq}V(R)[\Delta] \simeq V^{pq}(\text{gr } R)[\Delta] .$$

In particular the natural embedding $R = \text{gr } R \subset \text{gr } V(R)$ is such that $R \cap \text{gr}^{pq}V(R)[\Delta] = R^{pq}[\Delta]$.

Definition 4.16. Let $d : V(R) \rightarrow V(R)$ be a differential of the vertex algebra $V(R)$. We assume that d has \mathbb{Z}_+ -degree +1, that it preserves the Γ -filtration and the conformal weight Δ . In other words

$$d(F^pV(R)^n[\Delta]) \subset F^pV(R)^{n+1}[\Delta] . \quad (4.22)$$

As before, we denote by $d^{\text{gr}} : \text{gr } V(R) \rightarrow \text{gr } V(R)$ the induced differential of the associated graded vertex algebra. It clearly has bidegree $(0, 1)$, it preserves the conformal weight Δ , and it is an odd derivation of all n^{th} products of $\text{gr } V(R)$. We say that d is an *almost linear differential* of the non-linear Lie conformal algebra R if d^{gr} preserves $\mathfrak{g} \subset \text{gr } V(R)$, namely $d^{\text{gr}}(\mathfrak{g}^{pq}[\Delta]) \subset \mathfrak{g}^{p,q+1}[\Delta]$; or equivalently, in terms of d ,

$$d(\mathfrak{g}^{pq}[\Delta]) \subset \mathfrak{g}^{p,q+1}[\Delta] \oplus F^{p+\epsilon}V(R)^{p+q+1}[\Delta] . \quad (4.23)$$

In particular $d^{\text{gr}}|_R$ is a differential of the non-linear Lie conformal algebra $\text{gr } R = \mathbb{C}[T]\mathfrak{g}$, and d^{gr} is the induced differential of the universal enveloping vertex algebra $V(\text{gr } R) \simeq \text{gr } V(R)$.

The corresponding cohomology is $H(\text{gr } R, d^{\text{gr}}) = \mathbb{C}[T]H(\mathfrak{g}, d^{\text{gr}})$, with $H(\mathfrak{g}, d^{\text{gr}}) = \bigoplus_{p,q,\Delta} H^{pq}(\mathfrak{g}, d^{\text{gr}})[\Delta] = \bigoplus_{p,q,\Delta} \text{Ker}(d^{\text{gr}}|_{\mathfrak{g}^{pq}[\Delta]})/\text{Im}(d^{\text{gr}}|_{\mathfrak{g}^{p,q-1}[\Delta]})$. We say that d is a *good almost linear differential* of R if the cohomology $H(R, d^{\text{gr}})$ (or equivalently $H(\mathfrak{g}, d^{\text{gr}})$) is concentrated at \mathbb{Z}_+ -degree zero, namely $H^{pq}(\mathfrak{g}, d^{\text{gr}}) = 0$ if $p+q \neq 0$, or equivalently, in terms of d ,

$$\mathfrak{g}^{pq}[\Delta] \cap d^{-1}(F^{p+\epsilon}V(R)^{p+q+1}[\Delta]) \subset d(\mathfrak{g}^{p,q-1}[\Delta]) + F^{p+\epsilon}V(R)^{p+q}[\Delta] . \quad (4.24)$$

It follows that $H(\text{gr } R, d^{\text{gr}})$ has the decomposition

$$H(\text{gr } R, d^{\text{gr}}) = \bigoplus_{p, \Delta} H^{p, -p}(\text{gr } R, d^{\text{gr}})[\Delta] , \quad \text{where} \quad (4.25)$$

$$H^{p, -p}(\text{gr } R, d^{\text{gr}})[\Delta] = R^{p, -p}[\Delta] \cap \text{Ker } d^{\text{gr}} ,$$

in particular it is embedded in $\text{gr } R$, hence it has the induced structure of a non-linear Lie conformal algebra. We can thus apply Theorem 4.7 to conclude

$$H(\text{gr } V(R), d^{\text{gr}}) \simeq V(H(\text{gr } R, d^{\text{gr}})) . \quad (4.26)$$

From now on we let d be a good almost linear differential of R .

As we did before, for every $p \in \Gamma$, $\Delta \in \Gamma'_+$, we fix a basis $\mathcal{A}_{p\Delta} = \{e_i, i \in \mathcal{I}_{p\Delta}\}$ of $\mathfrak{g}^{p, -p}[\Delta] \cap \text{Ker } d^{\text{gr}}$, and we let $\mathcal{A} = \{e_i, i \in \mathcal{I}\} = \bigsqcup_{p\Delta} \mathcal{A}_{p\Delta}$. We extend these bases to $\tilde{\mathcal{A}}_{p\Delta} = \{e_{\tilde{\mathbf{i}}}, \tilde{\mathbf{i}} \in \tilde{\mathcal{I}}_{p\Delta}\}$, a basis of $H^{p, -p}(\text{gr } R, d^{\text{gr}})[\Delta] = R^{p, -p}[\Delta] \cap \text{Ker } d^{\text{gr}}$, and $\tilde{\mathcal{A}} = \{e_{\tilde{\mathbf{i}}}, \tilde{\mathbf{i}} \in \tilde{\mathcal{I}}\} = \bigsqcup_{p\Delta} \tilde{\mathcal{A}}_{p\Delta}$, a basis of $H(\text{gr } R, d^{\text{gr}})$. For example we can take $\tilde{\mathcal{I}}_{p\Delta} = \bigsqcup_{k \geq 0} (\mathcal{I}_{p, \Delta-k}, k)$ and $e_{\tilde{\mathbf{i}}} = T^{(k)} e_i$ for $\tilde{\mathbf{i}} = (i, k) \in (\mathcal{I}_{p, \Delta-k}, k) \subset \tilde{\mathcal{I}}_{p, \Delta}$. We then assume $\tilde{\mathcal{I}}$ is totally ordered and we denote

$$\tilde{\Sigma}_{p\Delta} = \left\{ : e_{\tilde{\mathbf{i}}_1} \dots e_{\tilde{\mathbf{i}}_s} : \left| \begin{array}{l} \tilde{\mathbf{i}}_k \in \tilde{\mathcal{I}}_{p_k \Delta_k}, \tilde{\mathbf{i}}_1 \leq \dots \leq \tilde{\mathbf{i}}_s \\ \tilde{\mathbf{i}}_k < \tilde{\mathbf{i}}_{k+1} \text{ if } p(\tilde{\mathbf{i}}_k) = \tilde{\mathbf{i}} \\ \sum_k p_k = p, \sum_k \Delta_k = \Delta \end{array} \right. \right\} .$$

We also let $\tilde{\Sigma}_p = \bigsqcup_{\Delta} \tilde{\Sigma}_{p\Delta}$, $\tilde{\Sigma}_{\Delta} = \bigsqcup_p \tilde{\Sigma}_{p\Delta}$ and $\tilde{\Sigma} = \bigsqcup_{p, \Delta} \tilde{\Sigma}_{p\Delta}$.

Lemma 4.17. (a) For $p, q \in \Gamma$ such that $p+q \neq 0$ we have $H^{pq}(\text{gr } V(R), d^{\text{gr}}) = 0$. Hence, for every $\Delta \in \Gamma_+$, the complex $(V(R)[\Delta], d)$ is good.

(b) For $p \in \Gamma$, $\Delta \in \Gamma_+$, the set $\tilde{\Sigma}_{p\Delta}$ is a basis of $H^{p, -p}(\text{gr } V(R), d^{\text{gr}})[\Delta]$.

Proof. By Theorem 4.7 and Lemma 4.15 we have an isomorphism $H(\text{gr } V(R), d^{\text{gr}}) \simeq V(H(\text{gr } R, d^{\text{gr}}))$. Moreover by assumption (4.25), $H(\text{gr } R, d^{\text{gr}})$ is concentrated in zero \mathbb{Z}_+ -degree, so must be its enveloping vertex algebra $V(H(\text{gr } R, d^{\text{gr}}))$. We thus have $H^{pq}(\text{gr } V(R), d^{\text{gr}}) = 0$ if $p+q \neq 0$, and $H^{p, -p}(\text{gr } V(R), d^{\text{gr}}) \simeq V^{p, -p}(H(\text{gr } R, d^{\text{gr}}))$. Part (a) of the lemma follows immediately by the fact that d^{gr} preserves the conformal weight Δ . Part (b) follows by (4.26) and the PBW Theorem. \square

By the first part of Lemma 4.17, the complex $(V(R)[\Delta], d)$ (which is finite dimensional, hence locally finite) satisfies all the assumptions of Lemma 4.2. We thus have by (4.1) and (4.2)

$$F^p V(R)^n[\Delta] \cap \text{Ker } d = F^p V(R)^n[\Delta] \cap \text{Im } d , \quad \text{for } n \neq 0 , \quad (4.27)$$

$$F^p V(R)^0[\Delta] \cap d^{-1}(F^{p+\epsilon} V(R)^1[\Delta]) = F^{p+\epsilon} V(R)^0[\Delta] \quad (4.28)$$

$$+ F^p V(R)^0[\Delta] \cap \text{Ker } d .$$

Thanks to (4.28), for each $i \in \mathcal{I}_{p\Delta}$ we can write $e_i = E_i - \epsilon_i \in \mathfrak{g}^{p,-p}[\Delta] \cap \text{Ker} d^{\text{gr}}$, with $\epsilon_i \in F^{p+\epsilon}V(R)^0[\Delta]$ and $E_i \in F^pV(R)^0[\Delta] \cap \text{Ker} d$. Moreover, for $\tilde{i} = (i, k) \in \tilde{\mathcal{I}}_{p\Delta}$, we let $E_{\tilde{i}} = T^{(k)}E_i$ and $\epsilon_{\tilde{i}} = T^{(k)}\epsilon_i$, so that again $e_{\tilde{i}} = E_{\tilde{i}} - \epsilon_{\tilde{i}}$. We then let

$$\begin{aligned} H^{p,-p}(\mathfrak{g}, d)[\Delta] &= \text{span}_{\mathbb{C}}\{E_i, i \in \mathcal{I}_{p\Delta}\} \subset H(\mathfrak{g}, d) = \bigoplus_{p, \Delta} H^{p,-p}(\mathfrak{g}, d)[\Delta], \\ H^{p,-p}(R, d)[\Delta] &= \text{span}_{\mathbb{C}}\{E_{\tilde{i}}, \tilde{i} \in \tilde{\mathcal{I}}_{p\Delta}\} \subset H(R, d) = \bigoplus_{p, \Delta} H^{p,-p}(R, d)[\Delta]. \end{aligned}$$

Finally we define the collection of ordered monomials in the elements $E_{\tilde{i}}$,

$$\tilde{S}_{p\Delta} = \left\{ : E_{\tilde{i}_1} \dots E_{\tilde{i}_s} : \left| \begin{array}{l} \tilde{i}_k \in \tilde{\mathcal{I}}_{p_k \Delta_k}, \tilde{i}_1 \leq \dots \leq \tilde{i}_s \\ \tilde{i}_k < \tilde{i}_{k+1} \text{ if } p(\tilde{i}_k) = \bar{1} \\ \sum_k p_k = p, \sum_k \Delta_k = \Delta \end{array} \right. \right\},$$

and we let $\tilde{S}_p = \bigsqcup_{\Delta} \tilde{S}_{p\Delta}$, $\tilde{S}_{\Delta} = \bigsqcup_p \tilde{S}_{p\Delta}$ and $\tilde{S} = \bigsqcup_{p, \Delta} \tilde{S}_{p\Delta}$. The analogue of Lemma 4.13 holds also in the vertex algebra set up, with the same proof (using Lemma 4.17 instead of Lemma 4.12). We thus have the following

Lemma 4.18. (a) *The cohomology of the complex $(V(R), d)$ is concentrated in degree 0: $H^n(V(R), d) = 0$, for $n \neq 0$.*

(b) *$H^0(V(R), d)$ is freely generated by $H(R, d) = \mathbb{C}[T]H(\mathfrak{g}, d)$. In particular \tilde{S} is a basis of $H^0(V(R), d)$ compatible with the Γ -filtration and the conformal weight.*

Theorem 4.19. (a) *The space $H(R, d) = \mathbb{C}[T]H(\mathfrak{g}, d)$ admits a structure of non-linear Lie conformal algebra whose enveloping vertex algebra is canonically isomorphic to $H(V(R), d)$:*

$$H(V(R), d) \simeq V(H(R, d)).$$

(b) *The decomposition $H(\mathfrak{g}, d) = \bigoplus_{p, \Delta} H^{p,-p}(\mathfrak{g}, d)[\Delta]$, given by the subspaces $H^{p,-p}(\mathfrak{g}, d)[\Delta] = \text{span}_{\mathbb{C}}\{E_i, i \in \mathcal{I}_{p\Delta}\}$, induces a vertex algebra Γ -filtration and a Γ_+ -valued conformal weight Δ on $H(V(R), d)$, given by the subspaces $F^p H(V(R), d)[\Delta] = F^p V(R)^0[\Delta] \cap \text{Ker} d$. In other words we have*

$$\begin{aligned} &F^{p_1} H(V(R), d)[\Delta_1]_{(n)} F^{p_2} H(V(R), d)[\Delta_2] \\ &\subset F^{p_1+p_2} H(V(R), d)[\Delta_1 + \Delta_2 - n - 1]. \end{aligned}$$

Proof. By Lemma 4.18, the vertex algebra $H(V(R), d)$ is freely generated by the free $\mathbb{C}[T]$ -module $H(R, d) = \mathbb{C}[T]H(\mathfrak{g}, d)$. Moreover, the Γ'_+ -grading of $H(\mathfrak{g}, d)$ induces a conformal weight Δ on $H(V(R), d)$. Hence, by the observations in Example 3.12, $H(V(R), d)$ is pregraded in the sense of Definition 3.11. Part (a) is now an immediate corollary of Theorem 3.14. Part (b) is obvious. \square

4.5 Zhu algebra and cohomology

We make here the same assumptions we made in the previous section. More precisely, we let \mathfrak{g} be a finite dimensional vector superspace graded as in (4.4), we assume the free $\mathbb{C}[T]$ -module $R = \mathbb{C}[T] \otimes \mathfrak{g}$, graded according to (4.19), is a non-linear Lie conformal algebra, and its λ -bracket satisfies the grading conditions (4.20). Therefore the universal enveloping vertex algebra $V(R)$ has induced \mathbb{Z}_+ -grading $V(R)^n$, Γ -filtration $F^p V(R)$, and Γ_+ -valued conformal weight Δ , defined by a Hamiltonian operator H , satisfying the grading conditions (4.21). We also fix $d : V(R) \rightarrow V(R)$, a good almost linear differential of R , so that equations (4.22), (4.23) and (4.24) hold.

Theorem 4.20. *There is a canonical associative algebra isomorphism*

$$\phi : Zhu_H H(V(R), d) \xrightarrow{\sim} H(Zhu_H V(R), d^Z) ,$$

where d^Z is a differential of $Zhu_H V(R)$ induced by $d : V(R) \rightarrow V(R)$.

The remaining part of this section will be devoted to the proof of Theorem 4.20. First, let us fix some notation. We denote by π_H the natural quotient map

$$\pi_H : \text{Ker} d \rightarrow \text{Ker} d / \text{Im} d = H(V(R), d) ,$$

and, as before, by π_Z the natural quotient map

$$\pi_Z : V(R) \rightarrow V(R) / J_{\hbar=1}(V(R)) = Zhu_H V(R) .$$

The vertex algebra $H(V(R), d)$ has the induced Hamiltonian operator H , so that it makes sense to consider its Zhu algebra. We then let π_Z^H be the natural quotient map

$$\pi_Z^H : H(V(R), d) \rightarrow H(V(R), d) / J_{\hbar=1}(H(V(R), d)) = Zhu_H H(V(R), d) .$$

By assumption the differential d is an odd derivation of all n^{th} products of $V(R)$, and it preserves the conformal weight Δ . Hence it is a derivation of all products $a_{(n, \hbar)} b$. In particular d preserves the subspace $J_{\hbar=1}(V(R)) \subset V(R)$, and there is an induced differential d^Z of the quotient space $Zhu_H V(R) = V(R) / J_{\hbar=1}(V(R))$, given by $d^Z(\pi_Z(A)) = \pi_Z(dA)$. We can then consider the cohomology $H(Zhu_H V(R), d^Z)$, and we let π_H^Z be the natural quotient map

$$\pi_H^Z : \text{Ker} d^Z \rightarrow \text{Ker} d^Z / \text{Im} d^Z = H(Zhu_H V(R), d^Z) .$$

An equivalent way to state Theorem 4.20 is to say that the map

$$\phi : Zhu_H H(V(R), d) \rightarrow H(Zhu_H V(R), d^Z) , \quad (4.29)$$

given by $\phi(\pi_Z^H \pi_H(A)) = \pi_H^Z \pi_Z(A)$, is a well defined associative algebra isomorphism.

As in the previous section, for each $p \in \Gamma$, $\Delta \in \Gamma'_+$, we fix a basis $\{e_i, \mathcal{I}_{p\Delta}\}$ of $\mathfrak{g}^{p,-p}[\Delta] \cap d^{-1}(F^{p+\epsilon}V(R)^1[\Delta])$, and elements $\{E_i, i \in \mathcal{I}_{p\Delta}\}$ such that $E_i \in F^pV(R)^0[\Delta] \cap \text{Ker } d$ and $E_i - e_i \in F^{p+\epsilon}V(R)^0[\Delta]$. We then let

$$H(\mathfrak{g}, d) = \bigoplus_{p, \Delta} H^{p,-p}(\mathfrak{g}, d)[\Delta], \quad H^{p,-p}(\mathfrak{g}, d)[\Delta] = \text{span}_{\mathbb{C}}\{\pi_H(E_i), i \in \mathcal{I}_{p\Delta}\},$$

$$H(R, d) = \mathbb{C}[T]H(\mathfrak{g}, d) \subset H(V(R), d).$$

By Theorem 4.19, $H(R, d)$ admits a structure of non-linear Lie conformal algebra such that $H(V(R), d) \simeq V(H(R, d))$. In particular $H(V(R), d)$ is PBW generated by the set $\{\pi_H(T^k E_i) \mid k \geq 0, i \in \mathcal{I} = \bigsqcup_{p\Delta} \mathcal{I}_{p\Delta}\}$. Moreover the natural Γ'_+ -grading of $H(\mathfrak{g}, d)$ induces a conformal weight Δ of the universal enveloping vertex algebra $H(V(R), d)$. Hence the vertex algebra $H(V(R), d)$ satisfies all assumptions of Theorem 3.25. We thus get that the space $Zhu_H H(R, d) = \pi_Z^H H(\mathfrak{g}, d) \simeq H(\mathfrak{g}, d)$ is naturally endowed with a structure of non-linear Lie superalgebra and $Zhu_H H(V(R), d) \simeq U(\pi_Z^H H(\mathfrak{g}, d))$. Hence $Zhu_H H(V(R), d)$ is PBW generated as associative algebra by the space $\pi_Z^H(H(\mathfrak{g}, d))$. We thus proved the following

Lemma 4.21. *The associative superalgebra $Zhu_H H(V(R), d)$ is PBW generated by the set $\{\pi_Z^H \pi_H(E_i) \mid i \in \mathcal{I}\}$.*

Consider now the associative algebra $Zhu_H V(R)$. By Theorem 3.25 it is isomorphic to $U(Zhu_H R)$, where $Zhu_H R = \pi_Z R \simeq \mathfrak{g}$ is a Lie superalgebra with Lie bracket $[\pi_Z(a), \pi_Z(b)] = \pi_Z([a, b]_{\hbar=1})$, for $a, b \in R$. By our assumptions the space $\pi_Z \mathfrak{g} \simeq \mathfrak{g}$ has a decomposition

$$\pi_Z \mathfrak{g} = \bigoplus_{\substack{p, q \in \Gamma, \ p+q \in \mathbb{Z}_+ \\ \Delta \in \Gamma'_+}} \pi_Z \mathfrak{g}^{pq}[\Delta], \quad \pi_Z \mathfrak{g}^{pq}[\Delta] \simeq \mathfrak{g}^{pq}[\Delta].$$

Lemma 4.22. (a) *We have the following inclusion*

$$\pi_Z R^{pq}[\Delta] \subset \bigoplus_{\Delta' \leq \Delta} \pi_Z \mathfrak{g}^{pq}[\Delta'].$$

(b) *The non-linear Lie bracket on $\pi_Z \mathfrak{g}$ satisfies the grading condition (4.6), namely*

$$[\pi_Z \mathfrak{g}^{p_1 q_1}[\Delta_1], \pi_Z \mathfrak{g}^{p_2 q_2}[\Delta_2]] \subset \bigoplus_{\substack{l \geq 0 \\ \Delta < \Delta_1 + \Delta_2}} \mathcal{T}^{p_1 + p_2 + l, q_1 + q_2 - l}(\pi_Z \mathfrak{g})[\Delta].$$

Proof. Since T increases the conformal weight Δ by 1, $R^{pq}[\Delta] = \bigoplus_{k \geq 0} T^k \mathfrak{g}^{pq}[\Delta]$. On the other hand we have $\pi_Z(T^k a) = \binom{-\Delta_a}{k} \pi_Z(a)$. The first part of the lemma

follows immediately. For part (b) we have, by the definition of the product $[\cdot, \cdot]_{\hbar=1}$ and the assumption (4.20),

$$\begin{aligned}
[\pi_Z \mathfrak{g}^{p_1 q_1} [\Delta_1], \pi_Z \mathfrak{g}^{p_2 q_2} [\Delta_2]] &= \pi_Z \left([\mathfrak{g}^{p_1 q_1} [\Delta_1], \mathfrak{g}^{p_2 q_2} [\Delta_2]]_{\hbar=1} \right) \\
&\subset \pi_Z \left(\bigoplus_{l \geq 0, \Delta < \Delta_1 + \Delta_2} \mathcal{T}^{p_1 + p_2 + l, q_1 + q_2 - l}(R) [\Delta] \right) \\
&\subset \bigoplus_{l \geq 0, \Delta < \Delta_1 + \Delta_2} \mathcal{T}^{p_1 + p_2 + l, q_1 + q_2 - l}(\pi_Z \mathfrak{g}) [\Delta] .
\end{aligned}$$

□

Lemma 4.23. *Let*

$$F^p U_{\Delta}(\pi_Z \mathfrak{g})^n = \text{span}_{\mathbb{C}} \left\{ \pi_Z(a_1) \cdots \pi_Z(a_s) \mid \begin{array}{l} a_i \in \mathfrak{g}^{p_i, q_i} [\Delta_i] \\ \sum_i (p_i + q_i) = n \\ \sum_i p_i \geq p, \sum_i \Delta_i \leq \Delta \end{array} \right\} ,$$

Then we have the inclusion

$$\pi_Z(F^p V(R)^n [\Delta]) \subset F^p U_{\Delta}(\pi_Z \mathfrak{g})^n .$$

Proof. It follows by Lemma 3.18 that every element in $F^p V(R)^n [\Delta]$ is a linear combination of monomials $A = (T^{k_1} a_1)_{(-1, \hbar=1)} \cdots_{(-1, \hbar=1)} (T^{k_s} a_s)$, with $a_i \in \mathfrak{g}^{p_i, q_i} [\Delta_i]$ and $\sum_i (p_i + q_i) = n$, $\sum_i p_i \geq p$, $\sum_i (k_i + \Delta_i) = \Delta$. On the other hand $\pi_Z(A)$ is, up to a constant coefficient, equal to $\pi_Z(a_1) \cdots \pi_Z(a_s)$, which belongs to $F^p U_{\Delta}(\pi_Z \mathfrak{g})^n$. □

As noticed above, $d : V(R) \rightarrow V(R)$ is a derivation of all (n, \hbar) products. Hence the induced map

$$d^Z : U(\pi_Z \mathfrak{g}) \simeq \text{Zhu}_H V(R) \longrightarrow U(\pi_Z \mathfrak{g}) ,$$

given by $d^Z(\pi_Z(A)) = \pi_Z(dA)$ is a differential of the associative algebra $U(\pi_Z \mathfrak{g}) \simeq \text{Zhu}_H V(R)$.

Lemma 4.24. *We have*

$$d^Z(F^p U_{\Delta}(\pi_Z \mathfrak{g})^n) \subset F^p U_{\Delta}(\pi_Z \mathfrak{g})^{n+1} .$$

Proof. Let $\bar{A} = \pi_Z(a_1) \cdots \pi_Z(a_s) \in F^p U_{\Delta}(\pi_Z \mathfrak{g})^n$, with $a_i \in \mathfrak{g}^{p_i, q_i} [\Delta_i]$, and $\sum_i (p_i + q_i) = n$, $\sum_i p_i \geq p$, $\sum_i \Delta_i = \Delta$. We have $\bar{A} = \pi_Z(A)$, with $A = a_{(-1, \hbar=1)}^1 \cdots_{(-1, \hbar=1)} a^s$, and it follows by Lemma 3.18 that $A \in F^p V(R)^n [\Delta]$. By assumption (4.22) $dA \in F^p V(R)^{n+1} [\Delta]$, hence by Lemma 4.23 above, we conclude that $d^Z(\bar{A}) = \pi_Z(dA) \in F^p U_{\Delta}(\pi_Z \mathfrak{g})^{n+1}$, as we wanted. □

So far we proved that $d^Z : U(\pi_Z \mathfrak{g}) \rightarrow U(\pi_Z \mathfrak{g})$ satisfies the conditions (4.8). We want to prove that d^Z is a good almost linear differential of the non-linear Lie algebra $\pi_Z \mathfrak{g}$. For this, we will need the following

Lemma 4.25. *We have*

$$\pi_Z \mathfrak{g}^{pq}[\Delta] \cap d^{Z-1}(F^{p+\epsilon} U_\Delta(\pi_Z \mathfrak{g})^{p+q+1}) = \pi_Z(g^{pq}[\Delta] \cap d^{-1}(F^{p+\epsilon} V(R)^{p+q+1}[\Delta])) .$$

Proof. The inclusion \supset follows immediately by Lemma 4.23 above, so we only need to prove the opposite inclusion. Let $a \in \mathfrak{g}^{pq}[\Delta]$ be such that $d^Z(\pi_Z(a)) = \pi_Z(da) \in F^{p+\epsilon} U_\Delta(\pi_Z \mathfrak{g})^{p+q+1}$. By assumption (4.23) on d , we have $da = \alpha + A \in \mathfrak{g}^{p,q+1}[\Delta] \oplus F^{p+\epsilon} V(R)^{p+q+1}[\Delta]$. We then have $\pi_Z(\alpha) = \pi_Z(da) - \pi_Z(A) \in F^{p+\epsilon} U_\Delta(\pi_Z \mathfrak{g})^{p+q+1}$. But $\pi_Z(\alpha) \in \pi_Z \mathfrak{g}^{p,q+1}[\Delta]$ and $\pi_Z \mathfrak{g}^{p,q+1}[\Delta] \cap F^{p+\epsilon} U_\Delta(\pi_Z \mathfrak{g})^{p+q+1} = 0$. Hence $\pi_Z(\alpha) = 0$, and $\alpha = 0$. In conclusion $da \in F^{p+\epsilon} V(R)^{p+q+1}[\Delta]$, which is what we wanted. \square

We now use assumption (4.23) and Lemma 4.23 to get

$$\begin{aligned} d^Z(\pi_Z \mathfrak{g}^{pq}[\Delta]) &= \pi_Z(d\mathfrak{g}^{pq}[\Delta]) \subset \pi_Z(\mathfrak{g}^{p,q+1}[\Delta] \oplus F^{p+\epsilon} V(R)^{p+q+1}[\Delta]) \\ &\subset \pi_Z \mathfrak{g}^{p,q+1}[\Delta] \oplus F^{p+\epsilon} U_\Delta(\pi_Z \mathfrak{g})^{p+q+1} , \end{aligned}$$

hence d^Z is an almost linear differential of $\pi_Z \mathfrak{g}$. Moreover, by assumption (4.24), Lemma 4.23 and Lemma 4.25, we have, for $p+q \neq 0$,

$$\pi_Z \mathfrak{g}^{pq}[\Delta] \cap d^{Z-1}(F^{p+\epsilon} U_\Delta(\pi_Z \mathfrak{g})^{p+q+1}) \subset d^Z(\pi_Z \mathfrak{g}^{p,q-1}[\Delta]) + F^{p+\epsilon} U_\Delta(\pi_Z \mathfrak{g})^{p+q} ,$$

hence the differential d^Z is good. We showed so far that π_Z, d^Z satisfy the conditions of Theorem 4.14. Let e_i, E_i be as above; namely $\{e_i, i \in \mathcal{I}_{p\Delta}\}$ is a basis of $\mathfrak{g}^{p,-p}[\Delta] \cap d^{-1}(F^{p+\epsilon} V(R)^1[\Delta])$ and $E_i \in F^p V(R)^0[\Delta] \cap \text{Kerd}$ is such that $E_i - e_i \in F^{p+\epsilon} V(R)^0[\Delta]$. By Lemma 4.25, $\{\pi_Z(e_i) \mid i \in \mathcal{I}_{p\Delta}\}$ is then a basis of $\pi_Z \mathfrak{g}^{p,-p}[\Delta] \cap d^{Z-1}(F^{p+\epsilon} U_\Delta(\pi_Z \mathfrak{g})^1)$, and, for $i \in \mathcal{I}_{p\Delta}$, $\pi_Z(E_i) \in F^p U_\Delta(\pi_Z \mathfrak{g})^0 \cap \text{Kerd}^Z$ is such that $\pi_Z(E_i) - \pi_Z(e_i) \in F^{p+\epsilon} U_\Delta(\pi_Z \mathfrak{g})^0$. Notice that by Lemma 4.13, the quotient map π_H^Z restricts to an isomorphism $U(\pi_Z \mathfrak{g})^0 \cap \text{Kerd}^Z \xrightarrow{\sim} H(\text{Zhu}_H V(R), d^Z)$. As in (4.16) we then let

$$\begin{aligned} H^{p,-p}(\pi_Z \mathfrak{g}, d^Z)[\Delta] &= \text{span}_{\mathbb{C}}\{\pi_H^Z \pi_Z(E_i), i \in \mathcal{I}_{p\Delta}\} , \\ H(\pi_Z \mathfrak{g}, d^Z) &= \bigoplus_{p,\Delta} H^{p,-p}(\pi_Z \mathfrak{g}, d^Z)[\Delta] \subset H(\text{Zhu}_H V(R), d^Z) . \end{aligned}$$

By Theorem 4.14, $H(\pi_Z \mathfrak{g}, d^Z)$ is a non-linear Lie algebra and $H(\text{Zhu}_H V(R), d^Z) \simeq U(H(\pi_Z \mathfrak{g}, d^Z))$. The following lemma follows immediately by the above observations.

Lemma 4.26. *The associative algebra $H(\text{Zhu}_H V(R), d^Z)$ is PBW generated by the set $\{\pi_H^Z \pi_Z(E_i) \mid i \in \mathcal{I}\}$.*

Proof of Theorem 4.20. First, we need to show that the map ϕ , given by (4.29), is well defined. The natural quotient map $\pi_Z : V(R) \rightarrow \text{Zhu}_H V(R)$ is such

that $\pi_Z(\text{Ker}d) \subset \text{Ker}d^Z$ and $\pi_Z(d(V(R))) = d^Z(Zhu_H V(R))$. Hence it induces a canonical map

$$H(V(R), d) \longrightarrow H(Zhu_H V(R), d^Z), \quad (4.30)$$

given by $\pi_H(A) \mapsto \pi_H^Z \pi_Z(A)$. Recall that $\pi_H : V(R) \rightarrow H(V(R), d)$ is a vertex algebra homomorphism which preserves the conformal weight. Hence $\pi_H(A)_{(-2, \hbar)} \pi_H(B) = \pi_H(A_{-2, \hbar} B)$. And obviously $\pi_H^Z \pi_Z(A_{(-2, \hbar=1)} B) = 0$. Therefore $J_{\hbar=1}(H(V(R), d)) = H(V(R), d)_{(-2, \hbar=1)} H(V(R), d)$ is in the kernel of (4.30), which of course guarantees that $\phi : \pi_H^Z \pi_H(A) \mapsto \pi_H^Z \pi_Z(A)$ is well defined. The fact that ϕ is an algebra homomorphism is immediate:

$$\begin{aligned} \phi(\pi_H^Z \pi_H(A) \cdot \pi_H^Z \pi_H(B)) &= \phi(\pi_H^Z(\pi_H(A_{-1, \hbar=1} B))) = \pi_H^Z(\pi_Z(A_{-1, \hbar=1} B)) \\ &= \pi_H^Z(\pi_Z(A)) \cdot \pi_H^Z(\pi_Z(B)) = \phi(A) \cdot \phi(B). \end{aligned}$$

Finally, ϕ is an isomorphism since, by Lemmas 4.21 and 4.26, it maps a PBW generating set of $Zhu_H H(V(R), d)$ to a PBW generating set of $H(Zhu_H V(R), d^Z)$. \square

5 Finite and affine W-algebras

5.1 Definition of the affine W-algebra $W_k(\mathfrak{g}, x, f)$

Let \mathfrak{g} is a simple finite dimensional Lie superalgebra, with a non-degenerate even supersymmetric invariant bilinear form $(\cdot | \cdot)$, and let $x, f \in \mathfrak{g}$ be a pair of even elements such that $\text{ad } x$ is diagonalizable on \mathfrak{g} with half-integer eigenvalues and $[x, f] = -f$. Let

$$\mathfrak{g} = \oplus_{j \in \frac{1}{2}\mathbb{Z}} \mathfrak{g}_j \quad (5.1)$$

be the eigenspace decomposition with respect to $\text{ad } x$. Let $\mathfrak{g}_+, \mathfrak{g}_-, \mathfrak{g}_{\geq}$ and \mathfrak{g}_{\leq} denote the sums of all eigenspaces \mathfrak{g}_j with j positive, negative, non-negative, and non-positive, respectively, and let $\pi_+, \pi_-, \pi_{\geq}, \pi_{\leq}$, denote the corresponding projections of \mathfrak{g} on $\mathfrak{g}_+, \mathfrak{g}_-,$ etc. We shall assume that the pair x, f is “good”, i.e. the centraliser \mathfrak{g}^f of f in \mathfrak{g} lies in \mathfrak{g}_{\leq} . In this case the grading (5.1) is also called good. It is easy to see that for a given x , all f such that the pair x, f is good, form a single $\exp(\mathfrak{g}_0)_{\bar{0}}$ -orbit, which is Zariski dense in $(\mathfrak{g}_{-1})_{\bar{0}}$. A case of special interest is when the pair x, f is a part of an \mathfrak{sl}_2 -triple $\mathfrak{sl}_2 \simeq \text{span}_{\mathbb{C}}\{e, h = 2x, f\} \subset \mathfrak{g}$. In this case the spectrum of $\text{ad } x$ is automatically in $\frac{1}{2}\mathbb{Z}$, and the pair x, f is good by representation theory of \mathfrak{sl}_2 . One can find a detailed study and classification of good gradings of simple Lie algebras in [EK].

Following [KRW, KW1], we recall here the construction of the family of vertex algebras $W_k(\mathfrak{g}, x, f)$, depending on a parameter $k \in \mathbb{C}$. We define a skew-supersymmetric bilinear form $\langle \cdot | \cdot \rangle_{ne}$ on $\mathfrak{g}_{\frac{1}{2}}$ by (see 0.6)

$$\langle a | b \rangle_{ne} = (f, [a, b]).$$

This form is non-degenerate since the map $\text{ad } f : \mathfrak{g}_{-1/2} \rightarrow \mathfrak{g}_{1/2}$ is a vector space isomorphism.

For each $j \in \frac{1}{2}\mathbb{Z}$ we fix a basis $\{u_\alpha \mid \alpha \in S_j\}$ of \mathfrak{g}_j compatible with the parity of \mathfrak{g} . We denote $p(\alpha) = p(u_\alpha)$, the parity of u_α , let $s(\alpha) = s(u_\alpha) = (-1)^{p(\alpha)}$, and let $j_\alpha = j$ if $u_\alpha \in \mathfrak{g}_j$. Furthermore, we let $S_+ = \bigsqcup_{j>0} S_j$ and $S_- = \bigsqcup_{j\leq 0} S_j$, so that $\{u_\alpha \mid \alpha \in S_+\}$ is a basis of \mathfrak{g}_+ and $\{u_\alpha \mid \alpha \in S_-\}$ is a basis of \mathfrak{g}_- . For $j \in \frac{1}{2}\mathbb{Z}$, we also let $\{u^\alpha \mid \alpha \in S_j\}$ be the basis of \mathfrak{g}_{-j} , dual to $\{u_\alpha \mid \alpha \in S_j\}$ with respect to the bilinear form $(\cdot | \cdot)$, i.e.

$$(u_\alpha | u^\beta) = s(\alpha)(u^\beta | u_\alpha) = \delta_\alpha^\beta, \quad \alpha, \beta \in S_j.$$

In particular $\{u^\alpha\}_{\alpha \in S_+}$ is a basis of \mathfrak{g}_- dual to the basis $\{u_\alpha\}_{\alpha \in S_+}$ of \mathfrak{g}_+ . We shall identify \mathfrak{g}_- and \mathfrak{g}_+^* via the bilinear form $(\cdot | \cdot)$. Finally, we let $\{u_\alpha\}$ and $\{v_\alpha\}$, $\alpha \in S_{\frac{1}{2}}$, be a pair of dual bases of $\mathfrak{g}_{\frac{1}{2}}$ with respect to the skew-supersymmetric form $\langle \cdot | \cdot \rangle_{ne}$, i.e.

$$\langle u_\alpha | v_\beta \rangle_{ne} = (f | [u_\alpha, v_\beta]) = \delta_{\alpha, \beta}, \quad \alpha, \beta \in S_{1/2}.$$

It is immediate to check that, with the above notation, we have

$$u^\alpha = [v_\alpha, f], \quad \alpha \in S_{1/2}. \quad (5.2)$$

The building blocks of the complex that defines the affine W -algebra are the universal enveloping vertex algebras of current and fermionic Lie conformal algebras, described in Subsection 1.8. However, we shall use a slightly more convenient language of non-linear Lie conformal algebras (cf. Example 3.9). First, the *current Lie conformal algebra* $Cur_k \mathfrak{g}$ of level k is, by definition, the free $\mathbb{C}[T]$ -module $\mathbb{C}[T] \otimes \mathfrak{g}$, together with the non-linear λ -bracket

$$[a \lambda b] = [a, b] + \lambda k(a|b), \quad a, b \in \mathfrak{g}.$$

Second, let $\varphi_{\mathfrak{g}_+} = \{\varphi_a \mid a \in \mathfrak{g}_+\}$ be a vector superspace isomorphic to $\Pi \mathfrak{g}_+$, where Π denotes parity-reversing, via $a \mapsto \varphi_a$, and we let $\varphi^{\mathfrak{g}_-} = \{\varphi^a \mid a \in \mathfrak{g}_-\}$ be a vector superspace isomorphic to $\Pi \mathfrak{g}_-$, via $a \mapsto \varphi^a$. The projections π_+ and π_- allow us to extend the notation φ_a, φ^a to all $a \in \mathfrak{g}$, by letting

$$\varphi_a := \varphi_{\pi_+ a}, \quad \varphi^a := \varphi^{\pi_- a}, \quad a \in \mathfrak{g}.$$

The superspace $\varphi_{\mathfrak{g}_+} \oplus \varphi^{\mathfrak{g}_-}$ is endowed with the following non-degenerate skew-supersymmetric bilinear form $(a, b \in \mathfrak{g})$:

$$\langle \varphi_a | \varphi_b \rangle_{ch} = \langle \varphi^a | \varphi^b \rangle_{ch} = 0, \quad \langle \varphi_a | \varphi^b \rangle_{ch} = s(a) \langle \varphi^b | \varphi_a \rangle_{ch} = (\pi_+ a | \pi_- b).$$

Here, as before, $s(a) = (-1)^{p(a)}$, where $p(a)$ denotes the parity of $a \in \mathfrak{g}$. We will denote $\varphi_\alpha = \varphi_{u_\alpha}$, $\varphi^\alpha = \varphi^{u^\alpha}$, $\alpha \in S_+$. Then $\{\varphi_\alpha, \varphi^\alpha \mid \alpha \in S_+\}$ is a basis of $\varphi_{\mathfrak{g}_+} \oplus \varphi^{\mathfrak{g}_-}$, such that $\langle \varphi_\alpha | \varphi_\beta \rangle_{ch} = \langle \varphi^\alpha | \varphi^\beta \rangle_{ch} = 0$, $\langle \varphi_\alpha | \varphi^\beta \rangle_{ch} = \delta_\alpha^\beta$. The corresponding *charged free fermion* non-linear Lie conformal algebra R_{ch} is, by definition, the free $\mathbb{C}[T]$ -module $\mathbb{C}[T] \otimes (\varphi_{\mathfrak{g}_+} \oplus \varphi^{\mathfrak{g}_-})$, with the λ -bracket

$$[\phi \lambda \psi] = \langle \phi | \psi \rangle_{ch}, \quad \phi, \psi \in \varphi_{\mathfrak{g}_+} \oplus \varphi^{\mathfrak{g}_-}.$$

Finally, let $\Phi_{\mathfrak{g}_{1/2}} = \{\Phi_a | a \in \mathfrak{g}_{1/2}\}$ be a vector superspace isomorphic to $\mathfrak{g}_{1/2}$, via $a \mapsto \Phi_a$, together with the skew-supersymmetric bilinear form $\langle \cdot | \cdot \rangle_{ne}$ defined above. We will denote $\Phi_\alpha = \Phi_{u_\alpha}$, $\Phi^\alpha = \Phi_{v_\alpha}$, $\alpha \in S_{1/2}$, so that $\langle \Phi_\alpha | \Phi^\beta \rangle_{ne} = -s(\alpha) \langle \Phi^\beta | \Phi_\alpha \rangle_{ne} = \delta_\alpha^\beta$. As before, the projection $\pi_{1/2}: \mathfrak{g} \twoheadrightarrow \mathfrak{g}_{1/2}$ allows us to extend the notation Φ_a to all $a \in \mathfrak{g}$ by letting

$$\Phi_a = \Phi_{\pi_{1/2}a}, \quad a \in \mathfrak{g}.$$

The corresponding *neutral free fermion* non-linear Lie conformal algebra R_{ne} is, by definition, the free $\mathbb{C}[T]$ -module $\mathbb{C}[T] \otimes \Phi_{\mathfrak{g}_{1/2}}$, with the λ -bracket

$$[\Phi_a \lambda \Phi_b] = \langle \Phi_a | \Phi_b \rangle_{ne}, \quad a, b \in \mathfrak{g}_{1/2}.$$

Now we define the (non-linear) Lie conformal algebra $R_k(\mathfrak{g}, x)$ as the direct sum of the non-linear Lie conformal algebras $Cur_k \mathfrak{g}$, R_{ch} and R_{ne} :

$$R_k(\mathfrak{g}, x) = \mathbb{C}[T]r(\mathfrak{g}, x), \text{ where } r(\mathfrak{g}, x) = \mathfrak{g} \oplus \varphi_{\mathfrak{g}^+} \oplus \varphi^{\mathfrak{g}^-} \oplus \Phi_{\mathfrak{g}_{1/2}},$$

and we let $C_k(\mathfrak{g}, x) = V(R_k(\mathfrak{g}, x))$ be the corresponding universal enveloping vertex algebra. (Note that we are taking the complex, described in Subsection 0.4, for the choice $\mathfrak{s} = \mathfrak{g}_{1/2}$.)

The following relations are obvious and they are useful for many computations ($a \in \mathfrak{g}$),

$$\begin{aligned} a &= \sum_{\alpha \in S} (a | u^\alpha) u_\alpha = \sum_{\alpha \in S} (u_\alpha | a) u^\alpha, \\ \varphi_a &= \sum_{\alpha \in S_+} (a | u^\alpha) \varphi_\alpha, \quad \varphi^a = \sum_{\alpha \in S_+} (u_\alpha | a) \varphi^\alpha, \\ \Phi_a &= \sum_{\alpha \in S_{1/2}} (f | [a, v_\alpha]) \Phi_\alpha = \sum_{\alpha \in S_{1/2}} (f | [u_\alpha, a]) \Phi^\alpha. \end{aligned}$$

We define the *energy momentum* element $L \in C_k(\mathfrak{g}, x)$ by

$$L = L^{\mathfrak{g}} + Tx + L^{ne} + L^{ch} \in C_k(\mathfrak{g}, x),$$

where $L^{\mathfrak{g}}$ is defined by (1.60), $L^{ne} = L^A$ is defined by (1.62), where $A = \Phi_{\mathfrak{g}_{1/2}}$, and, finally, $L^{ch} = L^{A, \vec{m}}$ is defined by (1.64), where $A = \varphi_{\mathfrak{g}^+} \oplus \varphi^{\mathfrak{g}^-}$ and $\vec{m} = (j_\alpha)$. It follows that L is a Virasoro element, i.e.

$$[L \lambda L] = (T + 2\lambda)L + \frac{1}{12}c_k(\mathfrak{g}, x)\lambda^3,$$

where, due to (1.61), (1.63), and (1.65), the central charge $c_k(\mathfrak{g}, x)$ is given by

$$c_k(\mathfrak{g}, x) = \frac{k \text{ sdim } \mathfrak{g}}{k + h^\vee} - 12k(x|x) - \sum_{\alpha \in S_+} s(\alpha)(12j_\alpha^2 - 12j_\alpha + 2) - \frac{1}{2} \text{ sdim } \mathfrak{g}_{1/2}.$$

It is straightforward to compute the λ -bracket of L with the generators of $C_k(\mathfrak{g}, x)$:

$$\begin{aligned} [L \lambda a] &= Ta + \lambda(a - [x, a]) - k\lambda^2(x|a), \quad [L \lambda \Phi_a] = (T + \frac{1}{2}\lambda)\Phi_a, \\ [L \lambda \varphi_a] &= T\varphi_a + \lambda\varphi_{a-[x, a]}, \quad [L \lambda \varphi^a] = T\varphi^a + \lambda\varphi^{-[x, a]}. \end{aligned}$$

In other words, all generators $u_\alpha (\alpha \in S)$, $\varphi_\alpha, \varphi^\alpha (\alpha \in S_+)$, and $\Phi_\alpha (\alpha \in S_{1/2})$, are eigenvectors of the Hamiltonian operator $H = L_0$, and are primary elements, except for $u_\alpha (\alpha \in S_0)$ such that $(u_\alpha|x) \neq 0$, with the following conformal weights:

$$\begin{aligned} \Delta(u_\alpha) &= 1 - j_\alpha \quad (\alpha \in S), \quad \Delta(\Phi_\alpha) = 1/2 \quad (\alpha \in S_{1/2}), \\ \Delta(\varphi_\alpha) &= 1 - j_\alpha \quad (\alpha \in S_+), \quad \Delta(\varphi^\alpha) = j_\alpha \quad (\alpha \in S_+). \end{aligned} \quad (5.3)$$

Next we define the *charge decomposition* $C_k(\mathfrak{g}, x) = \bigoplus_{n \in \mathbb{Z}} C_k(\mathfrak{g}, x)^n$ by letting

$$\begin{aligned} \text{charge } a &= \text{charge } \Phi = 0, \quad a \in \mathfrak{g}, \quad \Phi \in \Phi_{\mathfrak{g}_{1/2}}, \quad \text{charge } T = 0, \\ \text{charge } \varphi_a &= -1, \quad a \in \mathfrak{g}_+, \quad \text{charge } \varphi^a = 1, \quad a \in \mathfrak{g}_-. \end{aligned} \quad (5.4)$$

This makes $C_k(\mathfrak{g}, x)$ a \mathbb{Z} -graded vertex algebra. Notice that we are choosing the sign in (5.4) opposite to that of [KW1].

Now we introduce the following element of $C_k(\mathfrak{g}, x)$ (cf. Section 0.4):

$$\begin{aligned} d &= \sum_{\alpha \in S_+} s(\alpha) : \varphi^\alpha u_\alpha : + \sum_{\alpha \in S_{1/2}} : \varphi^\alpha \Phi_\alpha : + \varphi^f \\ &+ \frac{1}{2} \sum_{\alpha, \beta \in S_+} s(\alpha) : \varphi^\alpha \varphi^\beta \varphi_{[u_\beta, u_\alpha]} :. \end{aligned} \quad (5.5)$$

Lemma 5.1. (a) *The element $d \in C_k(\mathfrak{g}, x)$ has parity $\bar{1}$ and charge 1. Moreover d is a primary element of conformal weight 1, i.e. $[L_\lambda d] = (T + \lambda)d$, or $[d_\lambda L] = \lambda L$. In particular, $d_{(0)}L = 0$.*

(b) *The element d satisfies the following λ -bracket relations with the generators of $C_k(\mathfrak{g}, x)$ ($a \in \mathfrak{g}$):*

$$\begin{aligned} [d \lambda a] &= \sum_{\alpha \in S_+} s(\alpha) : \varphi^\alpha [u_\alpha, a] : + k s(a)(T + \lambda)\varphi^a, \\ [d \lambda \varphi_a] &= \pi_+ a + (a|f) + s(a)\Phi_a + \sum_{\alpha \in S_+} : \varphi^\alpha \varphi_{[u_\alpha, \pi_+ a]} :, \\ [d \lambda \varphi^a] &= \frac{1}{2} \sum_{\alpha \in S_+} s(\alpha) : \varphi^\alpha \varphi^{[u_\alpha, a]} :, \quad [d \lambda \Phi_a] = \varphi^{[\pi_{1/2} a, f]}. \end{aligned} \quad (5.6)$$

Using (5.2) we can rewrite the last equation as $[d \lambda \Phi^\alpha] = \varphi^\alpha$, $\alpha \in S_{1/2}$.

(c) *The λ -bracket of d with itself is trivial:*

$$[d \lambda d] = 0.$$

- (d) $d_{(0)}$ is an odd derivation of all n^{th} -products in $C_k(\mathfrak{g}, x)$, it commutes with T and H , it increases the charge by 1, and $d_{(0)}^2 = 0$, hence it is a differential of the vertex algebra $C_k(\mathfrak{g}, x)$.

Proof. (a) and (b) follow by a straightforward computation. (c) follows from (b), after another long computation. (d) is immediate by Proposition 1.12(a) and, in view of (c), by the commutator formula (1.18). See [KRW, KW1] for the details of the proof. \square

Definition 5.2. We define the *affine W -algebra* $W_k(\mathfrak{g}, x, f)$ to be the vector superspace

$$W_k(\mathfrak{g}, x, f) = H(C_k(\mathfrak{g}, x), d_{(0)}) ,$$

together with the vertex algebra structure induced from $C_k(\mathfrak{g}, x)$ (see Proposition 1.12(a)), and the Hamiltonian operator H , defined by (5.3).

Remark 5.3. Since, given x , all elements f , such that the pair x, f is good, are conjugate by $\exp(\mathfrak{g}_0)_{\bar{0}}$, the W -algebra $W_k(\mathfrak{g}, x, f)$ is independent, up to isomorphism, of the choice of f , hence we shall drop f from its notation. Of course, $W_k(\mathfrak{g}, x)$ depends, up to isomorphism, only on the conjugacy class of x in \mathfrak{g} .

Remark 5.4. After the present paper was completed, the paper [BrG] came to our attention, where the authors study the finite W -algebras in their Whittaker model definition, attached to more general good \mathbb{R} -gradings: $\mathfrak{g} = \bigoplus_{j \in \mathbb{R}} \mathfrak{g}_j$, defined by the property that there exists $f \in \mathfrak{g}_{-1}$ for which $\mathfrak{g}^f \subset \mathfrak{g}_{<1/2}$. Using this, they prove that all the corresponding finite W -algebras with the same f are isomorphic. It is easy to see that if we let $\mathfrak{g}_+ = \mathfrak{g}_{\geq 1/2}$, $\mathfrak{g}_- = \mathfrak{g}_{\leq -1/2}$, then the definitions of the affine and finite W -algebras $W_k(\mathfrak{g}, x, f)$ and $W^{\text{fin}}(\mathfrak{g}, x, f)$ in Sections 5.1 and 5.2 are still valid. Moreover, using the results of [BrG] on good \mathbb{R} -gradings, it is not difficult to show that affine W -algebras $W_k(\mathfrak{g}, x, f)$ with given f are isomorphic. However, it is still useful to vary x , which defines a good \mathbb{R} -grading with given f , since this produces different energy-momentum fields, hence different Hamiltonians H and different categories of positive-energy $W_k(\mathfrak{g}, x, f)$ -modules. One such example is treated in detail in [KRW].

5.2 Definition of the finite W -algebra $W^{\text{fin}}(\mathfrak{g}, x)$

In order to define the finite W -algebra $W^{\text{fin}}(\mathfrak{g}, x)$ we look at the H -twisted Zhu algebra of the vertex algebra $C_k(\mathfrak{g}, x)$. According to Definition 3.27, the H -twisted Zhu non-linear Lie algebra associated to $R_k(\mathfrak{g}, x)$ is the vector superspace

$$Zhu_H R_k(\mathfrak{g}) = \bar{r}(\mathfrak{g}, x) \simeq \bar{\mathfrak{g}} \oplus \bar{\varphi}_{\mathfrak{g}_+} \oplus \bar{\varphi}^{\mathfrak{g}_-} \oplus \bar{\Phi}_{\mathfrak{g}_{1/2}} ,$$

endowed with the non-linear Lie bracket $[\ , \]$ induced by the bracket $[\ * \]$ on $R_k(\mathfrak{g}, x)$, namely

$$\begin{aligned} [\bar{a}, \bar{b}] &= \overline{[a, b]} - k([x, a]|b) , \quad \text{for } \bar{a}, \bar{b} \in \bar{\mathfrak{g}} , \\ [\bar{\phi}, \bar{\psi}] &= \langle \bar{\phi} | \bar{\psi} \rangle_{ch} , \quad \text{for } \bar{\phi}, \bar{\psi} \in \bar{\varphi}_{\mathfrak{g}_+} \oplus \bar{\varphi}^{\mathfrak{g}_-} , \\ [\bar{\Phi}, \bar{\Phi}'] &= \langle \bar{\Phi} | \bar{\Phi}' \rangle_{ne} , \quad \text{for } \bar{\Phi}, \bar{\Phi}' \in \bar{\Phi}_{\mathfrak{g}_{1/2}} , \end{aligned}$$

and all other brackets equal zero. Indeed, since $[a_*b] = \sum_{j \in \mathbb{Z}_+} \binom{\Delta_a - 1}{j} a_{(j)}b$, we have $[\overline{a}, \overline{b}] = \overline{[a, b]} + k(\Delta_a - 1)(a|b)$, which gives the first formula; all other formulas are obvious. Here and further we denote elements of $Zhu_H C_k(\mathfrak{g}, x)$ with an overbar, in order to distinguish them from the same elements in $C_k(\mathfrak{g}, x)$. By Corollary 3.26, the H -twisted Zhu algebra of $C_k(\mathfrak{g}, x)$, which we denote $C_k^{\text{fin}}(\mathfrak{g}, x)$, is isomorphic to the universal enveloping associative algebra of $\overline{\tau}(\mathfrak{g}, x)$,

$$C_k^{\text{fin}}(\mathfrak{g}, x) := Zhu_H C_k(\mathfrak{g}, x) \simeq U(\overline{\tau}(\mathfrak{g}, x)) .$$

Remark 5.5. Notice that the associative algebras $C_k^{\text{fin}}(\mathfrak{g}, x)$, with $k \in \mathbb{C}$, are all isomorphic. Indeed we have an explicit isomorphism $\sigma : C_k^{\text{fin}}(\mathfrak{g}, x) \xrightarrow{\sim} C_{k=0}^{\text{fin}}(\mathfrak{g}, x)$, given by $\sigma(a) = a + k(x|a)$, for $a \in \mathfrak{g}$, with all other generators unchanged: $\sigma(\varphi_a) = \varphi_a$, $\sigma(\varphi^a) = \varphi^a$, $\sigma(\Phi_a) = \Phi_a$. Hence we shall denote, from now on, $C^{\text{fin}}(\mathfrak{g}, x)$ without subscript k .

Let π_Z be the natural quotient map from $C_k(\mathfrak{g}, x)$ to $C^{\text{fin}}(\mathfrak{g}, x)$. It is immediate to check, using Remark 4.12(b), that the element $\overline{d} = \pi_Z(d) \in C^{\text{fin}}(\mathfrak{g}, x)$, corresponding to d in (5.5), is given by

$$\begin{aligned} \overline{d} &= \sum_{\alpha \in S_+} s(\alpha) \overline{\varphi}^\alpha \overline{u}_\alpha + \sum_{\alpha \in S_{1/2}} \overline{\varphi}^\alpha \overline{\Phi}_\alpha + \overline{\varphi}^f \\ &+ \frac{1}{2} \sum_{\alpha, \beta \in S_+} s(\alpha) \overline{\varphi}^\alpha \overline{\varphi}^\beta \overline{\varphi}_{[u_\beta, u_\alpha]} . \end{aligned} \quad (5.7)$$

Moreover, since $\Delta(d) = 1$, we have $[d_*A] = d_{(0)}A$ for every $A \in C_k(\mathfrak{g}, x)$, hence $[\overline{d}, \pi_Z A] = \pi_Z(d_{(0)}A)$. In other words, the action on $C^{\text{fin}}(\mathfrak{g}, x)$ induced by $d_{(0)}$ is given by the adjoint action of \overline{d} :

$$ad \overline{d} : C^{\text{fin}}(\mathfrak{g}, x) \rightarrow C^{\text{fin}}(\mathfrak{g}, x) .$$

It follows that $ad \overline{d}$ is an odd derivation of the associative superalgebra $C^{\text{fin}}(\mathfrak{g}, x)$, and $(ad \overline{d})^2 = 0$.

Definition 5.6. The *finite W -algebra* $W^{\text{fin}}(\mathfrak{g}, x)$ is, by definition, the associative superalgebra

$$W^{\text{fin}}(\mathfrak{g}, x) = H(C^{\text{fin}}(\mathfrak{g}, x), ad \overline{d}) .$$

By Remark 5.5, the finite W -algebra is, up to isomorphism, independent of k , namely the isomorphism of W_k^{fin} with W_0^{fin} is given by the change of generators, described in the Remark. Here we are using the fact that the isomorphism σ leaves \overline{d} unchanged.

5.3 Structure of the affine W -algebra $W_k(\mathfrak{g}, x)$

The following vertex operators of the vertex algebra $C_k(\mathfrak{g}, x)$, will be important “building blocks” for the affine W -algebra (cf. the second formula in (5.6)):

$$J_a = a + \sum_{\alpha \in S_+} : \varphi^\alpha \varphi_{[u_\alpha, a]} : , \quad a \in \mathfrak{g} . \quad (5.8)$$

In other words, we define an injective linear map $\mathfrak{g} \xrightarrow{\sim} J_{\mathfrak{g}} \subset C_k(\mathfrak{g})$, given by $a \mapsto J_a$. We will also denote $J_{\alpha} = J_{u_{\alpha}}$, $\alpha \in S$. Notice that J_a has the same parity, the same charge degree, and the same conformal weight as a . It is clear from the definitions that the space $J_{\mathfrak{g}} \oplus \varphi_{\mathfrak{g}^+} \oplus \varphi^{\mathfrak{g}^-} \oplus \Phi_{\mathfrak{g}_{1/2}}$ is a freely generating subspace of the vertex algebra $C_k(\mathfrak{g}, x)$. We can then rewrite all formulas in Section 5.1 in terms of this new set of generators.

First, we compute the λ -bracket of J_a , $a \in \mathfrak{g}$, with the generators of $C_k(\mathfrak{g}, x)$. We immediately get for $a, b \in \mathfrak{g}$,

$$[J_a \lambda \varphi_b] = s(a)\varphi_{[a, \pi_+ b]}, \quad [J_a \lambda \varphi^b] = \varphi^{[a, \pi_- b]}, \quad [J_a \lambda \Phi_b] = 0. \quad (5.9)$$

It follows by a straightforward but rather lengthy computation, that

$$\begin{aligned} [J_a \lambda J_b] &= J_{[a, b]} + \lambda(k(a|b) + \text{str}_{\mathfrak{g}^+}(\pi_+(ad a)\pi_+(adb))) \\ &+ \sum_{\alpha \in S_+} : \varphi^{\alpha}(\varphi_{[\pi_{\leq}[u_{\alpha}, a], b]} - p(a, b)\varphi_{[\pi_{\leq}[u_{\alpha}, b], a]}) : . \end{aligned} \quad (5.10)$$

Notice that the last term in the above equation is identically zero if a and b are both in \mathfrak{g}_{\geq} or both in \mathfrak{g}_{\leq} . We then denote

$$\psi_k(a|b) = k(a|b) + \text{str}_{\mathfrak{g}^+}((\pi_+ ad a)(\pi_+ adb)), \quad (5.11)$$

so that equation (5.10) implies, for $a, b \in \mathfrak{g}_{\leq}$ or $a, b \in \mathfrak{g}_{\geq}$:

$$[J_a \lambda J_b] = J_{[a, b]} + \psi_k(a|b). \quad (5.12)$$

We are also interested in the λ -bracket of d with the generators of $C_k(\mathfrak{g}, x)$. It immediately follows from (5.6) that

$$[d \lambda \varphi_a] = J_{\pi_+ a} + (a|f) + s(a)\Phi_a. \quad (5.13)$$

We are left to compute $[d \lambda J_a]$, $a \in \mathfrak{g}$. This is obtained by another somewhat lengthy computation:

$$\begin{aligned} [d \lambda J_a] &= \sum_{\alpha \in S_+} s(\alpha) : \varphi^{\alpha} J_{\pi_{\leq}[u_{\alpha}, a]} : - s(a) \sum_{\alpha \in S_+} : \varphi^{\alpha} \Phi_{[u_{\alpha}, a]} : \\ &+ (T + \lambda) \sum_{\alpha \in S_+} \psi_k(a|u_{\alpha}) \varphi^{\alpha} + s(a) \varphi^{[f, a]}. \end{aligned} \quad (5.14)$$

Notice that the first three terms in the RHS are identically zero for $a \in \mathfrak{g}_+$. In other words, we have

$$[d \lambda J_a] = s(a) \varphi^{[f, a]} = -[d \lambda s(a)\Phi_a], \quad a \in \mathfrak{g}_+.$$

We now let R_+ and R be the following free $\mathbb{C}[T]$ -submodules of $C_k(\mathfrak{g}, x)$,

$$\begin{aligned} R_+ &= \mathbb{C}[T]r_+, \quad r_+ = \varphi_{\mathfrak{g}^+} \oplus d_{(0)}\varphi_{\mathfrak{g}^+}, \\ R &= \mathbb{C}[T]r, \quad r = J_{\mathfrak{g}_{\leq}} \oplus \varphi^{\mathfrak{g}^-} \oplus \Phi_{\mathfrak{g}_{1/2}}, \end{aligned} \quad (5.15)$$

where $J_{\mathfrak{g}_{\leq}} = \{J_a, a \in \mathfrak{g}_{\leq}\}$. Recall that (cf. (5.6))

$$d_{(0)}\varphi_a = J_a + (a|f) + s(a)\Phi_a, a \in \mathfrak{g}_+.$$

Hence $R_+ \cap R = 0$, and $R_+ \oplus R$ is a freely generating subspace for the vertex algebra $C_k(\mathfrak{g}, x)$. Moreover, it follows from the λ -bracket relations (5.9) and (5.12) that both R_+ and $R \oplus \mathbb{C}$ are closed under the λ -bracket. More precisely, R_+ has the following structure of a Lie conformal algebra ($a, b \in \mathfrak{g}_+$):

$$\begin{aligned} [\varphi_a \lambda \varphi_b] &= 0, \quad [d_{(0)}\varphi_a \lambda \varphi_b] = s(a)\varphi_{[a,b]}, \\ [d_{(0)}\varphi_a \lambda d_{(0)}\varphi_b] &= d_{(0)}\varphi_{[a,b]}. \end{aligned} \quad (5.16)$$

The third equation is obtained by applying $d_{(0)}$ to both sides of the second equation and recalling that $d_{(0)}$ is an odd derivation of the λ -bracket and $d_{(0)}^2 = 0$. Similarly, the λ -bracket of $C_k(\mathfrak{g}, x)$ restricted to R gives the following structure of a non-linear Lie conformal algebra on R (for $J_a, J_b \in J_{\mathfrak{g}_{\leq}}, \varphi^a, \varphi^b \in \varphi^{\mathfrak{g}^-}, \Phi_a, \Phi_b \in \Phi_{\mathfrak{g}_{1/2}}$):

$$\begin{aligned} [J_a \lambda J_b] &= J_{[a,b]} + \lambda\psi_k(a|b), \quad [J_a \lambda \varphi^b] = \varphi^{[a,b]}, \\ [\Phi_a \lambda \Phi_b] &= (f|[a,b]), \quad [J_a \lambda \Phi^b] = 0 = [\varphi^a \lambda \varphi^b] = [\varphi^a \lambda \Phi_b]. \end{aligned} \quad (5.17)$$

In conclusion, the universal enveloping vertex algebras $V(R_+)$ and $V(R)$ are naturally vertex subalgebras of $C_k(\mathfrak{g}, x)$, and, since $C_k(\mathfrak{g}, x)$ is freely generated by $R_+ \oplus R$, we have a vector space isomorphism

$$C_k(\mathfrak{g}, x) \simeq V(R_+) \otimes V(R), \quad (5.18)$$

which extends the natural embeddings of $V(R_+)$ and $V(R)$ in $C_k(\mathfrak{g}, x)$. Notice that (5.18) is not a vertex algebra isomorphism since, for example, the λ -bracket of $J_a, a \in \mathfrak{g}_{\leq}$ and $\varphi_b, b \in \mathfrak{g}_+$, is not necessarily zero (see (5.9)).

We further observe that the action of the differential $d_{(0)}$ preserves both vertex subalgebras $V(R_+)$ and $V(R)$. More precisely, $d_{(0)}$ obviously restricts to a differential, which we denote by d_+ , of the Lie conformal algebra $R_+ = \mathbb{C}[T](\varphi_{\mathfrak{g}_+} \oplus d_{(0)}\varphi_{\mathfrak{g}_+}) = \mathbb{C}[T]r_+$, commuting with T , and given by:

$$d_+(\varphi_a) = d_{(0)}\varphi_a, \quad d_+(d_{(0)}\varphi_a) = 0.$$

The corresponding cohomology vanishes, since $\text{Ker } d_+ = \text{Im } d_+ = \mathbb{C}[T]d_{(0)}\varphi_{\mathfrak{g}_+}$. On the other hand, $d_{(0)}$ restricts to a vertex algebra differential, which we denote by d , of $V(R) \subset C_k(\mathfrak{g}, x)$, but it does not preserve $R = \mathbb{C}[T]r$. Namely the action

of d on $r = J_{\mathfrak{g}_{\leq}} \oplus \varphi^{\mathfrak{g}^-} \oplus \Phi_{\mathfrak{g}_{1/2}}$ is "non-linear", and it is given explicitly by

$$\begin{aligned}
d(J_a) &= \sum_{\alpha \in S_+} s(\alpha) : \varphi^\alpha J_{\pi_{\leq}[u_\alpha, a]} : - s(a) \sum_{\alpha \in S_+} : \varphi^\alpha \Phi_{[u_\alpha, a]} : \\
&+ \sum_{\alpha \in S_+} \psi_k(a|u_\alpha) T \varphi^\alpha - s(a) \varphi^{[a, f]}, \quad a \in \mathfrak{g}_{\leq}, \\
d(\varphi^a) &= \frac{1}{2} \sum_{\alpha \in S_+} s(\alpha) : \varphi^\alpha \varphi^{[u_\alpha, a]} : , \quad a \in \mathfrak{g}_-, \\
d(\Phi_a) &= \varphi^{[a, f]}, \quad a \in \mathfrak{g}_{1/2}.
\end{aligned} \tag{5.19}$$

Recall that the affine W-algebra is defined as $W_k(\mathfrak{g}, x) = H(C_k(\mathfrak{g}, x), d_{(0)})$. We thus get, by the above observations,

$$\begin{aligned}
W_k(\mathfrak{g}, x) &= H(C_k(\mathfrak{g}, x), d_{(0)}) \simeq H(V(R_+) \otimes V(R), d_+ \otimes 1 + 1 \otimes d) \\
&\simeq H(V(R_+), d_+) \otimes H(V(R), d) \simeq H(V(R), d).
\end{aligned} \tag{5.20}$$

In the third equality we used the Kunneth Lemma, and for the last equality just notice that, by Theorem 4.7, $H(V(R_+), d_+) \simeq V(H(R_+, d_+)) = \mathbb{C}$.

We finally observe that the decomposition (5.18) is compatible with the H -grading (5.3) of $C_k(\mathfrak{g}, x)$. In particular $R = \mathbb{C}[T]r$ has the induced $\frac{1}{2}\mathbb{N}$ -grading, given by

$$\Delta(J_\alpha) = 1 - j_\alpha, \quad \Delta(\varphi^\alpha) = j_\alpha, \quad \Delta(\Phi_\alpha) = 1/2, \tag{5.21}$$

which induces a $\frac{1}{2}\mathbb{Z}_+$ -valued H -grading of the universal enveloping vertex algebra $V(R)$. Moreover, since H commutes with d , we have an induced H -grading of the affine W-algebra $W_k(\mathfrak{g}, x) \simeq H(V(R), d)$.

We can summarize all the results obtained so far in the following

Theorem 5.7. *The vertex algebra $W_k(\mathfrak{g}, x)$ is isomorphic to the vertex algebra $H(V(R), d)$, where $R = \mathbb{C}[T]r$, $r = J_{\mathfrak{g}_{\leq}} \oplus \varphi^{\mathfrak{g}^-} \oplus \Phi_{\mathfrak{g}_{1/2}}$, is the non-linear Lie conformal algebra with λ -bracket given by (5.17), $V(R)$ is the universal enveloping vertex algebra of R , and d is a differential of $V(R)$, given by (5.19).*

Moreover, the H -grading on $W_k(\mathfrak{g}, x)$ is induced by the $\frac{1}{2}\mathbb{N}$ -grading of r given by (5.21).

5.4 Structure of the finite W-algebra $W^{\text{fin}}(\mathfrak{g}, x)$

We want to apply the arguments from the last section to the finite W-algebra $W^{\text{fin}}(\mathfrak{g}, x)$. As before, we denote by π_Z the natural quotient map from the vertex algebra $C_k(\mathfrak{g}, x)$ to its H -twisted Zhu algebra $C^{\text{fin}}(\mathfrak{g}, x)$. In particular, using notation introduced in Section 5.2, we have ($a \in \mathfrak{g}$)

$$\pi_Z(a) = \bar{a}, \quad \pi_Z(\varphi_a) = \bar{\varphi}_a, \quad \pi_Z(\varphi^a) = \bar{\varphi}^a, \quad \pi_Z(\Phi_a) = \bar{\Phi}_a.$$

We also denote by $\bar{J}_a \in C^{\text{fin}}(\mathfrak{g}, x)$ the image via π_Z of $J_a \in C_k(\mathfrak{g}, x)$, defined in (5.8). We have:

$$\bar{J}_a = \bar{a} + \sum_{\alpha \in S_+} \bar{\varphi}^\alpha \bar{\varphi}_{[u_\alpha, a]} + \text{str}_{\mathfrak{g}^+}((ad x)(ada)), \quad a \in \mathfrak{g}. \tag{5.22}$$

The following identities can be used to prove equation (5.22), and they will be useful to translate all other results from the last section from the affine to the finite case. We have, for $a, b, c \in \mathfrak{g}$:

$$\begin{aligned}
\pi_Z(\colon \varphi^a \varphi_b \colon) &= \overline{\varphi}^a \overline{\varphi}_b + (x|[\pi_- a, \pi_+ b]) , \\
\pi_Z(\colon \varphi_a \varphi^b \colon) &= \overline{\varphi}_a \overline{\varphi}^b + (x|[\pi_+ a, \pi_- b]) - (a|b) , \\
\pi_Z(\colon \varphi^a \varphi^b \varphi_c \colon) &= \overline{\varphi}^a \overline{\varphi}^b \overline{\varphi}_c + (x|[\pi_- b, \pi_+ c]) \overline{\varphi}^a \\
&\quad - s(a)s(b)p(a, b)(x|[\pi_- a, \pi_+ c]) \overline{\varphi}^b , \\
\pi_Z(\colon \varphi^a J_b \colon) &= \overline{\varphi}^a \overline{J}_b + s(b) \overline{\varphi}^{[x, \pi_- a], b]} .
\end{aligned} \tag{5.23}$$

We explain how to derive the first relation, all the others are obtained in a similar way. By definition $\overline{\varphi}^a \overline{\varphi}_b = \pi_Z(\varphi^a *_{-1} \varphi_b)$. Hence the first equation in (5.23) follows by the identity

$$\varphi^a *_{-1} \varphi_b = \colon \varphi^a \varphi_b \colon - (x|[\pi_- a, \pi_+ b]) ,$$

which is easy to check by the definition (2.37) of the $*_{-1}$ -product.

Notice that the space $\overline{J}_{\mathfrak{g}} \oplus \overline{\varphi}_{\mathfrak{g}^+} \oplus \overline{\varphi}^{\mathfrak{g}^-} \oplus \overline{\Phi}_{\mathfrak{g}_{1/2}}$ PBW generates $C^{\text{fin}}(\mathfrak{g}, x)$. The commutation relations among these generators immediately follow from equations (5.9) and (5.10), via the condition

$$[\pi_Z A, \pi_Z B] = \pi_Z[A * B] , \quad A, B \in C_k(\mathfrak{g}, x) . \tag{5.24}$$

Moreover, recall that the adjoint action of \overline{d} on $C^{\text{fin}}(\mathfrak{g}, x)$ is induced, via π_Z , by the action of $d_{(0)}$ on $C_k(\mathfrak{g}, x)$. Namely

$$[\overline{d}, \pi_Z A] = \pi_Z(d_{(0)} A) , \quad A \in C_k(\mathfrak{g}, x) . \tag{5.25}$$

As in the previous section, we then define the spaces

$$\overline{r}_+ = \overline{\varphi}_{\mathfrak{g}^+} \oplus [\overline{d}, \overline{\varphi}_{\mathfrak{g}^+}] , \quad \overline{r} = \overline{J}_{\mathfrak{g} \leq} \oplus \overline{\varphi}^{\mathfrak{g}^-} \oplus \overline{\Phi}_{\mathfrak{g}_{1/2}} ,$$

which are non-linear Lie subalgebras of $C^{\text{fin}}(\mathfrak{g}, x)$, with the following commutation relations. In \overline{r}_+ we have, for $a, b \in \mathfrak{g}_+$:

$$\begin{aligned}
[\overline{\varphi}_a \lambda \overline{\varphi}_b] &= 0 , \quad [[\overline{d}, \overline{\varphi}_a], \overline{\varphi}_b] = s(a) \overline{\varphi}_{[a, b]} , \\
[[\overline{d}, \overline{\varphi}_a], [\overline{d}, \overline{\varphi}_b]] &= [\overline{d}, \overline{\varphi}_{[a, b]}] .
\end{aligned} \tag{5.26}$$

The second equation follows by (5.24) and the identity

$$[\overline{d}, \overline{\varphi}_a] = \overline{J}_{\pi_+ a} + (a|f) + s(a) \overline{\Phi}_a .$$

The last equation follows by the fact that $(ad \overline{d})^2 = 0$. In \overline{r} we have, again from (5.24), (for $\overline{J}_a, \overline{J}_b \in \overline{J}_{\mathfrak{g} \leq}$, $\overline{\varphi}^a, \overline{\varphi}^b \in \overline{\varphi}^{\mathfrak{g}^-}$, $\overline{\Phi}_a, \overline{\Phi}_b \in \overline{\Phi}_{\mathfrak{g}_{1/2}}$),

$$\begin{aligned}
[\overline{J}_a, \overline{J}_b] &= \overline{J}_{[a, b]} - \psi_k([x, a]|b) , \quad [\overline{J}_a, \overline{\varphi}^b] = \overline{\varphi}^{[a, b]} , \\
[\overline{\Phi}_a, \overline{\Phi}_b] &= (f|[a, b]) , \quad [\overline{J}_a, \overline{\Phi}^b] = 0 = [\overline{\varphi}^a, \overline{\varphi}^b] = [\overline{\varphi}^a, \overline{\Phi}_b] .
\end{aligned} \tag{5.27}$$

The universal enveloping algebras $U(\bar{\tau}_+)$ and $U(\bar{\tau})$ are naturally subalgebras of $C^{\text{fin}}(\mathfrak{g}, x)$ and, by the PBW Theorem, we have a vector space isomorphism

$$C^{\text{fin}}(\mathfrak{g}, x) \simeq U(\bar{\tau}_+) \otimes U(\bar{\tau}) .$$

Moreover the adjoint action of \bar{d} restricts to a differential \bar{d}_+ of $\bar{\tau}_+$, hence of $U(\bar{\tau}_+)$, is given by

$$\bar{d}_+(\bar{\varphi}_a) = [\bar{d}, \bar{\varphi}_a] , \quad \bar{d}_+([\bar{d}, \bar{\varphi}_a]) = 0 ,$$

and it restricts to a "non-linear" differential of $U(\bar{\tau})$, an odd derivation of the associative product, which we denote again by \bar{d} . It can be computed explicitly using (5.25) and it is given by

$$\begin{aligned} \bar{d}(\bar{J}_a) &= \sum_{\alpha \in S_+} s(\alpha) \bar{\varphi}^\alpha \bar{J}_{\pi_{\leq}[u_\alpha, a]} + s(a) \bar{\varphi}^{\sum_{S_+} [[x, u^\alpha], \pi_{\leq}[u_\alpha, a]]} \\ &\quad - s(a) \sum_{\alpha \in S_+} \bar{\varphi}^\alpha \bar{\Phi}_{[u_\alpha, a]} + \sum_{\alpha \in S_+} \psi_k([x, a] | u_\alpha) \bar{\varphi}^\alpha - s(a) \bar{\varphi}^{[a, f]} , \quad a \in \mathfrak{g}_{\leq} , \\ \bar{d}(\bar{\varphi}^a) &= \frac{1}{2} \sum_{\alpha \in S_+} s(\alpha) \bar{\varphi}^\alpha \bar{\varphi}^{[u_\alpha, a]} , \quad a \in \mathfrak{g}_{-} , \\ \bar{d}(\bar{\Phi}_a) &= \bar{\varphi}^{[a, f]} , \quad a \in \mathfrak{g}_{1/2} . \end{aligned} \tag{5.28}$$

Recalling Definition 5.6 of the finite W-algebra, we can repeat the same arguments from last section, equation (5.20), to conclude that

$$W^{\text{fin}}(\mathfrak{g}, x) = H(C^{\text{fin}}(\mathfrak{g}, x), ad\bar{d}) \simeq H(U(\bar{\tau}), \bar{d}) .$$

Moreover, it is clear from the above definitions that the non-linear Lie algebra $\bar{\tau}$ coincides with the H -twisted Zhu algebra of the non-linear Lie conformal algebra R , defined in (5.15). Hence, by Theorem 3.25 we have

$$Zhu_H V(R) \simeq U(Zhu_H R) \simeq U(\bar{\tau}) .$$

Finally, the action of \bar{d} on $U(\bar{\tau})$ is induced by the action of d on $V(R)$, defined in (5.19), namely

$$\bar{d}(\pi_Z A) = \pi_Z(dA) , \quad A \in V(R) ,$$

where, as before, π_Z denotes the quotient map from the vertex algebra $V(R)$ to its H -twisted Zhu algebra $U(\bar{\tau})$.

We can summarize all the above results in the following

Theorem 5.8. *The finite W-algebra $W^{\text{fin}}(\mathfrak{g}, x)$ is isomorphic to the associative algebra $H(U(\bar{\tau}), \bar{d})$, where $\bar{\tau} = \bar{J}_{\mathfrak{g}_{\leq}} \oplus \bar{\varphi}^{\mathfrak{g}_{-}} \oplus \bar{\Phi}_{\mathfrak{g}_{1/2}} \simeq Zhu_H R$ is the H -twisted Zhu algebra of the (non-linear) Lie conformal algebra R (defined in Theorem 5.7), with the Lie bracket given explicitly by equations (5.27), and \bar{d} is the differential on $U(\bar{\tau}) \simeq Zhu_H V(R)$ induced by d on $V(R)$ (again defined in Theorem 5.7), given explicitly by equations (5.28).*

5.5 Relation between finite and affine W-algebras

Recall that, by Theorem 5.7, the affine W-algebra can be defined as $W_k(\mathfrak{g}, x) = H(V(R), d)$, where $R = \mathbb{C}[T]r$, $r = J_{\mathfrak{g}^{\leq}} \oplus \varphi^{\mathfrak{g}^-} \oplus \Phi_{\mathfrak{g}_{1/2}}$, is the non-linear Lie conformal algebra defined by (5.17), and the differential $d : V(R) \rightarrow V(R)$ is given by (5.19). We now want to show that the pair (R, d) satisfies all assumptions of Theorem 4.19 and Theorem 4.20. For this we need to decompose the generating space r as in (4.4):

$$r = \bigoplus_{\substack{p, q \in \frac{1}{2}\mathbb{Z} \\ \Delta \in \frac{1}{2}\mathbb{Z}_+}} r^{pq}[\Delta] .$$

This is obtained by assigning the following degrees

	p	q	Δ
$J_\alpha, \alpha \in S_{\leq}$	$j_\alpha - \frac{1}{2}$	$-j_\alpha + \frac{1}{2}$	$1 - j_\alpha$
$\varphi^\alpha, \alpha \in S_+$	$-j_\alpha + \frac{1}{2}$	$j_\alpha + \frac{1}{2}$	j_α
$\Phi_\alpha, \alpha \in S_{1/2}$	0	0	$\frac{1}{2}$

Notice that Δ coincides with the conformal weight (5.3), and $p + q$ coincides with the charge (5.4). It is immediate to check that the Lie conformal algebra structure (5.17) of R satisfies the grading conditions (4.20), namely

$$r^{p_1 q_1}[\Delta_1]_{(n)} r^{p_2 q_2}[\Delta_2] \subset \bigoplus_{l \geq 0} \mathcal{T}^{p_1 + p_2 + l, q_1 + q_2 - l}(R)[\Delta_1 + \Delta_2 - n - 1] .$$

In particular, the universal enveloping vertex algebra $V(R)$ has the induced \mathbb{Z} -charge grading, the increasing $\frac{1}{2}\mathbb{Z}$ -filtration given by p , and $\frac{1}{2}\mathbb{Z}_+$ -valued conformal weight Δ . We finally need to show that $d : V(R) \rightarrow V(R)$ is a good almost linear differential of R , according to Definition 4.9. Indeed, d leaves unchanged the conformal weight and changes the charge degree by $+1$. It is also immediate to check that the action of d preserves the increasing filtration given by p . In other words condition (4.22) holds. The associated graded differential is easy to write down:

$$d^{\text{gr}}(J_a) = -s(a)\varphi^{[a, f]} , \quad d^{\text{gr}}(\varphi^a) = 0 , \quad d^{\text{gr}}(\Phi_a) = \varphi^{[a, f]} .$$

In particular, it is a "linear" differential of r of bidegree $(0, 1)$, with $\text{Ker}(d^{\text{gr}}|_r) = J_{\mathfrak{g}^f} \oplus \varphi^{\mathfrak{g}^-}$ and $\text{Im}(d^{\text{gr}}|_r) = \varphi^{\mathfrak{g}^-}$. Hence the corresponding cohomology is concentrated at the zero charge:

$$H(r, d^{\text{gr}}) \simeq \text{Ker}(d^{\text{gr}}|_{r^0}) = J_{\mathfrak{g}^f} .$$

In other words, conditions (4.23) and (4.24) hold.

The following result is now an immediate consequence of Theorem 4.19 (see also [KW1]):

Theorem 5.9. Fix bases $\{u_i | i \in I_j\}$ of \mathfrak{g}_j^f , $j \leq 0$, and let $J_i = J_{u_i}$, $i \in I = \sqcup_j I_j$. For every $i \in I_j$ we can find an element $E_i \in F^{j-1/2}V(R)$ of zero charge and of conformal weight $1 - j$, such that $d(E_i) = 0$ and $E_i - J_i \in F^jV(R)$. Moreover the space

$$H(R, d) = \mathbb{C}[T]H(r, d), \text{ where } H(r, d) = \text{span}_{\mathbb{C}}\{E_i, i \in I\}, \quad (5.29)$$

admits a structure of a non-linear Lie conformal algebra such that $W_k(\mathfrak{g}, x) \simeq V(H(R, d))$. In particular $W_k(\mathfrak{g}, x) = H^0(V(R), d)$, i.e. the cohomology of the complex $(C_k(\mathfrak{g}, x), d_{(0)})$ is concentrated at zero charge.

We now recall that, by Theorem 5.8, the finite W -algebra is isomorphic as associative algebra to $H(\text{Zhu}_H V(R), d^Z)$. We thus immediately get from Theorem 4.20 and Theorem 5.7 the main result of this section:

Theorem 5.10. There is a canonical associative algebra isomorphism

$$\text{Zhu}_H W_k(\mathfrak{g}, x) \xrightarrow{\sim} W^{fn}(\mathfrak{g}, x) \simeq U(H(r, d)),$$

where $H(r, d)$ is the space in (5.29), with the structure of a non-linear Lie algebra induced by the bracket $[\cdot, \cdot]$ on $W_k(\mathfrak{g}, x)$.

Remark 5.11. Let K_0 denote the subspace of zero charge d -closed elements of $V(R)$. Then, obviously, the canonical map $K_0 \rightarrow W_k(\mathfrak{g}, x)$ is an isomorphism of vertex algebras. Thus we have the "formality" property, i.e. the complex $(W_k(\mathfrak{g}, x), d = 0)$ embeds in the complex $(C_k(\mathfrak{g}, x), d_{(0)})$ as the subcomplex consisting of zero-charge $d_{(0)}$ -closed elements of the subcomplex $(V(R), d)$. Similar "formality" property holds for the finite W -algebras

6 Quasiclassical limit: quantum and classical W -algebras

If we remove the integral terms ("quantum corrections") in formulas (1.37), (1.38), (1.39) and (1.40) of the fourth definition of a vertex algebra, we arrive at the definition of a Poisson vertex algebra (cf. [FB]).

Definition 6.1. A Poisson vertex algebra is a quintuple $(\mathcal{V}, |0\rangle, T, \{\cdot_\lambda \cdot\}, \cdot)$, where

- (i) $(\mathcal{V}, T, \{\cdot_\lambda \cdot\})$ is a Lie conformal superalgebra,
- (ii) $(\mathcal{V}, |0\rangle, T, \cdot)$ is a unital associative commutative differential superalgebra,
- (iii) the operations $\{\cdot_\lambda \cdot\}$ and \cdot are related by the Leibniz rules (= commutative Wick formulas):

$$\{a_\lambda bc\} = \{a_\lambda b\}c + p(a, b)b\{a_\lambda c\}, \quad (6.1)$$

$$\{ab_\lambda c\} = (e^{T\partial_\lambda} a)\{b_\lambda c\} + p(a, b)(e^{T\partial_\lambda} b)\{a_\lambda c\} \quad (6.2)$$

($= p(a, b)p(a, c)\{b_{\lambda+T}c\}_{\rightarrow}a + p(b, c)\{a_{\lambda+T}c\}_{\rightarrow}b$, where the arrow means that $\lambda + T$ should be moved to the right).

As for vertex algebras, given a Poisson vertex algebra \mathcal{V} we can define n^{th} products for every $n \in \mathbb{Z}$ as follows:

$$a_{(-n-1)}b = (T^{(n)}a)b, \quad a_{(n)}b = \frac{d^n}{d\lambda^n}\{a_{\lambda}b\}|_{\lambda=0}, \quad n \in \mathbb{Z}_+.$$

Example 6.2. Let $\mathcal{V} = \mathbb{C}[u_i^{(n)} \mid i \in I, n \in \mathbb{Z}_+]$, be the algebra of polynomials in the even indeterminates $u_i^{(n)}$, let T be the derivation of \mathcal{V} defined by $Tu_i^{(n)} = u_i^{(n+1)}$, and let $|0\rangle = 1$. Define a λ -bracket on \mathcal{V} by

$$\{P_{\lambda}Q\} = \sum_{i,j \in I, p,q \in \mathbb{Z}_+} \frac{\partial Q}{\partial u_j^{(q)}} (\lambda + T)^q \{u_i_{\lambda+T}u_j\}_{\rightarrow} (-\lambda - T)^p \frac{\partial P}{\partial u_i^{(p)}},$$

where the λ -bracket between the $u_i = u_i^{(0)}$ is defined such that skewsymmetry (resp. Jacobi identity) holds for any pair u_i, u_j (resp. triple u_i, u_j, u_k). As before, $\{a_{\lambda+T}b\}_{\rightarrow}$ means that the powers of $\lambda + T$ are placed on the right. Then it is not difficult to check that \mathcal{V} is a Poisson vertex algebra. For example we can take $\{u_i_{\lambda}u_j\} = P_{i,j}(\lambda)$ be arbitrary polynomials such that $P_{i,j}(\lambda) = -P_{j,i}(-\lambda)$ for every $i, j \in I$. The simplest example of this is $\mathcal{V} = \mathbb{C}[u^{(n)} \mid n \in \mathbb{Z}_+]$ where $\{u_{\lambda}u\} = \lambda$, called the Gardner–Faddeev–Zakharov (GFZ) bracket.

Remark 6.3. Lie conformal algebras provide a very convenient framework for both classical and quantum Hamiltonian systems. This is based on the obvious observation that, given a Lie conformal algebra $(\mathcal{V}, \{\cdot_{\lambda}\cdot\})$, the bracket $\{a, b\} = \{a_{\lambda}b\}|_{\lambda=0}$ on \mathcal{V} satisfies the Jacobi identity for Lie algebras, it is such that $\{T\mathcal{V}, \mathcal{V}\} = 0$, and the subspace $T\mathcal{V}$ is a two-sided ideal with respect to this bracket, so that $(\mathcal{V}/T\mathcal{V}, \{\cdot, \cdot\})$ is a Lie algebra. A "local functional" is an element of $\mathcal{V}/T\mathcal{V}$. Given $h \in \mathcal{V}/T\mathcal{V}$, the corresponding "Hamiltonian equation" is, by definition, the equation

$$\dot{u} = \{h, u\}, \quad u \in \mathcal{V}.$$

A local functional h_1 is said to be an "integral of motion" for this equation if $\dot{h}_1 = 0$, or, equivalently, if h and h_1 are "in involution", i.e. $\{h, h_1\} = 0$. If \mathcal{V} is a Poisson vertex algebra, we obtain in this way classical Hamiltonian equations. For example, if $\mathcal{V} = \mathbb{C}[u, u', u'', \dots]$ is the Poisson vertex algebra with the GFZ bracket $\{u_{\lambda}u\} = \lambda$, then the local functionals $h_0 = \int u$, $h_1 = \int u^2$, $h_2 = \frac{1}{2} \int (u^3 - u'^2)$, \dots are in involution (here the sign of the integral simply means taking the image in $\mathcal{V}/T\mathcal{V}$), and the Hamiltonian equation $\dot{u} = \{h_2, u\}$ is the classical KdV equation $\dot{u} = 3uu' + u'''$. In exactly the same way, given a vertex algebra V , one defines a "local functional" $h \in V/T\mathcal{V}$ and the corresponding "quantum Hamiltonian equation" $\dot{u} = [h, u]$, $u \in V$. In particular, taking the vertex algebra $V^k(\mathfrak{g})$, where \mathfrak{g} is the 1-dimensional (even) Lie algebra with a non-degenerate bilinear form (its quasi-classical limit, discussed below, is the GFZ Poisson vertex algebra), one can in the same way construct and study quantum Hamiltonian equations and their integrals of motion (see [FF3]).

We next want to define the (H, Γ) -twisted Zhu algebra of a Poisson vertex algebra (cf. Section 2.2). A *Hamiltonian operator* H on a Poisson vertex algebra \mathcal{V} is a diagonalizable operator on \mathcal{V} such that (cf. (0.2))

$$H(a_{(n)}b) = (Ha)_{(n)}b + a_{(n)}(Hb) - (n+1)a_{(n)}b, \quad a, b \in \mathcal{V}, \quad n \in \mathbb{Z}.$$

As usual, if a is an eigenvector of H , its eigenvalue is denoted by Δ_a . Furthermore, let Γ be an additive subgroup of \mathbb{R} containing \mathbb{Z} . A Poisson vertex algebra \mathcal{V} is Γ/\mathbb{Z} -graded if we have a decomposition $\mathcal{V} = \bigoplus_{\bar{\gamma} \in \Gamma/\mathbb{Z}} \bar{\gamma}\mathcal{V}$ by H -invariant subspaces, such that

$$\bar{\alpha}\mathcal{V}\bar{\beta}\mathcal{V} \subset \bar{\alpha}+\bar{\beta}\mathcal{V}, \quad \{\bar{\alpha}\mathcal{V} \lambda \bar{\beta}\mathcal{V}\} \subset \mathbb{C}[\lambda] \otimes \bar{\alpha}+\bar{\beta}\mathcal{V}.$$

We will use the notation introduced in Section 2.2: given $a \in \bar{\gamma}_a\mathcal{V}[\Delta_a]$, we let ϵ_a be the maximal non-positive real number in $\bar{\gamma}_a - \Delta_a$, we let $\gamma_a = \Delta_a + \epsilon_a$, and for homogeneous $a, b \in \mathcal{V}$ we let $\chi(a, b) = 1$ or 0 depending on whether $\epsilon_a + \epsilon_b \leq -1$ or not. We also let H' be the diagonalizable operator of \mathcal{V} such that $H'(a) = \gamma_a a$. Consider the space $\mathcal{V}[\hbar]$, with the $\mathbb{C}[\hbar]$ -bilinear commutative associative product induced by the product ab on \mathcal{V} , and define the following $\mathbb{C}[\hbar]$ -bilinear \hbar -bracket on $\mathcal{V}[\hbar]$:

$$\{a, b\}_{\hbar} = \sum_{j \in \mathbb{Z}_+} \binom{\gamma_a - 1}{j} \hbar^j a_{(j)} b, \quad a, b \in \mathcal{V}. \quad (6.3)$$

Let \mathcal{V}_{Γ} (resp. $\mathcal{V}_{\hbar, \Gamma}$) be the \mathbb{C} -span (resp. $\mathbb{C}[\hbar]$ -span in $\mathcal{V}[\hbar]$) of all elements $a \in \mathcal{V}$ such that $\epsilon_a = 0$. Moreover let $J_{\hbar, \Gamma} \subset \mathcal{V}_{\hbar, \Gamma}$ be the $\mathbb{C}[\hbar]$ -submodule generated by all the elements $a_{(-2+\chi(a, b), \hbar, \Gamma)}b$, with $a, b \in \mathcal{V}$ such that $\epsilon_a + \epsilon_b \in \mathbb{Z}$, where we let

$$a_{(-2, \hbar, \Gamma)}b = ((T + \hbar H')a)b, \quad a_{(-1, \hbar, \Gamma)}b = ab, \quad a, b \in \mathcal{V}.$$

If we put $b = 1$ in the above expression, we get that $(T + \hbar H')a \in J_{\hbar, \Gamma}$ for all $a \in \mathcal{V}_{\hbar, \Gamma}$. Hence, by induction,

$$T^{(n)}a \equiv \hbar^n \binom{-\gamma_a}{n} a \pmod{J_{\hbar, \Gamma}}, \quad \text{for } a \in \mathcal{V}_{\hbar, \Gamma}. \quad (6.4)$$

Theorem 6.4. $J_{\hbar, \Gamma}$ is a two-sided ideal of $\mathcal{V}_{\hbar, \Gamma}$ with respect to both the commutative associative product \cdot and the \hbar -bracket $\{\cdot, \cdot\}_{\hbar}$. Moreover,

$$\text{Zhu}_{\hbar, \Gamma}\mathcal{V} := \mathcal{V}_{\hbar, \Gamma}/J_{\hbar, \Gamma}$$

is a Poisson superalgebra over $\mathbb{C}[\hbar]$, with commutative associative product induced by \cdot and with Lie bracket induced by $\{\cdot, \cdot\}_{\hbar}$.

Proof. It is clear, by the definition, that $J_{\hbar, \Gamma}$ is a two-sided ideal with respect to the commutative associative product of $\mathcal{V}_{\hbar, \Gamma}$. Moreover, it immediately follows by the definition (6.3) of the \hbar -bracket and by the left Leibniz rule (6.1), that

$$\{a, bc\}_{\hbar} = \{a, b\}_{\hbar}c + p(a, b)\{a, c\}_{\hbar}, \quad a, b \in \mathcal{V}. \quad (6.5)$$

It is not hard to prove, by a direct computation, that the following identity holds for all $a, b \in \mathcal{V}_\Gamma$ (cf. (2.11):

$$\{a, (T + \hbar H)b\}_\hbar = (T + \hbar H)\{a, b\}_\hbar. \quad (6.6)$$

If we then replace b by $(T + \hbar H)b$ in (6.5) and we use (6.6), we get, for $a, b, c \in \mathcal{V}_\Gamma$,

$$\{a, b_{(-2, \hbar, \Gamma)}c\}_\hbar = \{a, b\}_{\hbar(-2, \hbar, \Gamma)}c + p(a, b)b_{(-2, \hbar, \Gamma)}\{a, c\}_\hbar. \quad (6.7)$$

Next, we want to prove that the \hbar -bracket is skewsymmetric in the quotient space $\mathcal{V}_{\hbar, \Gamma}/J_{\hbar, \Gamma}$, namely

$$\{b, a\}_\hbar \equiv -p(a, b)\{a, b\}_\hbar \pmod{J_{\hbar, \Gamma}}, \quad a, b \in \mathcal{V}_\Gamma. \quad (6.8)$$

By definition of the \hbar -bracket and by the skewsymmetry of the λ -bracket, or, equivalently, by (1.30), we have

$$\begin{aligned} \{b, a\}_\hbar &= \sum_{i \in \mathbb{Z}_+} \binom{\Delta_b - 1}{i} \hbar^i b_{(i)}a \\ &= p(a, b) \sum_{i, j \in \mathbb{Z}_+} \binom{\Delta_b - 1}{i} \hbar^i (-1)^{i+j+1} T^{(j)}(a_{(i+j)}b). \end{aligned}$$

We then use (6.4) to get, for $a, b \in \mathcal{V}_\Gamma$,

$$\{b, a\}_\hbar \equiv -p(a, b) \sum_{i, j \in \mathbb{Z}_+} \binom{\Delta_b - 1}{i} \binom{-\Delta_a - \Delta_b + i + j + 1}{j} (-\hbar)^{i+j} a_{(i+j)}b \pmod{J_{\hbar, \Gamma}}. \quad (6.9)$$

Thanks to (2.22) with $x = \Delta_b - 1$, $y = -\Delta_a - \Delta_b + n + 1$, the RHS of (6.9) is equal to

$$-p(a, b) \sum_{n \in \mathbb{Z}_+} \binom{-\Delta_a + n}{n} (-\hbar)^n a_{(n)}b,$$

which in turn is equal to $-p(a, b)\{a, b\}_\hbar$, thus proving (6.8). It follows by (6.7) and (6.8) that $J_{\hbar, \Gamma} \subset \mathcal{V}_{\hbar, \Gamma}$ is a two-sided ideal with respect to the \hbar -bracket. We are then left to prove that the \hbar -bracket satisfies the Jacobi identity in $\mathcal{V}_{\hbar, \Gamma}/J_{\hbar, \Gamma}$. In fact, we will prove more, that the Jacobi identity holds already in $\mathcal{V}_{\hbar, \Gamma}$, namely that, for $a, b, c \in \mathcal{V}_\Gamma$,

$$\{a, \{b, c\}_\hbar\}_\hbar - p(a, b)\{b, \{a, c\}_\hbar\}_\hbar = \{\{a, b\}_\hbar, c\}_\hbar. \quad (6.10)$$

By the definition of the \hbar -bracket, the LHS of (6.10) is

$$\sum_{i, j \in \mathbb{Z}_+} \binom{\Delta_a - 1}{i} \binom{\Delta_b - 1}{j} \hbar^{i+j} \left(a_{(i)}(b_{(j)}c) - p(a, b)b_{(j)}(a_{(i)}c) \right). \quad (6.11)$$

By the Jacobi identity for the λ -bracket, or, equivalently, using (1.31), we get that (6.11) is equal to

$$\sum_{i,j,k \in \mathbb{Z}_+} \binom{\Delta_a - 1}{i} \binom{\Delta_b - 1}{j} \binom{i}{k} \hbar^{i+j} (a_{(k)} b)_{(i+j-k)} c .$$

In the above expression we replace $\binom{\Delta_a - 1}{i} \binom{i}{k}$ by $\binom{\Delta_a - 1}{k} \binom{\Delta_a - k - 1}{i - k}$, and we make the change of variables $n = i + j - k$, $l = i - k$, to get

$$\sum_{n,k \in \mathbb{Z}_+} \binom{\Delta_a - 1}{k} \sum_{l=0}^n \binom{\Delta_a - k - 1}{l} \binom{\Delta_b - 1}{n - l} \hbar^{n+k} (a_{(k)} b)_{(n)} c .$$

By (2.22) with $x = \Delta_a - k - 1$ and $y = \Delta_b - 1$, we then get

$$\sum_{n,k \in \mathbb{Z}_+} \binom{\Delta_a - 1}{k} \binom{\Delta_a + \Delta_b - k - 2}{n} \hbar^{n+k} (a_{(k)} b)_{(n)} c ,$$

which is in turn equal to $\{\{a, b\}_{\hbar}, c\}_{\hbar}$, thus proving (6.10). \square

Definition 6.5. A family of associative algebras A_ϵ is an associative superalgebra over the algebra of polynomials $\mathbb{C}[\epsilon]$, such that

$$[A_\epsilon, A_\epsilon] \subset \epsilon A_\epsilon . \quad (6.12)$$

It is called *regular* if the operator of multiplication by ϵ has zero kernel. Then $A^{\text{cl}} = A_\epsilon / \epsilon A_\epsilon$ is a commutative associative superalgebra, with the Poisson bracket $\{\bar{a}, \bar{b}\}$ defined by taking preimages $a, b \in A_\epsilon$ of $\bar{a}, \bar{b} \in A^{\text{cl}}$, writing

$$[a, b] = \epsilon \{a, b\} \in \epsilon A_\epsilon , \quad (6.13)$$

and taking the image of $\{a, b\}$ in A^{cl} . It is easy to see that this bracket is well defined and it makes A^{cl} a Poisson superalgebra. The Poisson superalgebra A^{cl} is called the *quasiclassical limit* of the family A_ϵ , and we let $\pi_0 : A_\epsilon \twoheadrightarrow A^{\text{cl}}$ be the natural quotient map. Similarly, $A = A_\epsilon / (\epsilon - 1)A_\epsilon$ is an associative algebra over \mathbb{C} , and we let $\pi_1 : A_\epsilon \twoheadrightarrow A$ be the natural quotient map.

Remark 6.6. The algebra $Zhu_{\hbar, \Gamma} V$ over $\mathbb{C}[\hbar]$ (defined in Section 2.2) is a family of associative algebras, by Theorem 2.13(f). However $Zhu_{\hbar, \Gamma} V$ is not, in general, a regular family of associative algebras, in the sense of Definition 6.5, since the multiplication by \hbar might have a non-zero kernel. For example, if V is an in Example 2.18, then

$$Zhu_{\hbar, H} V \simeq \mathbb{C}[\hbar] V / \left(\mathbb{C} \oplus \left(\bigoplus_{n \geq 1} \hbar V[n] \right) \right) = \mathbb{C}[\hbar] \oplus \left(\bigoplus_{n \geq 1} V[n] \right) ,$$

and $\text{Ker } \hbar \simeq \bigoplus_{n \geq 1} V[n] \neq 0$.

However, by Proposition 2.17(a), $Zhu_{\hbar,\Gamma}V/\hbar Zhu_{\hbar,\Gamma}V \simeq V_\Gamma/(V_\Gamma \cap V_{(-2+\chi)}V)$ is still a Poisson algebra, that we will call, again, the *quasiclassical limit* of $Zhu_{\hbar,\Gamma}V$. We then let ψ_0 be the natural quotient map $\psi_0 : Zhu_{\hbar,\Gamma}V \twoheadrightarrow V_\Gamma/(V_\Gamma \cap V_{(-2+\chi)}V)$. More generally, recall (see Section 2.2) that we can specialize $Zhu_{\hbar,\Gamma}V$ at any value $\hbar = c \in \mathbb{C}$, to get the associative superalgebra $Zhu_{\hbar,\Gamma}V/(\hbar - c)Zhu_{\hbar,\Gamma}V \simeq V_\Gamma/J_{\hbar=c,\Gamma}$. Denote by ψ_c the quotient map $Zhu_{\hbar,\Gamma}V \twoheadrightarrow V_\Gamma/J_{\hbar=c,\Gamma}$, and recall the notation $Zhu_\Gamma V = V_\Gamma/J_{\hbar=1,\Gamma}$ for the (H, Γ) -twisted Zhu algebra, from Section 2.2.

We will use the same notation also for a Poisson vertex algebra \mathcal{V} . Namely we let again ψ_c be the quotient map $Zhu_{\hbar,\Gamma}\mathcal{V} \twoheadrightarrow Zhu_{\hbar,\Gamma}\mathcal{V}/(\hbar - c)Zhu_{\hbar,\Gamma}\mathcal{V} \simeq \mathcal{V}_\Gamma/J_{\hbar=c,\Gamma}$ for arbitrary $c \in \mathbb{C}$, and we let $Zhu_\Gamma\mathcal{V} = \mathcal{V}_\Gamma/J_{\hbar=1,\Gamma}$. Obviously $\mathcal{V}_\Gamma/J_{\hbar=c,\Gamma}$ is a Poisson superalgebra over \mathbb{C} for every choice of $c \in \mathbb{C}$.

Definition 6.7. A family of vertex algebras V_ϵ is a vertex algebra over $\mathbb{C}[\epsilon]$ such that

$$[V_\epsilon \lambda V_\epsilon] \subset \mathbb{C}[\lambda] \otimes \epsilon V_\epsilon. \quad (6.14)$$

As before, V_ϵ is called *regular* if the operator of multiplication by ϵ has zero kernel. Then the normally ordered product on V_ϵ induces the structure of a unital commutative associative differential superalgebra on $(V^\text{cl} = V_\epsilon/\epsilon V_\epsilon, |0\rangle, T, \cdot)$, endowed with the λ -bracket $\{\bar{a} \lambda \bar{b}\}$, defined by taking preimages $a, b \in V_\epsilon$ of $\bar{a}, \bar{b} \in V^\text{cl}$, writing

$$[a \lambda b] = \epsilon \{a \lambda b\}, \quad (6.15)$$

and taking the image of $\{a \lambda b\}$ in $\mathbb{C}[\lambda] \otimes V^\text{cl}$. It is easy to see, using the fourth definition of a vertex algebra, that $(V^\text{cl}, |0\rangle, T, \{\cdot \lambda \cdot\}, \cdot)$ is a Poisson vertex algebra. Likewise, this Poisson vertex algebra is called the *quasiclassical limit* of the family V_ϵ , and we let $\pi_0 : V_\epsilon \twoheadrightarrow V^\text{cl}$ be the natural quotient map. Moreover, $V = V_\epsilon/(\epsilon - 1)V_\epsilon$ is clearly a vertex algebra over \mathbb{C} , and we let $\pi_1 : V_\epsilon \twoheadrightarrow V$ be the corresponding quotient map.

Example 6.8. A general example of a regular family of associative algebras $U(\mathfrak{g})_\epsilon$ is obtained by taking a Lie superalgebra \mathfrak{g} , defining a new $\mathbb{C}[\epsilon]$ -bilinear Lie bracket on $\mathbb{C}[\epsilon] \otimes \mathfrak{g}$ by the formula $[a, b]_\epsilon = \epsilon[a, b]$ for $a, b \in \mathfrak{g}$, and denoting by $U(\mathfrak{g})_\epsilon$ the universal enveloping superalgebra over $\mathbb{C}[\epsilon]$ of $\mathbb{C}[\epsilon] \otimes \mathfrak{g}$. It is clear that the quasiclassical limit of the family $U(\mathfrak{g})_\epsilon$ is the unital commutative associative superalgebra $S(\mathfrak{g})$ with the Poisson bracket which coincides with the Lie bracket on \mathfrak{g} and is extended to $S(\mathfrak{g})$ by the Leibniz rule.

In a similar fashion one constructs a regular family of vertex algebras $V(R)_\epsilon$, starting from a Lie conformal superalgebra R , by defining the $\mathbb{C}[\epsilon]$ -bilinear λ -bracket $[a \lambda b] = \epsilon[a \lambda b]$ on $\mathbb{C}[\epsilon] \otimes R$, and denoting $V(R)_\epsilon$ the universal enveloping vertex algebra over $\mathbb{C}[\epsilon]$ of $\mathbb{C}[\epsilon] \otimes R$. It is easy to see that the quasiclassical limit of the family $V(R)_\epsilon$ is the unital commutative associative differential superalgebra $S(R)$ with the Poisson λ -bracket which coincides with the λ -bracket on R and extends it to $S(R)$ by the Leibniz rules (6.1) and (6.2).

Example 6.9. In exactly the same way one constructs a regular family of associative superalgebras (resp. vertex algebras) in a slightly more general case

of a non-linear Lie superalgebra (resp. Lie conformal superalgebra), which is a central extension of a Lie superalgebra (resp. Lie conformal superalgebra), as described in Example 3.9. Also, the quasiclassical limits are computed in the same way. However this construction does not work for arbitrary non-linear Lie superalgebras or Lie conformal superalgebras.

Example 6.10. The GFZ Poisson vertex algebra from Example 6.2 is the quasiclassical limit of the family $B_\epsilon = V(R)_\epsilon$, where $R = \mathbb{C}[T]u$ is a non-linear Lie conformal algebra with λ -bracket $[u \lambda u] = \lambda$. Note that $B_{\epsilon=1}$ is the vertex algebra of a free boson (cf. [K]).

Let V_ϵ be a regular family of vertex algebras over $\mathbb{C}[\epsilon]$, with a Hamiltonian operator H and a Γ/\mathbb{Z} -grading. According to the above observations, starting from V_ϵ we can build the following commutative diagram of fundamental objects:

$$\begin{array}{ccccc}
 V^{\text{cl}} & \xleftarrow{\pi_0} & V_\epsilon & \xrightarrow{\pi_1} & V \\
 \vdots \downarrow \text{Zhu}_{\hbar,\Gamma} & & \vdots \downarrow \text{Zhu}_{\hbar,\Gamma} & & \downarrow \text{Zhu}_{\hbar,\Gamma} \\
 \text{Zhu}_{\hbar,\Gamma} V^{\text{cl}} & \xleftarrow{\pi_0} & \text{Zhu}_{\hbar,\Gamma} V_\epsilon & \xrightarrow{\pi_1} & \text{Zhu}_{\hbar,\Gamma} V \\
 \downarrow \psi_1 & & \downarrow \psi_1 & & \downarrow \psi_1 \\
 \text{Zhu}_\Gamma V^{\text{cl}} & \xleftarrow{\pi_0} & \text{Zhu}_\Gamma V_\epsilon & \xrightarrow{\pi_1} & \text{Zhu}_\Gamma V \\
 \downarrow \psi_0 & & \downarrow \psi_0 & & \downarrow \psi_0 \\
 \frac{V_\Gamma^{\text{cl}}}{V_\Gamma^{\text{cl}} \cap (V_{(-2+\chi)}^{\text{cl}} V^{\text{cl}})} & \xleftarrow{\pi_0} & \frac{V_{\epsilon\Gamma}}{V_{\epsilon\Gamma} \cap (V_{\epsilon(-2+\chi)} V_\epsilon)} & \xrightarrow{\pi_1} & \frac{V_\Gamma}{V_\Gamma \cap (V_{(-2+\chi)} V)}
 \end{array} \quad (6.16)$$

In the above diagram, horizontal left arrows denote the canonical map π_0 from an ϵ -family of algebras to its quasiclassical limit, described in Definitions 6.5 and 6.7. Similarly, the horizontal right arrows denote the quotient map π_1 from an ϵ -family of algebras to its specialization at $\epsilon = 1$, described in Definitions 6.5 and 6.7. The dotted vertical arrows stand for taking the family of associative algebras $\text{Zhu}_{\hbar,\Gamma}$ for a vertex algebra or a Poisson vertex algebra. The straight (respectively curved) vertical arrows denote the canonical quotient maps ψ_1 (resp. ψ_0) to the specializations at $\hbar = 1$ (resp. $\hbar = 0$), as described after Remark 6.6.

Example 6.11. Consider the special case of a family of vertex algebras $V(R)_\epsilon$ associated to a Lie conformal algebra R , or to a non-linear Lie conformal algebra of the type discussed in Example 6.9. Suppose R has a Hamiltonian operator H , and take the Γ/\mathbb{Z} -grading of $V(R)_\epsilon$ induced by the action of H (see Example 2.12). Then we can build, starting from $V(R)_\epsilon$, the whole diagram (6.16). In particular, at the top right and left corners we have respectively $V = V(R)$, the universal enveloping vertex algebra of R , and $V^{\text{cl}} = S(R)$, the symmetric algebra of R , with λ -bracket extending the λ -bracket of R by the Leibniz rules (6.1) and (6.2). Moreover, at the bottom left and right corners we have isomorphic Poisson algebras:

$$V^{\text{cl}} / (V_{(-2)}^{\text{cl}} V^{\text{cl}}) \simeq S(R/TR) \simeq V / (V_{(-2)} V), \quad (6.17)$$

where $S(R/TR)$ has Lie bracket induced by the 0th product of R/TR and extended by the Leibniz rule. To prove (6.17), note that $S(R)_{(-2)}S(R)$ is the ideal of $S(R)$ generated by TR , which implies the first isomorphism. Similarly $V(R)_{(-2)}V(R)$ is the ideal of $V(R)$, with respect to the -1^{st} product, generated by TR . Hence the natural linear map $R \rightarrow V(R)/(V(R)_{(-2)}V(R))$ induces a surjective homomorphism of associative algebras $\varphi : S(R/TR) \rightarrow V(R)/(V(R)_{(-2)}V(R))$. To prove that this is an isomorphism, we construct the inverse map as follows. The quotient map $R \twoheadrightarrow R/TR$ extends to a surjective linear map $\mathcal{T}(R) \twoheadrightarrow S(R/TR)$. It is easy to see that the subspace $\mathcal{M}(R) \subset \mathcal{T}(R)$ (see (3.8)) is in the kernel, hence we get a surjective linear map $V(R) \twoheadrightarrow S(R/TR)$. It is also easy to see that $V(R)_{(-2)}V(R)$ is in the kernel of this map, hence we get a map $V(R)/(V(R)_{(-2)}V(R)) \rightarrow S(R/TR)$ which is the inverse of φ . It is clear that φ is a Poisson algebra isomorphism.

Example 6.12. We present here an example of a regular family of vertex algebras $V(R)_\epsilon$ for which the Poisson algebras at the bottom left and right corners of diagram (6.16) are not isomorphic. Let $U = \bigoplus_{n \in \mathbb{Z}_+} U[n]$ be a graded unital commutative associative algebra with $U[0] = \mathbb{C}1$, and let H be the diagonalizable operator on U such that $H|_{U[n]} = nI_{U[n]}$. Then $V_\epsilon = \mathbb{C}[\epsilon] \otimes U$ is a regular family of vertex algebras with $|0\rangle = 1$, $T = \epsilon H$ and $a_{(n)}b = \delta_{n,-1}ab$. In this case $V = (U, |0\rangle = 1, T = H, Y(a, z) = L_a)$ and $V/V_{(-2)}V \simeq \mathbb{C}$, while $V^{\text{cl}} = (U, |0\rangle = 1, T = 0, Y(a, z) = L_a)$ (cf. Example 2.18) and $V^{\text{cl}}/(V_{(-2)}^{\text{cl}}V^{\text{cl}}) \simeq U$.

Returning to W -algebras, consider the family of W -algebras $W_k(\mathfrak{g}, x)_\epsilon$ defined as follows. Let $C_k(\mathfrak{g}, x)_\epsilon = V(R(\mathfrak{g}, x))_\epsilon$ be the regular family of vertex algebras associated to the non-linear Lie conformal algebra $R_k(\mathfrak{g}, x)$, as explained in Remark 6.9. Let d_ϵ be the element of $C_k(\mathfrak{g}, x)$ given by (5.5). Then $(C_k(\mathfrak{g}, x)_\epsilon, d_{\epsilon(0)})$ is a regular family of vertex algebras with a differential, and $W_k(\mathfrak{g}, x)_\epsilon$ is, by definition, the corresponding cohomology. Notice that $W_k(\mathfrak{g}, x)_\epsilon$ is, by the above construction, a vertex algebra over $\mathbb{C}[\epsilon]$. Moreover, by formality (cf. Remark 5.11), it is a vertex subalgebra of $C_k(\mathfrak{g}, x)_\epsilon$, hence it is a regular family of vertex algebras, according to Definition 6.7. By the observations in Section 5.1, $W_k(\mathfrak{g}, x)_\epsilon$ has a Hamiltonian operator defined by (5.3), and we take the Γ/\mathbb{Z} -grading of $W_k(\mathfrak{g}, x)_\epsilon$, induced by H (see Example 2.12).

We then build, starting from $W_k(\mathfrak{g}, x)_\epsilon$, the whole diagram (6.16) of fundamental objects. In particular, in the top right corner we have the usual affine W -algebra $W_k(\mathfrak{g}, x)$, while in the top left corner we have the *classical affine W -algebra* $W_k^{\text{cl}}(\mathfrak{g}, x)$. (We thus recover, in the case when f is a principal nilpotent element, the so called *classical Drinfeld–Sokolov reduction* [DS, FF1].) In the third row we have on the right the usual finite W -algebra $W^{\text{fin}}(\mathfrak{g}, x)$, and on the left the *classical finite W -algebra* $W^{\text{fin}, \text{cl}}(\mathfrak{g}, x)$. Moreover, in the bottom right corner we have a Poisson algebra which, by Proposition 3.30, is isomorphic to the associated graded of the finite W algebra $W^{\text{fin}}(\mathfrak{g}, x)$ with respect to the filtration (2.40) induced by the conformal weight, which coincides with the so called Kazhdan filtration. We thus recover the Poisson algebra on the Slodowy slice (see [GG, P1, P2]). Finally, we claim that the Poisson algebra at the bottom left corner is isomorphic to the one at the bottom right corner. To

see this, first observe that, by Example 6.11, they are isomorphic at the level of complexes, and then just notice that, since the differentials are derivations of all n^{th} products, taking cohomology commutes with taking quotients by $V_{(-2)}V$.

Appendix. Three equivalent definitions of finite W -algebras

A. D'Andrea, C. De Concini, A. De Sole, R. Heluani and V. Kac

We prove here that the definition of finite W -algebras via the Whittaker models, which goes back to [Ko] (see [P1, GG]), is equivalent to the definition presented in Section 5 of this paper (in fact, to the slightly more general definition, indicated in Section 0.4, with arbitrary coisotropic subspace $\mathfrak{s} \subset \mathfrak{g}_{1/2}$). For simplicity, we consider here the construction for a Lie algebra \mathfrak{g} , as the definition of finite W -algebras via the Whittaker models is known in the even case. However most of our arguments can be generalized to the super case.

We will use the notation introduced in Section 5: \mathfrak{g} is a simple, finite dimensional Lie algebra, with a non-degenerate symmetric invariant bilinear form $(\cdot | \cdot)$, and $x, f \in \mathfrak{g}$ form a "good" pair of elements of \mathfrak{g} , i.e. $\text{ad } x$ is diagonalizable on \mathfrak{g} with half integral eigenvalues and eigenspace decomposition $\mathfrak{g} = \bigoplus_{j \in \frac{1}{2}\mathbb{Z}} \mathfrak{g}_j$, and $f \in \mathfrak{g}_{-1}$ is such that $\mathfrak{g}^f = \{a \in \mathfrak{g} \mid [f, a] = 0\} \subset \bigoplus_{j \leq 0} \mathfrak{g}_j$. We choose an isotropic subspace $\mathfrak{l} \subset \mathfrak{g}_{1/2}$ with respect to the skewsymmetric non-degenerate bilinear form $(f|[a, b])$ on $\mathfrak{g}_{1/2}$, and denote by $\mathfrak{l}^\perp (= \mathfrak{s}) \subset \mathfrak{g}_{1/2}$ the orthogonal complement of \mathfrak{l} with respect to this bilinear form. Obviously $\mathfrak{l} \subset \mathfrak{l}^\perp$, and $(f|[a, b])$ induces a non-degenerate skewsymmetric bilinear form on $\mathfrak{l}^\perp/\mathfrak{l}$. We then let $\mathfrak{m}_\mathfrak{l} = \mathfrak{l} \oplus \mathfrak{g}_{\geq 1}$, and $\mathfrak{n}_\mathfrak{l} = \mathfrak{l}^\perp \oplus \mathfrak{g}_{\geq 1} \subset \mathfrak{g}$. They are obviously subalgebras of \mathfrak{g} , and $\mathfrak{m}_\mathfrak{l}$ is an ideal of $\mathfrak{n}_\mathfrak{l}$, with $\mathfrak{n}_\mathfrak{l}/\mathfrak{m}_\mathfrak{l} \simeq \mathfrak{l}^\perp/\mathfrak{l}$. (More generally, one can take a good \mathbb{R} -grading and let $\mathfrak{m}_\mathfrak{l} = \mathfrak{l} + \mathfrak{g}_{>1/2}$, $\mathfrak{n}_\mathfrak{l} = \mathfrak{l}^\perp + \mathfrak{g}_{>1/2}$, see Remark 5.4.)

The first definition, via quantum Hamiltonian reduction

Recall the Definition 5.6 of the finite W -algebra $W^{\text{fin}}(\mathfrak{g}, x)$. We can generalize it to an arbitrary choice of the isotropic subspace $\mathfrak{l} \subset \mathfrak{g}_{1/2}$ as follows. In analogy with the notation introduced in Section 5, we let $\varphi_{\mathfrak{n}_\mathfrak{l}} = \{\varphi_a \mid a \in \mathfrak{n}_\mathfrak{l}\}$ be the vector space $\mathfrak{n}_\mathfrak{l}$ with odd parity, $\varphi_{\mathfrak{n}_\mathfrak{l}^*} = \{\varphi^\phi \mid \phi \in \mathfrak{n}_\mathfrak{l}^*\}$ be the vector space $\mathfrak{n}_\mathfrak{l}^*$ with odd parity, and $\Phi_{\mathfrak{l}^\perp/\mathfrak{l}} = \{\Phi_{[a]} \mid [a] \in \mathfrak{l}^\perp/\mathfrak{l}\}$ be a vector superspace isomorphic to $\mathfrak{l}^\perp/\mathfrak{l}$. The usual pairing of $\mathfrak{n}_\mathfrak{l}$ and $\mathfrak{n}_\mathfrak{l}^*$ induces a non-degenerate skewsymmetric bilinear form on $\varphi_{\mathfrak{n}_\mathfrak{l}^*} \oplus \varphi_{\mathfrak{n}_\mathfrak{l}}$, which we denote by $\langle \cdot | \cdot \rangle_{ch}$, and, as notice above, the inner product $(f|[a, b])$ on $\mathfrak{g}_{1/2}$ induces a non-degenerate skewsymmetric bilinear form on $\Phi_{\mathfrak{l}^\perp/\mathfrak{l}}$, which we denote by $\langle \cdot | \cdot \rangle_{ne}$. We then define the non-linear Lie superalgebra

$$r = \mathfrak{g} \oplus \varphi_{\mathfrak{n}_\mathfrak{l}^*} \oplus \varphi_{\mathfrak{n}_\mathfrak{l}} \oplus \Phi_{\mathfrak{l}^\perp/\mathfrak{l}},$$

with non-linear Lie bracket given by

$$\begin{aligned} [a, b] &= \text{Lie bracket of } \mathfrak{g}, \quad a, b \in \mathfrak{g}, \\ [\varphi_a, \varphi^\phi] &= [\varphi^\phi, \varphi_a] = \phi(a) =: \langle \varphi_a | \varphi^\phi \rangle_{ch}, \quad a \in \mathfrak{n}_l, \phi \in \mathfrak{n}_l^*, \\ [\Phi_{[a]}, \Phi_{[b]}] &= (f|[a, b]) =: \langle \Phi_{[a]} | \Phi_{[b]} \rangle_{ch}, \quad [a], [b] \in \mathfrak{l}^\perp / \mathfrak{l}, \end{aligned}$$

and all other Lie brackets among generators equal zero. We denote by $C^I = U(r)$ the universal enveloping algebra of r , and we let d be the following odd element of C^I :

$$d = \sum_{\alpha \in S} \varphi^\alpha (u_\alpha + (f|u_\alpha) + \Phi_{[u_\alpha]}) + \frac{1}{2} \sum_{\alpha, \beta \in S} \varphi^\alpha \varphi^\beta \varphi_{[u_\beta, u_\alpha]} \in U(R),$$

where $\{u_\alpha\}, \{v^\alpha\}, \alpha \in S$, are dual bases of \mathfrak{n}_l and \mathfrak{n}_l^* , namely $v^\beta(u_\alpha) = \delta_{\alpha, \beta}$, $\varphi_\alpha = \varphi_{u_\alpha}$, $\varphi^\alpha = \varphi^{v^\alpha}$ for $\alpha \in S$, and $\Phi_{[a]}$, for $a \in \mathfrak{n}_l$, is defined via the isomorphism $\mathfrak{n}_l / \mathfrak{m}_l \simeq \mathfrak{l}^\perp / \mathfrak{l}$. It is not hard to check that $[d, d] = 0$, so that $d^I = \text{ad}(d)$ is a differential of the associative superalgebra C^I .

Definition A.1. The finite W -algebra is $W^I = H(C^I, d^I)$.

It is clear that, in the special case $\mathfrak{l} = 0$, Definition A.1 reduces to Definition 5.6.

Remark A.2. One can easily compute the action of d^I on the generators of C^I :

$$\begin{aligned} d^I(a) &= \sum_{\alpha} \varphi^\alpha [u_\alpha, a] \\ d^I(\varphi^\phi) &= \frac{1}{2} \sum_{\alpha} \varphi^\alpha \varphi^{u_\alpha \cdot \phi} \\ d^I(\varphi_a) &= a + (f|a) + \Phi_{[a]} + \sum_{\alpha} \varphi^\alpha \varphi_{[u_\alpha, a]} \\ d^I(\Phi_{[a]}) &= \sum_{\alpha} \varphi^\alpha (f|[u_\alpha, a]) \end{aligned}$$

In the second equation $a \cdot \phi$ denotes the coadjoint action of \mathfrak{n}_l on \mathfrak{n}_l^* , i.e. $(a \cdot \phi)(b) = \phi([b, a])$.

Remark A.3. For $a \in \mathfrak{n}_l$, we let $X_a = a + (f|a) + \Phi_{[a]}$. Then the map $a \mapsto X_a$ defines a Lie algebra isomorphism $\mathfrak{n}_l \simeq X_{\mathfrak{n}_l} \subset C^I$, namely $X_{[a, b]} = [X_a, X_b]$, $\forall a, b \in \mathfrak{n}_l$. Moreover, the action of d^I on $X_{\mathfrak{n}_l}$ is given by

$$d^I(X_a) = \sum_{\alpha} \varphi^\alpha X_{[u_\alpha, a]}.$$

The second definition, via Whittaker models

Let $\mathbb{C}_{\pm\chi} \simeq \mathbb{C}$ be the 1-dimensional representations of \mathfrak{m}_l given by $m \cdot 1 = \pm(f|m)$. Let M_l^0 be the left \mathfrak{g} -module induced by $\mathbb{C}_{-\chi}$:

$$M_l^0 = \text{Ind}_{\mathfrak{m}_l}^{\mathfrak{g}} \mathbb{C}_{-\chi} = U(\mathfrak{g}) \otimes_{U(\mathfrak{m}_l)} \mathbb{C}_{-\chi}. \quad (\text{A.1})$$

$M_{\mathfrak{l}}^0$ is, by definition, a left \mathfrak{g} -module via left multiplication. However it is easy to check that $M_{\mathfrak{l}}^0$ is also a left $\mathfrak{n}_{\mathfrak{l}}$ -module with respect to the adjoint action of $\mathfrak{n}_{\mathfrak{l}}$ on $U(\mathfrak{g})$.

Theorem A.4. [GG] *The Lie algebra cohomology of $\mathfrak{n}_{\mathfrak{l}}$ with coefficients in $M_{\mathfrak{l}}^0$ is concentrated at degree 0:*

$$H(\mathfrak{n}_{\mathfrak{l}}, M_{\mathfrak{l}}^0) = (M_{\mathfrak{l}}^0)^{\mathfrak{n}_{\mathfrak{l}}} ,$$

where the space of $\text{ad } \mathfrak{n}_{\mathfrak{l}}$ -invariants $(M_{\mathfrak{l}}^0)^{\mathfrak{n}_{\mathfrak{l}}}$ is an associative algebra with product induced from $U(\mathfrak{g})$. Moreover this algebra is independent of the choice of \mathfrak{l} , up to isomorphism.

Definition A.5. The finite W -algebra is $W^{II} = (M_{\mathfrak{l}}^0)^{\mathfrak{n}_{\mathfrak{l}}}$.

By Theorem A.4, W^{II} is the cohomology of the complex

$$C^{II} = \Lambda(\mathfrak{n}_{\mathfrak{l}}^*) \otimes (U(\mathfrak{g}) \otimes_{U(\mathfrak{m}_{\mathfrak{l}})} \mathbb{C}_{-\chi}) ,$$

with differential

$$\begin{aligned} d^{II}(\Psi \otimes (A \otimes_{U(\mathfrak{m}_{\mathfrak{l}})} 1)) &= \frac{1}{2} \sum_{\alpha} v^{\alpha} \wedge u_{\alpha} \cdot \Psi \otimes (A \otimes_{U(\mathfrak{m}_{\mathfrak{l}})} 1) \\ &\quad + \sum_{\alpha} v^{\alpha} \wedge \Psi \otimes (\text{ad } u_{\alpha}(A) \otimes_{U(\mathfrak{m}_{\mathfrak{l}})} 1) . \end{aligned}$$

As before, $a \cdot \Psi$ denotes the coadjoint action of $\mathfrak{n}_{\mathfrak{l}}$ on $\mathfrak{n}_{\mathfrak{l}}^*$, extended to $\Lambda(\mathfrak{n}_{\mathfrak{l}}^*)$.

In the rest of the Appendix we will prove the following

Theorem A.6. *There is an associative algebra isomorphism $W^I \simeq W^{II}$.*

The third definition of finite W -algebras

We present here an equivalent definition of the complex (C^I, d^I) defining the algebra W^I , which better enlightens the relation to the complex (C^{II}, d^{II}) and the corresponding cohomology W^{II} . Let $U(\mathfrak{n}_{\mathfrak{l}})_{-\chi}$ be the space $U(\mathfrak{n}_{\mathfrak{l}})$ with left action of $\mathfrak{m}_{\mathfrak{l}}$ given by $m \cdot A = (m - (f|m))A$, namely, as $\mathfrak{m}_{\mathfrak{l}}$ -module, $U(\mathfrak{n}_{\mathfrak{l}})_{-\chi} \simeq U(\mathfrak{n}_{\mathfrak{l}}) \otimes \mathbb{C}_{-\chi}$. Consider the induced \mathfrak{g} -module,

$$M_{\mathfrak{l}} = \text{Ind}_{\mathfrak{m}_{\mathfrak{l}}}^{\mathfrak{g}} U(\mathfrak{n}_{\mathfrak{l}})_{-\chi} = U(\mathfrak{g}) \otimes_{U(\mathfrak{m}_{\mathfrak{l}})} U(\mathfrak{n}_{\mathfrak{l}})_{-\chi} .$$

$M_{\mathfrak{l}}$ is clearly a left \mathfrak{g} -module by left multiplication. Moreover, a simple computation shows that $M_{\mathfrak{l}}$ has also the structure of left $\mathfrak{n}_{\mathfrak{l}}$ -module, induced by the adjoint action of $\mathfrak{n}_{\mathfrak{l}}$ on $U(\mathfrak{g})$, and of right $\mathfrak{n}_{\mathfrak{l}}$ -module, given by the right multiplication of $\mathfrak{n}_{\mathfrak{l}}$ on $U(\mathfrak{n}_{\mathfrak{l}})$. Notice that

$$M_{\mathfrak{l}} \simeq U(\mathfrak{g})_{\chi} \otimes_{U(\mathfrak{m}_{\mathfrak{l}})} U(\mathfrak{n}_{\mathfrak{l}}) , \tag{A.2}$$

where $\mathfrak{m}_{\mathfrak{l}}$ acts on $U(\mathfrak{n}_{\mathfrak{l}})$ by left multiplication, and on $U(\mathfrak{g})_{\chi} \simeq U(\mathfrak{g}) \otimes \mathbb{C}_{\chi}$ by right action. Similarly

$$M_{\mathfrak{l}}^0 \simeq U(\mathfrak{g})_{\chi} \otimes_{U(\mathfrak{m}_{\mathfrak{l}})} \mathbb{C} . \tag{A.3}$$

We then define, starting from M_l , a mixed Lie algebra cohomology and homology complex as follows. We first define the complex (C^h, d^h) for the Lie algebra homology of \mathfrak{n}_l with coefficients on the right module M_l , namely

$$C^h = M_l \otimes \Lambda(\mathfrak{n}_l) ,$$

and d^h is the following differential on C^h :

$$\begin{aligned} d^h(m \otimes c_1 \wedge \cdots \wedge c_s) &= \sum_{1 \leq k \leq s} (-1)^k m \cdot c_k \otimes c_1 \wedge \cdots \wedge c_s \\ &+ \sum_{1 \leq j < k \leq s} (-1)^{j+k+1} m \otimes [c_j, c_k] \wedge c_1 \wedge \cdots \wedge c_s . \end{aligned} \quad (\text{A.4})$$

Clearly C^h is again a left \mathfrak{n}_l -module, with respect to the adjoint action of \mathfrak{n}_l , hence we can talk about the Lie algebra cohomology of \mathfrak{n}_l with coefficients on C^h . We thus get a new complex (C^c, d^c) , where

$$C^c = \Lambda(\mathfrak{n}_l^*) \otimes M_l \otimes \Lambda(\mathfrak{n}_l) ,$$

and d^c is the usual Lie algebra cohomology differential:

$$\begin{aligned} d^c(\Psi \otimes m \otimes C) &= \frac{1}{2} \sum_{\alpha} v^{\alpha} \wedge u_{\alpha} \cdot \Psi \otimes m \otimes C \\ &+ \sum_{\alpha} v^{\alpha} \wedge \Psi \otimes \text{ad } u_{\alpha}(m \otimes C) . \end{aligned} \quad (\text{A.5})$$

We finally define a new complex (C^{III}, d^{III}) , where C^{III} is the same as above,

$$C^{III} = \Lambda(\mathfrak{n}_l^*) \otimes (U(\mathfrak{g}) \otimes_{U(\mathfrak{m}_l)} U(\mathfrak{n}_l)_{-\chi}) \otimes \Lambda(\mathfrak{n}_l) ,$$

and d^{III} is a mixed cohomology-homology differential:

$$d^{III} = d^c + (-1)^{\delta-1} \otimes d^h ,$$

with d^c as in (A.5), d^h as in (A.4), and δ denotes the usual grading of $\Lambda(\mathfrak{n}_l^*)$.

Proposition A.7. *There is an isomorphism of complexes $(C^{III}, d^{III}) \rightarrow (C^I, d^I)$ given by the following linear map*

$$\begin{aligned} i &: (\phi^1 \wedge \cdots \wedge \phi^i) \otimes (a_1 \dots a_s \otimes_{U(\mathfrak{m}_l)} b_1 \dots b_t) \otimes (c_1 \wedge \cdots \wedge c_j) \\ &\mapsto \varphi^{\phi^1} \cdots \varphi^{\phi^i} a_1 \dots a_s X_{b_1} \dots X_{b_t} \varphi_{c_1} \cdots \varphi_{c_j} \end{aligned}$$

Proof. It is immediate to check that i is well defined. It is obviously surjective, and it is injective thanks to the PBW Theorem for $U(r)$. We are left to prove that $i \circ d^{III} = d^I \circ i$. This is a straightforward computation, which can be done using the formulas in Remark A.2. \square

Proof of Theorem A.6. Thanks to Proposition A.7, in order to prove Theorem A.6 we need to show that (C^{III}, d^{III}) and (C^{II}, d^{II}) are quasi-isomorphic complexes, and the induced isomorphism of cohomologies $W^{III} \rightarrow W^{II}$ is an isomorphism of associative algebras. By definition, (C^{III}, d^{III}) is a bicomplex:

$$C^{III} = \bigoplus_{p \geq 0, q \leq 0} C^{p,q}, \quad d^{III} = d_1 + d_2,$$

where

$$C^{p,q} = \Lambda^p(\mathfrak{n}_l^*) \otimes (U(\mathfrak{g}) \otimes_{U(\mathfrak{m}_l)} U(\mathfrak{n}_l)_{-\chi}) \otimes \Lambda^{-q}(\mathfrak{n}_l),$$

and $d_1 = d^c$, $d_2 = (-1)^{\delta-1} \otimes d^h$ so that

$$d_1^2 = d_2^2 = d_1 d_2 + d_2 d_1 = 0,$$

and $d_1 : C^{p,q} \rightarrow C^{p+1,q}$, $d_2 : C^{p,q} \rightarrow C^{p,q+1}$. Notice that d^c and d^h commute, since d^c is defined in terms of the left action of \mathfrak{n}_l on M_l , while d^h is defined in terms of the right action of \mathfrak{n}_l on M_l . We can then study the corresponding spectral sequence $(E_r, d^r; r \geq 1)$. The first term of the spectral sequence is

$$\begin{aligned} E_1^{p,q} &= H^{p,q}(C^{III}, d^c) \\ &= \Lambda^p(\mathfrak{n}_l^*) \otimes H^q((U(\mathfrak{g}) \otimes_{U(\mathfrak{m}_l)} U(\mathfrak{n}_l)_{-\chi}) \otimes \Lambda(\mathfrak{n}_l), d^h) \\ &= \Lambda^p(\mathfrak{n}_l^*) \otimes H^q((U(\mathfrak{g})_{\chi} \otimes_{U(\mathfrak{m}_l)} U(\mathfrak{n}_l)) \otimes \Lambda(\mathfrak{n}_l), d^h) \\ &= \Lambda^p(\mathfrak{n}_l^*) \otimes (U(\mathfrak{g})_{\chi} \otimes_{U(\mathfrak{m}_l)} H^q(U(\mathfrak{n}_l) \otimes \Lambda(\mathfrak{n}_l), d^h)) \\ &= \delta_{q,0} \Lambda^p(\mathfrak{n}_l^*) \otimes (U(\mathfrak{g})_{\chi} \otimes_{U(\mathfrak{m}_l)} \mathbb{C}) \\ &= \delta_{q,0} \Lambda^p(\mathfrak{n}_l^*) \otimes (U(\mathfrak{g}) \otimes_{U(\mathfrak{m}_l)} \mathbb{C}_{-\chi}) = \delta_{q,0} C^{II,p}. \end{aligned}$$

In the second and fifth equalities above we used (A.2) and (A.3) respectively. The third equality is obvious, since d^h is defined in terms of the right action of \mathfrak{n}_l on $U(\mathfrak{g})_{\chi} \otimes_{U(\mathfrak{m}_l)} U(\mathfrak{n}_l)$ given by right multiplication on $U(\mathfrak{n}_l)$. Finally, the fourth equality follows by the isomorphism $H_q(\mathfrak{n}, U(\mathfrak{n})) \simeq \delta_{q,0} \mathbb{C}$. The corresponding differential on $E_1^{p,q} = \delta_{q,0} C^{II,p}$ is $d_1 = d^c = d^{II}$.

Since the complex E_1 is concentrated at $q = 0$, it immediately follows that the spectral sequence stabilizes at $r = 2$, namely $E_{\infty} = E_2$. Moreover, since the bigrading of C^{III} is bounded, we also know that

$$E_{\infty} \simeq \text{gr } H(C^{III}, d^{III}) \simeq H(C^{III}, d^{III}).$$

In conclusion, we have $H(C^{III}, d^{III}) \simeq E_2 \simeq H(C^{II}, d^{II})$.

The above arguments, together with Proposition A.7, show that the complexes (C^I, d^I) and (C^{II}, d^{II}) are quasi-isomorphic. In fact, we can write down an explicit quasi-isomorphism $\pi : (C^I, d^I) \rightarrow (C^{II}, d^{II})$ as follows:

$$\begin{aligned} \pi(\varphi^{\phi^1} \cdots \varphi^{\phi^i} a_1 \cdots a_s X_{b_1} \cdots X_{b_t} \varphi_{c_1} \cdots \varphi_{c_j}) \\ = \delta_{t,0} \delta_{j,0} \phi^1 \wedge \cdots \wedge \phi^i \otimes (a_1 \cdots a_s \otimes_{U(\mathfrak{m}_l)} 1). \end{aligned}$$

We are left to prove that the corresponding isomorphism of cohomologies $H(C^I, d^I) \simeq H(C^{II}, d^{II})$ preserves the associative algebra structures. Recall that in $H(C^{II}, d^{II})$ the associative product is induced by $U(\mathfrak{g})$, namely if $(A \otimes_{U(\mathfrak{m}_I)} 1), (B \otimes_{U(\mathfrak{m}_I)} 1) \in (M_I^0)^{\mathfrak{n}_I} \simeq H(C^{II}, d^{II})$, then $(A \otimes_{U(\mathfrak{m}_I)} 1) \cdot (B \otimes_{U(\mathfrak{m}_I)} 1) = (AB) \otimes_{U(\mathfrak{m}_I)} 1$. On the other hand, if we pick appropriate preimages in the complex (C^I, d^I) , say A and B , their product is AB . The claim follows immediately.

□

References

- [A1] T. Arakawa, *Representation theory of superconformal algebras and the Kac–Roan–Wakimoto conjecture*, math-ph/0405015.
- [A2] T. Arakawa, *Representation theory of W -algebras*, math.QA/0506056.
- [AM] A. Alekseev and E. Meinrenken, *The non-commutative Weil algebra*, Invent. Math. 139 (2000), 135-172.
- [BK] B. Bakalov and V. G. Kac, *Field algebras*, IMRN 3 (2003), 123-159.
- [BrG] J. Brundan and S. M. Goodwin, *Good gradings polytopes*, math.QA/0510205.
- [BrK] J. Brundan and A. Kleshchev, *Shifted Yangians and finite W -algebras*, Adv. Math. (to appear)
- [B] R. Borchers, *Vertex algebras, Kac-Moody algebras and the Monster*, Proc. Natl. Acad. Sci. USA 83(1986), 3068-3071.
- [BT1] J. de Boer and T. Tjin, *Quantization and representation theory of finite W -algebras*, Comm. Math. Phys. 158 (1993), 485-516
- [BT] J. de Boer and T. Tjin, *The relation between quantum W -algebras and Lie algebras*, Comm. Math. Phys. 160 (1994), 317-332 .
- [BS] P. Bouwknegt and K. Schoutens, *W -symmetry*, Advanced ser. in Math. Phys. 22, World Sci., 1995.
- [D’K] A. D’Andrea and V. G. Kac, *Structure theory of finite conformal algebras*, Selecta Math 4 (1998), 377-418.
- [DK] A. De Sole and V.G. Kac, *Freely generated vertex algebras and non-linear Lie conformal algebras*, Comm. Math. Phys. 254(2005), 659-694.
- [DV] K. de Vos and P. van Driel, *The Kazhdan-Lusztig conjecture for finite W -algebras*, Lett. in Math. Physics 35 (1995), 333-344.

- [DLM] C. Dong, H. Li and G. Mason, *Twisted representations of vertex operator algebras*, Math. Ann. 310 (1998), 571-600.
- [DS] V. G. Drinfeld and V.V. Sokolov, *Lie algebra and the KdV type equations*, Soviet J. Math. 30(1985), 1975-2036.
- [EK] A. G. Elashvili and V.G. Kac, *Classification of good gradings of simple Lie algebras*, Amer. Math. Soc. Transl (2) vol 213 (2005), 85-104, math-ph/031203.
- [FF1] B.L. Feigin and E. Frenkel, *Quantization of Drinfeld-Sokolov reduction*, Phys. Lett., B 246 (1990), 75-81.
- [FF2] B.L. Feigin and E. Frenkel, *Affine Kac-Moody algebras, bosonization and resolutions*, Lett. Math. Phys. 19 (1990), 307-317.
- [FF3] B.L. Feigin and E. Frenkel, *Integrals of motions and quantum groups*, in Lect. Notes in Math 1620 (1995), 349-418.
- [FB] E. Frenkel and D. Ben-Zvi, *Vertex algebras and algebraic curves*, AMS monographs 88, 2001. Second edition, 2004.
- [FKW] E. Frenkel, V. G. Kac and M. Wakimoto, *Characters and fusion rules for W -algebras via quantized Drinfeld-Sokolov reduction*, Comm. Math. Phys. 147 (1992), 295-328.
- [FHL] I. Frenkel, Y.-Z. Huang and J. Lepowsky, *On axiomatic approach to vertex operator algebras and modules*, Mem. Amer. Math. Soc. 104, 1993.
- [GG] W.L. Gan and V. Ginzburg, *Quantization of Slodowy slices*, IMRN 5 (2002), 243-255.
- [GMS] V. Gorbunov, F. Malikov, and V. Schechtman, *Gerbres of chiral differential operators II*, Invent. Math. 155(2004), 605-680.
- [J] N. Jacobson, *Lie algebras*, Interscience, New York, 1962.
- [K] V. G. Kac, *Vertex algebras for beginners*, Providence: AMS, University Lecture Notes, Vol. 10, 1996. Second edition, 1998.
- [KT] V. G. Kac and I. Todorov, *Superconformal current algebras and their unitary representations*, Comm. Math. Phys. 102 (1985), 337-347.
- [KRW] V. G. Kac, S.-S. Roan and M. Wakimoto, *Quantum reduction for affine superalgebras*, Comm. Math. Phys., 241 (2003), 307-342.
- [KW1] V. G. Kac and M. Wakimoto, *Quantum reduction and representation theory of superconformal algebras*, Adv. Math. 185 (2004), 400-458. Corrigendum, Adv. Math. 193 (2005), 453-455.

- [KW2] V. G. Kac and M. Wakimoto, *Quantum reduction in the twisted case*, in Progress in Math. 237, 2005, pp 85-126.
- [KWa] V.G. Kac and W. Wang, *Vertex operator superalgebras and their representations*, Contemp. Math. 175 (1994), 161-191.
- [Ko] B. Kostant, *On Whittaker vectors and representation theory*, Inv. Math 48 (1978), 101-184.
- [Ko1] B. Kostant, *A cubic Dirac operator and the emergence of Euler number multiplets of representations of equal rank subgroups*, Duke Math. J. 100 (1999), 447-501.
- [KS] B. Kostant and S. Sternberg, *Symplectic reduction, BRS cohomology and infinite-dimensional Clifford algebras*, Ann. Phys. 176 (1987), 49-113.
- [M] H. Matumoto, *Whittaker modules associated with highest weight modules*, Duke Math. J. 60 (1990), 59-113.
- [P1] A. Premet, *Special transverse slices and their enveloping algebras*, Adv. Math. 170 (2002), 1-55.
- [P2] A. Premet, *Enveloping algebras of Slodowy slices and the Joseph ideal*, 2005, preprint
- [RS] E. Ragoucy and P. Sorba, *Yangian realizations from finite W -algebras*, Comm. Math. Phys. 203 (1999), 551-576.
- [ST] A. Severin and W. Troost, *Extensions of Virasoro algebra and gauged WZW models*, Phys. Lett. B315 (1993), 304-310.
- [Za] A. Zamolodchikov, *Infinite extra symmetries in two-dimensional conformal quantum field theory*, Teor. Mat. Fiz 65(3)(1985), 347-359.
- [Z] Y. Zhu, *Modular invariance of characters of vertex operator algebras*, Journal AMS 9(1996), 237-302.