### FEYNMAN CYCLES IN THE BOSE GAS

#### DAN EL UELT SCHI

A bstract. We study the lengths of the cycles formed by trajectories in the Feynman-Kacrepresentation of the Bosegas. We discuss the occurrence of in nite cycles and their relation to Bose-Einstein condensation.

Keywords: Bose-Einstein condensation. Feynman cycles. In nite cycles.

2000 M ath. Subj. C lass.: 82B 10, 82B 21, 82B 26, 82D 50.

PACS num bers: 03.75 Hh, 05.30.-d, 05.30 Jp, 05.70 Fh, 31.15 Kb

### 1. Introduction

Bose and E instein understood 80 years ago that a curious phase transition occurs in a gas of non-interacting bosons; it is now commonly referred to as Bose-E instein condensation. Real particles interact, however, and form any years there were doubts that this transition takes place in natural systems. London suggested in 1938 that super uid Helium undergoes a Bose-E instein condensation, and this idea is largely accepted now adays. Bogolubov considered interacting systems; careful approximations allowed him to get back to a non-interacting gas, but with a dierent dispersion relation. See [16] and [8] form ore discussion and partial justications of Bogolubov theory.

In 1953 Feynm an studied the system in the Feynm an-K ac representation [5]. The partition function can be expanded as a gas of trajectories living in (d+1) dimensions. The extra dimension is commonly referred to as the time", although it is not related to physical time. The situation is illustrated in Fig. 1. A nite system with N particles induces a probability on the group  $S_N$  of permutations of N elements. Feynm an considered the probability for a given particle to belong to a cycle of length n. In the therm odynam ic limit, there may be strictly positive probability for in nite cycles to be present, and Feynm an suggested to use this as an order parameter for Bose-E instein condensation.

A few years later, in 1956, Penrose and Onsager introduced the concept of  $\$  -diagonal long-range order" [9]. Form ally, it is a correlation between positions x and y given by  $(x;y) = hc^y(x)c(y)i$ . The system displays o -diagonal long-range order when this correlation is strictly positive, uniform ly in the size of the system and in  $\dot{y}$  yj! 1. One can write a Feynm an-K ac version of this correlation, and it involves a special cycle starting at x and ending at y; this cycle m ay wind m any times around the imaginary time direction. In the limit where x and y are in nitely distant there corresponds a notion of in nite open cycle that is reminiscent of Feynman's approach.

Feynm an's order parameter is simpler; it is often used in numerical simulations or in order to gain heuristic understanding. On the other hand, everybody agrees that Penrose and Onsager order parameter is the correct one. Surprisingly, the question of their equivalence is usually eluded, and many physicists implicitely assume equivalence to hold. The rst mathematical investigation of this question is due to Suto, who showed that equivalence holds in the ideal gas. Indeed, he proved that in nite cycles occur in the presence of condensation [12], and that no in nite cycles occur in the absence of

1




Figure 1. The Feynm an-K ac representation of the partition function for a gas of bosons. The horizontal plane represents the d spatial dimensions, and the vertical axis is the imaginary timed imension. The picture shows a situation with veparticles and two cycles, of respective lengths 4 and 1.

condensation [13]; the latter result uses probabilistic methods from the theory of large deviations. These results have been extended to mean-eld systems in [1] and [3].

In this paper we explore the links between Feynm an cycles and o -diagonal long-range order. Let (x) denote the o -diagonal correlation between the origin and  $x \ 2 \ R^d$ , and % (n) denote the density of particles in cycles of length  $n \cdot W$  e propose the following form ula that relates both concepts:

$$(x) = X$$
 $c_n (x)% (n) + c_1 (x)% (1)$ : (1.1)

M athem atically, the problem is not well posed. M any choices for the coe cients  $c_n$  are possible  $\mid$  a trivial choice is  $c_n(x) = (x) = \text{ for all } n$ , including n = 1. We will see, however, that there is a natural de nition for  $c_n(x)$  in terms of W iener trajectories. In any case, we conjecture that Eq. (1.1) holds with coe cients satisfying

$$0.6 c_n (x) 6.1;$$
  $0.6 c_1 (x) 6.1;$ 

for all n;x. In addition, we should have

$$\lim_{n \to 1} c_n(x) = c_1(x)$$

for any x, and

$$\lim_{j \in j!} c_n(x) = 0$$

for any niten, but not uniform by in n; c<sub>1</sub> (x) m ay converge to a strictly positive constant c. If c = 1, we get from the dominated convergence theorem that  $\lim_{|x|_1} (x) = %(1)$  in which case the o -diagonal long-range order parameter is equal to the density of cycles of in nite lengths.

We establish this formula and these properties in the case of the ideal gas, where we show that

$$c_n(x) = e^{x^2-4n}$$
;  $c_1(x) = 1$ : (1.2)

We discuss the validity of the formula (1.1) in the interacting gas, proving that these properties hold true in a regime without Bose-Einstein condensation. The two order parameters should not be always equivalent, however. It is argued in [15] that they dier

when the bosons undergo a regular condensation into a crystalline phase. There is no o -diagonal long-range order, but in nite cycles may be present.

We work in the Feynman-Kac representation of the Bose gas. This representation is standard, see e.g. [4] for a clear and concise review, and [6] for a complete introduction. We assume the reader to possess some familiarity with it and in Section 2 we directly dene the main expressions | partition functions, density of cycles, o -diagonal long-range order | in terms of space-time trajectories. But basic notions and properties are reviewed in Appendix A.

The situation simplies in absence of interactions; we consider the ideal gas in Section 3, where we state and prove the formula that relates the two order parameters. The ideal gas is best discussed in the canonical ensemble. Rigorous proofs of macroscopic occupation of the zero mode have been proposed and they involve the grand-canonical ensemble, with a chemical potential that depends on the volume. Appendix B proposes a simple proof in the canonical ensemble.

Interacting systems constitute a form idable challenge; they are discussed in Section 4, where partial results are obtained.

In this paper, we denote nite volume expressions in plain characters, and in nite volume expressions in bold characters. Further, we always consider the canonical and grand-canonical ensembles where the temperature 1= is xed; we alleviate the notation by om itting the dependence of all quantities.

# 2. Feynman cycles and off-diagonal long-range order

2.1. Partition functions. Our Bose gas occupies a d-dim ensional dom ain D , always a cubic box of size L and volum e V =  $L^d$ . W e consider periodic boundary conditions. Let denote the particle density, the inverse temperature, and the chemical potential. The canonical partition function in the Feynman-K ac representation is given by

$$Y (N) = \frac{X^{N}}{k!} \frac{1}{k!} X Z dx_{1} ::: dx_{k} dW_{x_{1}x_{1}}^{n_{1}} (!_{1}) ::: dW_{x_{k}x_{k}}^{n_{k}} (!_{k})$$

$$Y^{k} \frac{1}{n_{1}} e^{U(!_{j})} Y e^{U(!_{i};!_{j})} : (2.1)$$

$$y^{k} \frac{1}{n_{j}} e^{U(!_{j})} dx_{k} dx$$

This expression is illustrated in Fig. 1. In words, we sum over the number k of closed trajectories and over their respective winding numbers  $n_1; \ldots; n_k$ . We integrate over the initial positions  $x_1; \ldots; x_k$ . We integrate over trajectories  $!_j : [0; n_j] !$  D that start and end at  $x_j$ ; here, W  $_{xx}$  denotes the W iener measure. See Appendix A form ore information, and in particular Eq. (A 10) for the normalization condition. Trajectories wind around the time direction according to their winding numbers; because of periodic boundary conditions, they may also wind around space directions.

G iven a trajectory! with winding numbern, the function U(!) denotes the interactions between dierent legs; explicitly,

$$U(!) = X \frac{1}{0.6 \text{ is j 6 n 1}} \frac{1}{0} U(!) = (i + s) (j + s) ds: (2.2)$$

And U (!; !  $^{0}$ ) denotes the interactions between closed trajectories! and !  $^{0}$ , of respective winding numbers n and n  $^{0}$ :

$$U(!;!^{0}) = X \frac{1}{0.6 \text{ i.6 n.1}\atop 0.6 \text{ j.6 n}^{0.1}} U(!;!^{0}) = X \frac{1}{0.6 \text{ i.6 n}^{0.1}} U(!;!^{0}) + S) ds: (2.3)$$

The function U (x) represents the pair interaction potential between two particles separated by a distance  $j_{x}j_{y}$ . We suppose that U (x) is nonnegative and spherically sym metric. We can allow the value + 1; all that is needed is that  $e^{U(!)}$  and  $e^{U(!)}$  be measurable functions with respect to the Wiener measure any piecewise continuous function D! [0;1] can be considered at this point.

The grand-canonical partition function is

$$Z() = {\begin{array}{c} X \\ e {\phantom{-}}^{N} & Y(N) \\ N > 0 \end{array}}$$
 (2.4)

(with the understanding that Y (0) = 1). We also need partition functions where a given trajectory  $!_0$  is present | these will be needed in the expression for cycle densities, see (2.8) and (2.9). Namely, we do ne

$$Y (N;!_{0}) = \frac{X^{N}}{k!} \frac{1}{k!} X Z X_{1} X_{1} X_{1} X_{1} X_{2} X_{1} X_{2} X_{2} X_{3} X_{4} X_{1} X_{1} X_{2} X_{3} X_{4} X_{5} X_{5}$$

The dependence on  $!_0$  comes from the last term, where the product includes terms with i=0. Notice that

$$Y (N ;!) 6 Y (N);$$
 (2.6)

with equality i U(x) 0, i.e. in absence of interactions. Finally, we introduce

$$Z(;!) = X e^{N} Y(N;!)$$
 (2.7)

(we set Y  $(0;!_0) = 1$ ). We also have Z (;!) 6 Z (), with equality i U (x) 0.

2.2. Cycle lengths. We now introduce the density of particles in cycles of length n, both in the canonical and grand-canonical ensembles. We denote the particle density by  $=\frac{N}{V}$ . When discussing the canonical ensemble, we always suppose that and V are such that N=V is an integer. The number of particles in cycles of length n is given by the random variable  $\sum_{j=1}^k n_{n_j m}$ . A veraging over all congurations of space-time closed trajectories,

we get

The last line follows from the rst line by replacing  $\frac{P_k}{j=1} n_{n_j,n}$  with  $nk_{n_1,n}$ ; isolating the integral over  $!_1$ ; using the de nition (2.5); using translation invariance, and  $_{D}$  dx<sub>1</sub> = V. Sim ilarly, we have the grand-canonical expression

% 
$$(n) = e^{-n} dW_{00}^{n} (!) e^{U(!)} \frac{Z(;!)}{Z()}$$
: (2.9)

One easily checks that

$$X$$
 %  $(n) = \frac{N}{V}$  ;  $x > 1$  %  $(n) = h\frac{N}{V}i$  ( ):  $x > 1$ 

We consider the thermodynamic limits of % (n) and % (n). Since 0 6 % (n) 6 , and since n is a discrete index, C antor diagonal process yields the existence of a sequence of increasing volum es  $V_k$ , with  $V_k = N_k$  an integer, such that % (n) converges to some  $\lim$  it that we denote % (n). Sim ilarly, we also obtain the in nite volume lim it % (n). Fatou's lem m a im plies that

This suggests to de ne the density of particles in in nite cycles by

% (1) = 
$$\binom{n > 1}{X}$$
 % (n);  
% (1) = () % (n):

The main question is whether % (1) diers from zero at given temperature, and at given density or chem ical potential.

W e chose to discuss densities of particles in cycles of given length, but one m ay consider probabilities as well. Namely, we could introduce the probability for particle 1 to belong

Thus % (n) = P (n) in the canonical ensemble, and % (1) = P (1). Things are not so simple in the grand-canonical ensemble. The probability P (n) is

$$Z Z Z = \frac{Z Z}{D} = \frac{Z Z}{D} \times \frac{Z}{D} = \frac{Z^{0}(;!)}{D} = \frac{Z^{0}(;!)}{Z(;!)} = \frac{Z^$$

Here,  $Z^0$  (;!) is like Z (;!) given in Eqs (2.7) and (2.5), but with a factor  $\frac{1}{(k+1)!}$  instead of  $\frac{1}{k!}$ . Heuristically, we should have  $h^{\underline{V}}_{nk}i=1=$  (), and % (n) = ()P (n), but this does not seem easy to establish. The ratio of partition functions in (2.13) is more discuss densities.

2.3. O -diagonal long-range order. Let us turn to Penrose and Onsager o -diagonal long-range order. Its Feynm an-K ac representation involves an open trajectory that starts at x and ends at y, that possibly winds several times around the time direction. Precisely, we introduce

$$(x) = \begin{cases} x^{N} & Z \\ dW_{0x}^{n} & (!) e^{U(!)} & \frac{Y(N - n;!)}{Y(N)}; \\ x^{n=1} & Z \\ X & e^{n} & dW_{0x}^{n} & (!) e^{U(!)} & \frac{Z(;!)}{Z(;)}; \end{cases}$$

$$(2.14)$$

Therm odynam ic lim its are denoted (x) and (x), provided they exist! One may actually restrict (x) and (x) on rational x, and use the C antor diagonal process to get convergence on a subsequence of increasing volumes. This is not necessary in this paper, as the lim its will be shown to exist in the regimes of parameters under consideration.

Sim ilarities between Eqs (2.8), (2.9) on the one hand, and Eqs (2.14) on the other hand, are manifest. We can write

where the coe cients  $c_n$ ; ,  $c_n$ ; are given by

$$c_{n}; (x) = \begin{bmatrix} z \\ dW_{00}^{n} (!) e^{U(!)} & \frac{Y(N - n;!)}{Y(N)} \end{bmatrix}^{1} \begin{bmatrix} z \\ dW_{0x}^{n} (!) e^{U(!)} & \frac{Y(N - n;!)}{Y(N)} \end{bmatrix};$$

$$c_{n}; (x) = \begin{bmatrix} z \\ dW_{00}^{n} (!) e^{U(!)} & \frac{Z(;!)}{Z()} \end{bmatrix}^{1} \begin{bmatrix} z \\ dW_{0x}^{n} (!) e^{U(!)} & \frac{Z(;!)}{Z()} \end{bmatrix};$$

$$(2.16)$$

As above, we denote the therm odynam ic limits by  $c_n$ ; (x) and  $c_n$ ; (x), provided they exist. One should be careful when sending the volume to in nity in Eqs (2.15), because a \leak to in nity" may yield a term involving % (1 ) | this actually occurs in the ideal gas, as shown in the next section.

## 3. The ideal gas

The ideal gas of quantum bosons is fascinating. Particles do not interact, yet they manage to display a phase transition. Historically, the Bose-Einstein condensation is the rst theoretical description of a phase transition. The ideal gas has been the object of many studies over the years; let us mention [17, 7, 10]. A simple proof of macroscopic occupation of the zero Fourier mode is presented in Appendix B.

In this section we elucidate the relation between cycle lengths and o -diagonal long-range order, thus clarifying results that were previously obtained by Sut) [12, 13]. We work

in the canonical ensemble and establish the formula (1.1) explicitely, for any dimension d > 1.

Theorem 1. For any 0 < ; < 1 , there exists a sequence of increasing cubes for which the therm odynam ic lim its of (x),  $c_n$ ; (x), % (n) exist for all x 2  $R^d$  and n 2 N. Further, we have

$$(x) = X$$
 $e^{\frac{x^2}{4n}} % (n) + % (1)$ :

The rest of this section is devoted to the proof of Theorem 1. The coe cient  $c_n$ ; (x), de ned in Eq. (2.16), has a simpler expression in absence of interactions. Indeed, we have U (!) = 0 and Y (N;!) = Y (N). It follows from properties of the W iener measure in periodic boxes, see Eq. (A.10), that

$$c_{n}$$
;  $(x) = \begin{cases} X & e^{\frac{L^{2}}{4n}(\frac{x}{L}z)^{2}} & X \\ & z^{2}z^{d} \end{cases}$   $e^{\frac{L^{2}}{4n}z^{2}}$ : (3.1)

Notice that  $\lim_{L \to 1} c_n$ ;  $(x) = e^{\frac{x^2}{4n}}$ , but the  $\lim_{L \to 1} i = n$ . If the sum over n is restricted to n 6  $cL^2$ , w ith c any n ite constant, w e can use the dominated convergence theorem and w e get

$$\lim_{L \mid 1} C_{n}; (x)% (n) = \begin{cases} X & \frac{x^{2}}{4n} % (n): \end{cases}$$
 (3.2)

(The  $\lim$  it is taken along the subsequence of increasing volumes for which % (n) is known to converge for any n.)

We now consider the term s with  ${\rm cL}^2$  < n 6 N . We estimate the sum s in (3.1) using integrals; we have

$$Z_{1}$$
 $e^{a(sb)^{2}}ds$  16  $e^{a(kb)^{2}}6$   $e^{a(sb)^{2}}ds+1$ : (3.3)

The Gaussian integral is equal to p = a. Consequently,

$$\frac{p}{4 n} L d p \frac{p}{4 n} + L d p \frac{p}{4 n} + L d p \frac{p}{4 n} L$$
(3.4)

These bounds hold provided  $\frac{p}{4n} > L$ . Since  $\frac{n}{L^2} > c$ , we have

W e obtain

$$X^{N}$$
  $c_{n}$ ;  $(x)$ %  $(n)$  6  $\frac{p}{4 c} + 1$  d  $X^{L^{2}}$  %  $(n)$  :  $(3.6)$ 

Using (32) with x=0 and the de nition (2.11) of the density of in nite cycles, we see that the last term converges to % (1 ) as L ! 1 . It then follows from (32) and (3.6) that

$$\lim_{L \mid 1} \sup_{n > 1} (x) 6 \stackrel{X}{=} e^{\frac{x^2}{4n}} % (n) + \frac{p}{\frac{4 c}{4 c}} + \frac{1}{1} % (1); \tag{3.7}$$

This inequality holds for any c, and the fraction is arbitrarily close to 1 by taking c large. A lower bound can be derived in a similar fashion, and we obtain the formula stated in Theorem 1.

### 4. The interacting gas

The interacting gas is much more dicult to study. We prove in this section the absence of in nite cycles when the chemical potential is negative (Theorem 2). We then study the coecients  $c_n$ ; (x) at low density and high temperature, using cluster expansion techniques. Their thermodynamic limit can be established, and we show that  $c_n$ ; (x)! O as  $\dot{x}$ ; 1 (Theorem 3).

Theorem 2. Let 0 < 1 and < 0; then

and

$$\lim_{j \neq j!} \lim_{1 \to 1!} \sup_{1 \to 1!} (x) = 0$$
:

Proof. Since U(!) > 0 and Z(;!) 6 Z(), the nite volume density % (n), Eq. (2.9), is less than

% (n) 6 e 
$$^{n}$$
  $^{2}$  dW  $^{n}_{00}$  (!) =  $\frac{e^{-n}}{(4 \text{ n})^{d=2}} \frac{X}{z^{2} z^{d}} e^{-\frac{L^{2}z^{2}}{4n}}$ : (4.1)

The right side is smaller than  $e^{-n}$  for all L large enough. We can therefore apply the dominated convergence theorem and we obtain

It follows that % (1 ) = 0. The statement about absence of o -diagonal long-range order can be treated similarly. We have the upper bound

(x) 6 
$$X = \frac{e^{-n}}{(4 - n)^{d=2}} X = \frac{(x - Lz)^2}{4n}$$
: (4.3)

By dominated convergence,

$$\lim_{x \to 0} \sup_{x \to 0} (x) 6 \sum_{n > 1}^{x} \frac{e^{-n}}{(4 \ n)^{d=2}} e^{x^{2} = 4n} : \tag{4.4}$$

We can again use the dominated convergence theorem for the  $\lim$  it ixj! 1, and we get the claim.

We continue the study of the interacting gas in the regime where cluster expansion converges. We assume that the chemical potential is negative, that the interaction potential U(x) is integrable, and that the temperature is high enough. The condition in Theorem 3 is stronger than necessary, but it is very explicit. We will invoke a weaker condition in the proof of the theorem that is based on the \K otecky-P reiss criterion" for the convergence of cluster expansion. Notice that G in ibre's survey [6] uses K irkwood-Salzburg equations; it applies to a broader range of potentials, but things are terribly intricate.

Theorem 3. Assume that , , and U satisfy  $\ \ \,$ 

$$\frac{1}{(4)^{d=2}} \sum_{R^d} X \times n^{d=2} 6$$

The therm odynam ic lim its of  $c_n$ ; (x) and % (n) exist, and we have

$$\lim_{\mathbf{j} \in \mathbb{N}^1, \ \mathbf{j} \in \mathbb{N}} c_n; \ (\mathbf{x}) = 0$$

for any n.

Proof. We need some notation in order to cast the grand-canonical partition function in a form suitable for the cluster expansion. Let us introduce a measure for trajectories that wind arbitrarily many times around the time direction. Namely, let  $X_n$  denote the measure space of continuous trajectories!: [0;n ]! D, and let  $X = [n > 1X_n]$  be the set of trajectories in D with arbitrary winding numbers. We introduce the measure on X whose integral means the following:

Z
$$F (!)d (!) = X \frac{e^{n}}{n} Z Z$$

$$dx dW_{xx}^{n} (!) e^{U(!)} F (!): (4.5)$$

It is clear that is a genuine measure on a reasonable measure space. But we describe the measure with more details for readers who are interested in analytic technicalities. The -algebra on  $X_n$  is the smallest -algebra that contains the sets  $f!\ 2\ X_n:!\ (t)\ 2\ B\ g$ , for any  $0\ 6\ t\ 6\ n$ , and any Borel set B D. Trajectories of  $X_1$  can be dilated in the time direction so as to yield trajectories with arbitrary winding numbers. One can then consider the product space  $X_1$  N with the product -algebra (the -algebra on N being the power set). The measure of a set of the kind A fng, with A a measurable subset of  $X_1$ , is de ned as

$$(A fng) = \frac{e^{-n}}{n} dx dW_{xx}^{n} (!^{0}) e^{-U(!^{0})} : (4.6)$$

Here, we introduced

$$A^0 = !^0 2 X_n : !^0(t) = ! (nt) \text{ for som } e ! 2 A :$$
 (4.7)

There is a unique extension to a measure on  $X_1$  N. There is a natural correspondence between X and  $X_1$  N, and we consider to be a measure on X.

W ith this notation, the grand-canonical partition function (2.4) is given by

$$Z() = \begin{cases} X & \frac{1}{k!} & X \\ x & X \end{cases}$$

$$X = \begin{cases} X & Y & h & i \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

$$X = \begin{cases} X & X \\ 0 & X \end{cases}$$

The term  $\,k=0$  is equal to 1 by de nition. Then Z ( ) has exactly the form assumed e.g. in [14]. The K otecky-P reiss criterion for the convergence of the cluster expansion requires the existence of a function a:X!  $R_+$  such that the following inequality holds for any ! 2 X:

<sup>Z</sup> h i 
$$e^{U(!;!^{0})} e^{a(!^{0})} d(!^{0}) 6 a(!)$$
: (4.9)

Choosing a (!) = n (with n the winding number of the trajectory !), it was shown in [14] that (4.9) is a consequence of the condition in Theorem 3.

The main result of the cluster expansion is that the partition function (4.8) is given by the exponential of a convergent series. Namely,

The combinatorial function '( $!_1; :::; !_k$ ) is equal to 1 if k = 1, and is otherwise equal to

$$(!_1; :::; !_k) = \frac{1}{k!} X Y h i$$

$$e^{U(!_i; !_j)} 1:$$
(4.11)

The sum is over connected graphs with k vertices, and the product is over edges of G. A proof for the relation (4.10) that directly applies here can be found in [14].

Observe now that the partition function Z (;!) is given by an expression similar to (4.8), where each d (! $_{\rm j}$ ) is replaced by e  $^{\rm U\,(!\,;!\,_{\rm j})}$  d (! $_{\rm j}$ ). Since U (! $_{\rm j}$ !) is positive, the criterion (4.9) is satisted with this new measure. It follows that Z (;!) has an expansion similar to (4.10), and we obtain the following expression for the ratio of partition functions,

$$\frac{Z(;!)}{Z()} = \exp \begin{pmatrix} X & Z & h & Y^{k} & i \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ &$$

It is not hard to check that

Then Equation (5) in [14] gives the necessary estimate for the exponent in (4.12), namely

This bound is uniform in the size of the domain, which is important. It follows that, as L ! 1 , the ratio Z ( ;!)=Z ( ) converges pointwise in and ! . The therm odynamic limits of  $c_n$ ; (x) and % (n) then clearly exist. Further,  $c_n$ ; (x) is bounded by

Z 
$$_{1}$$
 Z  $_{2}$   $_{3}$   $_{4.15}$   $_{5}$   $_{1}$   $_{5}$   $_{5}$   $_{6}$   $_{6}$   $_{6}$   $_{6}$   $_{6}$   $_{7}$   $_{1}$   $_{1}$   $_{1}$   $_{2}$   $_{2}$   $_{3}$   $_{1}$   $_{2}$   $_{3}$   $_{4.15}$   $_{1}$   $_{2}$   $_{3}$   $_{2}$   $_{3}$   $_{4}$   $_{2}$   $_{3}$   $_{4}$   $_{5}$   $_{5}$   $_{1}$   $_{2}$   $_{2}$   $_{3}$   $_{2}$   $_{3}$   $_{4}$   $_{2}$   $_{3}$   $_{3}$   $_{4}$   $_{5}$   $_{5}$   $_{5}$   $_{5}$   $_{6}$   $_{7}$   $_{1}$   $_{1}$   $_{2}$   $_{2}$   $_{3}$   $_{2}$   $_{3}$   $_{2}$   $_{3}$   $_{3}$   $_{4}$   $_{2}$   $_{3}$   $_{4}$   $_{2}$   $_{3}$   $_{3}$   $_{4}$   $_{2}$   $_{3}$   $_{3}$   $_{4}$   $_{2}$   $_{3}$   $_{3}$   $_{4}$   $_{2}$   $_{3}$   $_{3}$   $_{4}$   $_{2}$   $_{3}$   $_{3}$   $_{4}$   $_{2}$   $_{3}$   $_{3}$   $_{4}$   $_{2}$   $_{3}$   $_{3}$   $_{4}$   $_{2}$   $_{3}$   $_{3}$   $_{4}$   $_{2}$   $_{3}$   $_{3}$   $_{3}$   $_{3}$   $_{4}$   $_{2}$   $_{3}$   $_{3}$   $_{4}$   $_{3}$   $_{4}$   $_{4}$   $_{4}$   $_{5}$   $_{5}$   $_{5}$   $_{5}$   $_{5}$   $_{5}$   $_{5}$   $_{5}$   $_{7}$ 

It is not hard to show that the bracket is bounded away from zero uniformly in L (but not uniformly in and n). From (A.10), we have

$$\lim_{\substack{\text{im}\\ \text{ix} \mid 1}} \lim_{\text{L} \mid 1} \quad \text{dW }_{0x}^{\text{n}} \text{ (!)} = 0:$$

This implies that  $c_n$ ; (x) vanishes in the lim it of in nite jx j.

### 5. Conclusion

We introduced the formula (1.1) that relates the o-diagonal correlation function and the densities of cycles of given length. This formula involves coe cients c n that have a natural de nition in terms of integrals of Wiener trajectories. We conjectured several properties for the coe cients | these properties can actually be proved in the ideal gas for all tem peratures, and in the interacting gas for high tem peratures. These results seem to indicate that the order param eters of Feynm an and Penrose-O nsager agree. However, heuristic considerations based on the present fram ework [15] suggest that, if the gas is in a crystalline phase, the coe cients satisfy  $c_n(x)$  6  $e^{ax}$  for some a > 0, and for all n (including n = 1). Besides, one expects that %(1) > 0 if the temperature is su ciently low. The order param eters are not equivalent in this case.

An open problem is to establish the equivalence of the order parameters in weakly interacting gases in presence of Bose-Einstein condensation. Another question is whether c<sub>1</sub> (x) converges, as x j! 1, to a number that is strictly between 0 and 1. The corresponding phase would display a Bose condensate whose density is less than the density of particles in in nite cycles.

A cknow ledgm ents: It is a pleasure to thank Valentin Zagrebnov and Joe Pule for encouragements, and for pointing out an error with thermodynamic potentials in the original vresion of the manuscript. This research was supported in part by the grant DMS-0601075 from the USN ational Science Foundation.

# Appendix A. Feynman-Kac representation of the Bose gas

In this appendix we recall some properties of the Wiener measure, and we review the derivation of the Feynm an-K ac representation of the partition functions and of the o diagonal long-range order param eter. A complete account can be found in the excellent notes of G in ibre [6]; Faris w rote a useful survey [4].

Let D be the d-dim ensional cubic box of size L and volume  $V = L^d$ . We work with periodic boundary conditions, meaning that D is the d-dimensionial torus  $T_{\tau}^d$ . The state space is the Hilbert space H  $_{\rm D,N}$  of square-sum m able complex functions on D  $^{\rm N}$  , that are symmetric with respect to their arguments. Let S denote the symmetric projector on  $L^2(\mathbb{D}^N)$ , i.e.

S 
$$(x_1; ...; x_N) = \frac{1}{N!} X (x_{(1)}; ...; x_{(N)});$$
 (A.1)

where  $x_1; :::; x_N$  2 D and the sum is over all permutations of N elements. The state space for N bosons in D is therefore  $H_{D,N} = SL^2(D^N)$ , the projection of  $L^2(D^N)$  onto sym m etric functions.

The H am iltonian of the system is the sum H = T + V of kinetic and interaction energies.  $_{j=1}^{N}$   $_{j}$ , where  $_{j}$  is the Laplacian for the j-th particle. The kinetic energy is T = Interactions are given by the m ultiplication operator  $V = \frac{1}{2}$ 

Recall that and denote the inverse tem perature and the chem ical potential, respectively. The canonical and grand-canonical partition functions are

$$Y (;V;N) = Tr_{H_D,N} e^{H} ;$$
 (A.2)

$$Y (;V;N) = Tr_{H_{D;N}} e^{H};$$
 (A 2)  
 $Z (;V;) = e^{N} Y (;V;N):$  (A 3)

Under the assumption that U(x) is a stable potential and that it decays faster than  $jx j^d$  as jx j! 1, one can establish the existence of the therm odynam ic potentials (see [11])

$$f(;) = \lim_{V \mid 1} \frac{1}{V} \log Y (N);$$
 (A.4)

$$p(;) = \lim_{V = 1} \frac{1}{V} \log Z();$$
 (A.5)

Further, f and p are related by a Legendre transform,

$$f(;) = \sup \frac{1}{p}(;)$$
: (A.6)

This equation is useful to nd f from p in the case of the ideal gas, where p can be computed explicitly.

The Feynman-Kac representation allows to express  $e^{H}$  in term sofW iener trajectories (Brownian motion). We brie y review the main properties of the Wiener measure. Let  $X_1$  be the set of continuous paths!: [0; ]! D. Consider a function  $F:X_1$ ! R of the kind

$$F(!) = f(t_1); ...; t_n;$$
 (A.7)

where f is a bounded m easurable function on D  $^n$ , and 0 < t<sub>1</sub> <  $_{\text{ff}} <$  ; we extend f on R  $^d$  by periodicity. The integral of F with respect to the W iener m easure W  $_{\text{xy}}$  is given by

$$f(x_1; ...; x_n) dx_1 ... dx_n;$$
 (A.8)

where  $g_t$  is the normalized Gaussian function with mean zero and variance  $2t_r$ 

$$g_t(x) = \frac{1}{(4 + t)^{d-2}} e^{-\frac{x^2}{4t}}$$
: (A.9)

The sum over z accounts for periodic boundary conditions. A special case of (A.8) is when the function F is the constant function F (!) 1; we get

$$dW_{xy}(!) = (4)^{d=2} X_{e^{\frac{(x-y+Lz)^2}{4}}}$$
: (A 10)

Only the term z=0 remains in the limit L! 1. It can be proved that such a measure exists and is unique [6]. The W iener measure W  $_{xy}^{n}$  is concentrated on Holder continuous trajectories (with any Holder constant less than  $\frac{1}{2}$ ) that start at x and end at y. Integration with respect to W  $_{xy}$  and W  $_{00}$  are related as follows. De ne!  $^{0}$ (t) = ! (t)  $t^{\underline{y} \times x}$ ; then

F (!) 
$$dW_{xy}$$
 (!) =  $e^{\frac{(y-x)^2}{4}}$  Z F (!)  $dW_{00}$  (!): (A.11)

The Feynm an-K ac formula states that  $e^{-H}$  is given by an integral operator [2, 4, 6]. We are actually dealing with bosonic particles, and it is more convenient to consider the operator  $e^{-H}$  S that also projects onto symmetric functions. We have

$$e^{H}$$
 S  $(x_1; ...; x_N) = K (x_1; ...; x_N; y_1; ...; y_N) (y_1; ...; y_N) dy_1 ... dy_N; (A 12)$ 

where the kernel K is given by

$$\begin{array}{l} K \ (x_1; \ldots; x_N \ ; y_1; \ldots; y_N \ ) = \frac{1}{N \ !} \ X \\ \\ & = \sum_{\substack{2 \ S_N \\ \\ \text{exp} \\ \\ \text{i} < \text{j}} \ ^0 \end{array} \begin{array}{l} Z \\ \\ \\ \text{exp} \\ \\ \\ \text{i} < \text{j} \ ^0 \end{array} (!_1) :::: dW_{x_N \ y_{(N)}} \ (!_N \ ) \\ \\ & \times X \\ \\ \text{exp} \\ \\ & \times X \\ \\ \text{exp} \\ \\ \text{i} < \text{j} \ ^0 \end{array} (!_N \ )$$

The canonical partition function is then given by

We now group the cycles into closed trajectories, that may wind several times around the time direction. The number of permutations of N elements with k cycles of lengths  $n_1; \ldots; n_k$  (with  $j_1 n_j = N$ ) is

$$\frac{N!}{k!}$$
:

Further, we have

ex, we have Z Z Z 
$$\mathbb{Z}$$
  $\mathbb{Z}$   $\mathbb{Z$ 

The trajectory !:[0;n]! D in the right side is the concatenation of  $!_1;:::;!_n$ . The partition function (A.14) can then be rewritten in the form (2.1).

Let us turn to Penrose and Onsager o -diagonal long-range order [9]. Given a single particle function '  $2\ L^2$  (D), we do not the operator N, that represents the number of particles in the state '. The action of this operator is given by

$$N \cdot (x_1; :::; x_N) = \begin{cases} x^N & Z \\ & \checkmark (x) \end{cases} (x_1; :::; |x_X|, |x_2|; :::; x_N) \cdot (x_1) dx : \qquad (A.16)$$

$$j = 1 \quad D \quad j = 1$$

It is clear that 0 6 N , 6 N , and that  $[N \cdot ;S] = 0$ . Let  $'_0(x) = \frac{1}{V}$  denote the single particle ground state in absence of interactions. It is also the Fourier function with mode k=0. The average occupation of the zero mode is given by

$$%^{(0)} = \lim_{V! \ 1} \frac{1}{Y(N)} Tr_{H_{D,N}} \frac{h_{N_{0}}}{V} e^{H} : \qquad (A.17)$$

We set N = V, and the lim it exists at least along a subsequence of increasing volumes. A criterion for Bose-E instein condensation is that  $%^{(0)}$  diers from zero. We can derive a Feynman-K ac expression for this order parameter. From (A.12), (A.13), and (A.16), we have

Here, we set  $x_1 = x$ ,  $x_1 = y$ , and  $x_j = x_j$  for 2 6 j 6 N . Then  $^{(0)}$  can be written as

$$%^{(0)} = \lim_{V = 1} \frac{1}{V^2} \sum_{p=2}^{Z} (x - y) dx dy$$
 (A.19)

w here

$$(x \quad y) = \frac{1}{Y(;V;N)} \frac{1}{(N-1)!} \sum_{D^{N-1}}^{Z} dx_2 ::: dx_N$$

$$Z \qquad \qquad X \qquad \qquad Z \qquad \qquad X \qquad Z \qquad \qquad Z \qquad \qquad Z \qquad \qquad X \qquad Z \qquad \qquad Z \qquad$$

This expression involves an open cycle from x to y, winding n times around the time direction, with  $n=1;\ldots;N$ . Using the concatenation property (A 15), and thanks to the combinatorial factor  $\frac{(N-1)!}{(N-n)!}$ , we obtain the expression (2.14) for (x-y). The system displays o -diagonal long-range order if (x) is strictly positive, uniform ly in V;x.

Appendix B. A simple proof of macroscopic occupation in the ideal gas

In this section, we give a proof of the macroscopic occupation of the zero mode at low temperature. This is usually established in the grand-canonical ensemble, using a chemical potential that varies with the volume and tends to zero in the them odynamic limit. This approach is rather unnatural, and requires large deviation techniques to control the uctuations of the number of particles. The present proof is simpler and stays within the canonical ensemble.

The computation of the pressure and of the density in the grand-canonical ensemble can be found in any textbook dealing with quantum statistical mechanics. The chemical potential must be strictly negative. The in nite volume pressure is

potential m ust be strictly negative. The in nite volume pressure is 
$$p(;) = \frac{1}{(2)^d} \log 1 \quad e^{-(k^2)} dk; \tag{B.1}$$

and the density is

$$(;) = \frac{1}{(2)^{d}} \sum_{\mathbb{R}^{d}} \frac{dk}{e^{(k^{2})}} = \frac{1}{(4)^{d-2}} X e^{-n} n^{d-2} :$$
 (B 2)

The limit of (;) as % 0 is nite for d>3, and gives the critical density of the ideal Bose gas,  $_{c}$ . The graph of p(;) in three dimensions is plotted in Fig. 2 (a). Its Legendre transform (A.6) gives f(;), see Fig. 2 (b); it is nonanalytic at  $_{c}$ . The value of a () is given by

a() = 
$$\lim_{k \to 0} \frac{1}{k} p(;) = \lim_{k \to 0} f(;)$$
: (B.3)

The very nature of the Bose-E instein condensation is that the occupation number for k=0 becomes macroscopic. The average occupation of the zero mode  $\S^{(0)}$ , see Eq. (A.17), can be rewritten as

$$%^{(0)} = \lim_{V \mid 1} \frac{1}{Y(N)} \frac{X}{Y(N)} \frac{n_0}{v} e^{-\frac{P}{k} n_k k^2} :$$
 (B.4)

Here, N = V , and the sum is over all occupation numbers  $n_k > 0$ , with indices k 2  $(\frac{2}{L}\,Z\,)^d$ , such that k = N. The heart of Bose-Einstein condensation is the following result.




Figure 2. The pressure and the free energy of the ideal gas in three dimensions.

Theorem 4. For d > 3, the single-particle ground state is macroscopically occupied if > c. More precisely,

$$%^{(0)} = m ax(0; _{C}):$$

Proof. It is clear that  $%^{(0)} > 0$ . We now establish that  $%^{(0)} >$  c. Let us introduce the average occupation of the mode k,

$$hn_k i = \frac{1}{Y(N)} \sum_{(n_k 0) \in N}^{X} n_k e^{-P_{k0} n_k 0 k^{0^2}}$$
:

Thanks to the sum  $\text{ rule N } = \Pr_{k}^{P} m_{k} i$ , we have

$$%^{(0)} = \lim_{\substack{V : 1 \\ k \neq 0}} \frac{X}{v} \frac{\ln_k i}{V}$$
 (B.5)

We can view  $n_k$  as a random variable taking positive integer values; its expectation is therefore given by

$$\mathbf{m}_{k}\mathbf{i} = \sum_{i>1}^{X} P \operatorname{rob} (\mathbf{n}_{k} > \mathbf{i});$$
 (B.6)

where we de ned

Prob 
$$(n_k > i) = \frac{1}{Y(N)} X_{(n_k 0) N, i} e^{P_{k^0 n_k 0} k^{0^2}} :$$
 (B.7)

The sum is restricted to  $(n_{k^0})$  such that  $n_{k^0} = N$  and  $n_k > i$ . The change of variable  $n_k \,! \, n_k$  i leads to

Prob 
$$(n_k > i) = e^{-ik^{-2}} \frac{Y(N-i)}{Y(N)}$$
: (B.8)

The ratio of partition functions is also equal to the probability  $P \operatorname{rob}(n_0 > i)$ , which is smaller than 1. Equations (B.6) and (B.8) give a bound for the occupation numbers of all modes  $k \in 0$ , namely,

$$hn_k i 6 \frac{1}{e^{k^2} 1}$$
: (B.9)

Notice that  $hn_k$  i 6  $\frac{1}{k^2}$  6  $\frac{L^2}{4^{-2}}$  for k 6 0. This shows that only the zero mode can be macroscopically occupied (for d > 3). Inserting this bound into (B.5), we obtain

$$%^{(0)} > \frac{1}{(2)^d} \lim_{v \in I} X \frac{2}{L} \frac{d}{e^{k^2}} \frac{1}{1}$$
 (B.10)

The  $\lim$  it converges to the expression (B 2) with = 0, which is equal to c.

There remains to show that  $%^{(0)}$  6 max (0; c). From (B.8) with k = 0, and using the equivalence of ensembles, we have for any xed a,

$$\lim_{V = 1} \frac{1}{V} \log P \text{ rob } (n_0 > V \text{ a}) = f(;) \quad f(;) \quad a):$$
 (B.11)

The right side of (B.11) is strictly negative when a > m ax (0; c). There exists > 0 such that for large enough volumes,

Prob 
$$(n_0 > Va) 6 e^V$$
 : (B.12)

Let us assume that  $_{c} > 0$ ; the case  $_{c}$  6 0 can be treated similarly. Using (B.6) with k = 0, together with (B.12), we get

$$\frac{hn_0i}{V} = \frac{1}{V} \sum_{1 \text{ 6 i 6 aV}}^{X} P \text{ rob } (n_0 > i) + \frac{1}{V} \sum_{aV < i \text{ 6 N}}^{X} P \text{ rob } (n_0 > i)$$

$$6 a + e^{V} :$$
(B 13)

It follows that  $%^{(0)}$  is less than any number a >  $_{\text{c.}}$  hence  $%^{(0)}$  6  $_{\text{c.}}$ 

### References

- [1] G. Benfatto, M. Cassandro, I. Merola, E. Presutti, Limit theorems for statistics of combinatorial partition functions with applications to mean eld Bose gas, J. Math. Phys. 46, 033303 1{38 (2005)
- [2] O.Bratteli, D.W. Robinson, Operator Algebras and Quantum Statistical Mechanics II, Springer (1981)
- [3] T.C.D orlas, Ph.A.M artin, J.V.Pule, Long cycles in a perturbed mean eld model of a boson gas, J.Stat.Phys. 121, 433(461 (2005)
- [4] W .G .Faris, The Feynman-Kac formula, Notes for the Arizona Math.Phys.sem inar (2004)
- [5] R.P.Feynman, Atom ic theory of the transition in Helium, Phys. Rev. 91, 1291 (1301 (1953)
- [6] J.G in ibre, Some applications of functional integration in statistical mechanics, in \M ecanique statistique et theorie quantique des champs", Les Houches 1970, C.DeW itt and R. Stora eds, 327{427 (1971)
- [7] J.L.Lebow itz, M. Lenci, H. Spohn, Large deviations for ideal quantum systems, J.M ath.Phys.41, 1224(1243 (2000)
- [8] E.H. Lieb, R. Seiringer, J.P. Solovej, J.Y. ngvason, The mathematics of the Bose gas and its condensation, Oberwohlfach Seminars, Birkhauser (2005); cond-mat/0610117
- [9] O. Penrose, L. Onsager, Bose-Einstein condensation and liquid Helium, Phys. Rev. 104, 576(584 (1956)
- [10] J. V. Pule, V. A. Zagrebnov, The canonical perfect Bose gas in Casim ir boxes, J. Math. Phys. 45, 3565 (3583 (2004)
- [11] D. Ruelle, Statistical Mechanics: Rigorous Results, W. A. Benjamin (1969)
- [12] A. Suto, Percolation transition in the Bose gas, J. Phys. A 26, 4689 (4710 (1993)
- [13] A. Suto, Percolation transition in the Bose gas II, J. Phys. A 35, 6995 (7002 (2002)
- [14] D. Ueltschi, Cluster expansions and correlation functions, Moscow Math. J. 4, 511{522 (2004); math-ph/0304003
- [15] D. Ueltschi, The relation between Feynman cycles and o -diagonal long-range order, Phys. Rev. Lett., in press; cond-m at/0604005
- [16] V.A. Zagrebnov, J.B. Bru, The Bogoliubov model of weakly imperfect Bose gas, Phys. Reports 350, 291(434 (2001)
- [17] R.M.Zi, G.E.Uhlenbeck, M.K. ac, The idealBose-Einstein gas, revisited, Phys. Reports 32, 169 (248 (1977)

Daniel Ueltschi Department of Mathematics University of Arizona Tucson, AZ 85721, USA

E-m ailaddress: ueltschi@email.arizona.edu