

ANOTHER PROOF OF GELL-MANN AND LOW'S THEOREM

Luca Guido Molinari

Dipartimento di Fisica, Università degli Studi di Milano,
and INFN, Sezione di Milano, Via Celoria 16, 20133 Milano, Italy
luca.molinari@mi.infn.it

Abstract

The theorem by Gell-Mann and Low is a cornerstone in QFT and zero-temperature many-body theory. The standard proof is based on Dyson's time-ordered expansion of the propagator; a proof based on exact identities for the time-propagator is here given.

PACS: 03.65.-w, 11.10.-z, 71.10.-w, 24.10.Cn.

KEYWORDS: Gell-Mann Low Theorem, Interaction picture, Adiabatic theorem.

In the appendix of their paper "Bound States in Quantum Field Theory", Murray Gell-Mann and Francis Low proved a fundamental theorem[1] that bridges the ground states $|\Psi_0\rangle$ and $|\Psi\rangle$ of Hamiltonians H_0 and $H = H_0 + gV$ by means of time-propagators, and makes the transition of time-ordered correlators from the Heisenberg to the Interaction picture possible:

$$\langle \Psi | T \psi(1) \dots \psi^\dagger(n) | \Psi \rangle = \frac{\langle \Psi_0 | T S \psi(1) \dots \psi^\dagger(n) | \Psi_0 \rangle}{\langle \Psi_0 | S | \Psi_0 \rangle} \quad (1)$$

The single operator $S = U_I(\infty, -\infty)$ contains all the effects of the interaction. The theorem borrows ideas from the scattering and the adiabatic theories and introduces the concept of *adiabatic switching* of the interaction, through the time-dependent operator

$$H_\epsilon(t) = H_0 + e^{-\epsilon|t|} gV. \quad (2)$$

that interpolates between the operators of interest, H at $t = 0$ and H_0 at $|t| \rightarrow \infty$. The adiabatic limit is obtained for $\epsilon \rightarrow 0^+$. With the operator H_0 singled out, the theorem requires the time-propagator in the interaction picture,

$$U_{\epsilon I}(t, s) = e^{\frac{i}{\hbar} t H_0} U_\epsilon(t, s) e^{-\frac{i}{\hbar} s H_0} \quad (3)$$

where $U_\epsilon(t, s)$ is the full propagator[2]. The statement of Gell-Mann and Low's theorem is:

Theorem. Let $|\Psi_0\rangle$ be an eigenstate of H_0 with eigenvalue E_0 , and consider the vectors

$$|\Psi_\epsilon^{(\pm)}\rangle = \frac{U_{\epsilon I}(0, \pm\infty)|\Psi_0\rangle}{\langle\Psi_0|U_{\epsilon I}(0, \pm\infty)|\Psi_0\rangle} \quad (4)$$

If the limit vectors $|\Psi^{(\pm)}\rangle$ for $\epsilon \rightarrow 0^+$ exist, then they are eigenstates of H with same eigenvalue E .

Despite the validity of the theorem beyond perturbation theory, in the original paper[1] and in textbooks[3, 4, 5] the proof makes use of Dyson's expansion of the interaction propagator, and is rather cumbersome. Mathematical proofs are based on the adiabatic theorem[6]. In this letter a simple alternative proof is given, based on exact identities for the propagator that are stated in the following Lemma.

Lemma. If $U_\epsilon(t, s)$ is the time-propagator for $H_\epsilon(t)$ then, for all positive ϵ , the following relations hold:

$$i\hbar\epsilon g \frac{\partial}{\partial g} U_\epsilon(t, s) = H_\epsilon(t)U_\epsilon(t, s) - U_\epsilon(t, s)H_\epsilon(s), \quad \text{if } 0 \geq t \geq s, \quad (5)$$

$$-H_\epsilon(t)U_\epsilon(t, s) + U_\epsilon(t, s)H_\epsilon(s), \quad \text{if } t \geq s \geq 0. \quad (6)$$

Proof: The trick is to make the g -dependence of the propagator explicit into the time-dependence of some related propagator. Schrödinger's equation

$$i\hbar\partial_t U_\epsilon(t, s) = H_\epsilon(t)U_\epsilon(t, s), \quad U_\epsilon(s, s) = 1 \quad (7)$$

corresponds to the integral one, where we put $g = e^{t\theta}$:

$$U_\epsilon(t, s) = I + \frac{1}{i\hbar} \int_s^t dt' (H_0 + e^{\epsilon(\theta-|t'|)} V) U_\epsilon(t', s) \quad (8)$$

Consider the g -independent operators $H^{(\pm)}(t) = H_0 + e^{\pm\epsilon t} V$, with corresponding propagators $U^{(\pm)}(t, s)$. For $0 \geq t \geq s$, a time-translation in eq.(8) gives

$$U_\epsilon(t, s) = I + \frac{1}{i\hbar} \int_{s+\theta}^{t+\theta} dt' H^{(+)}(t') U_\epsilon(t' - \theta, s) \quad (9)$$

Comparison with the equation for $U^{(+)}(t + \theta, s + \theta)$:

$$U^{(+)}(t + \theta, s + \theta) = I + \frac{1}{i\hbar} \int_{s+\theta}^{t+\theta} dt' H^{(+)}(t') U^{(+)}(t', s + \theta)$$

and unicity of the solution imply the identification

$$U_\epsilon(t, s) = U^{(+)}(t + \theta, s + \theta) \quad (10)$$

Since θ enters in the operator $U^{(+)}(t + \theta, s + \theta)$ only in its temporal variables, we obtain

$$\partial_\theta U_\epsilon(t, s) = \partial_t U_\epsilon(t, s) + \partial_s U_\epsilon(t, s) \quad (11)$$

By using eq.(7) and its adjoint, the first identity is proven.

If $t \geq s \geq 0$, the same procedure gives $U_\epsilon(t, s) = U^{(-)}(t - \theta, s - \theta)$ and therefore $\partial_\theta U_\epsilon(t, s) = -\partial_t U_\epsilon(t, s) - \partial_s U_\epsilon(t, s)$, which leads to the identity (6). An identity for $t \geq 0 \geq s$ can be obtained by writing $U_\epsilon(t, s) = U_\epsilon(t, 0)U_\epsilon(0, s)$.

In the interaction picture, eq.(3), the identities transform straightforwardly into the following ones:

$$\begin{aligned} i\hbar\epsilon g \frac{\partial}{\partial g} U_{\epsilon I}(t, s) = & \quad (12) \\ H_{\epsilon I}(t)U_{\epsilon I}(t, s) - U_{\epsilon I}(t, s)H_{\epsilon I}(s), & \quad \text{if } 0 \geq t \geq s, \\ -H_{\epsilon I}(t)U_{\epsilon I}(t, s) + U_{\epsilon I}(t, s)H_{\epsilon I}(s), & \quad \text{if } t \geq s \geq 0, \end{aligned}$$

where $H_{\epsilon I}(t) = e^{\frac{i}{\hbar}tH_0} H_\epsilon(t) e^{-\frac{i}{\hbar}tH_0}$. By applying eqs.(12) to an eigenstate $|\Psi_0\rangle$ of H_0 , we obtain

$$(H - E_0 \pm i\hbar\epsilon g \frac{\partial}{\partial g}) U_{\epsilon I}(0, \pm\infty) |\Psi_0\rangle = 0 \quad (13)$$

This same equation is proven in the literature by direct use of Dyson's expansion. From now on, the proof of the theorem proceeds in the standard path, and is here sketched for completeness.

Existence of a non-trivial adiabatic limit of eq.(13) requires the vector $U_{\epsilon I}(0, \pm\infty) |\Psi_0\rangle$ to develop a phase proportional to $1/\epsilon$; this has been checked in diagrammatic expansion[7, 8]. The singular phase is precisely removed by the denominator in the definition of the vectors $|\Psi_\epsilon^{(\pm)}\rangle$, before the limit is taken.

1) For finite ϵ , the two identities eq.(13) are projected on the vector $|\Psi_0\rangle$, and yield a formula for the *energy shift*

$$\mp i\hbar\epsilon g \frac{\partial}{\partial g} \log \langle \Psi_0 | U_{\epsilon I}(0, \pm\infty) | \Psi_0 \rangle = E_\epsilon^{(\pm)} - E_0 \quad (14)$$

where $E_\epsilon^{(\pm)} = \langle \Psi_0 | H | \Psi_\epsilon^{(\pm)} \rangle$.

2) By eliminating E_0 in (13) with the aid of (14), with simple steps one obtains

$$(H - E_\epsilon^{(\pm)} \pm i\hbar\epsilon g \frac{\partial}{\partial g})|\Psi_\epsilon^{(\pm)}\rangle = 0 \quad (15)$$

Then, for $\epsilon \rightarrow 0^+$, the limit vectors $|\Psi^{(\pm)}\rangle$ obtained by pulling onward or backward in time the same asymptotic eigenstate $|\Psi_0\rangle$, are eigenvectors of $H = H_0 + gV$ with eigenvalues $E^{(\pm)}$.

3) The time-reversal operator has the action $T^\dagger U_\epsilon(t, s)T = U_\epsilon(-t, -s)$. If H_0 commutes with T the relation extends to the interaction propagator and $T^\dagger U_{\epsilon I}(0, \infty)T = U_{\epsilon I}(0, -\infty)$. If $|\Psi_0\rangle$ is also an eigenstate of T , it follows that $T^\dagger|\Psi_\epsilon^{(+)}\rangle$ is parallel to $|\Psi_\epsilon^{(-)}\rangle$ and $E^{(+)} = E^{(-)}$. The proportionality factor equals one, since $\langle E_0|\Psi^{(+)}\rangle = \langle E_0|\Psi^{(-)}\rangle$.

The formula for the energy shift, eq.(14), can be recast in a form involving the S -operator (Sucher's formula)[9]. From eqs.(12) the following relation follows:

$$-i\hbar\epsilon g \partial_g S = H_0 S + S H_0 - 2U_{\epsilon I}(\infty, 0)H U_{\epsilon I}(0, -\infty)$$

The expectation value on the eigenstate $|\Psi_0\rangle$ and use of the theorem give Sucher's formula:

$$E - E_0 = \lim_{\epsilon \rightarrow 0} \frac{i\hbar\epsilon}{2} g \frac{d}{dg} \log \langle \Psi_0 | U_{\epsilon I}(\infty, -\infty) | \Psi_0 \rangle \quad (16)$$

References

- [1] M.Gell-Mann e F.Low, Phys. Rev. **84** (1951) 350.
- [2] M.Reed and B.Simon, *Methods of Mathematical Physics*, Vol.II, Academic Press 1975.
- [3] P.Nozieres, *Physics of interacting Fermi systems*, (Benjamin, New York, 1964).
- [4] A.L.Fetter and J.D.Walecka, *Quantum Theory of Many Particle Systems*, MacGraw Hill, New York, 1971.
- [5] E.K.U.Gross, E.Runge and O.Heinonen, *Many-Particle theory*, Adam Hilger, Bristol, 1991.
- [6] G.Nenciu and G.Rasche, Helvetica Physica Acta **62** (1989) 372.

- [7] J.Hubbard, Proc. Roy. Soc. (London) A **240** (1957) 539.
- [8] K. Takayanagi, Phys. Rev. A **44** (1991) 59
- [9] J.Sucher, Phys. Rev. **107** (1957) 1448.