

Homogeneous Multivariate Polynomials with the Half-Plane Property

Young-Bin Choe

Department of Combinatorics and Optimization

University of Waterloo

Waterloo, Ontario N2L 3G1, CANADA

ybchoe@math.uwaterloo.ca

James G. Oxley

Department of Mathematics

Louisiana State University

Baton Rouge, LA 70803-4918, USA

oxley@math.lsu.edu

Alan D. Sokal

Department of Physics

New York University

New York, NY 10003, USA

sokal@nyu.edu

David G. Wagner

Department of Combinatorics and Optimization

University of Waterloo

Waterloo, Ontario N2L 3G1, CANADA

dgwagner@math.uwaterloo.ca

February 4, 2002

revised November 23, 2002

Abstract

A polynomial P in n complex variables is said to have the “half-plane property” (or Hurwitz property) if it is nonvanishing whenever all the variables lie in the open right half-plane. Such polynomials arise in combinatorics, reliability theory, electrical circuit theory and statistical mechanics. A particularly important case is when the polynomial is homogeneous and multiaffine: then it is the (weighted) generating polynomial of an r -uniform set system. We prove that the support (set of nonzero coefficients) of a homogeneous multiaffine polynomial with the half-plane property is necessarily the set of bases of a matroid. Conversely, we ask: For which matroids M does the basis generating polynomial $P_{\mathcal{B}(M)}$ have the half-plane property? Not all matroids have the half-plane property, but we find large classes that do: all sixth-root-of-unity matroids, and a subclass of transversal (or cotransversal) matroids that we call “nice”. Furthermore, the class of matroids with the half-plane property is closed under minors, duality, direct sums, 2-sums, series and parallel connection, full-rank matroid union, and some special cases of principal truncation, principal extension, principal cotruncation and principal coextension. Our positive results depend on two distinct (and apparently unrelated) methods for constructing polynomials with the half-plane property: a determinant construction (exploiting “energy” arguments), and a permanent construction (exploiting the Heilmann–Lieb theorem on matching polynomials). We conclude with a list of open questions.

KEY WORDS: Graph, matroid, jump system, abstract simplicial complex, spanning tree, basis, generating polynomial, reliability polynomial, Brown–Colbourn conjecture, half-plane property, Hurwitz polynomial, positive rational function, Lee–Yang theorem, Heilmann–Lieb theorem, matching polynomial, Grace–Walsh–Szegő coincidence theorem, matrix-tree theorem, electrical network, nonnegative matrix, determinant, permanent.

2000 MATHEMATICS SUBJECT CLASSIFICATION: Primary 05B35; Secondary 05C99, 05E99, 15A15, 15A48, 30C15, 32A99, 82B20, 94C05.

RUNNING HEAD: Polynomials with the Half-Plane Property

Contents

1	Introduction	4
2	Polynomials and the half-plane property	10
2.1	Basic definitions	10
2.2	Shifted half-plane property and leading part	11
2.3	Real-part-positive rational functions	12
2.4	Derivatives	14
2.5	The Grace–Walsh–Szegő coincidence theorem	16
2.6	Algorithms	17
3	The local half-plane property	18
4	Constructions	21
4.1	Deletion and contraction	21
4.2	Duality	22
4.3	Direct sum	23
4.4	Parallel connection, series connection, and 2-sum	24
4.5	Principal truncation and principal extension	27
4.6	Principal cotruncation and principal coextension	29
4.7	Multiaffine part and full-rank matroid union	31
4.8	“Folding mod 2” and convolution	33
5	Necessary and sufficient condition	36
5.1	General case	36
5.2	Rank-2 case	37
6	Necessary conditions I: Same-phase property	41

7	Necessary conditions II: Matroidal support	45
7.1	The exchangeability theorem	45
7.2	The weak half-plane property	49
8	The determinant condition and $\sqrt[6]{1}$-matroids	50
8.1	The determinant condition	50
8.2	(F, G) -representability and $\sqrt[6]{1}$ -matroids	51
9	Uniform matroids	58
9.1	Half-plane property	58
9.2	Brown–Colbourn property	60
10	The permanent condition: Transversal and cotransversal matroids	61
10.1	Heilmann–Lieb theorem	61
10.2	Application to transversal matroids	63
10.3	Examples	65
10.4	Rank-3 transversal matroids	74
10.5	A wild speculation	75
11	Counterexamples	76
12	Numerical experiments	80
13	Open questions	82
13.1	The strong half-plane property	82
13.2	Same-phase theorem	83
13.3	Support of polynomials with the half-plane property	83
13.4	(F, G) -representability of matroids	84
13.5	Determinant class, permanent class, and the half-plane property	85
13.6	Half-plane property for transversal matroids	87
13.7	Half-plane property for 7-element rank-3 matroids	88
13.8	Algorithms	88
A	Matroids considered in this paper	89
A.1	Rank-3 matroids on 6 elements	89
A.2	Rank-3 matroids on 7 elements	90
A.2.1	Matroids with a 4-point line	91
A.2.2	The Fano matroid F_7 and its relaxations	93
A.2.3	P_7 and its relaxations	94
A.3	Some rank-3 matroids on 8 or 9 elements	94
A.4	Some rank-4 matroids on 8 elements	97
A.5	Some transversal matroids	98
B	$(F, \{1\})$-representability of matroids	98

1 Introduction

It seems that the theory of polynomials, linear in each variable, that do not have zeros in a given multidisk or a more general set, has a long way to go, and has so far unnoticed connections to various other concepts in mathematics.

— Aimo Hinkkanen [37, p. 288]

Let us consider a connected graph $G = (V, E)$ as an electrical network, by assigning to each edge $e \in E$ a complex number x_e , called its *conductance* (or *admittance*).¹ The node voltages $\{\varphi_i\}_{i \in V}$ and current inflows $\{J_i\}_{i \in V}$ then satisfy the linear system $L(x)\varphi = J$, where $L(x)$ is the edge-weighted Laplacian matrix for G [that is, $L(x) = BXB^T$ where B is the directed vertex-edge incidence matrix for any orientation of G , and $X = \text{diag}(\{x_e\})$]. On physical grounds we expect that if $\text{Re } x_e > 0$ for all e (i.e. every branch is dissipative), then the network is uniquely solvable once we fix the voltage at a single reference node $i_0 \in V$, or in other words that the i_0 th principal cofactor of $L(x)$ is nonzero. Now, by the matrix-tree theorem [9, 17, 16, 80], each principal cofactor of $L(x)$ equals the spanning-tree sum

$$T_G(x) = \sum_{\text{trees } T \subseteq E} x^T, \quad (1.1)$$

where we have used the shorthand $x = \{x_e\}_{e \in E}$ and $x^T = \prod_{e \in T} x_e$. We therefore conjecture:

Theorem 1.1 *Let G be a connected graph. Then the spanning-tree polynomial T_G has the “half-plane property”, i.e. $\text{Re } x_e > 0$ for all e implies $T_G(x) \neq 0$.*

The proof of Theorem 1.1 is not difficult: Consider any nonzero complex vector $\varphi = \{\varphi_i\}_{i \in V}$ satisfying $\varphi_{i_0} = 0$. Because G is connected, we have $B^T\varphi \neq 0$. Therefore, the quantity

$$\varphi^* L(x) \varphi = \varphi^* BXB^T \varphi = \sum_{e \in E} |(B^T\varphi)_e|^2 x_e \quad (1.2)$$

has strictly positive real part whenever $\text{Re } x_e > 0$ for all e ; so in particular $(BXB^T\varphi)_i \neq 0$ for some $i \neq i_0$. It follows that the submatrix of $L(x)$ obtained by suppressing the i_0 th row and column is nonsingular, and so has a nonzero determinant. Theorem 1.1 then follows from the matrix-tree theorem.²

¹ In this paper all graphs are finite and undirected. Loops and multiple edges are allowed unless specified otherwise.

² This proof is well known in the circuit-theory literature: see e.g. [18, Section 2.7] as well as the related results in [23, pp. 398–401, 430–431 and 850–851] [58, pp. 52–53 and 67–69]. It has, moreover, a natural physical interpretation: if $\varphi = \{\varphi_i\}_{i \in V}$ are the node voltages, then the real part of the quadratic form (1.2) is the total power dissipated in the circuit. We thank Charles Desoer and Paul Penfield for pointing out references [23, 58].

An immediate corollary of Theorem 1.1 is that the complementary spanning-tree polynomial

$$\tilde{T}_G(x) = \sum_{\text{trees } T \subseteq E} x^{E \setminus T} \quad (1.3)$$

also has the half-plane property, since

$$\tilde{T}_G(x) = x^E T_G(1/x) \quad (1.4)$$

and the map $x_e \mapsto 1/x_e$ takes the right half-plane onto itself.

From a combinatorial point of view, the noteworthy fact is that the spanning trees of G constitute the bases of the graphic matroid $M(G)$, and their complements constitute the bases of the cographic matroid $M^*(G)$. So T_G and \tilde{T}_G are the (multivariate) basis generating polynomials for $M(G)$ and $M^*(G)$, respectively. This naturally suggests generalizing Theorem 1.1 to more general matroids and, perhaps, to more general set systems. Before posing these questions precisely, we need to fix some notation and terminology.

A *set system* (or *hypergraph*) \mathcal{S} on the (finite) ground set E is simply a collection \mathcal{S} of subsets of E . Given any set system \mathcal{S} on E , we define its (*multivariate*) *generating polynomial* to be

$$P_{\mathcal{S}}(x) = \sum_{S \in \mathcal{S}} x^S, \quad (1.5)$$

where $x = \{x_e\}_{e \in E}$ are commuting indeterminates (which we shall usually take to be complex variables). The *rank* of a set system is the maximum cardinality of its members (by convention we set $\text{rank} = -\infty$ if $\mathcal{S} = \emptyset$); equivalently, it is the degree of the generating polynomial $P_{\mathcal{S}}$. A set system \mathcal{S} is *r-uniform* if $|S| = r$ for all $S \in \mathcal{S}$, or equivalently if its generating polynomial $P_{\mathcal{S}}$ is homogeneous of degree r .

An *abstract simplicial complex* (or *complex* for short) is a set system \mathcal{S} satisfying

$$(I1) \quad \emptyset \in \mathcal{S}.$$

$$(I2) \quad \text{If } S \in \mathcal{S} \text{ and } S' \subseteq S, \text{ then } S' \in \mathcal{S} \text{ ("}\mathcal{S} \text{ is hereditary downwards").}$$

The members S of a complex are called *faces*, and the maximal members (with respect to set-theoretic inclusion) are called *facets*. A complex is called *pure* (of rank r) if all its facets have the same cardinality r .

A complex \mathcal{S} is called a *matroid complex* if it satisfies the further condition

$$(I3) \quad \text{If } S_1, S_2 \in \mathcal{S} \text{ and } |S_1| < |S_2|, \text{ then there exists an element } e \in S_2 \setminus S_1 \text{ such that } S_1 \cup \{e\} \in \mathcal{S} \text{ ("independence augmentation axiom for matroids").}$$

The faces of a matroid complex are called the *independent sets* of the matroid; the facets (i.e. the maximal independent sets) are called *bases*. It is easy to prove that every matroid complex is pure, i.e. all bases have the same cardinality r , called the *rank* of the matroid. Moreover, it is not difficult to show that a collection \mathcal{B} of subsets of E is the collection of bases of a matroid on E if and only if it satisfies the following two conditions:

(B1) \mathcal{B} is nonempty.

(B2) If $B_1, B_2 \in \mathcal{B}$ and $x \in B_1 \setminus B_2$, then there exists $y \in B_2 \setminus B_1$ such that $(B_1 \setminus x) \cup \{y\} \in \mathcal{B}$ (“basis exchange axiom for matroids”).

For more information on matroid theory, see [55].

We shall be particularly interested in the *basis generating polynomial* of a matroid M ,

$$P_{\mathcal{B}(M)}(x) = \sum_{B \in \mathcal{B}(M)} x^B. \quad (1.6)$$

We can now pose the following questions concerning possible extensions of Theorem 1.1:

Question 1.2 *For which matroids M does the basis generating polynomial $P_{\mathcal{B}(M)}$ have the half-plane property?*

More generally:

Question 1.3 *For which r -uniform set systems \mathcal{S} does the generating polynomial $P_{\mathcal{S}}$ have the half-plane property?*

Our original conjecture was that all matroids (and no non-matroidal set systems) have the half-plane property. That would be nice and neat, but it turns out to be false; and the truth is considerably more interesting and subtle. Our conjecture is half right: an r -uniform set system with the half-plane property is necessarily the set of bases of a matroid (Theorem 7.1). But not every matroid has the half-plane property, and we do not yet have a complete characterization of those that do. Nevertheless, we can find large classes of matroids with the half-plane property:

- (a) Every sixth-root-of-unity matroid [78] has the half-plane property (Theorem 8.1 and Corollary 8.2). This class properly includes the regular matroids, which in turn properly include the graphic and cographic matroids. The proof of Theorem 8.1 is, in fact, a direct generalization of the proof just given for Theorem 1.1.
- (b) Every uniform matroid has the half-plane property, and indeed has the (stronger) “Brown–Colbourn property” (Section 9).
- (c) A significant subclass of transversal matroids — those we call “nice” — have the half-plane property (Section 10). Indeed, it may well be true that *all* transversal matroids have the half-plane property, but we have no idea how to prove this.
- (d) All matroids of rank or corank at most 2 have the half-plane property (Corollary 5.5), as do all matroids on a ground set of at most 6 elements (Proposition 10.4).

- (e) The class of matroids with the half-plane property is closed under minors, duality, direct sums, 2-sums, series and parallel connection, full-rank matroid union, and some special cases of principal truncation, principal extension, principal cotruncation and principal coextension (Section 4).

Moreover, we can show that certain matroids do *not* have the half-plane property: among these are the Fano matroid F_7 , the non-Fano matroid F_7^- , their relaxations F_7^{--} , F_7^{-3} and $M(K_4) + e$, the matroids P_8 , P'_8 and P''_8 , the Pappus and non-Pappus matroids, the free extension $(\text{non-Pappus} \setminus 9) + e$, and all their duals (Section 11). The first six of these examples are minor-minimal, and we conjecture that the others are as well; but we strongly suspect that this list is incomplete, and indeed we consider it likely that the set of minor-minimal non-half-plane-property matroids is infinite.

More generally, we shall consider homogeneous multiaffine polynomials $P(x) = \sum_{S \subseteq E, |S|=r} a_S x^S$ with arbitrary complex coefficients a_S (not necessarily 0 or 1). We shall prove two *necessary* conditions for $P \not\equiv 0$ to have the half-plane property:

- (a) P must have the “same-phase property”, i.e. all the nonzero coefficients a_S must have the same phase (Theorem 6.1). So without loss of generality we can assume that all the a_S are nonnegative.
- (b) The *support* $\text{supp}(P) = \{S \subseteq E : a_S \neq 0\}$ must be the collection of bases of a matroid (Theorem 7.1).

This latter fact is particularly striking: it shows that matroids arise *naturally* from a consideration of homogeneous multiaffine polynomials with the half-plane property. We do not know whether the converse of Theorem 7.1 is true, i.e. whether for every matroid M there exists a homogeneous multiaffine polynomial P with the half-plane property such that $\text{supp}(P) = \mathcal{B}(M)$. But it is true, at least, for all matroids representable over \mathbb{C} (Corollary 8.2).

We shall also give two *sufficient* conditions for a homogeneous multiaffine polynomial P to have the half-plane property (or be identically zero):

- (a) *Determinant condition* (Theorem 8.1): $a_S = |\det(A \upharpoonright S)|^2$ for some $r \times n$ complex matrix A [here $n = |E|$, and $A \upharpoonright S$ denotes the square submatrix of A using the columns indexed by the set S]. This corresponds to $P(x) = \det(AXA^*)$ where $X = \text{diag}(\{x_e\})$ and $*$ denotes Hermitian conjugate.
- (b) *Permanent condition* (Theorem 10.2): $a_S = \text{per}(\Lambda \upharpoonright S)$ for some $r \times n$ nonnegative matrix Λ . This corresponds to $P(x) = \text{per}(\Lambda X)$.

Unfortunately, the relationship between these sufficient conditions and the half-plane property looks complicated. Neither family of polynomials contains the other; their intersection is nonempty; and their union is a proper subset of the set of all homogeneous multiaffine polynomials with the half-plane property.

These questions also have a close connection with reliability theory [19]. Consider a finite set E of communication channels, which fail independently with probabilities

$\{q_e\}_{e \in E}$. Let \mathcal{S} be a set system on E , whose members we shall interpret as the sets of failed channels that allow the system as a whole to be operational. Then the probability that the system is operational is given by the *multivariate reliability polynomial*

$$\text{Rel}_{\mathcal{S}}(q) = \sum_{A \in \mathcal{S}} q^A (1 - q)^{E \setminus A} \quad (1.7)$$

where $q = \{q_e\}_{e \in E}$. This is easily related to the multivariate generating polynomial

$$P_{\mathcal{S}}(x) = \sum_{A \in \mathcal{S}} x^A \quad (1.8)$$

by

$$\text{Rel}_{\mathcal{S}}(q) = (1 - q)^E P_{\mathcal{S}}\left(\frac{q}{1 - q}\right) \quad (1.9)$$

$$P_{\mathcal{S}}(x) = (1 + x)^E \text{Rel}_{\mathcal{S}}\left(\frac{x}{1 + x}\right) \quad (1.10)$$

In the reliability context it is natural to assume that \mathcal{S} is a complex, i.e. that \mathcal{S} contains \emptyset and is closed under taking subsets. Indeed, the simplest case arises when $G = (V, E)$ is a connected graph and we declare the system to be operational if the non-failed edges form a connected spanning subgraph (this is the “all-terminal reliability”). In this case \mathcal{S} is the set of complements of connected spanning subgraphs of G , i.e. the family $\mathcal{I}(M^*(G))$ of independent sets of the cographic matroid $M^*(G)$. Our Holy Grail is the following:

Conjecture 1.4 (multivariate Brown–Colbourn conjecture [10, 70]) *Let G be a connected graph. If $|q_e| > 1$ for all e , then $\text{Rel}_{\mathcal{I}(M^*(G))}(q) \neq 0$. Equivalently, if G is loopless and $\text{Re } x_e < -1/2$ for all e , then $P_{\mathcal{I}(M^*(G))}(x) \neq 0$.*

(Note that a loop in G corresponds to a coloop in $M^*(G)$. Each loop in G has no effect on the reliability polynomial but multiplies the independent-set polynomial of $M^*(G)$ by a factor $1 + x_e$, leading to a root at $x_e = -1$. This is why, in discussing the roots of $P_{\mathcal{I}(M^*(G))}$, we need to assume that G is loopless.)

We are at present quite far from a proof of Conjecture 1.4. One of us has proven Conjecture 1.4 for the special case of series-parallel graphs [70, Remark 3 in Section 4.1]; earlier, another one of us had proven the corresponding univariate result, i.e. when all the q_e take the same value [73]. But series-parallel graphs are a small subset of planar graphs, and an even smaller subset of all graphs!

Nonetheless one can dream, and even lacking a proof of Conjecture 1.4 it is reasonable to ask whether stronger results might be true. For example:

Question 1.5 *Which (coloopless) matroids M have the “Brown–Colbourn property”, i.e. $\text{Re } x_e < -1/2$ for all e implies $P_{\mathcal{I}(M)}(x) \neq 0$?*

Aside from the cographic (or equivalently, graphic) matroids of series-parallel graphs, we can prove the Brown–Colbourn property for uniform matroids $U_{r,n}$ with $0 \leq r < n$ (the condition $r < n$ corresponds precisely to forbidding coloops): this follows immediately from the corresponding univariate result [73, Proposition 7.3] combined with the Grace–Walsh–Szegő coincidence theorem (see Section 9). On the other hand, it is easy to show (Corollary 2.3) that the Brown–Colbourn property for $P_{\mathcal{I}(M)}$ implies the half-plane property for $P_{\mathcal{B}(M)}$. So the results of this paper imply that the Brown–Colbourn property fails for many matroids (e.g. F_7 , F_7^- , ...) and for all pure complexes that are not matroidal.

One purpose of this paper is, therefore, to serve as a “warm-up” for an attack on the Brown–Colbourn property, by studying first a property (the half-plane property) that is a *necessary* condition for the Brown–Colbourn property and may be easier to characterize. But the results of our investigation show that the half-plane property is very interesting in its own right!

Our study of polynomials with the half-plane property can also be viewed in the wider context of theorems asserting that some combinatorially interesting class of multivariate polynomials are nonvanishing in some large domain of complex n -space. Theorems of this kind include the Lee–Yang theorem on ferromagnetic Ising models and generalizations thereof [46, 68, 47, 37] and the Heilmann–Lieb theorem on matching polynomials [35] (discussed in Section 10.1 below). Useful tools for manipulating such polynomials include the Grace–Walsh–Szegő coincidence theorem [31, 71, 74, 51] (see Section 2.5), the Asano contraction lemma [3, 63, 37] and the Hinkkanen composition theorem [37] (see Section 4.8).

The plan of this paper is as follows: In Section 2 we discuss the basic properties of multivariate polynomials with the half-plane property, including their important connection with real-part-positive rational functions. In Section 3 we discuss a key criterion that we call the “local half-plane property”. In Section 4 we describe a large number of constructions that preserve the half-plane property; most of these are motivated by standard operations on matroids. In Section 5 we provide a necessary and sufficient condition for a polynomial to have the half-plane property; this condition will play an important role in finding counterexamples. We also provide a simple explicit criterion in the rank-2 case. In Section 6 we prove that, in a homogeneous polynomial with the half-plane property, all the nonzero coefficients must have the same phase. In Section 7 we prove that the support of a homogeneous multiaffine polynomial must be the collection of bases of a matroid; we also prove a generalization to the non-multiaffine case. In Section 8 we give the “determinant construction”, and deduce as a corollary that every sixth-root-of-unity matroid has the half-plane property. This section also contains some results on (F, G) -representability of matroids that may be of independent interest. In Section 9 we prove that every uniform matroid has the half-plane property, and indeed has the Brown–Colbourn property. In Section 10 we give the “permanent construction”, and deduce as a corollary that a certain subclass of transversal matroids (those we call “nice”) have the half-plane property; we also give numerous examples of nice and non-nice transversal matroids.

In Section 11 we give several examples of matroids that do *not* have the half-plane property. In Section 12 we report our numerical experiments on matroids for which we have been unable to prove or disprove the half-plane property. We conclude, in Section 13, with a list of open questions. In Appendix A we provide, for the convenience of the reader, a list of the matroids considered in this paper, along with brief summaries of their properties. In Appendix B we analyze the $(F, \{1\})$ -representability of matroids.

Note Added (November 2002): Gordon Royle and one of the authors (A.D.S.) have recently discovered — to our great surprise — that the Brown–Colbourn conjecture is false! The multivariate Brown–Colbourn conjecture is false already for the simplest non-series-parallel graph, namely the complete graph K_4 . The univariate Brown–Colbourn conjecture is false for certain simple planar graphs that can be obtained from K_4 by parallel and series extension of edges. This work will be reported separately [61].

2 Polynomials and the half-plane property

Our main interest is in polynomials that are homogeneous of degree r and are multiaffine (i.e. of degree ≤ 1 in each variable separately): these include the generating polynomials of r -uniform set systems and, more specifically, the basis generating polynomials of rank- r matroids. However, many of our results will be valid for homogeneous polynomials that are not necessarily multiaffine, or for multiaffine polynomials that are not necessarily homogeneous, or (sometimes) for general polynomials. We shall endeavor to state our results in whatever degree of generality seems most natural, without being pedantic.

2.1 Basic definitions

Let $P(x) = \sum_{\mathbf{m}} a_{\mathbf{m}} x^{\mathbf{m}}$ be a polynomial with complex coefficients in the variables $x = \{x_e\}_{e \in E}$. We call the finite set $E = E(P)$ the *ground set* of P , and we call its members *elements*. We shall generally use the letter n to denote the cardinality of E ; often we shall simply take E to be $[n] = \{1, \dots, n\}$. In P , the sum ranges over all *multi-indices* \mathbf{m} (that is, functions $\mathbf{m}: E \rightarrow \mathbb{N}$), and only finitely many of the coefficients $a_{\mathbf{m}}$ are nonzero. The *degree* of the monomial $x^{\mathbf{m}} = \prod_{e \in E} x_e^{m(e)}$ is $|\mathbf{m}| = \sum_{e \in E} m(e)$, and the *degree* of P is $\deg P = \max\{|\mathbf{m}|: a_{\mathbf{m}} \neq 0\}$. We say that P is *homogeneous of degree* r if $a_{\mathbf{m}} = 0$ whenever $|\mathbf{m}| \neq r$. For $e \in E$, the *degree of* P in x_e is $\deg_e P = \max\{m(e): a_{\mathbf{m}} \neq 0\}$. We say that P is *affine in* e if $\deg_e P \leq 1$, and we say that P is *multiaffine* if it is affine in e for all $e \in E$. A multiaffine polynomial can be written in the form $P(x) = \sum_{S \subseteq E} a_S x^S$.

The *support* of a polynomial $P(x) = \sum_{\mathbf{m}} a_{\mathbf{m}} x^{\mathbf{m}}$ is the set of multi-indices with nonzero coefficients:

$$\text{supp}(P) = \{\mathbf{m}: a_{\mathbf{m}} \neq 0\}. \quad (2.1)$$

In the case of a multiaffine polynomial $P(x) = \sum_{S \subseteq E} a_S x^S$, we identify multi-indices with subsets of E , so that

$$\text{supp}(P) = \{S \subseteq E: a_S \neq 0\}. \quad (2.2)$$

If D is a domain (connected open set) in \mathbb{C}^n , we denote by \mathcal{F}_D the set of all functions analytic in D that are either nonvanishing in D or else identically zero. By Hurwitz's theorem³, \mathcal{F}_D is closed in the topology of uniform convergence on compact subsets of D . (That is why we included the phrase “or else identically zero” in the definition of \mathcal{F}_D : it takes care of a trivial degenerate case in a convenient way.)

Let us now suppose that $D = D_1 \times \cdots \times D_n$ where the D_i are domains in \mathbb{C} . If $f \in \mathcal{F}_D$ and we fix some of the variables (say x_1, \dots, x_m) to particular values $x_1^{(0)} \in D_1, \dots, x_m^{(0)} \in D_m$, then clearly $f(x_1^{(0)}, \dots, x_m^{(0)}, \cdot) \in \mathcal{F}_{D_{m+1} \times \cdots \times D_n}$. But more is true, at least if $f \in \mathcal{F}_D$ is also continuous in the closure $\overline{D} = \overline{D_1} \times \cdots \times \overline{D_n}$: then we can fix $x_1^{(0)} \in \overline{D_1}, \dots, x_m^{(0)} \in \overline{D_m}$ and we still have $f(x_1^{(0)}, \dots, x_m^{(0)}, \cdot) \in \mathcal{F}_{D_{m+1} \times \cdots \times D_n}$ (again by Hurwitz's theorem). That is, we can fix some variables even on the boundary of the domain, and as a function of the remaining variables, f must be either nonvanishing or else identically zero.

We denote by H the open right half-plane $\{x \in \mathbb{C}: \text{Re } x > 0\}$, and by \overline{H} the closed right half-plane $\{x \in \mathbb{C}: \text{Re } x \geq 0\}$. If the polynomial P in n variables belongs to \mathcal{F}_{H^n} (i.e. is either nonvanishing in the product of open right half-planes or else identically zero), we say that P has the *half-plane property*.⁴ The discussion of the previous paragraph has the following corollary:

Proposition 2.1 *Let P have the half-plane property, let E' be a subset of the ground set $E = E(P)$, and fix $x' \in \overline{H}^{E'}$. Then $P(x', \cdot)$, considered as a polynomial on the ground set $E \setminus E'$, has the half-plane property. In particular, this holds when we take $x' = 0$.*

2.2 Shifted half-plane property and leading part

Let us denote by $H_{\theta, K}$ the rotated translated open half-plane $\{x \in \mathbb{C}: \text{Re}(e^{-i\theta}x) > K\}$. If the polynomial P in n variables belongs to $\mathcal{F}_{H_{\theta, K}^n}$ for some θ and K , we say

³ Hurwitz's theorem states that if D is a domain in \mathbb{C}^n and (f_k) are nonvanishing analytic functions on D that converge to f uniformly on compact subsets of D , then f is either nonvanishing or else identically zero. Hurwitz's theorem for $n = 1$ is proved in most standard texts on the theory of analytic functions of a single complex variable (see e.g. [1, p. 176]). Surprisingly, we have been unable to find Hurwitz's theorem proven for general n in any standard text on several complex variables (but see [42, p. 306] and [68, p. 337]). So here, for completeness, is the sketch of a proof: Suppose that $f(c) = 0$ for some $c = (c_1, \dots, c_n) \in D$, and let $D' \subset D$ be a small polydisc centered at c . Applying the single-variable Hurwitz theorem, we conclude that $f(z_1, c_2, \dots, c_n) = 0$ for all z_1 such that $(z_1, c_2, \dots, c_n) \in D'$. Applying the same argument repeatedly in the variables z_2, \dots, z_n , we conclude that f is identically vanishing on D' and hence, by analytic continuation, also on D .

⁴ In the engineering literature, polynomials $P \not\equiv 0$ with the half-plane property are termed *widest-sense Hurwitz polynomials*: see e.g. [26].

that P has the *shifted half-plane property*. In particular, if $P \in \mathcal{F}_{H_{\pi, 1/2}^n}$, we say that P has the *Brown–Colbourn property*.

If P is a polynomial of degree r , we denote by P^\sharp the polynomial consisting of those terms in P that have degree r , and we call it the *leading part* of P . Clearly P^\sharp is homogeneous of degree r . Moreover, it is easy to see that

$$P^\sharp(x) = \lim_{\zeta \rightarrow \infty} \zeta^{-r} P(\zeta x) \quad (2.3)$$

as ζ tends to infinity in \mathbb{C} , uniformly for x in compact subsets of \mathbb{C}^n . Using Hurwitz’s theorem, we deduce immediately that:

Proposition 2.2 *If P has the shifted half-plane property, then P^\sharp has the half-plane property [and more generally belongs to $\mathcal{F}_{H_{\theta', 0}^n}$ for all θ'].*

Corollary 2.3 *Let M be a matroid. If the independent-set generating polynomial $P_{\mathcal{I}(M)}$ has the Brown–Colbourn property, then the basis generating polynomial $P_{\mathcal{B}(M)}$ has the half-plane property.*

Henceforth we shall say that a set system \mathcal{S} (resp. a matroid M) “has the half-plane property” if its generating polynomial $P_{\mathcal{S}}$ (resp. its basis generating polynomial $P_{\mathcal{B}(M)}$) has the half-plane property, and that a matroid M “has the Brown–Colbourn property” if its independent-set generating polynomial $P_{\mathcal{I}(M)}$ does.

2.3 Real-part-positive rational functions

Let D be a domain in \mathbb{C}^n , and let f be a complex-valued function that is analytic on D . We say that f is *real-part-positive on D* (resp. *strictly real-part-positive on D*) if $\operatorname{Re} f(x) \geq 0$ (resp. $\operatorname{Re} f(x) > 0$) for all $x \in D$.

Lemma 2.4 *Let f be real-part-positive on D . Then either f is strictly real-part-positive on D , or else f is a pure imaginary constant. (In particular, f is either nonvanishing on D or else identically zero.)*

PROOF. By the open mapping theorem, either the image $f[D]$ is open in \mathbb{C} or else f is constant. Since f is real-part-positive, $f[D]$ is contained in the closed right half-plane \overline{H} . If $f[D]$ is open, then it is contained in the interior of \overline{H} , i.e. in the open right half-plane H , so that f is strictly real-part-positive. If f is constant, this constant value is either pure imaginary or else has strictly positive real part. \square

Lemma 2.5 *Let D be a domain in \mathbb{C}^n , and let k be a positive integer. Let f be real-part-positive (resp. strictly real-part-positive) on H^k , and let g_1, \dots, g_k be strictly real-part-positive on D . Then $f \circ \mathbf{g}$ is real-part-positive (resp. strictly real-part-positive) on D . [Here we have used the obvious shorthand $\mathbf{g} = (g_1, \dots, g_k)$.]*

PROOF. Trivial. \square

Lemma 2.6 *Let D be a domain in \mathbb{C}^n , and let g and h be analytic functions on D . Suppose that h is nonvanishing on D , and that g/h is real-part-positive on D . Then g is either nonvanishing on D or else identically zero.*

In particular, let P and Q be polynomials in n variables, with $Q \not\equiv 0$. Suppose that Q has the half-plane property, and that the rational function P/Q is real-part-positive on H^n . Then P has the half-plane property.

PROOF. By Lemma 2.4, g/h is either a (pure imaginary) constant function or else is strictly real-part-positive on D . If $g/h = c$, then g is either identically zero (if $c = 0$) or nonvanishing on D (if $c \neq 0$). If g/h is strictly real-part-positive on D , then g is manifestly nonvanishing on D . \square

In this lemma we have *assumed* that h is nonvanishing on D in order to guarantee without fuss that g/h is analytic on D . If we drop this assumption, we can still consider g/h as an analytic function on $D \setminus \mathcal{Z}(h)$, where $\mathcal{Z}(h) = \{x \in D : h(x) = 0\}$ is the zero set of h . [Note that $\mathcal{Z}(h)$ is a closed set, and has empty interior whenever $h \not\equiv 0$.] It turns out that no generality is gained by this maneuver, at least when g and h are polynomials and we exclude the trivial possibility that g and h contain a common factor:

Lemma 2.7 *Let $P \not\equiv 0$ and $Q \not\equiv 0$ be polynomials in n complex variables, with P and Q relatively prime (over \mathbb{C}). Let D be a domain in \mathbb{C}^n . Suppose that the rational function P/Q is real-part-positive on $D \setminus \mathcal{Z}(Q)$. Then in fact $\mathcal{Z}(Q) \cap D = \emptyset$.*

PROOF. Suppose that for some $z^{(0)} \in D$ we have $Q(z^{(0)}) = 0$.

(a) If $P(z^{(0)}) \neq 0$, then Q/P is analytic in some neighborhood $U \ni z^{(0)}$ and is nonconstant, so by the open mapping theorem $(Q/P)[U]$ contains a neighborhood $V \ni 0$. Therefore, $(Q/P)[U \setminus \mathcal{Z}(Q)]$ contains $V \setminus \{0\}$, in violation of the hypothesis that P/Q is real-part-positive on $D \setminus \mathcal{Z}(Q)$.

(b) If $P(z^{(0)}) = 0$, then it is known [62, Theorem 1.3.2] that for every neighborhood $U \ni z^{(0)}$ we have $(P/Q)[U \setminus \mathcal{Z}(Q)] = \mathbb{C}$, which again violates the hypothesis that P/Q is real-part-positive on $D \setminus \mathcal{Z}(Q)$. \square

The most important case for us will be $D = H^n$. As we shall see, there is a close interplay between polynomials with the half-plane property and rational functions that are real-part-positive on H^n .⁵ One direction of this interplay is given by Lemmas 2.6 and 2.7; the other direction will be given in Proposition 2.8 below.

⁵ In the engineering literature, rational functions that are real-part-positive on H^n are called *positive* (or *positive real* if their numerator and denominator polynomials have real coefficients): see e.g. [12, 57, 26]. We feel, however, that these terms are likely to cause confusion, so we prefer the more precise term “real-part-positive”.

2.4 Derivatives

If P is a polynomial with ground set E and $\mathbf{m} = \{m_e\}_{e \in E}$ is a multi-index, we define the polynomial $\partial^{\mathbf{m}}P$ in the obvious way: $(\partial^{\mathbf{m}}P)(x) = (\prod_{e \in E} (\partial/\partial x_e)^{m_e})P(x)$. We have the following easy but fundamental result:

Proposition 2.8 *Let P be a polynomial with ground set E , and let $\{\lambda_e\}_{e \in E}$ be non-negative real numbers. Suppose that P has the half-plane property. Then:*

- (a) *The polynomial $\sum_{e \in E} \lambda_e \partial P / \partial x_e$ has the half-plane property.*
- (b) *Provided that $P \not\equiv 0$, the rational function $P^{-1} \sum_{e \in E} \lambda_e \partial P / \partial x_e$ is real-part-positive on H^E . Indeed, it is strictly real-part-positive on H^E except when $\sum_{e \in E} \lambda_e \partial P / \partial x_e \equiv 0$.*

PROOF. If $P \equiv 0$ the proposition is trivial, so assume $P \not\equiv 0$. We shall prove (b); Lemma 2.6 then implies (a).

Consider first the univariate case $|E| = 1$. If P is a constant function, then $P' \equiv 0$ and the theorem is trivial. So let P be a univariate polynomial of degree $k \geq 1$ with the half-plane property, i.e. $P(x) = C \prod_{i=1}^k (x - \alpha_i)$ with $C \neq 0$ and $\operatorname{Re} \alpha_i \leq 0$. Then

$$\frac{P'(x)}{P(x)} = \sum_{i=1}^k \frac{1}{x - \alpha_i} \quad (2.4)$$

is strictly real-part-positive.

Now consider the general multivariate case. Applying the univariate result to x_e with $\{x_f\}_{f \neq e}$ held fixed in the open right half-plane, we conclude that $P^{-1} \partial P / \partial x_e$ is real-part-positive on H^E . The same therefore holds for $P^{-1} \sum_{e \in E} \lambda_e \partial P / \partial x_e$. Moreover, by Lemma 2.4 this function is either strictly real-part-positive on H^E or else a (pure imaginary) constant. But it cannot be a nonzero constant, because $\deg(\sum_{e \in E} \lambda_e \partial P / \partial x_e) < \deg P$. This proves (b) in the general case. \square

Remarks. 1. Part (b) of Proposition 2.8 was proven by Koga [41, Theorem 11], but it is probably older.

2. The same proof shows that the differential operator

$$\mathcal{O} = \alpha + \sum_{e \in E} \beta_e \frac{\partial}{\partial x_e} + \sum_{e \in E} \gamma_e x_e \quad (2.5)$$

with $\operatorname{Re} \alpha \geq 0$ and $\beta_e, \gamma_e \geq 0$ preserves the half-plane property, since the function

$$\frac{\mathcal{O}P}{P} = \frac{\alpha P + \sum_{e \in E} \beta_e \frac{\partial P}{\partial x_e} + \sum_{e \in E} \gamma_e x_e P}{P} \quad (2.6)$$

is real-part-positive.

3. More generally, let $\mathcal{O}_1, \dots, \mathcal{O}_n$ be differential operators of the type (2.5) with $\operatorname{Re} \alpha \geq 0$ and $\beta_e, \gamma_e \geq 0$, and let P_1, \dots, P_n have the half-plane property. For each index i , let Q_i be one of the two functions P_i and $\mathcal{O}_i P_i$, and let R_i be the other one. Then

$$S = \sum_{i=1}^n \left(\prod_{j \neq i} R_j \right) Q_i \quad (2.7)$$

has the half-plane property, because

$$\frac{S}{\prod_{i=1}^n R_i} = \sum_{i=1}^n \frac{Q_i}{R_i} \quad (2.8)$$

is real-part-positive. [If exactly one of the R_i is identically zero, then $S = Q_i$. If two or more of the R_i are identically zero, then $S \equiv 0$.] This construction was inspired by our reading of the brief paper of Bose [7].

Corollary 2.9 *If the polynomial P belongs to $\mathcal{F}_{H_{\theta,K}^n}$, then so does $\partial^{\mathbf{m}} P$ for every multi-index \mathbf{m} . In particular, if P has the half-plane property, then so does $\partial^{\mathbf{m}} P$ for every multi-index \mathbf{m} .*

Suppose we partition the ground set E into two disjoint subsets E' and E'' , and write

$$P(x) = \sum_{\mathbf{m}'} P_{\mathbf{m}'}(x'') (x')^{\mathbf{m}'}, \quad (2.9)$$

where of course $x' = \{x_e\}_{e \in E'}$ and $x'' = \{x_e\}_{e \in E''}$. We then have:

Corollary 2.10 *Let P be written in the form (2.9). If P has the half-plane property, then so do all of the coefficient functions $P_{\mathbf{m}'}$.*

PROOF. We have $P_{\mathbf{m}'}(x'') = (1/\mathbf{m}'!) \partial_{x'}^{\mathbf{m}'} P(x', x'')|_{x'=0}$ where $\mathbf{m}'! = \prod_{e \in E'} m'_e!$. The result then follows immediately from Corollary 2.9 and Proposition 2.1. \square

In fact, a result much stronger than Corollary 2.9 was proven two decades ago by Lieb and one of the authors [47, Proposition 2.2]:

Theorem 2.11 (Lieb and Sokal [47]) *Let $\{P_i\}_{i=1}^k$ and $\{Q_i\}_{i=1}^k$ be polynomials in n complex variables, and define*

$$R(v, w) = \sum_{i=1}^k P_i(v) Q_i(w) \quad (2.10)$$

$$S(x) = \sum_{i=1}^k P_i(\partial/\partial x) Q_i(x) \quad (2.11)$$

If R has the half-plane property, then so does S .

In particular, we have the case $k = 1$: if P and Q have the half-plane property, then so does $P(\partial/\partial x)Q(x)$.

2.5 The Grace–Walsh–Szegő coincidence theorem

Finally, we shall need a version of the Grace–Walsh–Szegő coincidence theorem [31, 71, 74]. Let $P(x_1, \dots, x_n)$ be a multiaffine polynomial that is *symmetric* under all permutations of the x_1, \dots, x_n . Any such polynomial can be written in the form

$$P(x_1, \dots, x_n) = \sum_{k=0}^n a_k \binom{n}{k}^{-1} E_k(x_1, \dots, x_n) \quad (2.12)$$

where the E_k are the elementary symmetric polynomials, i.e. $E_0 = 1$ and

$$E_k(x_1, \dots, x_n) = \sum_{1 \leq i_1 < i_2 < \dots < i_k \leq n} x_{i_1} x_{i_2} \cdots x_{i_k} ; \quad (2.13)$$

and conversely, any polynomial of the form (2.12) is symmetric and multiaffine. Now let us define a *closed circular region* in \mathbb{C} to be a closed disc (including the degenerate case of a single point), a closed half-plane, or the closed exterior of a disc. And let us define an *open circular region* in \mathbb{C} to be an open disc, an open half-plane, or the open exterior of a disc. We then have:

Theorem 2.12 (Grace–Walsh–Szegő) *Let P be a symmetric multiaffine polynomial in n complex variables, let C be an open or closed circular region in \mathbb{C} , and let $x_1^{(0)}, \dots, x_n^{(0)}$ be points in the region C . Suppose, further, that either $\deg P = n$ (i.e. $a_n \neq 0$) or C is convex (or both). Then there exists at least one point $\xi \in C$ such that*

$$P(x_1^{(0)}, \dots, x_n^{(0)}) = P(\xi, \dots, \xi) . \quad (2.14)$$

PROOF. Note first that it suffices to prove the theorem for closed circular regions: for if C is an open circular region, then there exists a closed circular region $C' \subset C$ that still contains the points $x_1^{(0)}, \dots, x_n^{(0)}$. Now, the standard proof of the Grace–Walsh–Szegő coincidence theorem (see e.g. [51, Theorem 15.4]) applies whenever $a_n \neq 0$. If $a_n = 0$ and C is a closed disc, then by taking limits from the case $a_n \neq 0$ and using the compactness of C , we can obtain the desired result.⁶

Finally, suppose that $a_n = 0$ and C is a closed half-plane. Without loss of generality we can suppose that C is the closed right half-plane \overline{H} . Let $\alpha = \max\{|\operatorname{Im} x_i^{(0)}| : 1 \leq$

⁶ **PROOF.** Let $P_\epsilon(x_1, \dots, x_n) = P(x_1, \dots, x_n) + \epsilon x_1 \cdots x_n$. Applying the case $a_n \neq 0$, we conclude that for each $\epsilon \neq 0$ there exists $\xi_\epsilon \in C$ such that $P_\epsilon(x_1^{(0)}, \dots, x_n^{(0)}) = P_\epsilon(\xi_\epsilon, \dots, \xi_\epsilon)$. Since C is compact, the family $\{\xi_\epsilon\}$ must have at least one limit point $\xi \in C$ as $\epsilon \rightarrow 0$. This limit point satisfies (2.14).

$i \leq n\}$ and $\beta = \max\{\operatorname{Re} x_i^{(0)} : 1 \leq i \leq n\}$. Let C_R be the closed disc of radius $\sqrt{R^2 + \alpha^2}$ centered at R . Note that C_R contains the points $x_1^{(0)}, \dots, x_n^{(0)}$ whenever $R \geq \beta/2$. Now apply the theorem to C_R : we obtain a point $\xi_R \in C_R$ satisfying $P(x_1^{(0)}, \dots, x_n^{(0)}) = P(\xi_R, \dots, \xi_R)$. If, for at least one R , the point ξ_R lies in $C_R \cap \overline{H}$, we are done; and if not, then the points ξ_R all lie in $C_R \setminus \overline{H}$, and as $R \rightarrow \infty$ they must have a limit point ξ in the interval $[-i\alpha, i\alpha]$ of the imaginary axis. \square

This theorem is usually applied as follows: Starting from a univariate polynomial $Q(x) = \sum_k a_k x^k$ and an integer $n \geq \deg Q$, we construct the unique symmetric multiaffine polynomial $P(x_1, \dots, x_n)$ satisfying $P(x, \dots, x) = Q(x)$, namely (2.12). In this situation, Theorem 2.12 implies that if Q is nonvanishing in a circular region C , then P is nonvanishing in C^n (provided that $n = \deg Q$ or C is convex).

More generally, starting from a multivariate polynomial Q in variables $\{x_e\}_{e \in E}$ and integers $n_e \geq \deg_e Q$, we can introduce variables $\{x_e^{(j)}\}_{e \in E, 1 \leq j \leq n_e}$ and form the unique multiaffine polynomial P in all these variables that is symmetric in each family $X_e = \{x_e^{(j)}\}_{1 \leq j \leq n_e}$ separately and reduces to Q when $x_e^{(j)} = x_e$ for all j, e . We refer to P as being obtained by the n_e -fold *polarization* of x_e in Q , for all $e \in E$. Then Theorem 2.12 implies that if $\{C_e\}_{e \in E}$ are circular regions and $Q(x)$ is nonvanishing when $x_e \in C_e$ for all e , then P is nonvanishing when $x_e^{(j)} \in C_e$ for all j, e (provided that, for each e , either $n_e = \deg_e Q$ or C_e is convex). By this technique, we can often prove theorems for general polynomials Q by reducing them to the special case of multiaffine polynomials P (albeit in a larger number of variables).

2.6 Algorithms

It is worth remarking that the half-plane property is algorithmically decidable, using quantifier-elimination methods for the theory of real closed fields [15]. Indeed, let $P \not\equiv 0$ be a polynomial in complex variables x_1, \dots, x_n . Setting $x_k = a_k + ib_k$ and separating out real and imaginary parts

$$R(a_1, \dots, a_n, b_1, \dots, b_n) = \operatorname{Re} P(\{a_k + ib_k\}) \quad (2.15a)$$

$$I(a_1, \dots, a_n, b_1, \dots, b_n) = \operatorname{Im} P(\{a_k + ib_k\}) \quad (2.15b)$$

the half-plane property for P is immediately seen to be equivalent to the assertion

$$\neg(\exists a_1, \dots, a_n, b_1, \dots, b_n \in \mathbb{R}) (a_1 > 0) \wedge \dots \wedge (a_n > 0) \wedge (R(a, b) = 0) \wedge (I(a, b) = 0) \quad (2.16)$$

in the first-order theory of the real numbers, which is decidable according to a classic result of Tarski [72]. Indeed, by the method of cylindrical algebraic decomposition (CAD) [20], this computation can be performed in time $c_1^{c_2^n}$ for suitable constants c_1 and c_2 (“doubly exponential time”). Moreover, some more recent algorithms [34, 36, 60, 5] require only a time c^n (“singly exponential time”). Unfortunately, this computation seems at present to be unfeasible in practice even for $n = 4$.

3 The local half-plane property

For any element $e \in E$, the *deletion of e from P* is the polynomial $P^{\setminus e}$ on ground set $E \setminus e$ that is obtained from P by setting $x_e = 0$. The *contraction of e from P* is the polynomial $P^{/e}$ on ground set E defined by $P^{/e} = \partial P / \partial x_e$. Note that if P is affine in e , then $P^{/e}$ can also be considered as a polynomial on ground set $E \setminus e$ (and we shall usually do so). We say that an element $e \in E$ is a *loop* of P in case $P^{/e} \equiv 0$, and is a *coloop* of P in case $P^{\setminus e} \equiv 0$.

Proposition 3.1 (Deletion/contraction) *Let P be a polynomial with the half-plane property. Then, for every $e \in E(P)$, both $P^{\setminus e}$ and $P^{/e}$ have the half-plane property.*

PROOF. Proposition 2.1 implies that $P^{\setminus e}$ has the half-plane property. Proposition 2.8(a) implies that $P^{/e}$ has the half-plane property. \square

Remark. If P is affine in e , then the half-plane property for the contraction $P^{/e}$ can alternatively be proven using

$$P^{/e}(x_{\neq e}) = \lim_{x_e \rightarrow +\infty} \frac{P(x)}{x_e} \quad (3.1)$$

[see (3.2) below] and Hurwitz's theorem.

Let P be a polynomial and $e \in E(P)$, and suppose that P is affine in e . Then P can be written in the form

$$P(x) = P^{\setminus e}(x_{\neq e}) + x_e P^{/e}(x_{\neq e}), \quad (3.2)$$

where we have used the shorthand $x_{\neq e} = \{x_f\}_{f \in E \setminus e}$. In this situation, we say that the pair (P, e) satisfies the *local half-plane property* if, whenever $\operatorname{Re} x_f > 0$ for all $f \in E \setminus e$, we have $P^{\setminus e}(x_{\neq e}) \neq 0$, $P^{/e}(x_{\neq e}) \neq 0$ and

$$\operatorname{Re} \frac{P^{\setminus e}(x_{\neq e})}{P^{/e}(x_{\neq e})} \geq 0. \quad (3.3)$$

Otherwise put, (P, e) has the local half-plane property in case

- (a) e is neither a loop nor a coloop of P ,
- (b) both $P^{\setminus e}$ and $P^{/e}$ have the half-plane property, and
- (c) $P^{\setminus e}/P^{/e}$ is real-part-positive on $H^{E \setminus e}$.

Note also that by Lemma 2.4, there is either strict inequality in (3.3) or else $P^{\setminus e}/P^{/e}$ is a nonzero pure imaginary constant [i.e. $P(x) = (1 + i\alpha x_e)P^{\setminus e}(x_{\neq e})$ for some nonzero real number α].

We have the following fundamental result:

Theorem 3.2 *Let P be affine in e . Then the following are equivalent:*

- (a) P has the half-plane property.
- (b) One of the following four mutually exclusive possibilities holds:
 - (i) $P^{/e} \equiv 0$ and $P^{\setminus e} \equiv 0$;
 - (ii) $P^{/e} \equiv 0$, and $P^{\setminus e} \not\equiv 0$ has the half-plane property;
 - (iii) $P^{\setminus e} \equiv 0$, and $P^{/e} \not\equiv 0$ has the half-plane property;
 - (iv) (P, e) has the local half-plane property, i.e. $P^{\setminus e} \not\equiv 0$ and $P^{/e} \not\equiv 0$ both have the half-plane property, and

$$\operatorname{Re} \frac{P^{\setminus e}(x_{\neq e})}{P^{/e}(x_{\neq e})} \geq 0 \quad (3.4)$$

for all $x_{\neq e} \in H^{E \setminus e}$.

- (c) One of the following three mutually exclusive possibilities holds:
 - (i) $P^{/e} \equiv 0$ and $P^{\setminus e} \equiv 0$;
 - (ii) $P^{/e} \equiv 0$, and $P^{\setminus e} \not\equiv 0$ has the half-plane property;
 - (iii) $P^{/e} \not\equiv 0$ has the half-plane property, and

$$\operatorname{Re} \frac{P(x)}{P^{/e}(x_{\neq e})} > 0 \quad (3.5)$$

for all $x \in H^E$.

PROOF. (a) \implies (b): By Proposition 3.1, both $P^{\setminus e}$ and $P^{/e}$ have the half-plane property. If one or both of them is identically zero, we are in one of the cases (i)–(iii). If neither is identically zero, then both $P^{\setminus e}(x_{\neq e})$ and $P^{/e}(x_{\neq e})$ are nonzero for all $x_{\neq e} \in H^{E \setminus e}$. Therefore, for each such $x_{\neq e}$, there is a unique value for x_e such that $P(x) = 0$, namely

$$x_e = -\frac{P^{\setminus e}(x_{\neq e})}{P^{/e}(x_{\neq e})}. \quad (3.6)$$

Since P has the half-plane property, we must have $\operatorname{Re} x_e \leq 0$, proving (3.4).

(b) \implies (c): This is trivial if either (b)(i) or (b)(ii) holds; so suppose that either (b)(iii) or (b)(iv) holds, and let $x \in H^E$. Then $P^{/e}(x_{\neq e}) \neq 0$, and

$$\operatorname{Re} \frac{P(x)}{P^{/e}(x_{\neq e})} = \operatorname{Re} \left(\frac{P^{\setminus e}(x_{\neq e})}{P^{/e}(x_{\neq e})} + x_e \right) > 0 \quad (3.7)$$

since $\operatorname{Re} [P^{\setminus e}(x_{\neq e})/P^{/e}(x_{\neq e})] \geq 0$ and $\operatorname{Re} x_e > 0$.

(c) \implies (a): In cases (c)(i) and (c)(ii) this is trivial. And in case (c)(iii), we clearly have $P(x) \neq 0$ for all $x \in H^E$, so it is also trivial. \square

Corollary 3.3 *Let P be a multiaffine polynomial. Then the following are equivalent:*

- (a) *P has the half-plane property.*
- (b) *There exists at least one $e \in E(P)$ for which condition (b) of Theorem 3.2 holds.*
- (b') *For all $e \in E(P)$, condition (b) of Theorem 3.2 holds.*
- (c) *There exists at least one $e \in E(P)$ for which condition (c) of Theorem 3.2 holds.*
- (c') *For all $e \in E(P)$, condition (c) of Theorem 3.2 holds.*

Now let us drop the hypothesis that P is affine in e . We can still write P in the form

$$P(x) = \sum_{k=0}^M P_k(x_{\neq e}) x_e^k \quad (3.8)$$

where $M = \deg_e P$. There is no longer any simple necessary and sufficient condition, analogous to Theorem 3.2, for P to have the half-plane property. But Fettweis and Basu [26, Theorem 18] have proven a very interesting *necessary* condition:

Theorem 3.4 (Fettweis and Basu [26]) *Let P be a polynomial with the half-plane property, written in the form (3.8) for some $e \in E(P)$. Then each coefficient function P_k has the half-plane property, and moreover for each $r \geq 0$ we have:*

- (a) *It is impossible to have $P_r \neq 0$, $P_{r+1} \equiv P_{r+2} \equiv \dots \equiv P_{r+\ell} \equiv 0$, $P_{r+\ell+1} \neq 0$ with $\ell \geq 2$.*
- (b) *If $P_r \neq 0$, $P_{r+1} \equiv 0$ and $P_{r+2} \neq 0$, then P_{r+2}/P_r is a strictly positive constant.*
- (c) *If $P_r \neq 0$ and $P_{r+1} \neq 0$, then P_{r+1}/P_r is real-part-positive on $H^{E \setminus e}$.*

To prove Theorem 3.4, we start with a lemma [26, Lemma 4]:

Lemma 3.5 *Let $P(x) = \sum_{k=0}^M P_k(x_{\neq e}) x_e^k$ be a polynomial. Let $0 \leq r \leq s \leq M$ and define*

$$Q(x) = \sum_{k=r}^s \frac{k!}{(k-r)!} \frac{(M-k)!}{(s-k)!} P_k(x_{\neq e}) x_e^k. \quad (3.9)$$

If P has the half-plane property, then so does Q .

PROOF. Define

$$P_1(x) = \frac{\partial^r}{\partial x_e^r} P(x) \quad (3.10)$$

$$P_2(x) = x_e^{M-r} P_1(x_{\neq e}, 1/x_e) \quad (3.11)$$

$$P_3(x) = \frac{\partial^{M-s}}{\partial x_e^{M-s}} P_2(x) \quad (3.12)$$

$$Q(x) = x_e^s P_3(x_{\neq e}, 1/x_e) \quad (3.13)$$

By Proposition 2.8 and the fact that $x_e \mapsto 1/x_e$ maps the open right half-plane onto itself, all these operations preserve the half-plane property. \square

PROOF OF THEOREM 3.4. We have already proven (Corollary 2.10) that all the coefficient functions P_k have the half-plane property, Now apply Lemma 3.5 with $s = r + \ell + 1$ [using $\ell = 1$ in (b) and $\ell = 0$ in (c)], yielding a polynomial

$$Q(x) = \alpha P_r(x_{\neq e}) x_e^r + \beta P_{r+\ell+1}(x_{\neq e}) x_e^{r+\ell+1} \quad (3.14)$$

with $\alpha, \beta > 0$ and $P_r, P_{r+\ell+1} \not\equiv 0$ having the half-plane property. If $\ell \geq 2$, this polynomial Q never has the half-plane property [just choose any $x_{\neq e} \in H^{E \setminus e}$, then there exists a solution to $Q(x) = 0$ with $\operatorname{Re} x_e > 0$]. If $\ell = 1$, then Q has the half-plane property if and only if $P_{r+2}(x_{\neq e})/P_r(x_{\neq e}) > 0$ for all $x_{\neq e} \in H^{E \setminus e}$; but by the open mapping theorem, this implies that P_{r+2}/P_r is a constant function. Finally, if $\ell = 0$, then Q has the half-plane property if and only if $\operatorname{Re} [P_{r+1}(x_{\neq e})/P_r(x_{\neq e})] \geq 0$ for all $x_{\neq e} \in H^{E \setminus e}$. \square

4 Constructions

In the previous section we saw that if a polynomial P has the half-plane property, then so do the deletion $P^{\setminus e}$ and the contraction $P^{/e}$, for every element $e \in E$. In this section we describe some other constructions that preserve the half-plane property.

Let us recall that the *support* of a multiaffine polynomial $P(x) = \sum_{S \subseteq E} a_S x^S$ is $\operatorname{supp}(P) = \{S \subseteq E : a_S \neq 0\}$. We shall see later (Theorem 7.1) that if $\bar{P} \not\equiv 0$ is a homogeneous multiaffine polynomial with the half-plane property, then $\operatorname{supp}(P)$ is necessarily the set $\mathcal{B}(M)$ of bases of some matroid M on the ground set $E(P)$. (Theorem 7.2 generalizes this result to homogeneous polynomials that are not necessarily multiaffine.) Therefore, while describing these constructions on polynomials, we will also explain, in the homogeneous multiaffine case, the corresponding operations on matroids. Indeed, our choice of terminology for these constructions on polynomials is motivated by analogy with standard terms in matroid theory.

4.1 Deletion and contraction

The following proposition is evident from the definitions:

Proposition 4.1 *Let P be a homogeneous multiaffine polynomial with $\operatorname{supp}(P) = \mathcal{B}(M)$ for a matroid M , and let $e \in E(P)$. Then:*

- (a) *e is a loop of P if and only if e is a loop of M .*
- (b) *e is a coloop of P if and only if e is a coloop of M .*

- (c) If e is not a loop of P , then $\text{supp}(P^{/e}) = \mathcal{B}(M/e)$, the contraction of M by e .
- (d) If e is not a coloop of P , then $\text{supp}(P^{\setminus e}) = \mathcal{B}(M \setminus e)$, the deletion of e from M .
- (e) If e is not a loop of P , and P is the basis generating polynomial for M , then $P^{/e}$ is the basis generating polynomial for M/e .
- (f) If e is not a coloop of P , and P is the basis generating polynomial for M , then $P^{\setminus e}$ is the basis generating polynomial for $M \setminus e$.

4.2 Duality

Given a multiaffine polynomial $P(x) = \sum_{S \subseteq E} a_S x^S$ on the ground set E , we define the *dual polynomial*

$$P^*(x) = x^E P(1/x) = \sum_{S \subseteq E} a_S x^{E \setminus S}. \quad (4.1)$$

We have already encountered (in the Introduction) a special case of the following principle:

Proposition 4.2 (Duality) *Let P be a multiaffine polynomial.*

If P has the half-plane property, then the dual polynomial P^ also has the half-plane property.*

If $\text{supp}(P) = \mathcal{B}(M)$ for a matroid M , then $\text{supp}(P^) = \mathcal{B}(M^*)$, where M^* is the matroid dual to M .*

If P is the basis generating polynomial for a matroid M , then P^ is the basis generating polynomial for M^* .*

PROOF. The half-plane property for P^* follows immediately from the fact that $x_e \mapsto 1/x_e$ maps the open right half-plane onto itself. The remainder follows immediately from the definition of the dual matroid M^* . \square

Let us also record, for future reference, some easy identities relating duality to deletion and contraction:

Lemma 4.3 *Let P be a multiaffine polynomial on the ground set E , and let $e \in E$. Then*

$$(P^{\setminus e})^* = (P^*)^{/e} \quad (4.2)$$

$$(P^{/e})^* = (P^*)^{\setminus e} \quad (4.3)$$

where $P^{\setminus e}$, $P^{/e}$, $(P^*)^{/e}$ and $(P^*)^{\setminus e}$ are all considered as having ground set $E \setminus e$.

4.3 Direct sum

The next proposition is trivial, and has already been used implicitly:

Proposition 4.4 *Let P and Q be polynomials with the half-plane property. Then the pointwise product PQ [considered on the ground set $E(P) \cup E(Q)$] also has the half-plane property.*

The special case where P and Q have disjoint ground sets has a simple matroidal interpretation:

Proposition 4.5 (Direct sum) *Let P and Q be homogeneous multiaffine polynomials with $E(P) \cap E(Q) = \emptyset$.*

If $\text{supp}(P) = \mathcal{B}(M)$ and $\text{supp}(Q) = \mathcal{B}(N)$ for matroids M and N , then $\text{supp}(PQ) = \mathcal{B}(M \oplus N)$, the direct sum of M and N .

If P and Q are the basis generating polynomials for matroids M and N , then PQ is the basis generating polynomial for $M \oplus N$.

Clearly, if a matroid M is the direct sum of two matroids M_1, M_2 of nonzero rank, then its basis generating polynomial $P_{\mathcal{B}(M)}$ is reducible. A very strong version of the converse is true as well:

Proposition 4.6 *Let M be a connected matroid with ground set E , and let P be a multiaffine polynomial in the indeterminates $\{x_e\}_{e \in E}$ with coefficients in an integral domain R . If $\text{supp}(P) = \mathcal{B}(M)$, then P is irreducible over R .*

We begin with an easy but crucial lemma:

Lemma 4.7 *Let P_1 and P_2 be nonzero polynomials in the indeterminates $\{x_e\}_{e \in E}$ with coefficients in an integral domain R . Suppose that $P_1 P_2$ is multiaffine. Then:*

- (a) *There exist disjoint subsets $E_1, E_2 \subseteq E$ such that P_i uses only the variables $\{x_e\}_{e \in E_i}$ ($i = 1, 2$).*
- (b) *P_1 and P_2 are both multiaffine.*

PROOF. Suppose there exists $e \in E$ such that both P_1 and P_2 use the variable x_e . For $i = 1, 2$, let $d_i \geq 1$ be the degree of P_i in the variable x_e , and let $Q_i \neq 0$ be the coefficient of $x_e^{d_i}$ in P_i , considered as an element of the polynomial ring $R[x_{\neq e}]$. Then $Q_1 Q_2 \neq 0$ because $R[x_{\neq e}]$ is an integral domain [39, Theorem III.5.1 and Corollary III.5.7]. But this shows that the coefficient of $x_e^{d_1 + d_2}$ in $P_1 P_2$ is nonzero, contradicting the hypothesis that $P_1 P_2$ is multiaffine (since $d_1 + d_2 \geq 2$). This proves (a); and (b) is an easy consequence. \square

PROOF OF PROPOSITION 4.6. If $|E| = 1$, the result is trivial, so assume henceforth that $|E| \geq 2$. Suppose that $P = P_1 P_2$ where P_1 and P_2 are non-constant polynomials over R . By Lemma 4.7, there exist disjoint subsets $E_1, E_2 \subseteq E$ such that P_i uses only the variables $\{x_e\}_{e \in E_i}$. Moreover, each term x^S in P arises from a unique pair of terms in P_1 and P_2 (namely, $x^{S \cap E_1}$ and $x^{S \cap E_2}$); and conversely, each pair of nonzero terms in P_1 and P_2 gives rise to a nonzero term in P (because R is an integral domain). Since every term in P has degree $r(M)$ [the rank of M], it follows that there exist integers $r_1, r_2 \geq 1$ such that every term in P_i has degree r_i ($i = 1, 2$).

Let B be a basis of M . For each element $i \in E \setminus B$, there is a unique circuit $C(i, B)$ of M contained in $B \cup \{i\}$. Consider the bipartite graph G with vertex classes B and $E \setminus B$ where a vertex $i \in E \setminus B$ is adjacent to a vertex $j \in B$ if and only if $j \in C(i, B) \setminus i$. By a theorem of Cunningham [21] and Krogdahl [43] (see [55, Lemma 10.2.8]), since $|E| \geq 2$ and M is connected, G is a connected graph. Thus, by interchanging E_1 and E_2 if necessary, we may assume that $E_1 \setminus B$ contains an element i such that $C(i, B)$ meets $E_2 \cap B$. Suppose $j \in C(i, B) \cap E_2 \cap B$. Then $(B \setminus \{j\}) \cup \{i\}$ is a basis of M . But $|(B \setminus \{j\}) \cup \{i\} \cap E_1| > |B \cap E_1|$, so P_1 has a term of degree exceeding r_1 : a contradiction. \square

Remark. This proof is a natural extension of the proof given by Duffin and Morley [24, Theorem 8] of the irreducibility of the basis generating polynomial of a connected *binary* matroid.

4.4 Parallel connection, series connection, and 2-sum

Let P and Q be polynomials such that $E(P) \cap E(Q) = \{e\}$. The *parallel connection* of P and Q is defined to be

$$(P \parallel Q)(x) = P^{\setminus e}(x) Q^{/e}(x) + P^{/e}(x) Q^{\setminus e}(x) + x_e P^{/e}(x) Q^{/e}(x). \quad (4.4)$$

Note that if both P and Q are affine in e , then so is $P \parallel Q$; and if both P and Q are multiaffine, then so is $P \parallel Q$ (the latter assertion uses the fact that $E(P)$ and $E(Q)$ meet only in e).

Proposition 4.8 (Parallel connection) *Let P and Q be polynomials such that $E(P) \cap E(Q) = \{e\}$, and suppose further that both P and Q are affine in e . If both P and Q have the half-plane property, then $P \parallel Q$ also has the half-plane property.*

If, in addition, both P and Q are homogeneous and multiaffine, with $\text{supp}(P) = \mathcal{B}(M)$ and $\text{supp}(Q) = \mathcal{B}(N)$ for matroids M and N , and e is not a loop in at least one of P and Q , then $\text{supp}(P \parallel Q) = \mathcal{B}(M \parallel N)$, the parallel connection of M and N .⁷

If P and Q are the basis generating polynomials for matroids M and N , and e is not a loop in at least one of P and Q , then $P \parallel Q$ is the basis generating polynomial for $M \parallel N$.

⁷ The standard notation for the parallel connection of M and N is $P(M, N)$, but in order to avoid confusion with polynomials, we write $M \parallel N$ instead.

FIRST PROOF. If e is a loop of P , then $(P\|Q) = P^{\setminus e}Q^{/e}$; while if e is a coloop of P , then $P\|Q = P^{/e}Q$. Propositions 3.1 and 4.4 imply that $P^{\setminus e}Q^{/e}$ and $P^{/e}Q$ have the half-plane property. If e is a loop or coloop of Q , we argue analogously. Hence we may assume that e is not a loop or coloop in either P or Q .

From Theorem 3.2(a) \implies (b), both (P, e) and (Q, e) have the local half-plane property. Let x be such that $\operatorname{Re} x_f > 0$ for all $f \in (E(P) \cup E(Q)) \setminus e$. Then $P^{\setminus e}(x)$, $P^{/e}(x)$, $Q^{\setminus e}(x)$ and $Q^{/e}(x)$ are all nonzero, by the local half-plane property. If $(P\|Q)(x) = 0$, we can solve this equation for x_e , yielding

$$x_e = -\frac{P^{\setminus e}(x)}{P^{/e}(x)} - \frac{Q^{\setminus e}(x)}{Q^{/e}(x)}, \quad (4.5)$$

which has nonpositive real part as a consequence of the local half-plane property for (P, e) and (Q, e) . Therefore, $P\|Q$ has the half-plane property.

According to [55, Proposition 7.1.13(ii)], the bases of $M\|N$ are exactly those sets that can be obtained from some $B_1 \in \mathcal{B}(M)$ and some $B_2 \in \mathcal{B}(N)$ in one of the following ways:

- (i) $(B_1 \cup B_2) \setminus e$, where $e \notin B_1$ and $e \in B_2$;
- (ii) $(B_1 \cup B_2) \setminus e$, where $e \in B_1$ and $e \notin B_2$;
- (iii) $B_1 \cup B_2$, where $e \in B_1$ and $e \in B_2$.

These three classes correspond precisely to the three terms of (4.4). Moreover, it is easy to see that each basis B of $M\|N$ is obtained in this result from a unique pair $(B_1, B_2) \in \mathcal{B}(M) \times \mathcal{B}(N)$. The claims about support and about basis generating polynomials follow immediately. \square

SECOND PROOF. Since P and Q are affine in e , we have

$$P(x) = P^{\setminus e}(x_{\neq e}) + x_e P^{/e}(x_{\neq e}) \quad (4.6)$$

$$Q(x) = Q^{\setminus e}(x_{\neq e}) + x_e Q^{/e}(x_{\neq e}) \quad (4.7)$$

and hence

$$\frac{\partial}{\partial x_e}(PQ)(x) = P^{\setminus e}(x)Q^{/e}(x) + P^{/e}(x)Q^{\setminus e}(x) + 2x_e P^{/e}(x)Q^{/e}(x). \quad (4.8)$$

By Propositions 4.4 and 2.8(a), the polynomial (4.8) has the half-plane property. Replacing x_e by $x_e/2$ (which preserves the half-plane property), we obtain $(P\|Q)(x)$. The rest of the proposition is proven as before. \square

Remark. The assertion in Proposition 4.8 that $P\|Q$ has the half-plane property does not require the hypothesis that $E(P) \cap E(Q) = \{e\}$. But without this hypothesis, $P\|Q$ may not be multiaffine, even if P and Q both are.

Corollary 4.9 (2-sum) *Let P and Q be polynomials such that $E(P) \cap E(Q) = \{e\}$, and suppose further that P and Q are both affine in e . If both P and Q have the half-plane property, then $(P \parallel Q)^{\setminus e}$ also has the half-plane property.*

If, in addition, both P and Q are homogeneous and multiaffine, with $\text{supp}(P) = \mathcal{B}(M)$ and $\text{supp}(Q) = \mathcal{B}(N)$ for matroids M and N , and e is not a loop or coloop in either P or Q , then $\text{supp}((P \parallel Q)^{\setminus e}) = \mathcal{B}(M \oplus_2 N)$, the 2-sum of M and N .

If P and Q are the basis generating polynomials for matroids M and N , respectively, and e is not a loop or coloop in either P or Q , then $(P \parallel Q)^{\setminus e}$ is the basis generating polynomial for $M \oplus_2 N$.

PROOF. This follows immediately from Propositions 3.1 and 4.8 and the definition of 2-sum. \square

Let P and Q be polynomials such that $E(P) \cap E(Q) = \{e\}$. The *series connection* of P and Q is defined to be

$$(P \& Q)(x) = P^{\setminus e}(x)Q^{\setminus e}(x) + x_e P^{\setminus e}(x)Q^{/e}(x) + x_e P^{/e}(x)Q^{\setminus e}(x). \quad (4.9)$$

Note that if both P and Q are affine in e , then so is $P \& Q$, and indeed we have $P \& Q = (P^{*e} \parallel Q^{*e})^{*e}$ where *e denotes duality with respect to x_e only [that is, $P^{*e}(x_e, x_{\neq e}) = x_e P(1/x_e, x_{\neq e})$; this operation preserves the half-plane property, by the same argument used in the proof of Proposition 4.2]. If both P and Q are multiaffine, then so is $P \& Q$, and we have $P \& Q = (P^* \parallel Q^*)^*$.

Proposition 4.10 (Series connection) *Let P and Q be polynomials such that $E(P) \cap E(Q) = \{e\}$, and suppose further that P and Q are both affine in e . If both P and Q have the half-plane property, then $P \& Q$ also has the half-plane property.*

If, in addition, both P and Q are homogeneous and multiaffine, with $\text{supp}(P) = \mathcal{B}(M)$ and $\text{supp}(Q) = \mathcal{B}(N)$ for matroids M and N , and e is not a coloop in at least one of P and Q , then $\text{supp}(P \& Q) = \mathcal{B}(M \& N)$, the series connection of M and N .⁸

If P and Q are the basis generating polynomials for matroids M and N , and e is not a coloop in at least one of P and Q , then $P \& Q$ is the basis generating polynomial for $M \& N$.

PROOF. The proof is analogous to that of Proposition 4.8. \square

⁸ The standard notation for the series connection of M and N is $S(M, N)$, but in order to avoid confusion with polynomials, we write $M \& N$ instead.

4.5 Principal truncation and principal extension

Let P be a polynomial with ground set E , and let $\{\lambda_e\}_{e \in E}$ be nonnegative real numbers. Then the polynomial

$$\mathrm{tr}_\lambda P(x) = \sum_{e \in E} \lambda_e P^{/e}(x) \quad (4.10)$$

is called a *(weighted) principal truncation* of P . If a is a new element (not occurring in E), then

$$\mathrm{ext}_\lambda P(x_{\cup a}) = P(x) + x_a \mathrm{tr}_\lambda P(x) \quad (4.11)$$

is called a *(weighted) principal extension* of P ; here we have used the shorthand $x_{\cup a} = \{x_e\}_{e \in E \cup a}$.

Proposition 4.11 (Principal truncation/extension) *Let P be a polynomial with ground set E , and let $\{\lambda_e\}_{e \in E}$ be nonnegative real numbers. If P has the half-plane property, then both $\mathrm{tr}_\lambda P$ and $\mathrm{ext}_\lambda P$ have the half-plane property.*

PROOF. Proposition 2.8(a) asserts that $\mathrm{tr}_\lambda P$ has the half-plane property. Proposition 2.8(b) together with Theorem 3.2(b) \implies (a) imply that $\mathrm{ext}_\lambda P$ has the half-plane property. \square

The terms “principal truncation” and “principal extension” are taken from matroid theory; they are justified by the following result:

Proposition 4.12 *Let P be a homogeneous multiaffine polynomial with ground set E , and let $\{\lambda_e\}_{e \in E}$ be nonnegative real numbers, not all of which are zero. Suppose that $\mathrm{supp}(P) = \mathcal{B}(M)$ for a matroid M of nonzero rank, and let F be the closure in M of the set $S = \{e: \lambda_e > 0\}$. Then $\mathrm{supp}(\mathrm{tr}_\lambda P) = \mathcal{B}(\mathrm{tr}_F(M))$, where $\mathrm{tr}_F(M)$ is the principal truncation of M with respect to F . Also, $\mathrm{supp}(\mathrm{ext}_\lambda P) = \mathcal{B}(M +_F a)$, where $M +_F a$ is the principal extension of M on F .*

PROOF. The bases of the truncation $\mathrm{tr}_F(M)$ are all the sets of the form $B \setminus f$ where $B \in \mathcal{B}(M)$ and $f \in B \cap F$. The sets that appear in the support of $\mathrm{tr}_\lambda P$ are all the sets of the form $B \setminus g$ where $B \in \mathcal{B}(M)$ and $g \in B \cap S$. Since $S \subseteq F$, we clearly have $\mathrm{supp}(\mathrm{tr}_\lambda P) \subseteq \mathcal{B}(\mathrm{tr}_F(M))$. Conversely, suppose $B \in \mathcal{B}(M)$ and $f \in B \cap F$, and let r be the rank of M . Since $\mathrm{rank}(B \setminus f) = r - 1$ and $\mathrm{rank}((B \setminus f) \cup S) = r$, the independence augmentation axiom implies that there is an element $g \in S$ such that $B' = (B \setminus f) \cup \{g\}$ is a basis of M . Since $B \setminus f = B' \setminus g$, we conclude that $\mathcal{B}(\mathrm{tr}_F(M)) \subseteq \mathrm{supp}(\mathrm{tr}_\lambda P)$, as desired.

The bases of the extension $M +_F a$ are given by

$$\mathcal{B}(M +_F a) = \mathcal{B}(M) \cup \{B \cup \{a\}: B \in \mathcal{B}(\mathrm{tr}_F(M))\}. \quad (4.12)$$

From this and the previous paragraph, the claim about the support of $\text{ext}_\lambda P$ follows readily. \square

In the situation of Propositions 4.11 and 4.12, even if P is the basis generating polynomial of the matroid M , it need not be the case that $\text{tr}_\lambda P$ is the basis generating polynomial of $\text{tr}_F(M)$, since the coefficients need not all equal 1. Rather, for this to be the case we must choose the weights $\{\lambda_f\}_{f \in F}$ so as to have

$$\sum_{f \in F: B \cup \{f\} \in \mathcal{B}(M)} \lambda_f = 1 \quad (4.13)$$

for all $B \in \mathcal{B}(\text{tr}_F(M))$. Since $\mathcal{B}(\text{tr}_F(M))$ is generally much larger than F , this system of equations is usually overdetermined. However, if this system does have a nonnegative solution $\{\lambda_f\}_{f \in F}$, then for this choice of λ it follows that $\text{tr}_\lambda P$ is the basis generating polynomial for $\text{tr}_F(M)$, and that $\text{ext}_\lambda P$ is the basis generating polynomial for $M +_F a$. When this happens we say that the principal truncation (or extension) of M by F is *nice*.

Proposition 4.13 *For every $1 \leq r \leq n$, the uniform matroid $U_{r-1,n}$ is a nice principal truncation of $U_{r,n}$. Therefore, for all $0 \leq r \leq n$, the basis generating polynomial of $U_{r,n}$ has the half-plane property.*

PROOF. Fix n , and let $E = \{1, \dots, n\}$ be the common ground set of all the matroids $U_{r,n}$. We have $U_{r-1,n} = \text{tr}_E(U_{r,n})$. Setting $\lambda_e = 1/(n-r+1)$ for all $e \in E$ gives a solution to the system (4.13), so $U_{r-1,n}$ is a nice principal truncation of $U_{r,n}$. The basis generating polynomial of $U_{n,n}$ is x^E , which clearly has the half-plane property. Reverse induction on r (from n down to 0) using Proposition 4.11 now shows that the basis generating polynomial of $U_{r,n}$ has the half-plane property for all $0 \leq r \leq n$. \square

Here are a few more examples:

Example 4.1. The matroid $M(K_4)^+$ shown in Figure 5 (see Appendix A) is the principal extension of the graphic matroid $M(K_4)$ obtained by adding one point freely to one of the 3-point lines of $M(K_4)$, say 123. An easy computation shows that this principal extension is nice: it suffices to take $\lambda_1 = \lambda_2 = \lambda_3 = 1/2$ and $\lambda_4 = \lambda_5 = \lambda_6 = 0$. It follows (by Theorem 1.1 and Proposition 4.11) that $M(K_4)^+$ has the half-plane property. \square

Example 4.2. The matroid $M(K_4)+e$ (Figure 5) is the free extension of the graphic matroid $M(K_4)$, i.e. the principal extension of $M(K_4)$ obtained by adding one point freely to the flat $F = E$. A straightforward computation shows that this principal extension is not nice: the system (4.13) has no solution.

Let us remark that $M(K_4) + e$ fails the half-plane property (Example 11.4). This shows that non-nice principal extension does *not* in general preserve the half-plane property. \square

Example 4.3. The matroid $F_7^{-4} + e$ (Figure 8) is the free extension of the matroid F_7^{-4} (Figure 6), i.e. the principal extension of F_7^{-4} obtained by adding one point freely to the flat $F = E$. An easy computation shows that this free extension is nice: it suffices to take $\lambda_1 = 0$ and $\lambda_2 = \dots = \lambda_7 = 1/4$. It follows that $F_7^{-4} + e$ has the half-plane property if F_7^{-4} does. Unfortunately, we have been unable to prove whether or not F_7^{-4} has the half-plane property; but our numerical tests (Section 12) suggest that it does (and that $F_7^{-4} + e$ does also). \square

Example 4.4. The matroid Q_7 (Figure 5) is the free extension of the matroid $Q_7 \setminus 7$. Curiously, the system (4.13) does have a solution, but this (unique) solution fails to be nonnegative: it is $\lambda_1 = -1/6$, $\lambda_2 = \lambda_3 = 1/2$, $\lambda_4 = \lambda_5 = \lambda_6 = 1/3$. So this free extension is not nice. Nevertheless, it turns out that Q_7 has the half-plane property (Corollary 10.3 and Example 10.7). \square

We have written a MATHEMATICA program `niceprincipal.m` to test a principal extension for niceness by solving the linear system (4.13); it is available as part of the electronic version of this paper at arXiv.org.

4.6 Principal cotruncation and principal coextension

Let P be a *multiaffine* polynomial with ground set E , and let $\{\lambda_e\}_{e \in E}$ be nonnegative real numbers. Then the polynomial

$$\text{cotr}_\lambda P(x) = \sum_{e \in E} \lambda_e x_e P^{\setminus e}(x_{\neq e}) \quad (4.14)$$

is called a (*weighted*) *principal cotruncation* of P . If a is a new element (not occurring in E), then

$$\text{coext}_\lambda P(x_{\cup a}) = \text{cotr}_\lambda P(x) + x_a P(x) \quad (4.15)$$

is called a (*weighted*) *principal coextension* of P .

Note that by Lemma 4.3, we have

$$(\text{cotr}_\lambda P)^* = \sum_{e \in E} \lambda_e (P^*)^{/e}(x_{\neq e}) = \text{tr}_\lambda P^* \quad (4.16)$$

(to understand the first equality, observe that $P^{\setminus e}$ is considered as a polynomial on $E \setminus e$, while the rest are considered as polynomials on E). Likewise,

$$(\text{coext}_\lambda P)^* = x_a (\text{cotr}_\lambda P)^* + P^* = x_a \text{tr}_\lambda P^* + P^* = \text{ext}_\lambda P^* \quad (4.17)$$

(taking care again to notice that some of these are polynomials on $E \cup a$ and others on E). So principal cotruncation and coextension are indeed the duals of principal truncation and extension.

Proposition 4.14 (Principal cotruncation/coextension) *Let P be a multiaffine polynomial with ground set E , and let $\{\lambda_e\}_{e \in E}$ be nonnegative real numbers. If P has the half-plane property, then both $\text{cotr}_\lambda P$ and $\text{coext}_\lambda P$ have the half-plane property.*

PROOF. The result is trivial if $P \equiv 0$, so assume $P \not\equiv 0$. We have

$$\frac{\text{cotr}_\lambda P(x)}{P(x)} = \sum_{e \in E} \lambda_e \frac{x_e P^{\setminus e}(x_{\neq e})}{P(x)} \quad (4.18a)$$

$$= \sum_{e \in E} \lambda_e \frac{x_e P^{\setminus e}(x_{\neq e})}{P^{\setminus e}(x_{\neq e}) + x_e P^{/e}(x_{\neq e})} \quad (4.18b)$$

$$= \sum_{e \in E} \lambda_e \left(\frac{1}{x_e} + \frac{P^{/e}(x_{\neq e})}{P^{\setminus e}(x_{\neq e})} \right)^{-1}, \quad (4.18c)$$

which is real-part-positive by Theorem 3.2. [Use (4.18b) in place of (4.18c) for those elements e having $P^{\setminus e} \equiv 0$ or $P^{/e} \equiv 0$.] So $\text{cotr}_\lambda P$ has the half-plane property. Moreover,

$$\frac{\text{coext}_\lambda P(x)}{P(x)} = \frac{\text{cotr}_\lambda P(x)}{P(x)} + x_a \quad (4.19)$$

is real-part-positive, so $\text{coext}_\lambda P$ has the half-plane property. \square

The principal cotruncation of a matroid M on a set D , which appears in the next proposition, is also called the principal lift of M on D [14, pp. 160–161].

Proposition 4.15 *Let P be a homogeneous multiaffine polynomial with ground set E , and let $\{\lambda_e\}_{e \in E}$ be nonnegative real numbers, not all of which are zero. Suppose that $\text{supp}(P) = \mathcal{B}(M)$ for a matroid M of rank less than $|E|$, let $S = \{e: \lambda_e > 0\}$, and let D be the subset of E such that $E \setminus D$ is the union of all circuits contained in $E \setminus S$. Then $\text{supp}(\text{cotr}_\lambda P) = \mathcal{B}(\text{cotr}_D(M))$, where $\text{cotr}_D(M)$ is the principal cotruncation of M on D . Also, $\text{supp}(\text{coext}_\lambda P) = \mathcal{B}(M \times_D a)$, where $M \times_D a$ is the principal coextension of M on D .*

PROOF. The bases of the cotruncation $\text{cotr}_D(M)$ are all the sets of the form $B \cup d$ where $B \in \mathcal{B}(M)$ and $d \in D \setminus B$. The sets that appear in the support of $\text{cotr}_\lambda P$ are all the sets of the form $B \cup f$ where $B \in \mathcal{B}(M)$ and $f \in S \setminus B$. Since $S \subseteq D$, it follows that $\text{supp}(\text{cotr}_\lambda P) \subseteq \mathcal{B}(\text{cotr}_D(M))$.

Now suppose that $B \in \mathcal{B}(M)$ and $f \in D \setminus B$. If $f \in S$, then $B \cup f \in \mathcal{B}(\text{cotr}_D(M))$. Now suppose that $f \notin S$. Clearly $B \cup f$ contains a circuit C of M and $f \in C$. If C avoids S , then C is a circuit contained in $E \setminus S$, so $C \subseteq E \setminus D$, which is a contradiction. Thus C contains an element g of S . Then $(B \setminus g) \cup f \in \mathcal{B}(M)$ and $(B \setminus g) \cup f \cup g = B \cup f \in \mathcal{B}(\text{cotr}_D(M))$. Hence $\text{supp}(\text{cotr}_\lambda P) \supseteq \mathcal{B}(\text{cotr}_D(M))$ and so equality holds.

The bases of the principal coextension $M \times_D a$ are given by

$$\mathcal{B}(M \times_D a) = \{B \cup a : B \in \mathcal{B}(M)\} \cup \mathcal{B}(\text{cotr}_D(M)). \quad (4.20)$$

From this and the previous paragraph, the claim about the support of $\text{coext}_\lambda P$ follows readily. \square

Note that even if P is the basis generating polynomial of the matroid M , it need not be the case that $\text{cotr}_\lambda P$ is the basis generating polynomial of $\text{cotr}_D(M)$, since the coefficients need not all equal 1. Rather, for this to be the case we must choose the weights $\{\lambda_d\}_{d \in D}$ so as to have

$$\sum_{d \in D : B \setminus d \in \mathcal{B}(M)} \lambda_d = 1 \quad (4.21)$$

for all $B \in \mathcal{B}(\text{cotr}_D(M))$. When this system of equations has a nonnegative solution $\{\lambda_d\}_{d \in D}$, we say that the principal cotruncation (or coextension) of M by D is *nice*.

Proposition 4.16 *For every $1 \leq r \leq n$, the uniform matroid $U_{r,n}$ is a nice principal cotruncation of $U_{r-1,n}$. Therefore, for all $0 \leq r \leq n$, the basis generating polynomial of $U_{r,n}$ has the half-plane property.*

PROOF. Fix n , and let $E = \{1, \dots, n\}$ be the common ground set of all the matroids $U_{r,n}$. We have $U_{r,n} = \text{cotr}_E(U_{r-1,n})$. Setting $\lambda_e = 1/r$ for all $e \in E$ gives a solution to the system (4.21), so $U_{r,n}$ is a nice principal cotruncation of $U_{r-1,n}$. The basis generating polynomial of $U_{0,n}$ is 1, which clearly has the half-plane property. Induction on r using Proposition 4.14 shows that the basis generating polynomial of $U_{r,n}$ has the half-plane property for all $0 \leq r \leq n$. \square

Remark. The construction in this section was inspired by our analysis of Fettiweis' [25] proof of the half-plane property for the uniform matroids $U_{r,n}$ (see Section 9).

4.7 Multiaffine part and full-rank matroid union

Let $P(x) = \sum_{\mathbf{m}} a_{\mathbf{m}} x^{\mathbf{m}}$ be a polynomial with ground set E . For any subset $A \subseteq E$, define

$$P^{bA}(x) = \sum_{\mathbf{m} : m_e \leq 1 \forall e \in A} a_{\mathbf{m}} x^{\mathbf{m}}. \quad (4.22)$$

When $A = E$ we shall write simply P^b , and shall call it the *multiaffine part* of P . Clearly the map $P \mapsto P^{bA}$ is linear and idempotent.

Proposition 4.17 *If P has the half-plane property, then for any $A \subseteq E$, P^{bA} has the half-plane property.*

PROOF. Consider first the univariate case $|E| = 1$ with $A = E$. The case $P \equiv 0$ is trivial, so let $P \not\equiv 0$ be a univariate polynomial of degree k with the half-plane property: we have $P(x) = C \prod_{i=1}^k (x + \alpha_i)$ with $C \neq 0$ and $\operatorname{Re} \alpha_i \geq 0$. The cases $k = 0, 1$ are trivial, so assume $k \geq 2$. Then $P^b(x) = a_0 + a_1 x$ where

$$a_0 = C \prod_{i=1}^k \alpha_i \quad (4.23a)$$

$$a_1 = C \sum_{j=1}^k \prod_{i \neq j} \alpha_i \quad (4.23b)$$

If one or more of the α_i are zero, then $a_0 = 0$ and P^b has the half-plane property. Otherwise,

$$\frac{a_1}{a_0} = \sum_{j=1}^k \frac{1}{\alpha_j} \quad (4.24)$$

has nonnegative real part, so that P^b again has the half-plane property.

Consider next the multivariate case with $|A| = 1$, i.e. $A = \{a\}$. Let us write $P(x) = \sum_k P_k(x_{\neq a}) x_a^k$, so that $P^{ba}(x) = P_0(x_{\neq a}) + P_1(x_{\neq a}) x_a$. Applying the univariate case to the variable x_a , we conclude that for each $x_{\neq a} \in H^{E \setminus a}$, the univariate polynomial $P^{ba}(\cdot, x_{\neq a})$ is either identically zero or else nonvanishing on H . By Corollary 2.10, both P_0 and P_1 have the half-plane property, hence each one is either identically zero or else nonvanishing on $H^{E \setminus a}$. If both P_0 and P_1 are identically zero, then P^{ba} is identically zero and hence has the half-plane property. If one or both of them is nonvanishing on $H^{E \setminus a}$, it follows that $P^{ba}(\cdot, x_{\neq a})$ cannot be identically zero for any $x_{\neq a} \in H^{E \setminus a}$, so P^{ba} must be nonvanishing on H^E .

The general case with $|A| = \{a_1, \dots, a_N\}$ is now obtained by successively applying the case $|A| = 1$ to each a_i . \square

Remark. Note that P^b can be identically zero even if P is not identically zero (and has the half-plane property), e.g. $P(x) = x^2$.

For matroids M and N , the *union* $M \vee N$ is the matroid on ground set $E(M) \cup E(N)$ that has as its collection of independent sets all sets of the form $I_1 \cup I_2$, with I_1 independent in M and I_2 independent in N . The bases of $M \vee N$ are its maximal independent sets, as always. It is easy to see that $\operatorname{rank}(M \vee N) = \operatorname{rank}(M) + \operatorname{rank}(N)$ holds if and only if there exist $B_1 \in \mathcal{B}(M)$ and $B_2 \in \mathcal{B}(N)$ with $B_1 \cap B_2 = \emptyset$; and in this case the bases of $M \vee N$ are precisely the sets $B_1 \cup B_2$ with $B_1 \in \mathcal{B}(M)$, $B_2 \in \mathcal{B}(N)$ and $B_1 \cap B_2 = \emptyset$. When $\operatorname{rank}(M \vee N) = \operatorname{rank}(M) + \operatorname{rank}(N)$ holds, we call this *full-rank matroid union*.

Proposition 4.18 (Matroid union) *Let P and Q be homogeneous multiaffine polynomials with the same-phase property such that $\operatorname{supp}(P) = \mathcal{B}(M)$ and $\operatorname{supp}(Q) = \mathcal{B}(N)$ for matroids M and N .*

(a) If $\text{rank}(M \vee N) = \text{rank}(M) + \text{rank}(N)$, then $\text{supp}((PQ)^{\flat}) = \mathcal{B}(M \vee N)$.

(b) If $\text{rank}(M \vee N) < \text{rank}(M) + \text{rank}(N)$, then $(PQ)^{\flat} \equiv 0$.

PROOF. If $P(x) = \sum_S a_S x^S$ and $Q(x) = \sum_S b_S x^S$, then $(PQ)^{\flat}(x) = \sum_S c_S x^S$ where

$$c_S = \sum_{\substack{T \cup U = S \\ T \cap U = \emptyset}} a_T b_U. \quad (4.25)$$

If $\text{rank}(M \vee N) < \text{rank}(M) + \text{rank}(N)$, then there are no pairs $T \in \text{supp}(P)$ and $U \in \text{supp}(Q)$ with $T \cap U = \emptyset$, so $(PQ)^{\flat} \equiv 0$. If $\text{rank}(M \vee N) = \text{rank}(M) + \text{rank}(N)$, then the sets S for which (4.25) contains at least one nonzero summand are precisely the bases of $M \vee N$; and the same-phase property for P and Q ensures that all the contributing terms $a_T b_U$ have the same phase, so that $c_S \neq 0$. \square

Note, however, that even if P and Q are the basis generating polynomials of M and N , it need not be the case that $(PQ)^{\flat}$ is a multiple of the basis generating polynomial of $M \vee N$, since the coefficients need not all be equal. For example, let M and N be matroids on the set $\{1, 2, 3, 4\}$ with basis generating polynomials $P(x) = x_1 + x_2 + x_3$ and $Q(x) = x_1 + x_2 + x_4$. Then $(PQ)^{\flat}(x) = 2x_1x_2 + x_1x_3 + x_1x_4 + x_2x_3 + x_2x_4 + x_3x_4$.

4.8 “Folding mod 2” and convolution

Let $P(x) = \sum_{\mathbf{m}} a_{\mathbf{m}} x^{\mathbf{m}}$ be a polynomial with ground set E . For any subset $A \subseteq E$, define

$$P^{\natural A}(x) = \sum_{\mathbf{m}} a_{\mathbf{m}} \prod_{e \in A} x_e^{m_e \bmod 2} \prod_{e \in E \setminus A} x_e^{m_e} \quad (4.26)$$

where $m_e \bmod 2 = 0$ or 1 . When $A = E$ we shall write simply P^{\natural} ; note that P^{\natural} is multiaffine. Clearly the map $P \mapsto P^{\natural A}$ is linear and satisfies $(P_1 \cdots P_k)^{\natural A} = (P_1^{\natural A} \cdots P_k^{\natural A})^{\natural A}$.

Proposition 4.19 *If P has the half-plane property and is not identically zero, then for any $A \subseteq E$, $P^{\natural A}$ has the half-plane property and is not identically zero.*

PROOF. Consider first the univariate case $|E| = 1$. Let $P \neq 0$ be a univariate polynomial of degree k with the half-plane property, so that $P(x) = C \prod_{i=1}^k (x + \alpha_i)$ with $C \neq 0$ and $\text{Re } \alpha_i \geq 0$. The cases $k = 0, 1$ are trivial, so assume $k \geq 2$. By repeated use of the identity $(P_1 P_2)^{\natural} = (P_1^{\natural} P_2^{\natural})^{\natural}$ [where P_1 is a linear factor and P_2 is the product of the remaining factors], it suffices to prove the case $k = 2$. In this case we have $P^{\natural}(x) = C[(1 + \alpha_1 \alpha_2) + (\alpha_1 + \alpha_2)x]$. It is easy to see that $1 + \alpha_1 \alpha_2$ and

$\alpha_1 + \alpha_2$ cannot both be zero, so $P^\natural \not\equiv 0$. If one of them is zero, then P^\natural clearly has the half-plane property. If both of them are nonzero, then

$$\frac{1 + \alpha_1 \alpha_2}{\alpha_1 + \alpha_2} = \frac{1}{\alpha_1 + \alpha_2} + \left(\frac{1}{\alpha_1} + \frac{1}{\alpha_2} \right)^{-1} \quad (4.27)$$

has nonnegative real part [this happens also if one of the α_i is zero], so P^\natural again has the half-plane property.

Consider now the multivariate case, i.e. suppose that $P(x) \neq 0$ whenever $\operatorname{Re} x_e > 0$ for all $e \in E$. Let $A = \{a_1, \dots, a_N\}$ and define $A_j = \{a_1, \dots, a_j\}$ for $1 \leq j \leq N$. Applying the univariate case successively to each x_{a_i} (with all other x_e held fixed in the open right half-plane), we produce a sequence of polynomials $P^{\natural A_1}, P^{\natural A_2}, \dots, P^{\natural A_N} \equiv P^{\natural A}$, each of which has the half-plane property and is not identically zero. \square

Remarks. 1. There obviously exist polynomials $P \not\equiv 0$ for which $P^\natural \equiv 0$, e.g. $P(x) = 1 - x^2$. Proposition 4.19 asserts that this cannot happen when P has the half-plane property.

2. Suppose that in the definition of $P^{\natural A}$ we replace “mod 2” by “mod n ”. Then the map $P \mapsto P^{\natural A}$ is still linear and satisfies $(P_1 \cdots P_k)^{\natural A} = (P_1^{\natural A} \cdots P_k^{\natural A})^{\natural A}$, but for $n \geq 3$ it does not preserve the half-plane property. A univariate counterexample is $P(x) = (1 + i\epsilon x)^n$ with ϵ real, for which $P^\natural(x) = (1 + i\epsilon x)^n - (i\epsilon)^n(x^n - 1)$. For small $\epsilon \neq 0$, the roots of P^\natural are $x = \epsilon^{-1}i/(1 - \omega) + O(\epsilon^{n-1})$ where $\omega \neq 1$ is an n th root of unity; and for $n \geq 3$ at least one of these roots has strictly positive real part.

3. If $P(z_1, z_2) = a + bz_1 + cz_2 + dz_1z_2$, the *Asano contraction* of P is the univariate polynomial $\tilde{P}(z) = a + dz$. The Asano contraction lemma [3, 63, 37] states that if P is nonvanishing in the unit bidisc $|z_i| < 1$ ($i = 1, 2$), then \tilde{P} is nonvanishing in the unit disc $|z| < 1$; this lemma and its generalizations play an important role in the derivation of Lee–Yang theorems in statistical mechanics [63, 64, 47, 37]. The Asano lemma can be obtained as a corollary of Proposition 4.19, using the Möbius transformation

$$z_i = \frac{1 - x_i}{1 + x_i} \iff x_i = \frac{1 - z_i}{1 + z_i} \quad (4.28)$$

that maps the right half-plane $\operatorname{Re} x_i > 0$ onto the unit polydisc $|z_i| < 1$. It suffices to note that

$$\begin{aligned} Q(x_1, x_2) &\equiv (1 + x_1)(1 + x_2)P\left(\frac{1 - x_1}{1 + x_1}, \frac{1 - x_2}{1 + x_2}\right) \\ &= (a + b + c + d) + (a - b + c - d)x_1 + (a + b - c - d)x_2 + (a - b - c + d)x_1x_2 \\ &\equiv \alpha + \beta x_1 + \gamma x_2 + \delta x_1x_2 \end{aligned} \quad (4.29)$$

while

$$R(x) \equiv (1 + x)\tilde{P}\left(\frac{1 - x}{1 + x}\right) = (a + d) + (a - d)x = \frac{\alpha + \delta}{2} + \frac{\beta + \gamma}{2}x, \quad (4.30)$$

so that $2R$ can be obtained from Q by setting $x_1 = x_2 = x$ and applying the \natural operation.

Now let $P(x) = \sum_S a_S x^S$ and $Q(x) = \sum_S b_S x^S$ be multiaffine polynomials on the same ground set E . The *convolution* $P * Q$ of P and Q is defined to be

$$(P * Q)(x) = \sum_{S, T} a_S b_T x^{S \Delta T}, \quad (4.31)$$

where Δ denotes the symmetric difference of sets.⁹ It is easily seen that convolution is a bilinear, commutative and associative operation, and that $P_1 * \dots * P_k = (P_1 \dots P_k)^\natural$. It therefore follows immediately from Propositions 4.4 and 4.19 that:

Proposition 4.20 (Convolution) *Let P and Q be multiaffine polynomials with the half-plane property. Then $P * Q$ also has the half-plane property. Moreover, if P and Q are not identically zero, then $P * Q$ is not identically zero.*

Remarks. 1. The ring of multiaffine polynomials with respect to convolution does have divisors of zero, e.g. $P(x) = 1 + x$ and $Q(x) = 1 - x$ have $P * Q \equiv 0$. But Proposition 4.20 asserts that this cannot happen when P and Q both have the half-plane property.

2. Proposition 4.2 (duality) is a special case of Proposition 4.20, obtained by taking $Q(x) = x^E$.

3. Proposition 4.20 is closely related to Hinkkanen's composition theorem [37], which states that if $P(z) = \sum_S a_S z^S$ and $Q(z) = \sum_S b_S z^S$ are multiaffine polynomials (on the same ground set E) that are nonvanishing in the unit polydisc, then the *Schur–Hadamard product*

$$(P \circ Q)(z) = \sum_S a_S b_S z^S \quad (4.32)$$

is either nonvanishing in the unit polydisc or else identically zero. Indeed, Proposition 4.20 and Hinkkanen's composition theorem are interderivable, using the Möbius transformation (4.28) that maps the right half-plane $\operatorname{Re} x_i > 0$ onto the unit polydisc $|z_i| < 1$. To see this, note first that, given a polynomial $P(x) = \sum_S a_S x^S$, we can define a polynomial

$$\widehat{P}(z) = (1 + z)^E P\left(\frac{1 - z}{1 + z}\right) = \sum_T \widehat{a}_T z^T \quad (4.33)$$

where the coefficients $\{\widehat{a}_T\}$ are obtained from $\{a_S\}$ by Fourier transformation on the group 2^E :

$$\widehat{a}_T = \sum_{S \subseteq E} a_S (-1)^{|S \cap T|}. \quad (4.34)$$

⁹ Strictly speaking, it is the *coefficients* $a = \{a_S\}$ and $b = \{b_S\}$ — which are complex-valued functions on the group 2^E of subsets of E with respect to symmetric difference — that are being convoluted here.

Clearly, P has the half-plane property if and only if \widehat{P} has the “unit polydisc property”; and since Fourier transformation carries elementwise product onto convolution (and vice versa), we have $\widehat{P * Q} = \widehat{P} \circ \widehat{Q}$. In this way, Proposition 4.20 can be given an alternate proof as a corollary of Hinkkanen’s composition theorem; and, conversely, our proof of Proposition 4.20 yields an alternate proof of Hinkkanen’s composition theorem.

In Proposition 4.18 we showed that matroid union can be obtained via the \flat operation in case $\text{rank}(M \vee N) = \text{rank}(M) + \text{rank}(N)$. Let us now show that, in this same full-rank situation, matroid union can also be obtained via convolution (followed by taking the leading part):

Corollary 4.21 (Matroid union, revisited) *Let P and Q be homogeneous multiaffine polynomials with the same-phase property such that $\text{supp}(P) = \mathcal{B}(M)$ and $\text{supp}(Q) = \mathcal{B}(N)$ for matroids M and N . If $\text{rank}(M \vee N) = \text{rank}(M) + \text{rank}(N)$, then $(P * Q)^\sharp = (PQ)^\flat$ and hence $\text{supp}((P * Q)^\sharp) = \mathcal{B}(M \vee N)$.*

PROOF. As noted earlier, the equality $\text{rank}(M \vee N) = \text{rank}(M) + \text{rank}(N)$ implies that there exist pairs (S, T) with $S \in \mathcal{B}(M)$, $T \in \mathcal{B}(N)$ and $S \cap T = \emptyset$; moreover, the bases of $M \vee N$ are precisely the corresponding sets $S \cup T$. Now, for each $B \subseteq E$ of cardinality $\text{rank}(M \vee N)$, the coefficient c_B of B in $P * Q$ is the sum of $a_S b_T$ over all pairs $(S, T) \in \mathcal{B}(M) \times \mathcal{B}(N)$ for which $S \cup T = B$ (note that this implies $S \cap T = \emptyset$); and the same-phase property for P and Q ensures that all the terms $a_S b_T$ have the same phase, so that $c_B \neq 0$ whenever $B \in \mathcal{B}(M \vee N)$. It follows that $P * Q$ indeed has degree $\text{rank}(M \vee N)$ [rather than some lower degree], so that $(P * Q)^\sharp = (PQ)^\flat$. \square

5 Necessary and sufficient condition

5.1 General case

Let P be a polynomial in n complex variables with complex coefficients. Gregor [33, Lemma 6] has given a necessary and sufficient condition for P to have the half-plane property:

Proposition 5.1 (Gregor [33]) *Let P be a polynomial in n complex variables, and fix $k \geq \deg P$. For $x, y \in \mathbb{R}^n$, define the univariate polynomial $\widehat{p}_{x,y}(\zeta) = \zeta^k P(\zeta x + \zeta^{-1}y)$. Then the following are equivalent:*

- (a) *P has the half-plane property and is not identically zero.*
- (b) *For all $x, y \geq 0$ with $x + y > 0$, $\widehat{p}_{x,y}$ has the half-plane property and is not identically zero [that is, all the roots of $\widehat{p}_{x,y}$ lie in the closed left half-plane].*

(c) For all $x, y > 0$, $\widehat{p}_{x,y}$ has the half-plane property and is not identically zero.

PROOF. (a) \implies (b): Let $x, y \geq 0$ with $x+y > 0$. Then $\zeta \in H$ implies $\zeta x + \zeta^{-1}y \in H^n$, so $\widehat{p}_{x,y}(\zeta) \neq 0$.

(b) \implies (c) is trivial.

(c) \implies (a): Let $z = (z_1, \dots, z_n)$ with $\operatorname{Re} z_i > 0$ for all i . Choose $\zeta \in H$ in such a way that all the z_i lie in the interior of the convex cone generated by ζ and ζ^{-1} [i.e. choose ζ so that $\max |\arg z_i| < |\arg \zeta| < \pi/2$]; then there exist numbers $x_i, y_i > 0$ with $z_i = \zeta x_i + \zeta^{-1}y_i$. It follows that $P(z) = \zeta^{-k} \widehat{p}_{x,y}(\zeta) \neq 0$. \square

If P is homogeneous, this result can be simplified slightly:

Proposition 5.2 *Let P be a homogeneous degree- r polynomial in n complex variables. For $x, y \in \mathbb{R}^n$, define the univariate polynomial $p_{x,y}(\zeta) = P(\zeta x + y)$. Then the following are equivalent:*

(a) P has the half-plane property and is not identically zero.

(b) For all $x, y \geq 0$ with $x + y > 0$, all the roots (real or complex) of $p_{x,y}$ lie in $(-\infty, 0]$.

(c) For all $x, y > 0$, all the roots (real or complex) of $p_{x,y}$ lie in $(-\infty, 0]$.

PROOF. (a) \implies (b): Let $x, y \geq 0$ with $x+y > 0$. If $\zeta_0 \in \mathbb{C} \setminus (-\infty, 0]$, then we can find $\alpha, \beta \in H$ such that $\alpha/\beta = \zeta_0$. But this means that $p_{x,y}(\alpha/\beta) = \beta^{-r} P(\alpha x + \beta y) \neq 0$.

(b) \implies (c) is trivial.

(c) \implies (a): Let $z = (z_1, \dots, z_n)$ with $\operatorname{Re} z_i > 0$ for all i . Choose $\alpha, \beta \in H$ in such a way that all the z_i lie in the interior of the convex cone generated by α and β , i.e. such that there exist numbers $x_i, y_i > 0$ with $z_i = \alpha x_i + \beta y_i$. Then $P(z) = P(\alpha x + \beta y) = \beta^r p_{x,y}(\alpha/\beta) \neq 0$. \square

Remark. The necessary and sufficient conditions given in Propositions 5.1 and 5.2 are algorithmically testable in time $c_1^{c_2^n}$ using cylindrical algebraic decomposition, and in time c^n using more recent algorithms (see Section 2.6). But this computation seems thoroughly unfeasible in practice, at least at present.

5.2 Rank-2 case

The drawback of the necessary-and-sufficient characterization contained in Proposition 5.2 is, of course, that the sufficient condition (b) or (c) is not easy to verify, because of the universal quantification over $x, y \geq 0$. However, in the rank-2 case we can obtain an easily-checkable alternative condition. Let $A = \{a_{ij}\}_{i,j=1}^n$ be a symmetric matrix with nonnegative real entries, i.e. $a_{ij} = a_{ji} \geq 0$. (We shall see in

Theorem 6.1 that the restriction to nonnegative real entries is no loss of generality.) Define the homogeneous degree-2 polynomial in n variables,

$$P_A(z) = \frac{1}{2} \sum_{i,j=1}^n a_{ij} z_i z_j = \frac{1}{2} z^T A z. \quad (5.1)$$

Let $\lambda_1(A) \geq \lambda_2(A) \geq \dots \geq \lambda_n(A)$ be the eigenvalues of A . We then have:

Theorem 5.3 (due largely to [33, 4]) *Let A be a symmetric matrix with nonnegative real entries. Then the following are equivalent:*

- (a) *Either $P_A(H^n) = \{0\}$ (i.e. $P_A \equiv 0$) or else $P_A(H^n) = \mathbb{C} \setminus (-\infty, 0]$.*
- (b) *P_A has the half-plane property.*
- (c) *$\lambda_2(A) \leq 0$.*
- (d) *If $x, y \in \mathbb{R}^n$ with $y^T A y \geq 0$, then $(x^T A y)^2 \geq (x^T A x)(y^T A y)$.*
- (e) *If $x, y \in \mathbb{R}^n$ with $x, y \geq 0$, then $(x^T A y)^2 \geq (x^T A x)(y^T A y)$.*

PROOF. If $A = 0$, then (a)–(e) are all true; so let us consider the nontrivial case $A \neq 0$ [hence $\lambda_1(A) > 0$ because A is symmetric]. By the Perron–Frobenius theorem [27, p. 66], there exists an eigenvector $x^{(1)} \geq 0$ corresponding to the eigenvalue λ_1 . Now extend this to an orthonormal basis of real eigenvectors $x^{(1)}, \dots, x^{(n)}$ satisfying $Ax^{(k)} = \lambda_k x^{(k)}$.

(a) \implies (b) is trivial.

(b) \implies (c): Suppose that $\lambda_2(A) > 0$. Perturb $x^{(1)}, x^{(2)}$ slightly to get vectors $\tilde{x}^{(1)} > 0$ and $\tilde{x}^{(2)}$ satisfying $\tilde{x}^{(1)T} A \tilde{x}^{(2)} = 0$, $\tilde{x}^{(1)T} A \tilde{x}^{(1)} > 0$ and $\tilde{x}^{(2)T} A \tilde{x}^{(2)} > 0$.¹⁰ Consider the vector $z = \tilde{x}^{(1)} + i\alpha \tilde{x}^{(2)}$ where $\alpha \in \mathbb{R}$. Then

$$P_A(z) = \frac{1}{2} (\tilde{x}^{(1)T} A \tilde{x}^{(1)} - \alpha^2 \tilde{x}^{(2)T} A \tilde{x}^{(2)}), \quad (5.2)$$

which vanishes when $\alpha = \pm \sqrt{(\tilde{x}^{(1)T} A \tilde{x}^{(1)}) / (\tilde{x}^{(2)T} A \tilde{x}^{(2)})}$. So P_A does not have the half-plane property.

(c) \implies (d): If $x^T A x \leq 0$ or $y^T A y = 0$, the assertion is trivial; so we can assume that $x^T A x > 0$ and $y^T A y > 0$. We must have $x^{(1)} \cdot x \neq 0$, since otherwise [thanks to the negative-semidefiniteness of A on the orthogonal complement of $x^{(1)}$] we would have $x^T A x \leq 0$; and likewise we must have $x^{(1)} \cdot y \neq 0$. Now define

$$g(\alpha) = (x + \alpha y)^T A (x + \alpha y). \quad (5.3)$$

¹⁰ For example, let v be any vector with strictly positive components, and define $\tilde{x}^{(1)} = x^{(1)} + \epsilon v$, $\tilde{x}^{(2)} = x^{(2)} + \delta v$ with $\delta = -\epsilon(\tilde{x}^{(2)T} A v) / (\tilde{x}^{(1)T} A v + \epsilon v^T A v)$ and $\epsilon > 0$ sufficiently small.

We have $g(0) = x^T A x > 0$. On the other hand, for $\alpha = -(x^{(1)} \cdot x)/(x^{(1)} \cdot y)$ we have $x^{(1)} \cdot (x + \alpha y) = 0$ and hence $g(\alpha) \leq 0$ [again by the negative-semidefiniteness of A on $x^{(1)\perp}$]. So the quadratic equation $g(\alpha) = 0$ has a real solution, which implies that its discriminant is nonnegative, i.e. that $(x^T A y)^2 \geq (x^T A x)(y^T A y)$.

(d) \implies (e) is trivial.

(e) \implies (a): Suppose that $P_A(z) = -C \leq 0$ for some vector $z = (z_1, \dots, z_n)$ with $\operatorname{Re} z_i > 0$ for all i . Choose θ so that $|\arg z_i| < \theta < \pi/2$ for all i . Then all the z_i lie in the interior of the convex cone generated by $e^{i\theta}$ and $e^{-i\theta}$, i.e. there exist vectors $x, y > 0$ such that $z = e^{i\theta}x + e^{-i\theta}y$. Then

$$P_A(z) = \frac{1}{2}[(x^T A x)e^{2i\theta} + 2(x^T A y) + (y^T A y)e^{-2i\theta}] = -C. \quad (5.4)$$

Defining $\zeta = e^{2i\theta}$, it follows that the quadratic equation

$$(x^T A x)\zeta^2 + 2[(x^T A y) + C]\zeta + (y^T A y) = 0 \quad (5.5)$$

has a root $\zeta \in \mathbb{C} \setminus (-\infty, 0]$. Since all the coefficients of (5.5) are positive, this root ζ cannot be positive, so it must have a nonzero imaginary part. This means that the discriminant of (5.5) is negative, i.e. $[(x^T A y) + C]^2 < (x^T A x)(y^T A y)$. Since $x^T A y \geq 0$ and $C \geq 0$, it follows that $(x^T A y)^2 < (x^T A x)(y^T A y)$, contradicting hypothesis (e). Therefore $P_A(H^n) \subseteq \mathbb{C} \setminus (-\infty, 0]$.

On the other hand, $P_A(1, \dots, 1) > 0$ and $P_A(\lambda, \dots, \lambda) = \lambda^2 P_A(1, \dots, 1)$; so as λ runs over the open right half-plane, $P_A(\lambda, \dots, \lambda)$ runs over all of $\mathbb{C} \setminus (-\infty, 0]$. \square

Remarks. 1. The proof of (c) \implies (d) is taken from [4, Theorem 4.4.6], where it is also proven that (d) \implies (c). After completing this proof, we learned that the equivalence of (b), (c) and (e) had already been proven by Fiedler and Gregor [33, Theorems 5 and 8]. Numerous equivalent characterizations are given in [4, Theorem 4.4.6].

2. The reference in Theorem 5.3(c) to the eigenvalues of the matrix A (i.e. treating A as a *linear operator*) is in fact quite misleading. What we really have here is a *quadratic form* $Q(x) = \sum_{i,j=1}^n a_{ij}x_i x_j$ on a real vector space $V (= \mathbb{R}^n)$ equipped with a distinguished convex cone C (i.e. the vectors with nonnegative components). We then extend Q bilinearly to the complexification $V + iV$, and say that it has the “half-plane property” if it is either identically zero or else is nonvanishing on the set $C^\circ + iV$ (here C° denotes the interior of C). Condition (c) is the statement that the inertia (n_+, n_0, n_-) of the real quadratic form Q satisfies $n_+ \leq 1$.

Let us now specialize to the 0-1-valued case: we are given a 2-uniform set system \mathcal{S} on $[n]$ — that is, a simple graph G with vertex set $[n]$ and edge set \mathcal{S} — and we define

$$a_{ij} = a_{ji} = \begin{cases} 1 & \text{if } \{i, j\} \in \mathcal{S} \\ 0 & \text{if } \{i, j\} \notin \mathcal{S} \end{cases} \quad (5.6a)$$

$$a_{ii} = 0 \quad \text{for all } i \quad (5.6b)$$




Figure 1: Forbidden induced subgraphs in Corollary 5.4(d).

Thus A is simply the adjacency matrix of G , and P_A is the generating polynomial $P_{\mathcal{S}}$. Theorem 5.3 then becomes:

Corollary 5.4 *Let G be a simple graph on the vertex set $[n]$ with edge set \mathcal{S} , and let A be the corresponding adjacency matrix. Then the following are equivalent:*

- (a) P_A has the half-plane property.
- (b) $\lambda_2(A) \leq 0$.
- (c) G is a complete multipartite graph (possibly empty) plus possible isolated vertices.
- (d) G contains none of the graphs in Figure 1 as induced subgraphs.
- (e) \mathcal{S} is the set of bases for a rank-2 matroid on the ground set $[n]$.

PROOF. If $\mathcal{S} = \emptyset$ (i.e. $A = 0$), then (a)–(e) are all true; so let us consider the nontrivial case $\mathcal{S} \neq \emptyset$. The equivalence of (a) and (b) follows from Theorem 5.3. The equivalence of (b), (c) and (d) is a result of Smith [69, Theorem 1] (see also [22, Theorem 6.7] and [38, Theorem 2.1]).

Consider, finally, the basis exchange property for a matroid:

- (B2) If $B_1, B_2 \in \mathcal{S}$ and $i \in B_1 \setminus B_2$, then there exists $j \in B_2 \setminus B_1$ such that $(B_1 \setminus i) \cup j \in \mathcal{S}$.

For a rank-2 matroid, this property holds trivially when $|B_1 \cap B_2| = 1$ or 2 . When $B_1 \cap B_2 = \emptyset$ (so that $|B_1 \cup B_2| = 4$), straightforward checking of cases shows that property (B2) holds if and only if none of the configurations in Figure 1 occurs on $B_1 \cup B_2$. This proves the equivalence of (d) and (e). \square

Corollary 5.5 *All rank-2 matroids have the half-plane property, as do all corank-2 matroids.*

PROOF. This is an immediate consequence of Corollary 5.4(e) \implies (a) and Proposition 4.2. \square

6 Necessary conditions I: Same-phase property

In this section and the next, we derive some *necessary* conditions that a homogeneous polynomial with the half-plane property must satisfy.

Let us begin with a bit of motivation: Many “counting” results in combinatorics find a more natural context when they are generalized to allow nonnegative real weights, rather than restricting all weights to be 0 or 1. For example, the max-flow-min-cut theorem arises by generalizing Menger’s theorem in this way. Now, in *some* cases it is natural to go further and allow complex weights, but in some cases it is not. For example, in the Heilmann–Lieb theorem (see Section 10.1 below), the complex vertex weights $\{x_i\}_{i \in V}$ are the main point, but it is essential that the edge weights $\{\lambda_e\}_{e \in E}$ be nonnegative. Roughly speaking, it seems natural that *variables* in multivariate polynomials be promoted to complex variables, but it is often necessary that the *coefficients* in the polynomial be nonnegative real numbers.¹¹

With this background in mind, we shall prove that if a *homogeneous* polynomial $P(x) = \sum_{\mathbf{m}} a_{\mathbf{m}} x^{\mathbf{m}}$ has the half-plane property, then *necessarily* all the coefficients $a_{\mathbf{m}}$ are nonnegative modulo an overall multiplicative phase factor.

Let us say that a pair of nonzero complex numbers a, b *have the same phase* if $a/b \in (0, \infty)$. Obviously, having the same phase is an equivalence relation on $\mathbb{C} \setminus 0$; and a collection $\{a_\alpha\}$ of nonzero complex numbers all have the same phase if and only if there exists $\theta \in \mathbb{R}$ such that $e^{-i\theta} a_\alpha > 0$ for all α . We say that a polynomial $P(x) = \sum_{\mathbf{m}} a_{\mathbf{m}} x^{\mathbf{m}}$ has the *same-phase property* if all the nonzero coefficients $a_{\mathbf{m}}$ have the same phase.

Theorem 6.1 *Let $P(x) = \sum_{\mathbf{m}} a_{\mathbf{m}} x^{\mathbf{m}}$ be a polynomial in n complex variables that is homogeneous of degree r . If P has the half-plane property, then all the nonzero coefficients $a_{\mathbf{m}}$ have the same phase.*

PROOF. The proof is by induction on n . If $n = 1$, we must have $P(x) = a_r x^r$, so the theorem is trivial. So assume $n \geq 2$. Let M ($\leq r$) be the degree of x_n in $P(x)$, and let us write

$$P(x) = \sum_{k=0}^M x_n^k P_k(x_1, \dots, x_{n-1}) \quad (6.1)$$

where of course

$$P_k(x_1, \dots, x_{n-1}) = \frac{1}{k!} (\partial/\partial x_n)^k P(x) \Big|_{x_n=0}. \quad (6.2)$$

Clearly each coefficient of P corresponds to exactly one coefficient of exactly one P_k , and vice versa. Now, each P_k is a homogeneous polynomial (of degree $r - k$) in $n - 1$

¹¹ Of course, given a polynomial expression, it is not always obvious which objects should be considered to be “variables” and which should be absorbed into the “coefficients”. So this admittedly vague principle may in some cases be best understood backwards, i.e. as an injunction to treat as “variables” those objects that can naturally be made complex, and as “coefficients” those that cannot.

variables; and by Corollary 2.10 it has the half-plane property. So, by the inductive hypothesis, all its nonzero coefficients have the same phase.

To complete the proof, we need only show that these phases are the same for all k . Applying Proposition 5.2 with $x = (0, \dots, 0, 1)$ and $y = (1, \dots, 1, 0)$, we get

$$p_{x,y}(\zeta) = P(1, \dots, 1, \zeta) = \sum_{k=0}^M P_k(1, \dots, 1) \zeta^k. \quad (6.3)$$

Since $p_{x,y}$ has only real nonpositive zeros, it can be written as $p_{x,y}(\zeta) = a \prod_{k=1}^M (\zeta + c_k)$ for some complex number a and nonnegative real numbers c_1, c_2, \dots, c_M ; so its coefficients $P_k(1, \dots, 1)$ must all have the same phase. \square

Remark. It is easy to see that if P is homogeneous of degree 1, then P has the half-plane property *if and only if* all its nonzero coefficients have the same phase.

We can generalize Theorem 6.1 to a class of not-necessarily-homogeneous polynomials. Let us say that a polynomial $P(x) = \sum_{\mathbf{m}} a_{\mathbf{m}} x^{\mathbf{m}}$ has *definite parity* if all the nonzero monomials have total degree of the same parity (i.e. $a_{\mathbf{m}}, a_{\mathbf{m}'} \neq 0$ implies that $|\mathbf{m}| \equiv |\mathbf{m}'| \pmod{2}$); this is equivalent to saying that the polynomial P is either even or odd. We then have:

Theorem 6.2 *Let $P(x) = \sum_{\mathbf{m}} a_{\mathbf{m}} x^{\mathbf{m}}$ be a polynomial in n complex variables that has definite parity and the half-plane property. Then all the nonzero coefficients $a_{\mathbf{m}}$ have the same phase.*

We begin the proof of Theorem 6.2 with a lemma:

Lemma 6.3 *Let P and Q be polynomials in n complex variables, and suppose that $P + cQ$ has the same-phase property for all real $c > 0$. Then either*

(a) *all the nonzero coefficients of P and Q have the same phase*

or else

(b) *P and Q each have the same-phase property, and $P = \alpha Q$ for some nonzero complex number α .*

PROOF. Letting $c \rightarrow 0$ (resp. $c \rightarrow +\infty$), we deduce that P (resp. Q) has the same-phase property. Let us write $P(x) = \sum_{\mathbf{m}} a_{\mathbf{m}} x^{\mathbf{m}}$ and $Q(x) = \sum_{\mathbf{m}} b_{\mathbf{m}} x^{\mathbf{m}}$. Now there are two cases:

- 1) There is a multi-index \mathbf{m} such that both $a_{\mathbf{m}}$ and $b_{\mathbf{m}}$ are nonzero.
- 2) There is no such multi-index \mathbf{m} .

In case 2, the fact that $P+Q$ has the same-phase property implies that all the nonzero coefficients of P and Q have the same phase.

In case 1, choose a multi-index \mathbf{m} such that both $a_{\mathbf{m}}$ and $b_{\mathbf{m}}$ are nonzero, and define $r_{\mathbf{m}} = a_{\mathbf{m}}/b_{\mathbf{m}}$. If there is no multi-index $\mathbf{n} \neq \mathbf{m}$ with either $a_{\mathbf{n}}$ or $b_{\mathbf{n}}$ nonzero, then $P(x) = a_{\mathbf{m}}x^{\mathbf{m}}$ and $Q(x) = b_{\mathbf{m}}x^{\mathbf{m}}$, which proves assertion (b) [with $\alpha = r_{\mathbf{m}}$]. So assume that there is at least one such multi-index \mathbf{n} , and consider an arbitrary one of them. Then, for all positive real numbers c , the complex numbers $a_{\mathbf{m}} + cb_{\mathbf{m}}$ and $a_{\mathbf{n}} + cb_{\mathbf{n}}$ have the same phase whenever they are both nonzero. If one of $a_{\mathbf{n}}$ and $b_{\mathbf{n}}$ is zero, then by taking $c \rightarrow 0$ or $c \rightarrow +\infty$ we can conclude (using the fact that P and Q have the same-phase property) that $a_{\mathbf{m}}$, $b_{\mathbf{m}}$ and the nonzero member of $\{a_{\mathbf{n}}, b_{\mathbf{n}}\}$ all have the same phase, which implies (again using the fact that P and Q have the same-phase property) that all the nonzero coefficients of P and Q have the same phase. If, on the other hand, neither $a_{\mathbf{n}}$ nor $b_{\mathbf{n}}$ is zero, let us define $r_{\mathbf{n}} = a_{\mathbf{n}}/b_{\mathbf{n}}$. Then $(r_{\mathbf{m}} + c)b_{\mathbf{m}}$ and $(r_{\mathbf{n}} + c)b_{\mathbf{n}}$ have the same phase for all positive real numbers $c \notin \{-r_{\mathbf{m}}, -r_{\mathbf{n}}\}$. Since $b_{\mathbf{m}}$ and $b_{\mathbf{n}}$ have the same phase, so do $r_{\mathbf{m}} + c$ and $r_{\mathbf{n}} + c$. Since this is valid for all positive real $c \notin \{-r_{\mathbf{m}}, -r_{\mathbf{n}}\}$, it follows that either $r_{\mathbf{m}}$ and $r_{\mathbf{n}}$ are both real and positive, or else that $r_{\mathbf{m}} = r_{\mathbf{n}}$. If the former case holds for at least one multi-index \mathbf{n} for which $a_{\mathbf{n}}$ and $b_{\mathbf{n}}$ are both nonzero, then all the nonzero coefficients of P and Q have the same phase. If the latter case holds for all \mathbf{n} for which at least one of $a_{\mathbf{n}}$ and $b_{\mathbf{n}}$ is nonzero, then $P = \alpha Q$ with $\alpha = r_{\mathbf{m}}$. \square

PROOF OF THEOREM 6.2. We proceed by induction on n . For $n = 1$, since P has definite parity, its zeros are symmetric with respect to the origin. Since P also has the half-plane property, these zeros all lie symmetrically on the imaginary axis, so that $P(x) = ax^m \prod_{i=1}^k (x^2 + c_i)$ for some complex number a , nonnegative integers m and k , and positive real numbers c_1, c_2, \dots, c_k . This implies the same-phase property for P .

For the induction step, suppose that $P(x_1, \dots, x_{n+1})$ has definite parity and the half-plane property. Let $P_j(x)$ be the coefficient of x_{n+1}^j in $P(x)$. Each P_j has definite parity (this parity alternates with j) and has the half-plane property (by Corollary 2.10); so by the inductive hypothesis each P_j has the same-phase property. We now want to show that the nonzero coefficients of the various P_j all have the same phase.

Define $j_{\min} = \min\{j: P_j \not\equiv 0\}$ and $j_{\max} = \max\{j: P_j \not\equiv 0\}$, and suppose that $j_{\min} \leq j < j_{\max}$. If $P_j \equiv 0$ (which of course can happen only for $j > j_{\min}$), then Theorem 3.4 states that P_{j+1}/P_{j-1} is a strictly positive constant, so that the nonzero coefficients of P_{j-1} and P_{j+1} have the same phase. If $P_j \not\equiv 0$ and $P_{j+1} \not\equiv 0$, then Theorem 3.4 states that P_{j+1}/P_j is real-part-positive on H^n . Thus, by applying Theorem 3.2 to the polynomial $P_j(x) + yP_{j+1}(x)$, where y is a new variable, we deduce that this polynomial has the half-plane property (in the variables x_1, \dots, x_n, y). It follows that, for any nonnegative real numbers c_1, \dots, c_n , the polynomial $P_j(x) + (c_1x_1 + \dots + c_nx_n)P_{j+1}(x)$ also has the half-plane property (in the variables x_1, \dots, x_n).

Since this latter polynomial has definite parity, by the inductive hypothesis it has the same-phase property. Now apply Lemma 6.3 with $P = P_j$ and $Q(x) = x_1 P_{j+1}(x)$. In case (a), we conclude that all the nonzero coefficients of P_j and P_{j+1} have the same phase. In case (b), we conclude that $P_j/P_{j+1} = \alpha x_1$ for some nonzero complex number α ; but since P_j/P_{j+1} is real-part-positive on H^n , α must be a positive real number, so we again conclude that all the nonzero coefficients of P_j and P_{j+1} have the same phase.

The foregoing observations show that the nonzero coefficients of all the P_j have the same phase, and hence that P has the same-phase property. This completes the induction step, and hence the proof. \square

Finally, let us take this opportunity to discuss the image of H^n under a homogeneous polynomial P .

Proposition 6.4 *Let $P \not\equiv 0$ be a homogeneous polynomial of degree $r \geq 1$ in n complex variables. If P does not have the half-plane property, then $P(H^n) = \mathbb{C}$. If P does have the half-plane property, then there exists $\theta \in \mathbb{R}$ such that*

$$P(H^n) = e^{i\theta} H^r = \begin{cases} e^{i\theta} H & \text{if } r = 1 \\ \mathbb{C} \setminus e^{i\theta}(-\infty, 0] & \text{if } r = 2 \\ \mathbb{C} \setminus 0 & \text{if } r \geq 3 \end{cases} \quad (6.4)$$

[We apologize for the inconsistent notations $H^n = H \times \dots \times H$ and $H^r = \{z^r : z \in H\}$. We trust that this will not cause any confusion.]

PROOF. Clearly P is nonconstant, so by the open mapping theorem, $P(H^n)$ is open. By homogeneity, $P(H^n)$ is a cone (i.e. invariant under multiplication by any $\lambda > 0$). If P does not have the half-plane property, then $P(H^n) \ni 0$; and an open cone containing 0 must be all of \mathbb{C} . Now suppose that P does have the half-plane property, so that $P(H^n) \not\ni 0$. Then $P(1, \dots, 1) \neq 0$, so let $\theta = \arg P(1, \dots, 1)$. Since $P(\zeta, \dots, \zeta) = \zeta^r P(1, \dots, 1)$, it follows that $P(\zeta, \dots, \zeta)$ covers the set $e^{i\theta} H^r$ as ζ ranges over H . Therefore, $P(H^n) \supseteq e^{i\theta} H^r$.

Let us now prove the reverse containment. Suppose first that $r = 1$ and $P(x) = \sum_{i=1}^n a_i x_i$. Then, as remarked above, P has the half-plane property if and only if all the nonzero a_i have the same phase, i.e. there exists $\theta \in \mathbb{R}$ such that $\operatorname{Re}(e^{-i\theta} a_i) \geq 0$ for all i . But in this case $P(H^n) = e^{i\theta} H$.

Suppose next that $r = 2$ and $P(x) = \sum_{i,j=1}^n a_{i,j} x_i x_j$. By Theorem 6.1, there exists $\theta \in \mathbb{R}$ such that $\operatorname{Re}(e^{-i\theta} a_{i,j}) \geq 0$ for all i, j . But then Theorem 5.3(b) \implies (a) tells us that $P(H^n) = \mathbb{C} \setminus e^{i\theta}(-\infty, 0]$.

If $r \geq 3$, the containment $P(H^n) \subseteq \mathbb{C} \setminus 0$ is trivial. \square

Corollary 6.5 *Fix $N \geq 2$. Let $\{E_i\}_{i=1}^N$ be disjoint finite sets, and for each i let P_i be a polynomial of degree r_i on the ground set E_i . [Note that P_i need not be homogeneous.] Suppose that $P = \sum_{i=1}^N P_i$, considered as a polynomial on the ground set $E = \bigcup_{i=1}^N E_i$, has the half-plane property. Then there is at most one index i with $r_i \geq 2$ and $P_i \neq 0$.*

PROOF. Suppose that there are two such indices i, j . By setting $x_e = 0$ for all $e \in E \setminus (E_i \cup E_j)$, we can assume without loss of generality that $N = 2$ and $E = E_1 \cup E_2$. Let us write $x^{(1)} = \{x_e\}_{e \in E_1}$ and $x^{(2)} = \{x_e\}_{e \in E_2}$, and let us define

$$Q_\lambda(x) = \lambda^{-1} P(\lambda^{1/r_1} x^{(1)}, \lambda^{1/r_2} x^{(2)}) . \quad (6.5)$$

Then Q_λ has the half-plane property for all $\lambda > 0$; and by Hurwitz's theorem, so does

$$\lim_{\lambda \rightarrow +\infty} Q_\lambda(x) = P_1^\#(x^{(1)}) + P_2^\#(x^{(2)}) \quad (6.6)$$

(recall from Section 2.2 that $^\#$ denotes leading part). So we can assume without loss of generality that P_1 and P_2 are homogeneous. Applying Proposition 6.4, we conclude that there exist $\theta_1, \theta_2 \in \mathbb{R}$ such that $P_i(H^{E_i}) \supseteq \mathbb{C} \setminus e^{i\theta_i}(-\infty, 0]$ for $i = 1, 2$. But then we can choose $x^{(1)} \in H^{E_1}$ and $x^{(2)} \in H^{E_2}$ so that $P_1(x^{(1)}) + P_2(x^{(2)}) = 0$, showing that P does not have the half-plane property. \square

Remark. We can also allow the sets E_i to be non-disjoint, provided that $\deg P_1, \deg P_2 \geq |E_1 \cap E_2| + 2$: it suffices to fix $x_e \in H$ for $e \in E_1 \cap E_2$ and then apply Corollary 6.5 in the remaining variables.

Let us conclude by mentioning an alternate proof of Theorem 6.1, which is more lengthy than the one given here but also more elementary (because it does not require Proposition 5.2). First one proves Theorem 6.1 in the multiaffine case by induction on r , deducing the same-phase property for P from that of its contractions $P^{/i}$; the key tool here (ensuring the necessary “connectedness”) is Corollary 6.5. Then one uses a Grace–Walsh–Szegő construction to reduce the case of a non-multiaffine polynomial to that of a multiaffine polynomial in a larger number of variables.

7 Necessary conditions II: Matroidal support

7.1 The exchangeability theorem

In this section we shall prove that if $P \neq 0$ is a homogeneous multiaffine polynomial with the half-plane property, then $\text{supp}(P)$ is necessarily the collection of bases of a matroid.

Theorem 7.1 *Let $P(x) = \sum_{S \subseteq E, |S|=r} a_S x^S$ be a multiaffine homogeneous degree- r polynomial (on the ground set E) that has the half-plane property and is not identically zero. Then there exists a rank- r matroid M on the ground set E such that $\text{supp}(P) = \mathcal{B}(M)$.*

When $\text{supp}(P) = \mathcal{B}(M)$, we call the (obviously unique) matroid M the *support matroid* of P .

Indeed, with only a little more work we can prove a generalization to homogeneous polynomials that are not necessarily multiaffine. First, a bit of notation: For multi-indices \mathbf{m} and \mathbf{p} , let $\mathbf{m} \wedge \mathbf{p}$ be the multi-index defined by $(\mathbf{m} \wedge \mathbf{p})(e) = \min[\mathbf{m}(e), \mathbf{p}(e)]$ for all $e \in E$. For $e \in E$, let us define the multi-index δ_e by

$$\delta_e(f) = \begin{cases} 1 & \text{if } f = e \\ 0 & \text{if } f \neq e \end{cases} \quad (7.1)$$

If \mathcal{S} is a set of multi-indices and $\mathbf{m}, \mathbf{p} \in \mathcal{S}$, we say that \mathbf{m} is *exchangeable towards* \mathbf{p} in \mathcal{S} (and write “ $\mathbf{m} \rightarrow \mathbf{p}$ in \mathcal{S} ”) to denote the following condition: for every $e \in E$ such that $\mathbf{m}(e) > \mathbf{p}(e)$, there exists an $f \in E$ such that $\mathbf{m}(f) < \mathbf{p}(f)$ and $\mathbf{m} - \delta_e + \delta_f \in \mathcal{S}$. This is an obvious generalization of the basis-exchange property for matroids, to which it reduces if all the multi-indices are 0-1-valued.

Theorem 7.2 *Let $P(x) = \sum_{|\mathbf{m}|=r} a_{\mathbf{m}} x^{\mathbf{m}}$ be a homogeneous degree- r polynomial (on the ground set E) that has the half-plane property and is not identically zero. Then, for every $\mathbf{m}, \mathbf{m}' \in \text{supp}(P)$, \mathbf{m} is exchangeable towards \mathbf{m}' in $\text{supp}(P)$.*

Clearly, by specializing Theorem 7.2 to the multiaffine case, we obtain Theorem 7.1.

PROOF OF THEOREM 7.2. Since P is homogeneous, by Theorem 6.1 we may assume without loss of generality that $a_{\mathbf{m}} > 0$ for all $\mathbf{m} \in \text{supp}(P)$.

We proceed by induction on r . The cases $r = 0$ and $r = 1$ are trivial. The induction step fails when passing from $r = 1$ to $r = 2$, for reasons that will be explained below. Therefore, we treat $r = 2$ as the base case of the induction, and deduce it from Theorem 5.3 as follows: Let $A = (a_{ij})_{i,j \in E}$ be the unique symmetric real matrix such that $P = P_A$. If the conclusion of the theorem fails, then there exist $\mathbf{m}, \mathbf{m}' \in \text{supp}(P)$ and $e \in E$ with $\mathbf{m}(e) > \mathbf{m}'(e)$ such that for any $f \in E$ with $\mathbf{m}(f) < \mathbf{m}'(f)$ we have $\mathbf{m} - \delta_e + \delta_f \notin \text{supp}(P)$. Since $r = 2$ we may write $\mathbf{m} = \delta_e + \delta_g$ and $\mathbf{m}' = \delta_b + \delta_c$ (some of b, c, e, g may coincide). Since neither $\delta_g + \delta_b$ nor $\delta_g + \delta_c$ is in $\text{supp}(P)$, we have $a_{bg} = a_{gb} = a_{cg} = a_{gc} = 0$. Let $x = \delta_e + \rho \delta_g$ for some $\rho > 0$, and let $y = \delta_b + \delta_c$. We see that

$$x^T A y = (a_{eb} + a_{ec})/2 \quad (7.2a)$$

$$y^T A y = a_{bc} + (a_{bb} + a_{cc})/2 > 0 \quad (7.2b)$$

$$x^T A x = \rho a_{eg} + (a_{ee} + \rho^2 a_{gg})/2 > 0 \quad (7.2c)$$

where $a_{eg}, a_{bc} > 0$ since $\mathbf{m}, \mathbf{m}' \in \text{supp}(P)$]. As $\rho \rightarrow \infty$, inequality (e) of Theorem 5.3 eventually fails, so P does not have the half-plane property.

For the induction step, assume that $r \geq 3$ and that the theorem holds for homogeneous polynomials of degree $r - 1$. We shall employ an inner induction on $n = |E|$. The base case $n = 1$ is trivial. So assume the result for homogeneous polynomials of degree r and ground sets of size strictly less than n . Let P be a homogeneous polynomial of degree r on an n -element ground set E , and let $\mathbf{m}, \mathbf{m}' \in \text{supp}(P)$. We shall show that $\mathbf{m} \rightarrow \mathbf{m}'$ in $\text{supp}(P)$, i.e. that for every $e \in E$ such that $\mathbf{m}(e) > \mathbf{m}'(e)$, there exists an $f \in E$ such that $\mathbf{m}(f) < \mathbf{m}'(f)$ and $\mathbf{m} - \delta_e + \delta_f \in \text{supp}(P)$. We divide the argument into three cases:

1. There exists $b \in E$ such that $\mathbf{m}(b) = \mathbf{m}'(b) = 0$.
2. There exists $b \in E$ such that $\mathbf{m}(b) > 0$ and $\mathbf{m}'(b) > 0$.
3. For every $b \in E$, exactly one of $\mathbf{m}(b) > 0$ or $\mathbf{m}'(b) > 0$ holds.

Case 1. Suppose there exists an element b such that $\mathbf{m}(b) = \mathbf{m}'(b) = 0$. Then consider the polynomial $P^{\setminus b}$, which by Proposition 3.1 has the half-plane property. The ground set $E \setminus b$ of $P^{\setminus b}$ is strictly smaller than the ground set E of P , so the hypothesis of the inner induction applies. Both \mathbf{m} and \mathbf{m}' are in the support of $P^{\setminus b}$, so we have $\mathbf{m} \rightarrow \mathbf{m}'$ in $\text{supp}(P^{\setminus b})$. Since $\text{supp}(P^{\setminus b}) \subseteq \text{supp}(P)$, it follows that $\mathbf{m} \rightarrow \mathbf{m}'$ in $\text{supp}(P)$.

Case 2. Suppose there exists an element b such that $\mathbf{m}(b) > 0$ and $\mathbf{m}'(b) > 0$. Then consider the polynomial $P^{\setminus b}$, which by Proposition 3.1 has the half-plane property. This polynomial is homogeneous of degree $r - 1$, so the hypothesis of the outer induction applies. Both $\mathbf{m} - \delta_b$ and $\mathbf{m}' - \delta_b$ are in the support of $P^{\setminus b}$, so we have $(\mathbf{m} - \delta_b) \rightarrow (\mathbf{m}' - \delta_b)$ in $\text{supp}(P^{\setminus b})$. Now let $e \in E$ be such that $\mathbf{m}(e) > \mathbf{m}'(e)$. Since $(\mathbf{m} - \delta_b)(e) > (\mathbf{m}' - \delta_b)(e)$, there is an $f \in E$ with $\mathbf{m}(f) < \mathbf{m}'(f)$ and $\mathbf{m} - \delta_b - \delta_e + \delta_f \in \text{supp}(P^{\setminus b})$. This implies that $\mathbf{m} - \delta_e + \delta_f \in \text{supp}(P)$.

This argument shows that whenever $\mathbf{p}, \mathbf{q} \in \text{supp}(P)$ are such that $\mathbf{p} \wedge \mathbf{q} > 0$, we have $\mathbf{p} \rightarrow \mathbf{q}$ in $\text{supp}(P)$. This fact will be used repeatedly in the remainder of the proof.

Case 3. Suppose that for every $b \in E$, exactly one of $\mathbf{m}(b) > 0$ or $\mathbf{m}'(b) > 0$ holds. We proceed in a series of steps:

(a) We claim that there is a multi-index $\mathbf{k} \in \text{supp}(P)$ such that $\mathbf{m} \wedge \mathbf{k} \neq 0$ and $\mathbf{m}' \wedge \mathbf{k} \neq 0$. Suppose not, and specialize $x_b = y$ if $\mathbf{m}(b) > 0$ and $x_b = z$ if $\mathbf{m}'(b) > 0$. The result is a polynomial of the form $Ay^r + Bz^r$ for some numbers $A, B > 0$, which has the half-plane property. However, upon substituting the values $y = B^{1/r}e^{i\pi/2r}$ and $z = A^{1/r}e^{-i\pi/2r}$ this polynomial vanishes, a contradiction (since $r \geq 2$). Hence there must exist a multi-index $\mathbf{k} \in \text{supp}(P)$ as claimed.

(b) Next, we claim that there is a multi-index $\mathbf{k}' \in \text{supp}(P)$ such that $|\mathbf{m} \wedge \mathbf{k}'| \geq 2$ and $|\mathbf{m}' \wedge \mathbf{k}'| \geq 1$. (This is what fails in the case $r = 2$.) Perhaps the \mathbf{k} found in step

(a) already satisfies this condition. If not, then $|\mathbf{m} \wedge \mathbf{k}| = 1$ and $|\mathbf{m}' \wedge \mathbf{k}| \geq 2$. Let $b \in E$ be such that $(\mathbf{m}' \wedge \mathbf{k})(b) > 0$. Since $\mathbf{k} \rightarrow \mathbf{m}$ in $\text{supp}(P)$ [as shown in Case 2], there is a $g \in E$ with $\mathbf{k}(g) < \mathbf{m}(g)$ such that $\mathbf{k}' = \mathbf{k} - \delta_b + \delta_g \in \text{supp}(P)$. This \mathbf{k}' meets the required conditions.

(c) Next, we claim that there is a multi-index $\mathbf{p} \in \text{supp}(P)$ such that $|\mathbf{m} \wedge \mathbf{p}| \geq 2$ and $|\mathbf{m}' \wedge \mathbf{p}| \geq 1$ and, moreover, for all $b \in E$ such that $\mathbf{m}(b) > 0$ we have $\mathbf{p}(b) \leq \mathbf{m}(b)$. To show this, we define

$$\nu(\mathbf{p}, \mathbf{m}) = \sum_{b \in E: \mathbf{m}(b) > 0} \max[0, \mathbf{p}(b) - \mathbf{m}(b)] \quad (7.3)$$

and choose $\mathbf{k}' \in \text{supp}(P)$ as in step (b). If $\nu(\mathbf{k}', \mathbf{m}) = 0$ then we are done, so assume that $\nu(\mathbf{k}', \mathbf{m}) > 0$. Let $b \in E$ be such that $\mathbf{m}(b) > 0$ and $\mathbf{k}'(b) > \mathbf{m}(b)$. Since $\mathbf{k}' \rightarrow \mathbf{m}$ in $\text{supp}(P)$, there is a $g \in E$ with $\mathbf{k}'(g) < \mathbf{m}(g)$ such that $\mathbf{p}' = \mathbf{k}' - \delta_b + \delta_g \in \text{supp}(P)$. We have $\nu(\mathbf{p}', \mathbf{m}) < \nu(\mathbf{k}', \mathbf{m})$ and $|\mathbf{m} \wedge \mathbf{p}'| = |\mathbf{m} \wedge \mathbf{k}'| \geq 2$ and $|\mathbf{m}' \wedge \mathbf{p}'| = |\mathbf{m}' \wedge \mathbf{k}'| \geq 1$. Repeating this argument inductively as required produces a multi-index $\mathbf{p} \in \text{supp}(P)$ satisfying the claim.

(d) Let $e \in E$ be such that $\mathbf{m}(e) > \mathbf{m}'(e)$, and consider two subcases:

(i) $\mathbf{m}(e) > \mathbf{p}(e)$.

(ii) $\mathbf{m}(e) = \mathbf{p}(e)$.

Subcase (i): If $\mathbf{m}(e) > \mathbf{p}(e)$, then since $\mathbf{m} \rightarrow \mathbf{p}$ in $\text{supp}(P)$, there is an $f \in E$ with $\mathbf{m}(f) < \mathbf{p}(f)$ such that $\mathbf{m} - \delta_e + \delta_f \in \text{supp}(P)$. Since $\mathbf{m}(f) < \mathbf{p}(f)$ we must have $\mathbf{m}(f) = 0$, so that $\mathbf{m}(f) < \mathbf{m}'(f)$.

Subcase (ii): Here $\mathbf{m}(e) = \mathbf{p}(e) > 0 = \mathbf{m}'(e)$. Therefore, since $\mathbf{p} \rightarrow \mathbf{m}'$ in $\text{supp}(P)$, there is a $g \in E$ with $\mathbf{p}(g) < \mathbf{m}'(g)$ such that $\mathbf{q} = \mathbf{p} - \delta_e + \delta_g \in \text{supp}(P)$. Now we have $\mathbf{m}(e) > \mathbf{q}(e)$ and $\mathbf{m} \wedge \mathbf{q} > 0$ and, moreover, for all $b \in E$ such that $\mathbf{m}(b) > 0$ we have $\mathbf{q}(b) \leq \mathbf{m}(b)$. We may therefore repeat the argument of subcase (i) with \mathbf{q} in place of \mathbf{p} . \square

Remark. In step (a) of Case 3 we needed to invoke the same-phase property (Theorem 6.1) in order to ensure that $A, B \neq 0$. In the multiaffine case this is unnecessary, because only \mathbf{m} and \mathbf{m}' contribute to A and B , so $A, B \neq 0$ is guaranteed; and this fact is enough to imply that $Ay^r + Bz^r$ cannot have the half-plane property (by Corollary 6.5). In most other respects, however, the proof of Theorem 7.2 is not much more complicated than the proof for the multiaffine case.

Theorem 7.2 shows that the support of a homogeneous polynomial with the half-plane property satisfies a multi-analogue of the matroid basis-exchange axiom. Such structures were introduced by Bouchet and Cunningham [8] and are called “jump systems” (see also [49, 28]). For $\mathbf{p} \in \mathbb{Z}^E$, let $\|\mathbf{p}\| = \sum_{e \in E} |\mathbf{p}(e)|$. A nonempty subset $\mathcal{J} \subseteq \mathbb{Z}^E$ is called a *jump system* (with ground set E) in case it satisfies the following condition:

- (J) For any $\mathbf{m}, \mathbf{m}' \in \mathcal{J}$ and $\mathbf{k} \in \mathbb{Z}^E$ such that $\|\mathbf{m} - \mathbf{k}\| = 1$ and $\|\mathbf{k} - \mathbf{m}'\| = \|\mathbf{m} - \mathbf{m}'\| - 1$, either $\mathbf{k} \in \mathcal{J}$ or there exists $\mathbf{k}' \in \mathcal{J}$ with $\|\mathbf{k} - \mathbf{k}'\| = 1$ and $\|\mathbf{k}' - \mathbf{m}'\| = \|\mathbf{m} - \mathbf{m}'\| - 2$.

A jump system \mathcal{J} is said to have *constant sum* when $|\mathbf{m}| = |\mathbf{m}'|$ for all $\mathbf{m}, \mathbf{m}' \in \mathcal{J}$. In this case, the condition defining a jump system is equivalent to requiring that for all $\mathbf{m}, \mathbf{m}' \in \mathcal{J}$, both $\mathbf{m} \rightarrow \mathbf{m}'$ in \mathcal{J} and $(-\mathbf{m}) \rightarrow (-\mathbf{m}')$ in $-\mathcal{J} \equiv \{-\mathbf{k} : \mathbf{k} \in \mathcal{J}\}$. (The definition of exchangeability extends in the natural way from \mathbb{N}^E to \mathbb{Z}^E .)

Corollary 7.3 *Let $P \neq 0$ be a homogeneous polynomial with the half-plane property. Then $\text{supp}(P)$ is a jump system with constant sum.*

PROOF. Since P is homogeneous, $\text{supp}(P)$ has constant sum. For any $\mathbf{m}, \mathbf{m}' \in \text{supp}(P)$, we have $\mathbf{m} \rightarrow \mathbf{m}'$ in $\text{supp}(P)$ by Theorem 7.2. For each $e \in E$, let $\mathbf{p}(e) = \deg_e P$, and consider $Q(x) = x^{\mathbf{p}} P(1/x)$. Since the open right half-plane is invariant under the transformation $z \mapsto 1/z$, Q is also a homogeneous polynomial with the half-plane property. Thus, by Theorem 7.2 again, $(\mathbf{p} - \mathbf{m}) \rightarrow (\mathbf{p} - \mathbf{m}')$ in $\text{supp}(Q)$. But this means that $(-\mathbf{m}) \rightarrow (-\mathbf{m}')$ in $-\text{supp}(P)$. \square

A nonempty intersection of a jump system in \mathbb{Z}^E with the set $\{0, 1\}^E$ is known as a *delta-matroid*, and is usually regarded as a set system (by identifying a set with its characteristic vector). When a delta-matroid is uniform (i.e. all sets have the same size), then it is the collection of bases of a matroid (as is easily seen), so that Theorem 7.1 is a special case of Corollary 7.3.

It is natural to wonder what can be said about the support of a polynomial P with the half-plane property, if one omits (or weakens) the assumption that P is homogeneous. The fact that the same-phase property need not hold in this case causes great complications; one might start by considering the subclass of polynomials P that have the half-plane property *and* the same-phase property.

Question 7.4 *If P is multiaffine and has the half-plane property, is $\text{supp}(P)$ a delta-matroid? What if P also has the same-phase property?*

Question 7.5 *Assume that P has the half-plane property and has definite parity (i.e. $\mathbf{m}, \mathbf{m}' \in \text{supp}(P)$ implies that $|\mathbf{m}| \equiv |\mathbf{m}'| \pmod{2}$). Is P then a jump system? (Recall from Theorem 6.2 that all such polynomials have the same-phase property.)*

7.2 The weak half-plane property

Recall that a set system \mathcal{S} (resp. a matroid M) is said to have the half-plane property if its generating polynomial $P_{\mathcal{S}}$ (resp. its basis generating polynomial $P_{\mathcal{B}(M)}$) does. Let us now say that a set system \mathcal{S} (resp. a matroid M) has the *weak half-plane property* if there exists a polynomial P with the half-plane property for which

$\text{supp}(P) = \mathcal{S}$ [resp. $\text{supp}(P) = \mathcal{B}(M)$]. This is obviously a weaker condition, since the nonzero coefficients of P need not all be equal. (By Theorem 6.1 they must, however, all have the same phase in the matroid case or, more generally, when the set system is r -uniform.) Theorem 7.1 can then be rephrased as saying that if an r -uniform set system has the weak half-plane property, it is necessarily the collection of bases of a matroid. A central open question is the converse:

Question 7.6 *Does every matroid M have the weak half-plane property? And if not, which ones do?*

We shall soon show (Corollary 8.2) that all matroids representable over \mathbb{C} have the weak half-plane property. But we are totally in the dark about matroids not representable over \mathbb{C} , such as the Fano matroid F_7 and its coextensions $AG(3, 2)$ and S_8 (which are representable only over fields of characteristic 2), the matroids T_8 , R_9 , $S(5, 6, 12)$ and $PG(2, 3)$ (which are representable only over fields of characteristic 3), or the Vámos and non-Pappus matroids (which are not representable over any field).

Remark. The weak half-plane property for an n -element matroid M is algorithmically testable in time $c_1^{c_2^{2n+|\mathcal{B}(M)|}}$ using cylindrical algebraic decomposition, and in time $c^{n|\mathcal{B}(M)|}$ using more recent algorithms (see Section 2.6). But this is thoroughly unfeasible in practice!

8 The determinant condition and $\sqrt[6]{1}$ -matroids

8.1 The determinant condition

Let A be an $r \times n$ matrix with complex entries, and define

$$Q_A(x) = \det(AXA^*) \quad (8.1)$$

where $X = \text{diag}(x_1, \dots, x_n)$ and $*$ denotes Hermitian conjugate. Clearly Q_A is multiaffine and is homogeneous of degree r . We have the following straightforward generalization of Theorem 1.1:

Theorem 8.1 *Let A be an arbitrary complex $r \times n$ matrix, and define Q_A by (8.1). Then:*

- (a) Q_A has the half-plane property.
- (b) $Q_A(x) = \sum_{\substack{S \subseteq [n] \\ |S| = r}} |\det(A \upharpoonright S)|^2 x^S$, where $A \upharpoonright S$ denotes the (square) submatrix of A using the columns indexed by the set S .

PROOF. Note first that if $\text{rank } A < r$, then $Q_A \equiv 0$ and so trivially has the half-plane property; so let us assume that $\text{rank } A = r$. Then $\ker A^* = 0$, i.e. for every nonzero $\psi \in \mathbb{C}^r$, we have $A^*\psi \neq 0$ in \mathbb{C}^n . It follows that, for each $\psi \neq 0$, the quantity

$$\psi^* A X A^* \psi = \sum_{i=1}^n |(A^*\psi)_i|^2 x_i \quad (8.2)$$

has strictly positive real part whenever $\text{Re } x_i > 0$ for all i ; so in particular $A X A^* \psi \neq 0$. Therefore the matrix $A X A^*$ is nonsingular, and so has a nonzero determinant.

The identity (b) is an immediate consequence of the Cauchy–Binet theorem.¹²

□

Remark. This proof is a direct generalization of the proof of Theorem 1.1, to which it reduces if we take A to be the directed vertex-edge incidence matrix for any orientation of G with the i_0 th row suppressed (here i_0 is an arbitrary vertex of G).

Corollary 8.2 (a) *Every matroid representable over \mathbb{C} has the weak half-plane property.*

(b) *Let M be a rank- r matroid on n elements that can be represented over \mathbb{C} by an $r \times n$ matrix A for which every $r \times r$ subdeterminant is either zero or else of modulus 1. Then M has the half-plane property.*

PROOF. (a) This is an immediate consequence of Theorem 8.1, since $\det(A \upharpoonright S) \neq 0$ if and only if $S \in \mathcal{B}(M)$.

(b) Clearly $|\det(A \upharpoonright S)|^2 = 1$ if $S \in \mathcal{B}(M)$ and 0 if $S \notin \mathcal{B}(M)$. So, by Theorem 8.1(b) we have $Q_A = P_{\mathcal{B}(M)}$, and by Theorem 8.1(a) Q_A has the half-plane property. □

8.2 (F, G) -representability and $\sqrt[6]{1}$ -matroids

Which matroids have the property of Corollary 8.2(b)? Certainly regular matroids do, as they can be represented over \mathbb{R} (hence also over \mathbb{C}) by a totally unimodular

¹² The Cauchy–Binet theorem states that if A is an $m \times n$ matrix and B is an $n \times m$ matrix, where $m \leq n$, then

$$\det(AB) = \sum_{\substack{S \subseteq [n] \\ |S| = m}} (\det A[[m]|S]) (\det B[S|[m]]) .$$

Here the sum runs over all m -element subsets $S \subseteq [n] \equiv \{1, 2, \dots, n\}$, $A[[m]|S]$ denotes the submatrix of A consisting of the columns from S (taken in order), and $B[S|[m]]$ denotes the submatrix of B consisting of the rows from S (taken in order). See e.g. [50, pp. 128–129] for a proof.

$r \times n$ matrix A (i.e. one for which every square subdeterminant is either 0, $+1$ or -1). But, in fact, Corollary 8.2(b) applies to a larger class of matroids. Let us pose these questions more generally in the following context.

Let F be a field, let F^* be the multiplicative group $F \setminus 0$, and let G be a subgroup of F^* . If A is a matrix over F , let us call A an (F, G) -matrix if every nonzero subdeterminant of A lies in G (note, in particular, that every nonzero entry of A must lie in G). And let us call A a *weak* (F, G) -matrix if every nonzero $r \times r$ subdeterminant of A lies in G , where A has rank r . Finally, let us call a matroid (F, G) -representable if it is representable over F by an (F, G) -matrix, and *weakly* (F, G) -representable if it is representable over F by a weak (F, G) -matrix.

The concept of an (F, G) -representable matroid was introduced by Whittle [78] and studied further, under the hypothesis that $-1 \in G$, by Semple and Whittle [67] within the more general framework of partial fields.¹³ (We shall not need this more general notion here, and we refer the reader to Semple and Whittle’s paper for a discussion of it.) As these authors note, many important classes of matroids are special cases of (F, G) -representable matroids:

- F -representable matroids: $G = F^*$.
- Regular matroids (also known as unimodular matroids [76]): Let F be any field of characteristic zero (e.g. $F = \mathbb{Q}, \mathbb{R}$ or \mathbb{C}), and let $G = \{-1, 1\}$.
- k -regular matroids [77, 78, 65, 56]: Let F be the field $\mathbb{Q}(\alpha_1, \dots, \alpha_k)$ obtained by extending the rationals by k algebraically independent transcendental elements $\alpha_1, \dots, \alpha_k$, and let G be the set of all products of integer powers of differences of distinct members of $\{0, 1, \alpha_1, \dots, \alpha_k\}$. Thus, 0-regular matroids are just the regular matroids; 1-regular matroids are also called near-regular. A matroid is ω -regular if it is k -regular for some k .
- Dyadic matroids [77, 78]: Let F be any field of characteristic zero, and let $G = \{\pm 2^k : k \in \mathbb{Z}\}$.
- Sixth-root-of-unity matroids ($\sqrt[6]{1}$ -matroid for short) [78]: Let $F = \mathbb{C}$ and G be the multiplicative group \mathbb{Z}_6 of complex sixth roots of unity.
- Complex unimodular matroids: Let $F = \mathbb{C}$ and G be the multiplicative group $U(1)$ of complex numbers of modulus 1.

Most “naturally arising” examples of (F, G) -representability — including those relevant to this paper — have $-1 \in G$, and (as we shall see) the theory takes a simpler form under this hypothesis. Nevertheless, there is also some interesting theory that can be developed for the case $-1 \notin G$: for example, in Appendix B, we characterize $(F, \{1\})$ -representable matroids. We shall therefore comment briefly on which results seem to need $-1 \in G$ and which do not.

¹³ These authors call it a (G, F) -matroid.

We begin with some easy lemmas:

Lemma 8.3 *Let A be an (F, G) -matrix, and let B be an F -matrix.*

- (a) *If B is obtained from A by a sequence of row or column deletions, then B is an (F, G) -matrix.*
- (b) *If B is obtained from A by a sequence of row or column scalings by factors lying in G , then B is an (F, G) -matrix.*
- (c) *If $-1 \in G$, and B is obtained from A by a sequence of row or column interchanges, then B is an (F, G) -matrix.*
- (d) *If $-1 \in G$, and B is obtained from A by a sequence of pivots, then B is an (F, G) -matrix.*

PROOF. (a)–(c) are trivial. The proof of (d) is essentially identical to that of [67, Proposition 3.3]. \square

Lemma 8.4 *The class of (F, G) -representable matroids is closed under deletion and direct sum. Moreover, if $-1 \in G$, then the class is closed under contraction, duality, 2-sum, and series and parallel connection.*

PROOF. The proof of the first sentence is straightforward. When $-1 \in G$, closure under contraction follows from Lemma 8.3(d) by mimicking the argument of [55, Proposition 3.2.6]. As noted in [67, Proposition 4.2], closure under duality, 2-sum, and series and parallel connection follows by mimicking the standard arguments for fields that appear in [55, Theorem 2.2.8 and Proposition 7.1.21]. \square

It follows from the characterization of $(F, \{1\})$ -representable matroids in Appendix B that, when $-1 \notin G$, the class of (F, G) -representable matroids need not be closed under duality, 2-sum, series connection, or parallel connection. One question we have been unable to answer is the following:

Question 8.5 *When $-1 \notin G$, is the class of (F, G) -representable matroids always closed under contraction?*

It follows from Theorem B.1 that the answer is affirmative when $G = \{1\}$, but we do not know whether this is so in general.

Our first nontrivial result is that weak (F, G) -representability is no more general than (F, G) -representability, at least when $-1 \in G$. We do not know whether this result holds without the assumption that $-1 \in G$.

Proposition 8.6 *Let F be a field, and let G be a subgroup of the multiplicative group F^* that contains -1 . Let A be a rank- r $m \times n$ matrix over F that is a weak (F, G) -matrix. Then there is a rank- r $r \times n$ matrix B over F that is an (F, G) -matrix and for which $M[A] = M[B]$.*

PROOF. Let $M = M[A]$. Since A has rank r , we may delete $m - r$ rows from A leaving an $r \times n$ matrix A_1 whose rows are linearly independent. Moreover, A_1 is a weak (F, G) -matrix representing M . Now the following row and column operations leave a determinant unchanged except for possibly multiplying it by -1 :

- (i) Interchanging two rows or two columns.
- (ii) Adding a multiple of one row to another.

By operations (i) and (ii), A_1 can be transformed into a matrix A_2 of the form $[D|Z]$ where D is a diagonal matrix all of whose diagonal entries are nonzero.

The matrix $[D|Z]$ can be further transformed as follows. Let the top-left 2×2 submatrix of D be $\begin{bmatrix} d_1 & 0 \\ 0 & d_2 \end{bmatrix}$. In $[D|Z]$, perform the following operations:

- (i) replace row 2 by row 2 plus d_1^{-1} times row 1;
- (ii) replace row 1 by row 1 minus d_1 times row 2;
- (iii) replace row 2 by row 2 plus d_1^{-1} times row 1; and
- (iv) interchange rows 1 and 2.

We now have, in the top-left corner, the submatrix $\begin{bmatrix} 1 & 0 \\ 0 & -d_1 d_2 \end{bmatrix}$. We can repeat this process until all the diagonal entries in D except possibly the last, d_r , are ones. The resulting matrix $[D'|Z']$ is a weak (F, G) -matrix representing M . Since $\det D' = d_r$, it follows that $d_r \in G$. Thus, we can multiply the last row of $[D'|Z']$ by d_r^{-1} to obtain a weak (F, G) -matrix A_3 representing M of the form $[I_r|Y]$.

We shall show next that A_3 is an (F, G) -matrix. Let A'_3 be a $k \times k$ submatrix of A_3 for some k with $1 \leq k \leq r - 1$. The identity submatrix I_r of A_3 has an $r \times (r - k)$ submatrix B_1 each of whose columns contains all zeros in the rows of A_3 that meet A'_3 . Let A''_3 be the submatrix of A_3 consisting of those columns that meet some column of A'_3 . As A_3 is a weak (F, G) matrix, the determinant of the matrix $[B_1|A''_3]$ is in G . But the value of this determinant is $\pm \det A''_3$. Since $-1 \in G$, we deduce that $\det A''_3 \in G$. We conclude that A_3 is indeed an (F, G) -matrix representing M . \square

Proposition 8.7 *Let F be a field, and let G be a subgroup of F^* that contains -1 . Let A be an (F, G) -matrix of rank r . Then there is an (F, G) -matrix $B = [I_r|D]$ such that $M[A] \simeq M[B]$ and B is obtainable from A by a sequence of row or column interchanges, pivots, and deletions of zero rows.*

PROOF. This follows from Lemma 8.3 by well-known arguments: see e.g. [67, Propositions 3.5 and 4.1]. \square

If D is a matrix over F , let us denote by $\mathcal{G}(D^\sharp)$ the simple bipartite graph whose vertices correspond to the rows and columns of D and whose edges correspond to the nonzero entries of D [55, pp. 190, 194]. If H is a subgroup of F^* , let us call D *H-normalized* if there exists a basis B for the cycle matroid of $\mathcal{G}(D^\sharp)$ [i.e. a maximal spanning forest in $\mathcal{G}(D^\sharp)$] such that each entry in D corresponding to an edge in B lies in H .

We aim next to show that, given an (F, G) -matrix $[I_r|D]$, we can perform a sequence of row and column scalings by elements of G (thereby maintaining an (F, G) -matrix) to yield a matrix $[I_r|D']$ for which D' is $\{1\}$ -normalized. In fact, a much stronger result is true:

Proposition 8.8 *Let F be a field, and let G be a subgroup of F^* . Consider a matrix $[I_r|D]$ over F , in which all the nonzero elements of D lie in G . Let $B = \{b_1, \dots, b_k\}$ be a basis for the cycle matroid of $\mathcal{G}(D^\sharp)$, and let $\{\theta_1, \dots, \theta_k\}$ be elements of G . Then, by a sequence of row and column scalings with scale factors in G , one can obtain from $[I_r|D]$ a matrix $[I_r|D']$ in which the entry of D' corresponding to b_i is θ_i . In particular, every subdeterminant of $[I_r|D']$ differs from the corresponding subdeterminant of $[I_r|D]$ by a factor lying in G .*

PROOF. The proof of [55, Theorem 6.4.7] carries through with very minor modifications to prove this proposition. \square

By definition, every $\sqrt[6]{1}$ -matroid is a complex unimodular matroid. In fact, the two classes coincide:

Theorem 8.9 *The classes of complex unimodular matroids and $\sqrt[6]{1}$ -matroids are equal.*

PROOF. Evidently we need only prove that every complex unimodular matroid is a $\sqrt[6]{1}$ -matroid. By Propositions 8.7 and 8.8, we can represent any complex unimodular matroid M by a matrix $[I_r|D]$ over \mathbb{C} in which all nonzero subdeterminants have modulus one and in which D is $\{1\}$ -normalized. The desired result then follows from the following two lemmas:

Lemma 8.10 *Let D be a \mathbb{Z}_6 -normalized matrix over \mathbb{C} in which all nonzero subdeterminants have modulus one. Then all the nonzero entries of D are sixth roots of unity.*

Lemma 8.11 *Let A be a square matrix over \mathbb{C} , whose determinant has modulus 1, such that all nonzero entries are sixth roots of unity. Then $\det(A)$ is in fact a sixth root of unity.*

Let us prove these two lemmas in reverse order:

PROOF OF LEMMA 8.11. Since $\det(A)$ is a linear combination, with coefficients ± 1 , of products of entries in A , it follows that $\det(A)$ belongs to the additive subgroup of \mathbb{C} generated by the sixth roots of unity. In the complex plane, the elements of this subgroup are the vertices of a triangular lattice, which intersects the unit circle precisely at the sixth roots of unity. \square

PROOF OF LEMMA 8.10. Let B be a basis for the cycle matroid of $\mathcal{G}(D^\sharp)$ such that each entry in D corresponding to an edge in B is a sixth root of unity. For each nonzero entry d of D , let e_d be the corresponding edge of $\mathcal{G}(D^\sharp)$. If $e_d \notin B$, then adding e_d to the subgraph of $\mathcal{G}(D^\sharp)$ induced by B creates a unique cycle C_d , whose length is some even integer $2k \geq 4$. Assume that there exists a nonzero entry d that is not a sixth root of unity, and choose one for which k is minimal.

Let D_d be the $k \times k$ submatrix of D induced by the vertices of C_d . By construction, all the entries of D_d corresponding to edges in $C_d \setminus e_d$ are sixth roots of unity. Moreover, if D_d contains any nonzero entry d' besides those of C_d , then the corresponding edge $e_{d'}$ of $\mathcal{G}(D^\sharp)$ is a diagonal of C_d (note that this cannot occur in the minimal case $2k = 4$). It follows that $|C_{d'}| < |C_d|$, so the choice of d implies that d' is a sixth root of unity. Hence every nonzero entry of D_d except possibly d is a sixth root of unity.

Consider now the subgraph of $\mathcal{G}(D^\sharp)$ induced by the vertices of C_d ; among the cycles of this graph containing e_d , choose a cycle C' of shortest length $2j \leq 2k$. Let D'_d be the $j \times j$ submatrix of D_d induced by the vertices of C' . Then each row and column of D'_d has exactly two nonzero entries corresponding to edges of C' , and no other nonzero entries (by the minimality of C'). Moreover, all the nonzero entries of D'_d , except possibly d , are sixth roots of unity.

Because each row and column of D'_d has exactly two nonzero entries, the expansion of $\det(D'_d)$ in permutations has exactly two nonzero terms, exactly one of which contains d . We conclude that $\det(D'_d) = a - bd$ where a and b are sixth roots of unity. If $\det(D'_d) = 0$, then d is a sixth root of unity. If $\det(D'_d)$ has modulus 1, then so does $c \equiv a/b - d$. Hence d is a complex number of modulus 1, which differs from a sixth root of unity (a/b) by another complex number (c) of modulus 1. It follows by simple geometry that d is a sixth root of unity. \square

It is well known that the regular matroids are characterized by the following list of equivalent properties [76]:

Theorem 8.12 *The following are equivalent for a matroid M :*

- (a) M is $(F, \{\pm 1\})$ -representable for some field F of characteristic 0.
- (a) M is weakly $(F, \{\pm 1\})$ -representable for some field F of characteristic 0.
- (c) M is $(\mathbb{Q}, \{\pm 1\})$ -representable.

- (d) M is weakly $(\mathbb{Q}, \{\pm 1\})$ -representable.
- (e) M is representable over $GF(2)$ and over at least one field of characteristic not equal to 2.
- (f) M is representable over all fields.

More recently, Whittle [78] has given an analogous characterization of the $\sqrt[6]{1}$ -matroids. Combining his (deep) results with our (comparatively elementary) Proposition 8.6 and Theorem 8.9, we obtain:

Theorem 8.13 (Whittle [78]) *The following are equivalent for a matroid M :*

- (a) M is representable over $GF(3)$ and at least one field of characteristic 2.
- (b) M is representable over $GF(3)$ and $GF(4)$.
- (c) M is representable over $GF(3)$ and all fields $GF(2^{2^k})$ for k integer.
- (d) M is representable over all fields F that contain a root of the polynomial $x^2 - x + 1$. [In particular, M is representable over all fields $GF(q)$ for which q is a power of 3, q is a square, or $q \equiv 1 \pmod{3}$.]
- (e) M is representable over all fields $GF(q)$ for which $q \not\equiv 2 \pmod{3}$.
- (f) M is $(\mathbb{C}, \mathbb{Z}_6)$ -representable.
- (g) M is $(\mathbb{C}, U(1))$ -representable.
- (h) M is weakly $(\mathbb{C}, U(1))$ -representable.

The excluded-minor characterization of regular matroids is well known [55, Theorems 13.1.1 and 13.1.2]:

Theorem 8.14 (Tutte)

- (a) A binary matroid M is regular if and only if it has no minor isomorphic to F_7 or F_7^* .
- (b) A matroid M is regular if and only if it has no minor isomorphic to $U_{2,4}$, F_7 or F_7^* .

More recently, Geelen, Gerards and Kapoor [29], as a corollary of their important work in determining the excluded minors for $GF(4)$ -representable matroids, have given an analogous excluded-minor characterization of $\sqrt[6]{1}$ -matroids:

Theorem 8.15 (Geelen–Gerards–Kapoor [29])

- (a) A ternary matroid M is a $\sqrt[6]{1}$ -matroid if and only if M has no minor isomorphic to F_7^- , $(F_7^-)^*$ or P_8 .
- (b) A matroid M is a $\sqrt[6]{1}$ -matroid if and only if M has no minor isomorphic to $U_{2,5}$, $U_{3,5}$, F_7 , F_7^* , F_7^- , $(F_7^-)^*$ or P_8 .

It will be shown in Section 11 that none of F_7 , F_7^* , F_7^- , $(F_7^-)^*$ and P_8 has the half-plane property. The next two corollaries come from combining this fact with the last two theorems and Corollary 8.2:

Corollary 8.16 *A binary matroid has the half-plane property if and only if it is regular.*

Corollary 8.17 *A ternary matroid has the half-plane property if and only if it is a $\sqrt[6]{1}$ -matroid.*

To conclude this section, we remark that a much less elementary proof of Theorem 8.9 than the one given above comes from using Theorem 8.15(b) together with the fact, which is not difficult to verify, that none of $U_{2,5}$, $U_{3,5}$, F_7 , F_7^* , F_7^- , $(F_7^-)^*$ and P_8 is a complex unimodular matroid.

9 Uniform matroids

9.1 Half-plane property

The basis generating polynomial of the uniform matroid $U_{r,n}$ is the elementary symmetric polynomial

$$E_{r,n}(x_1, \dots, x_n) = \sum_{1 \leq i_1 < i_2 < \dots < i_r \leq n} x_{i_1} x_{i_2} \cdots x_{i_r} \quad (9.1)$$

(we set $E_{0,n} \equiv 1$). We have the following fundamental result:

Theorem 9.1 *Let $0 \leq r \leq n$. Then:*

- (a) *The elementary symmetric polynomial $E_{r,n}$ has the half-plane property.*
- (b) *For $r \geq 1$, the rational function $F_{r,n} \equiv E_{r,n}/E_{r-1,n}$ is strictly real-part-positive on H^n .*

Note first that (a) implies (b), by Proposition 2.8(b) with $\lambda_i = 1/(n - r + 1)$ for all i ; and (b) implies (a), by Lemmas 2.7 and 2.6 and the fact (which is a special case of Proposition 4.6) that the elementary symmetric polynomials $E_{r,n}$ with $r < n$ are irreducible over any field. [Alternatively, the truth of (b) for *all* $r \geq 1$ allows one to

prove (a) by induction on r , starting from $E_{0,n} \equiv 1$ and using Lemma 2.6.] So we can try to prove whichever half seems more convenient.

We have already given two proofs of Theorem 9.1(a): one based on nice principal truncation (i.e. differentiation) starting from the easy case $r = n$ (Proposition 4.13), and another based on nice principal cotruncation starting from the easy case $r = 0$ (Proposition 4.16). We shall give a third proof in Section 10, based on showing that $U_{r,n}$ is a nice transversal matroid (Corollary 10.3 and Example 10.1 following it). Let us here give two more proofs: one based on the Grace–Walsh–Szegő coincidence theorem, and one based on “series-parallel” identities for the rational functions $F_{r,n}$.

FOURTH PROOF. We have $E_{r,n}(\xi, \dots, \xi) = \binom{n}{r} \xi^r$, which is manifestly nonvanishing for ξ in the open right half-plane H . Since $E_{r,n}$ is symmetric and multiaffine, the Grace–Walsh–Szegő coincidence theorem (Theorem 2.12) implies that $E_{r,n}$ is nonvanishing in H^n . \square

FIFTH PROOF [25]. We shall prove (b) by induction on r . The case $r = 1$ is trivial for all n . Now the following identities are easy to prove (for $1 \leq r \leq n$):

$$E_{r,n}(x) = E_{r,n-1}(x_{\neq i}) + x_i E_{r-1,n-1}(x_{\neq i}) \quad \text{for any index } i \quad (9.2)$$

$$E_{r,n}(x) = \frac{1}{r} \sum_{i=1}^n x_i E_{r-1,n-1}(x_{\neq i}) \quad (9.3)$$

We therefore have

$$F_{r,n}(x) = \frac{1}{r} \sum_{i=1}^n \frac{x_i E_{r-1,n-1}(x_{\neq i})}{E_{r-1,n}(x)} \quad (9.4a)$$

$$= \frac{1}{r} \sum_{i=1}^n \frac{x_i E_{r-1,n-1}(x_{\neq i})}{E_{r-1,n-1}(x_{\neq i}) + x_i E_{r-2,n-1}(x_{\neq i})} \quad (9.4b)$$

$$= \frac{1}{r} \sum_{i=1}^n \frac{x_i F_{r-1,n-1}(x_{\neq i})}{F_{r-1,n-1}(x_{\neq i}) + x_i} \quad (9.4c)$$

$$= \frac{1}{r} \sum_{i=1}^n \left(\frac{1}{x_i} + \frac{1}{F_{r-1,n-1}(x_{\neq i})} \right)^{-1}, \quad (9.4d)$$

so that the strict real-part-positivity of $F_{r,n}$ follows from that of $F_{r-1,n-1}$. \square

Remarks. 1. The computation (9.4) is really just (4.18) specialized to the case of uniform matroids. So this is just an explicit version of the “nice cotruncation” proof of Proposition 4.16. Indeed, it was our analysis of the proof (9.4) that led us to abstract the idea of weighted principal cotruncation (Section 4.6).

2. The identity (9.4d) shows how $F_{r,n}(x)$ can be synthesized as the admittance of a 2-terminal series-parallel network whose elementary branch admittances are positive

multiples of the x_i . To begin with, $F_{1,n}(x) = x_1 + \cdots + x_n$ is the admittance of branches x_1, \dots, x_n placed in parallel. Then $F_{r,n}(x)$ is the admittance of n branches in parallel, the i th of which consists of admittances x_i/r and $F_{r-1,n-1}(x_{\neq i})/r$ in series. [Of course, the admittance $F_{r-1,n-1}(x_{\neq i})/r$ is obtained from the network giving $F_{r-1,n-1}(x_{\neq i})$ by dividing each branch admittance by r .]

Now let $\mathcal{G}_{r,n}$ be the graph obtained by this construction: it contains approximately n^r edges, and each edge e is assigned an admittance $\alpha_e x_{i(e)}$ for some positive constant α_e and some index $i(e)$. Then the spanning-tree polynomial $T_{\mathcal{G}_{r,n}}(\{x_e\})$, with its arguments specialized as $x_e \rightarrow \alpha_e x_{i(e)}$, contains the polynomial $E_{r,n}(x_1, \dots, x_n)$ as a factor. In this way, Theorem 9.1(a) can be obtained as a corollary of Theorem 1.1.

More generally, one might try to prove the half-plane property for a polynomial P by starting from the spanning-tree polynomial T_G of a large graph G , specializing to fewer variables, and then extracting a factor. We do not know whether this method can be applied fruitfully in other cases.

9.2 Brown–Colbourn property

Let us now prove that for $r < n$, the uniform matroids $U_{r,n}$ not only have the half-plane property but have the Brown–Colbourn property (which is stronger, by virtue of Corollary 2.3).

Theorem 9.2 *Let $0 \leq r \leq n - 1$. Then the uniform matroid $U_{r,n}$ has the “Brown–Colbourn property”, i.e. $\operatorname{Re} x_i < -1/2$ for all i implies $P_{\mathcal{I}(U_{r,n})}(x) \neq 0$, where $\mathcal{I}(U_{r,n}) = \{S \subseteq \{1, \dots, n\} : |S| \leq r\}$ is the collection of independent sets of $U_{r,n}$.*

PROOF. Since $P_{\mathcal{I}(U_{r,n})}(x_1, \dots, x_n) = \sum_{k=0}^r E_{k,n}(x_1, \dots, x_n)$ is manifestly symmetric and multiaffine, it follows from the Grace–Walsh–Szegő coincidence theorem (Theorem 2.12) that it suffices to prove the Brown–Colbourn property for the univariate polynomial

$$I_{r,n}(z) = P_{\mathcal{I}(U_{r,n})}(z, \dots, z) = \sum_{k=0}^r \binom{n}{k} z^k. \quad (9.5)$$

A slightly sharper result than this is given in Proposition 9.3 below. \square

As preparation for the statement and proof of Proposition 9.3, let us note that the univariate reliability polynomial [cf. (1.7)] for the uniform matroid $U_{r,n}$ is given by

$$\operatorname{Rel}_{r,n}(q) = (1 - q)^n I_{r,n}\left(\frac{q}{1 - q}\right) = \sum_{k=0}^r \binom{n}{k} q^k (1 - q)^{n-k} \quad (9.6)$$

and hence can be written as $\operatorname{Rel}_{r,n}(q) = (1 - q)^{n-r} H_{r,n}(q)$ with

$$H_{r,n}(q) = \sum_{k=0}^r \binom{n}{k} q^k (1 - q)^{r-k} \quad (9.7a)$$

$$= \sum_{k=0}^r \sum_{j=0}^{r-k} (-1)^j \binom{n}{k} \binom{r-k}{j} q^{j+k} \quad (9.7b)$$

$$= \sum_{\ell=0}^r q^\ell \sum_{j=0}^{\ell} (-1)^j \binom{n}{\ell-j} \binom{r-\ell+j}{j} \quad (9.7c)$$

$$= \sum_{\ell=0}^r \binom{n-r-1+\ell}{\ell} q^\ell \quad (9.7d)$$

where we define $\binom{n-r-1+\ell}{\ell} = \delta_{\ell 0}$ in case $r = n$.¹⁴

Proposition 9.3 (Wagner [73]) *Let $0 \leq r \leq n-1$. Then all the zeros of $H_{r,n}(q)$ lie in the annulus*

$$\frac{1}{n-r} \leq |q| \leq \frac{r}{n-1}. \quad (9.8)$$

In particular, all the zeros of $H_{r,n}(q)$ lie in $|q| \leq 1$, so that all the zeros of $I_{r,n}(z)$ lie in $\operatorname{Re} z \geq -1/2$.

PROOF. The ratios of successive coefficients of $H_{r,n}$ are $\lambda_\ell = \binom{n-r-1+\ell}{\ell} / \binom{n-r+\ell}{\ell+1} = (\ell+1)/(n-r+\ell)$, which is nondecreasing as ℓ runs from 0 to $r-1$. Thus, by the Eneström–Kakeya theorem [51, Theorem 30.3 and Exercise 2] (see also [2]) it follows that all the zeros of $H_{r,n}$ lie in the annulus $\lambda_0 \leq |q| \leq \lambda_{r-1}$. \square

10 The permanent condition: Transversal and co-transversal matroids

10.1 Heilmann–Lieb theorem

We begin by recalling the Heilmann–Lieb [35] theorem on the zeros of matching polynomials. This theorem is most often quoted in its univariate version (see e.g. [48, Section 8.5]), but it is the multivariate result [35, Theorem 4.6 and Lemma 4.7] that is truly fundamental.

Let $G = (V, E)$ be a loopless graph, and let us define the matching polynomial with edge weights $\{\lambda_e\}_{e \in E}$ and vertex weights $\{x_i\}_{i \in V}$:

$$M_G(x; \lambda) = \sum_{\text{matchings } M} \prod_{e=ij \in M} \lambda_e x_i x_j. \quad (10.1)$$

(Here $e = ij$ means that the endpoints of e are i, j ; if G is not simple, this is a slight abuse of notation but unlikely to cause any confusion. Note also that we could,

¹⁴ The binomial-coefficient identity used in the last step can be deduced from [32, eqns. (1.7) and (1.9)]. Alternatively, it is a specialization of [79, Exercise 4.15(a)] followed by [32, eqn. (1.7)].

if we wanted, restrict attention to simple graphs by replacing each set of parallel edges e_1, \dots, e_k with a single edge e' of weight $\lambda_{e'} = \sum_{i=1}^k \lambda_{e_i}$.) Define also the complementary matching polynomial

$$\widetilde{M}_G(x; \lambda) = x^V M_G(1/x; \lambda) \quad (10.2)$$

where $x^V = \prod_{i \in V} x_i$. If $e = ij \in E$, we have the fundamental recursion relation

$$M_G(x; \lambda) = M_{G \setminus e}(x; \lambda) + \lambda_e x_i x_j M_{G-i-j}(x; \lambda) \quad (10.3)$$

or equivalently

$$\widetilde{M}_G(x; \lambda) = \widetilde{M}_{G \setminus e}(x; \lambda) + \lambda_e \widetilde{M}_{G-i-j}(x; \lambda). \quad (10.4)$$

Repeated application of (10.4) to all edges incident on a vertex i , followed by deletion of i , yields the key identity

$$\widetilde{M}_G(x; \lambda) = x_i \widetilde{M}_{G-i}(x; \lambda) + \sum_{\substack{e \sim i \\ e = ij}} \lambda_e \widetilde{M}_{G-i-j}(x; \lambda) \quad (10.5)$$

where $e \sim i$ denotes that e is incident on i . The Heilmann–Lieb theorem asserts that if the edge weights are nonnegative, then the polynomials M_G and \widetilde{M}_G have the half-plane property:

Theorem 10.1 (Heilmann and Lieb [35]) *Let $G = (V, E)$ be a loopless graph, and let $\{\lambda_e\}_{e \in E}$ be nonnegative edge weights. If $\operatorname{Re} x_i > 0$ for all $i \in V$, then*

- (a) $\widetilde{M}_G(x; \lambda) \neq 0$.
- (b) For every $i \in V$, $\widetilde{M}_{G-i}(x; \lambda) \neq 0$.
- (c) For every $i \in V$, $\operatorname{Re} \frac{\widetilde{M}_G(x; \lambda)}{\widetilde{M}_{G-i}(x; \lambda)} > 0$.
- (d) $M_G(x; \lambda) \neq 0$.

PROOF. We shall prove (a)–(c) by induction on $|V|$. The theorem is trivial if $|V| = 0$ or 1, since $\widetilde{M}_G = 1$ for $|V| = 0$ and $\widetilde{M}_G(x_1) = x_1$ for $|V| = 1$. So assume that (a)–(c) hold for all graphs with fewer than n vertices, and let $|V| = n$. Then (b) holds for G because (a) holds for $G - i$. It then follows from (10.5) that

$$\frac{\widetilde{M}_G(x; \lambda)}{\widetilde{M}_{G-i}(x; \lambda)} = x_i + \sum_{\substack{e \sim i \\ e = ij}} \lambda_e \frac{\widetilde{M}_{G-i-j}(x; \lambda)}{\widetilde{M}_{G-i}(x; \lambda)}. \quad (10.6)$$

The right-hand side has positive real part because $\operatorname{Re} x_i > 0$, $\lambda_e \geq 0$ and $\operatorname{Re} [\widetilde{M}_{G-i-j}(x; \lambda)/\widetilde{M}_{G-i}(x; \lambda)] > 0$ by hypothesis (c) applied to $G - i$ (using the fact

that the function $z \mapsto 1/z$ maps the right half-plane to itself). We conclude that (c) holds for G . Hence so does (a).

Finally, (d) follows from (a) by Proposition 4.2. \square

Remarks. 1. The same conclusion obviously holds [with a reversal of sign in (c)] if $\operatorname{Re} x_i < 0$ for all i . But the proof does not apply to other rotated half-planes, as we need invariance under the function $z \mapsto 1/z$. Indeed, the result for other half-planes is false already for $G = K_2$, which has $\widetilde{M}_G(x_1, x_2; \lambda) = \lambda_{12} + x_1 x_2$.

2. This proof is basically the same as that of Heilmann and Lieb [35, Lemma 4.7], but is slightly simpler and stronger: they prove (c) only for pairs $(G', G' - i)$ belonging to their set \mathcal{E}_G .

3. How could one have guessed the key inductive hypothesis (c)? It suffices to note, from (10.5), that $\partial \widetilde{M}_G / \partial x_i = \widetilde{M}_{G-i}$; and of course $\widetilde{M}_{G-i} \not\equiv 0$ because it has constant term 1. Therefore, by either Proposition 2.8(b) or Theorem 3.2(c), if \widetilde{M}_G has the half-plane property, then necessarily hypothesis (c) must hold.

4. Here is a simple alternate proof of Theorem 10.1(d), based on the concept of multiaffine part (Proposition 4.17): The polynomial $P_G(x) = \prod_{e=ij \in E} (1 + \lambda_e x_i x_j)$ manifestly has the half-plane property whenever $\lambda_e \geq 0$ for all e . But P_G^b (the multiaffine part of P_G) is precisely M_G .

10.2 Application to transversal matroids

Now let us specialize the Heilmann–Lieb theorem to the case of a bipartite graph $G = (V, E)$ with bipartition $V = A \cup B$, setting $x_j = 1$ for $j \in B$ and considering $M_G(x; \lambda)$ as a polynomial in $\{x_i\}_{i \in A}$. Then the restricted matching polynomial

$$\bar{M}_G(x; \lambda) = \sum_{\text{matchings } M} \prod_{\substack{e=ij \in M \\ i \in A \\ j \in B}} \lambda_e x_i \quad (10.7)$$

also has the half-plane property, provided that the edge weights are nonnegative.

Consider now the transversal matroid $M[G, A]$ with ground set A defined by the bipartite graph G , in which a subset $S \subseteq A$ is declared independent if it can be matched into B . Defining the weighted sum of such matchings,

$$c(S; \lambda) = \sum_{\substack{\text{matchings } M \\ V(M) \cap A = S}} \prod_{e \in M} \lambda_e, \quad (10.8)$$

we find immediately that

$$\bar{M}_G(x; \lambda) = \sum_{S \in \mathcal{I}(M[G, A])} c(S; \lambda) x^S. \quad (10.9)$$

So the restricted matching polynomial $\bar{M}_G(x; \lambda)$, which has the half-plane property, is *almost* the independent-set generating polynomial for $M[G, A]$: the only trouble comes from the weights $c(S; \lambda)$, which need not be equal. Likewise, the restricted *maximum*-matching polynomial

$$\bar{M}_G^\sharp(x; \lambda) = \sum_{\substack{S \in \mathcal{I}(M[G, A]) \\ |S| = \text{rank } M[G, A]}} c(S; \lambda) x^S \quad (10.10)$$

also has the half-plane property (by Proposition 2.2), and is almost the basis generating polynomial for $M[G, A]$, again modulo the problem of possibly unequal weights.

Note also that we may, if we wish, let G be the *complete* bipartite graph on the vertex set $A \cup B$, since any undesired edges can always be given weight $\lambda_e = 0$. In particular, if we take $A = [n]$, $B = [r]$ and $G = K_{n,r}$, we have the following permanental analogue of Theorem 8.1 (see [53] for the definition and properties of permanents):

Theorem 10.2 *Let Λ be an arbitrary nonnegative $r \times n$ matrix ($r \leq n$), and define P_Λ by*

$$P_\Lambda(x) = \text{per}(\Lambda X) \quad (10.11)$$

where $X = \text{diag}(x_1, \dots, x_n)$. Then:

(a) P_Λ has the half-plane property.

(b) $P_\Lambda(x) = \sum_{\substack{S \subseteq [n] \\ |S| = r}} \text{per}(\Lambda \upharpoonright S) x^S$, where $\Lambda \upharpoonright S$ denotes the (square) submatrix of Λ using the columns indexed by the set S .

PROOF. Part (b) is an immediate consequence of the definition of the permanent of a non-square matrix. Part (a) follows immediately from the Heilmann–Lieb theorem and the fact that

$$P_\Lambda(x) = \bar{M}_G^\sharp(x; \lambda) = \lim_{\alpha \rightarrow \infty} \alpha^{-r} \bar{M}_G(\alpha x; \lambda) \quad (10.12)$$

with $G = K_{n,r}$. \square

Let us now call the pair (G, A) *nice* if there exists a collection $\{\lambda_e\}_{e \in E}$ of non-negative edge weights so that $c(S; \lambda)$ has the same nonzero value for all bases S of $M[G, A]$. And let us call the transversal matroid M *nice* if there exists a nice pair (G, A) such that $M \simeq M[G, A]$. The foregoing results can be rephrased as follows:

Corollary 10.3

(a) *Every transversal matroid has the weak half-plane property.*

(b) *Every nice transversal matroid has the half-plane property.*

Unfortunately, Corollary 10.3(a) adds nothing to Corollary 8.2(a), since every transversal matroid is representable over all fields of sufficiently large cardinality, and hence in particular over \mathbb{C} [55, Corollary 12.2.17]. But Corollary 10.3(b) is powerful, as we shall soon see.

Our next task should be to characterize nice pairs (G, A) and nice transversal matroids M . Unfortunately, we are unable to do this in general, so we shall content ourselves with giving some examples of nice and non-nice transversal matroids. As a preliminary, let us observe that the restriction of a nice pair (G, A) to any subset $A' \subseteq A$ (and the same set B) yields a nice pair (G', A') . In particular, any restriction of a nice transversal matroid is a nice transversal matroid.

Let us also note that, to prove that a matroid M has the half-plane property, it suffices by Proposition 4.2 to show that either M or its dual is a nice transversal matroid. If M^* is transversal, we say that M is *cotransversal* (also called a *strict gammoid*); if M^* is nice transversal, we say that M is *co-nice cotransversal*.

10.3 Examples

In this subsection we give some examples of nice and non-nice transversal matroids. Sometimes, instead of specifying the pair (G, A) , we shall find it more convenient to specify the family $\mathcal{A} = \{A_j\}_{j \in B}$ of subsets of A defined by $i \in A_j$ if and only if $ij \in E$. The family \mathcal{A} is called a *presentation* for $M[G, A]$.

Example 10.1. Every uniform matroid $U_{r,n}$ is nice. Indeed, it suffices to take the obvious presentation, namely the one induced by the complete bipartite graph $G = K_{n,r}$ with $|A| = n$ and $|B| = r$, and to set $\lambda_e = 1$ for all e . \square

Example 10.2. Not all pairs (G, A) are nice. To see this, consider the bipartite graph G shown in Figure 2, with $A = \{1, 2, 3, 4\}$ and $B = \{x, y\}$. Then every 2-element subset of A is matchable into B , and the weights are

$$c(12) = ad \tag{10.13a}$$

$$c(13) = ae \tag{10.13b}$$

$$c(14) = af \tag{10.13c}$$

$$c(23) = be + cd \tag{10.13d}$$

$$c(24) = bf \tag{10.13e}$$

$$c(34) = cf \tag{10.13f}$$

where for conciseness we have used the name of the edge e in place of λ_e . Suppose that all these subset weights are equal and nonzero. Then all the edge weights a, \dots, f must be nonzero, and moreover we must have $d = e = f$ and $a = b = c$. But then $c(23) = 2c(24)$, a contradiction. So this pair (G, A) is not nice. Note, nevertheless,




Figure 2: A bipartite graph $G = (V, E)$ with bipartition $V = A \cup B$.

that $M[G, A] \simeq U_{2,4}$, which has an alternative (nonisomorphic) presentation that *is* nice (Example 10.1). \square

Example 10.3. Not all transversal matroids are nice. To see this, consider the rank-3 whirl \mathcal{W}^3 , which has ground set $\{1, 2, 3, 4, 5, 6\}$ and 3-point lines 123, 345 and 561. It is transversal with maximal presentation $\mathcal{A} = \{456, 126, 234\}$ (i.e. the sets of the presentation are the complements of the three 3-point lines). By a general result ([52] or [11, Theorem 5.2.6]), this is the unique maximal presentation. Now suppose that we could weight the edges in the bipartite graph to make \mathcal{W}^3 nice. The bases 413, 513 and 613 all occur exactly once as a transversal; therefore, all the edges associated with the first set in the presentation must receive the same weight, call it $\lambda(1)$. Likewise, 513, 523 and 563 all occur once, so all the edges associated with the second set in the presentation must receive the same weight, call it $\lambda(2)$. Finally, 512, 513 and 514 all occur once, so all the edges associated with the third set in the presentation must receive the same weight, call it $\lambda(3)$. Thus the basis 135 gets weight $\lambda(1)\lambda(2)\lambda(3)$. But the basis 246 gets weight $2\lambda(1)\lambda(2)\lambda(3)$. Therefore \mathcal{W}^3 is not nice.

It follows that any transversal matroid having \mathcal{W}^3 as a restriction is also non-nice: this includes \mathcal{W}^{3+} and $\mathcal{W}^3 + e$ (see Figures 5 and 6 in Appendix A) and their free extensions $\mathcal{W}^{3+} + e$ and $\mathcal{W}^3 + e + f$ (see Figure 8). \square

Example 10.4. For $n_1, n_2, n_3 \geq 3$, let L_{n_1, n_2, n_3} be the rank-3 matroid consisting of three nonintersecting lines containing n_1 , n_2 and n_3 points, respectively. This matroid has a presentation consisting of the complements of the three lines, in which each basis occurs exactly twice as a transversal. Therefore this matroid is nice (taking all weights λ_e equal).

It follows that the matroids P_6 , S_7 , F_7^{-6} and P_7''' (see Figures 4–7 in Appendix A) are nice, since they can be obtained by deleting elements from $L_{3,3,3}$ or $L_{4,3,3}$. \square

Example 10.5. More generally, let $L_{n_1, n_2, n_3; n'}$ (with $n_1, n_2, n_3 \geq 3$ and $n' \geq 0$) be the rank-3 matroid consisting of three nonintersecting lines plus n' freely added points. Let us show that the smallest example with freely added points, namely $M = L_{3,3,3;1}$, is *not* nice:

Let the 3-point lines be abc , def and ghi , and call the freely added point 0. Then M has a presentation consisting of the complements of the three 3-point lines, i.e. $\mathcal{A} = \{0abcdef, 0abcghi, 0defghi\}$. Indeed, this is the unique presentation.¹⁵ Now suppose that we could weight the edges in the bipartite graph to make M nice. For each element x of $0defghi$, the basis abx occurs exactly twice as a transversal: as abx and as bax . This means that all the edges associated with the third set in the presentation must receive the same weight, call it $\lambda(3)$. By the same argument, all the edges associated with the second set in the presentation must receive the same weight $\lambda(2)$, and all the edges associated with the first set in the presentation must receive the same weight $\lambda(1)$. Thus the basis abd gets weight $2\lambda(1)\lambda(2)\lambda(3)$. But the basis $0ae$ gets weight $3\lambda(1)\lambda(2)\lambda(3)$. Therefore M is not nice. \square

Example 10.6. Let $L_{n_1; n'}$ (with $n_1 \geq 3$ and $n' \geq 1$) be the rank-3 matroid consisting of one n_1 -point line plus n' freely added points. By deletion from $L_{n_1, 3, 3}$ (Example 10.4), we can conclude that $L_{n_1; n'}$ is nice whenever $n' \leq 4$. Let us show that the smallest remaining case, $L_{3;5}$, is *not* nice:

$L_{3;5}$ is transversal with unique maximal presentation $\mathcal{A} = \{12345678, 12345678, 45678\}$. Let us denote the weights associated with the first (resp. second, third) set in the presentation as λ_i (resp. λ'_i, λ''_i). We shall use the combinations $\alpha_{ij} = \lambda_i\lambda'_j + \lambda_j\lambda'_i$ and $\beta_{ij} = \lambda_i\lambda'_j - \lambda_j\lambda'_i$.

For each pair of distinct elements $i, j \in \{1, 2, 3\}$ and each $x \in \{4, 5, 6, 7, 8\}$, the basis ijx occurs exactly twice as a transversal, always with the same two ways of selecting i and j from the first two sets. Fixing i and j and letting x vary, we conclude that all the edges associated with the third set in the presentation must receive the same weight, which without loss of generality we may take equal to 1. It then follows that $\alpha_{ij} = 1$ for all distinct $i, j \in \{1, 2, 3\}$.

For each $i \in \{1, 2, 3\}$ and each triplet of distinct elements $x, y, z \in \{4, 5, 6, 7, 8\}$, the bases ixy , ixz and iyz each occur exactly four times as transversals. Thus $\alpha_{ix} + \alpha_{iy} = 1$, $\alpha_{ix} + \alpha_{iz} = 1$ and $\alpha_{iy} + \alpha_{iz} = 1$. By comparing the first equation with

¹⁵ PROOF. The presentation \mathcal{A} is a maximal presentation, since adding an element to any set will result in one of abc , def or ghi occurring as a transversal. By a general result ([52] or [11, Theorem 5.2.6]), this is the unique maximal presentation. The complements of the three sets in the presentation are the three flats ghi , def and abc . Any other presentation must come from the maximal one by deleting elements from the sets in the presentation. This will increase the complements of these sets. But the complements of the sets in any presentation must be flats in M , and there are no flats other than $E(M)$ that contain any one of abc , def or ghi . This proves the claim.




Figure 3: The matroid M_{n_1, n_2} .

the difference of the last two, we deduce that $\alpha_{ix} = 1/2$ for all $i \in \{1, 2, 3\}$ and $x \in \{4, 5, 6, 7, 8\}$.

For each triplet of distinct elements $x, y, z \in \{4, 5, 6, 7, 8\}$, the basis xyz occurs exactly six times as a transversal, so we have $\alpha_{xy} + \alpha_{xz} + \alpha_{yz} = 1$. Therefore, for each quadruplet of distinct elements $x, y, z, w \in \{4, 5, 6, 7, 8\}$, we have $\alpha_{xy} + \alpha_{xz} + \alpha_{yz} = 1$, $\alpha_{xy} + \alpha_{xw} + \alpha_{yw} = 1$, $\alpha_{xz} + \alpha_{xw} + \alpha_{zw} = 1$ and $\alpha_{yz} + \alpha_{yw} + \alpha_{zw} = 1$. Subtracting the sum of the last two equations from the sum of the first two, we conclude that $\alpha_{xy} = \alpha_{zw}$. Since there are *five* elements in $\{4, 5, 6, 7, 8\}$, we can conclude that $\alpha_{xy} = 1/3$ for all distinct $x, y \in \{4, 5, 6, 7, 8\}$.

Finally, for any distinct $i, j \in \{1, 2, 3\}$ and distinct $x, y \in \{4, 5, 6, 7, 8\}$, we have $\alpha_{ij}\alpha_{xy} - \alpha_{iy}\alpha_{jx} = 1/3 - 1/4 = 1/12$. Simplifying this, we find $\beta_{ix}\beta_{jy} = -1/12$ for all such i, j, x, y . But this implies that all the numbers β_{ix} ($1 \leq i \leq 3$, $4 \leq x \leq 8$) are equal and take the value $\pm\sqrt{-1/12}$. In particular, there are no *real* solutions. [The niceness equations do have *complex* solutions of the form $\lambda_1 = \lambda_2 = \lambda_3 = e^{\pm i\pi/6}/\sqrt{2}$, $\lambda'_1 = \lambda'_2 = \lambda'_3 = e^{\mp i\pi/6}/\sqrt{2}$, $\lambda_4 = \dots = \lambda_8 = 1/\sqrt{6}$, $\lambda'_4 = \dots = \lambda'_8 = 1/\sqrt{6}$ and $\lambda''_4 = \dots = \lambda''_8 = 1$.] \square

Example 10.7. For $n_1, n_2 \geq 2$, let M_{n_1, n_2} be the rank-3 matroid on the $(n_1 + n_2 + 2)$ -element ground set $\{0, 1, \dots, n_1 + n_2 + 1\}$ such that there are lines $\{0, 1, 2, \dots, n_1\}$ and $\{0, n_1 + 1, n_1 + 2, \dots, n_1 + n_2\}$ but no other 3-element circuits (see Figure 3). This matroid is transversal with $A = \{0, 1, \dots, n_1 + n_2 + 1\}$ and $B = \{x, y, z\}$, and has as a presentation

$$A_x = \{0, 1, 2, \dots, n_1 + n_2\} \quad (10.14a)$$

$$A_y = \{1, 2, \dots, n_1, n_1 + n_2 + 1\} \quad (10.14b)$$

$$A_z = \{n_1 + 1, n_1 + 2, \dots, n_1 + n_2, n_1 + n_2 + 1\} \quad (10.14c)$$

Now put weights

$$\lambda_{0x} = 1 \quad (10.15a)$$

$$\lambda_{ix} = 1/2 \quad \text{for } 1 \leq i \leq n_1 + n_2 \quad (10.15b)$$

$$\lambda_{iy} = 1 \quad \text{for all } i \in A_y \quad (10.15c)$$

$$\lambda_{iz} = 1 \quad \text{for all } i \in A_z \quad (10.15d)$$

It is easily checked that $c(S; \lambda) = 1$ for every basis S of M_{n_1, n_2} . So this is a nice presentation, and M_{n_1, n_2} is a nice transversal matroid. In particular, $Q_6 \simeq M_{2,2}$, $P_6 \simeq M_{2,3} \setminus 0$, $Q_7 \simeq M_{2,3}$, $S_7 \simeq M_{3,4} \setminus \{0, 1\}$ and $P_7''' \simeq M_{3,3} \setminus 0$ are nice transversal matroids. \square

Proposition 10.4 *All matroids on a ground set of at most 6 elements have the half-plane property.*

PROOF. The half-plane property is preserved by direct sums (Proposition 4.4) and 2-sums (Corollary 4.9). Therefore, we can restrict attention to 3-connected matroids. Since all rank-1 and rank-2 matroids have the half-plane property (Corollary 5.5), we can restrict attention to matroids of rank ≥ 3 . Finally, since the half-plane property is invariant under duality (Proposition 4.2), we can restrict attention to matroids of rank $\leq \lfloor n/2 \rfloor$, where n is the number of elements. So it suffices to consider 3-connected rank-3 matroids on 6 elements. All these matroids are shown in Figure 4 (see Appendix A). The uniform matroid $U_{3,6}$ has the half-plane property, as we have shown many times (Section 9). The graphic matroid $M(K_4)$ and the whirl \mathcal{W}^3 are $\sqrt[6]{1}$ -matroids, hence have the half-plane property by Corollary 8.2(b). Finally, we have just shown (Examples 10.4 and 10.7) that Q_6 and P_6 are nice transversal matroids, hence also have the half-plane property. \square

Example 10.8. The matroid F_7^{-5} (see Figure 6 in Appendix A) is transversal with unique maximal presentation $\mathcal{A} = \{2347, 4567, 1234567\}$. The bases 251, 351, 451 and 751 all occur exactly once as a transversal; therefore, all the edges associated with the first set in the presentation must receive the same weight, call it $\lambda(1)$. By symmetry, all the edges associated with the second set in the presentation must receive the same weight, call it $\lambda(2)$. But then 471 occurs twice as a transversal, so it gets twice the weight that 251 gets. It follows that F_7^{-5} is not nice. \square

Example 10.9. The matroid $(F_7^{-3})^*$ (see Figure 6) is transversal with unique maximal presentation $\mathcal{A} = \{123, 147, 156, 345\}$. The bases 1465, 2465 and 3465 all occur exactly once as a transversal, so all the edges associated with the first set in the presentation must receive the same weight, call it $\lambda(1)$. By symmetry, all the edges associated with the second set in the presentation must receive the same weight, call it $\lambda(2)$, and all the edges associated with the third set in the presentation must receive the same weight, call it $\lambda(3)$. Finally, the bases 1763, 1764 and 1765 all occur

exactly once as a transversal, so all the edges associated with the fourth set in the presentation must receive the same weight, call it $\lambda(4)$. But then 1345 occurs three times as a transversal, so it gets three times the weight that 1456 gets. So $(F_7^{-3})^*$ is not nice. \square

Example 10.10. The matroid $(F_7^{-4})^*$ (see Figure 6) is transversal with unique maximal presentation $\mathcal{A} = \{123, 147, 156, 1234567\}$. The bases 1456, 2456 and 3456 all occur exactly twice as transversals, all with the same two ways of selecting 456 from the last three sets in the presentation; so all the edges associated with the first set in the presentation must receive the same weight, call it $\lambda(1)$. By symmetry, all the edges associated with the second set in the presentation must receive the same weight, call it $\lambda(2)$, and all the edges associated with the third set in the presentation must receive the same weight, call it $\lambda(3)$. Without loss of generality we can set $\lambda(1) = \lambda(2) = \lambda(3) = 1$. Let us denote the weights associated with the fourth set in the presentation as λ_i ($1 \leq i \leq 7$). Bases then get weights as follows:

$$\begin{aligned} 1245: & \lambda_1 + \lambda_2 + \lambda_4 + \lambda_5 \\ 1345: & \lambda_1 + \lambda_3 + \lambda_4 + \lambda_5 \\ 1246: & \lambda_1 + \lambda_2 + \lambda_4 + \lambda_6 \\ 1346: & \lambda_1 + \lambda_3 + \lambda_4 + \lambda_6 \\ 1276: & \lambda_1 + \lambda_2 + \lambda_7 + \lambda_6 \\ 1376: & \lambda_1 + \lambda_3 + \lambda_7 + \lambda_6 \\ 1234: & \lambda_2 + \lambda_3 \\ 1475: & \lambda_4 + \lambda_7 \\ 1562: & \lambda_5 + \lambda_6 \end{aligned}$$

It follows from the first six bases that $\lambda_2 = \lambda_3$, $\lambda_5 = \lambda_6$ and $\lambda_4 = \lambda_7$. It then follows from the last three bases that $\lambda_2 = \lambda_3 = \dots = \lambda_7$. But then we must have $\lambda_1 = -\lambda_2$. Therefore, the equations have no *nonnegative* solution, so $(F_7^{-4})^*$ is not nice.

Now consider the matroid $(F_7^{-4}+e)^*$ (see Figure 8) which is transversal with unique maximal presentation $\mathcal{A} = \{123, 147, 156, 12345678, 12345678\}$. The first part of the argument just given for $(F_7^{-4})^*$ carries over with slight modification to $(F_7^{-4}+e)^*$: the bases 14568, 24568 and 34568 all occur exactly *four* times as transversals, all with the same four ways of selecting 4568 from the last four sets in the presentation, so all the edges associated with the first set in the presentation must receive the same weight $\lambda(1)$; and by symmetry, all the edges associated with the second (resp. third) set in the presentation must receive the same weight $\lambda(2)$ [resp. $\lambda(3)$]. Without loss of generality we can set $\lambda(1) = \lambda(2) = \lambda(3) = 1$. Let us denote the weights associated with the fourth (resp. fifth) set in the presentation as λ_i (resp. λ'_i) for $1 \leq i \leq 8$. Bases then get weights as follows:

$$\begin{aligned} 14568: & (\lambda_5 + \lambda_6)\lambda'_8 + (\lambda'_5 + \lambda'_6)\lambda_8 \\ 14578: & (\lambda_4 + \lambda_7)\lambda'_8 + (\lambda'_4 + \lambda'_7)\lambda_8 \\ 12458: & (\lambda_1 + \lambda_2 + \lambda_4 + \lambda_5)\lambda'_8 + (\lambda'_1 + \lambda'_2 + \lambda'_4 + \lambda'_5)\lambda_8 \\ 12678: & (\lambda_1 + \lambda_2 + \lambda_6 + \lambda_7)\lambda'_8 + (\lambda'_1 + \lambda'_2 + \lambda'_6 + \lambda'_7)\lambda_8 \end{aligned}$$

All bases must get equal (and nonzero) weight. Thus, by comparing the sum of the weights of the first two bases with the sum of the weights of the last two bases, we conclude that $\lambda_1\lambda'_8 = \lambda_2\lambda'_8 = \lambda'_1\lambda_8 = \lambda'_2\lambda_8 = 0$ (since the weights are nonnegative). But, by symmetry, the element 2 can be replaced by 3, 4, 5, 6 or 7. Hence $\lambda_i\lambda'_8 = \lambda'_i\lambda_8 = 0$ for $1 \leq i \leq 7$, a contradiction. Therefore $(F_7^{-4} + e)^*$ is not nice. \square

Example 10.11. The matroid $(\mathcal{W}^3 + e)^*$ (see Figure 6) is transversal with unique maximal presentation $\mathcal{A} = \{123, 345, 156, 1234567\}$. The bases 1467, 2467 and 3467 all occur exactly once as a transversal; so all the edges associated with the first set in the presentation must receive the same weight, call it $\lambda(1)$. By symmetry, all the edges associated with the second set in the presentation must receive the same weight, call it $\lambda(2)$, and all the edges associated with the third set in the presentation must receive the same weight, call it $\lambda(3)$. But then 1357 occurs twice as a transversal, so it gets twice the weight that 1467 does. So $(\mathcal{W}^3 + e)^*$ is not nice.

Now consider the matroid $(\mathcal{W}^3 + e + f)^*$ (see Figure 8) which is transversal with unique maximal presentation $\mathcal{A} = \{123, 345, 156, 12345678, 12345678\}$. The argument just given for $(\mathcal{W}^3 + e)^*$ carries over verbatim to $(\mathcal{W}^3 + e + f)^*$ if we replace each basis $abc7$ by $abc78$ and we change “all occur exactly once as a transversal” to “all occur exactly twice as a transversal, all with the same two ways of selecting 78 from the last two sets in the presentation”. It follows that $(\mathcal{W}^3 + e + f)^*$ is not nice. \square

Example 10.12. The matroid $(F_7^{-5})^*$ (see Figure 6) is transversal with (non-maximal) presentation $\mathcal{A} = \{123, 156, 2467, 3457\}$, in which each basis occurs exactly twice as a transversal. Therefore this matroid is nice (taking all weights λ_e equal). \square

Example 10.13. The matroid $(\mathcal{W}^{3+})^*$ (see Figure 5) is transversal with unique maximal presentation $\mathcal{A} = \{156, 345, 1237, 1237\}$. The bases 1427, 5427 and 6427 all occur exactly twice as transversals, all with the same two ways of selecting 27 from the last two sets in the presentation; so all the edges associated with the first set in the presentation must receive the same weight, call it $\lambda(1)$. By symmetry, all the edges associated with the second set in the presentation must receive the same weight, call it $\lambda(2)$. Let us denote the weights associated with the third (resp. fourth) set in the presentation as λ_i (resp. λ'_i) for $i = 1, 2, 3, 7$. The bases 6417, 6427 and 6437 all occur exactly twice as transversals, from which we can conclude that $\lambda_i\lambda'_7 + \lambda'_i\lambda_7 = \alpha > 0$ for $i = 1, 2, 3$. But then 1357 gets weight $2\lambda(1)\lambda(2)\alpha$, while 1427 gets weight $\lambda(1)\lambda(2)\alpha$. So $(\mathcal{W}^{3+})^*$ is not nice.

Now consider the matroid $(\mathcal{W}^{3+} + e)^*$ (see Figure 8) which is transversal with unique maximal presentation $\mathcal{A} = \{156, 345, 1237, 1237, 12345678\}$. The argument just given for $(\mathcal{W}^{3+})^*$ carries over verbatim to $(\mathcal{W}^{3+} + e)^*$ if we replace each basis

$abcd$ by $abcd8$. So $(\mathcal{W}^{3+} + e)^*$ is not nice. \square

Example 10.14. The matroid $(Q_7)^*$ (see Figure 5) is transversal with unique maximal presentation $\mathcal{A} = \{123, 1456, 1456, 1234567\}$. The bases 1457, 2457 and 3457 all occur exactly twice as transversals, all with the same two ways of selecting 457 from the last three sets in the presentation; so all the edges associated with the first set in the presentation must receive the same weight, which without loss of generality we take to be 1. Let us denote the weights associated with the second (resp. third, fourth) set in the presentation as λ_i (resp. λ'_i, λ''_i). For each $x \in \{4, 5, 6\}$, the basis $21x7$ occurs twice as a transversal (as $21x7$ and $2x17$); so $\lambda_1\lambda'_x + \lambda_x\lambda'_1$ takes the same value $\alpha \neq 0$ for $x = 4, 5, 6$. For each pair of distinct $x, y \in \{4, 5, 6\}$, the basis $1xy7$ occurs twice as a transversal (as $1xy7$ and $1yx7$); so $\lambda_x\lambda'_y + \lambda_y\lambda'_x = \alpha$ as well. Bases then get weights as follows:

$$\begin{aligned} 1452: & \alpha(\lambda''_1 + \lambda''_2 + \lambda''_4 + \lambda''_5) \\ 1453: & \alpha(\lambda''_1 + \lambda''_3 + \lambda''_4 + \lambda''_5) \\ 1463: & \alpha(\lambda''_1 + \lambda''_3 + \lambda''_4 + \lambda''_6) \\ 1563: & \alpha(\lambda''_1 + \lambda''_3 + \lambda''_5 + \lambda''_6) \\ 2143: & \alpha(\lambda''_2 + \lambda''_3) \\ 2456: & \alpha(\lambda''_4 + \lambda''_5 + \lambda''_6) \end{aligned}$$

Comparing the first four bases, we conclude that $\lambda''_2 = \lambda''_3 \equiv \beta$ and $\lambda''_4 = \lambda''_5 = \lambda''_6 \equiv \gamma$. But then $\lambda''_1 + \beta + 2\gamma = 2\beta = 3\gamma$, which implies that $\lambda''_1 = -\frac{1}{2}\gamma$. Therefore, the equations have no *nonnegative* solution, so $(Q_7)^*$ is not nice. \square

Example 10.15. The matroid P''_7 (see Figure 7) is transversal with unique maximal presentation $\mathcal{A} = \{1267, 1345, 4567\}$. The bases 134, 234, 634 and 734 all occur exactly once as a transversal; therefore, all the edges associated with the first set in the presentation must receive the same weight. Likewise, 216, 236, 246 and 256 (resp. 134, 135, 136 and 137) all occur once, so all the edges associated with the second (resp. third) set in the presentation must receive the same weight. But then 147 gets twice the weight of 134. So P''_7 is not nice. \square

Example 10.16. The matroid $(P'_7)^*$ (see Figure 7) is transversal with unique maximal presentation $\mathcal{A} = \{123, 156, 267, 345\}$. The bases 1574, 2574 and 3574 all occur exactly once as a transversal; therefore, all the edges associated with the first set in the presentation must receive the same weight. Analogous arguments using 2174, 2574, 2674 (resp. 3524, 3564, 3574 or 2673, 2674, 2675) show that all the edges associated with the second (resp. third or fourth) set in the presentation must receive the same weight. But then 1263 gets twice the weight of 1574. So $(P'_7)^*$ is not nice.

Now consider the matroid $(P'_7 + e)^*$ (see Figure 8) which is transversal with unique maximal presentation $\mathcal{A} = \{123, 156, 267, 345, 12345678\}$. The argument just given

for $(P'_7)^*$ carries over verbatim to $(P'_7 + e)^*$ if we replace each basis $abcd$ by $abcd8$. So $(P'_7 + e)^*$ is not nice. \square

Example 10.17. The matroid $(P''_7)^*$ (see Figure 7) is transversal with (non-maximal) presentation $\mathcal{A} = \{123, 267, 345, 14567\}$, in which each basis occurs exactly twice as a transversal. Therefore $(P''_7)^*$ is nice (taking all weights λ_e equal). \square

Example 10.18. More generally, for each $n \geq 1$, let N_n be the rank- $(n+1)$ transversal matroid on the ground set $\{1, \dots, 2n+1\}$ given by the presentation

$$A_{\hat{0}} = \{1, 2, 4, 6, 8, \dots, 2n-4, 2n-2, 2n, 2n+1\} \quad (10.16a)$$

$$A_{\hat{j}} = \{2j-1, 2j, 2j+1\} \quad \text{for } 1 \leq j \leq n \quad (10.16b)$$

Geometrically, the dual of N_n consists of 3-point lines $123, 345, 567, \dots, (2n-1)(2n)(2n+1)$ joined together in general position in rank n . Let S be a matchable subset of $\{1, \dots, 2n+1\}$. By using induction on n and considering separately the cases when S does and does not contain two consecutive members of $\{1, \dots, 2n+1\}$, we can show (after some non-trivial work) that S arises from exactly two matchings. So, putting equal weights on all edges, we conclude that this is a nice presentation, and N_n is a nice transversal matroid. We have $N_1 \simeq U_{2,3}$, $N_2 \simeq U_{3,5}$ and $N_3 \simeq (P''_7)^*$. \square

Example 10.19. The matroid $(\text{non-Pappus} \setminus 1)^*$ is transversal with maximal presentation $\mathcal{A} = \{247, 269, 348, 359, 456\}$. For each $x \in \{4, 5, 6\}$, the basis $7283x$ occurs exactly once as a transversal, so all edges associated with the last set in the presentation receive the same nonzero weight. For each $y \in \{3, 5, 9\}$, the basis $728y6$ occurs exactly once as a transversal, so all edges associated with the fourth set in the presentation receive the same nonzero weight. For each $z \in \{2, 6, 9\}$, the basis $7z834$ occurs exactly once as a transversal, so all edges associated with the second set in the presentation receive the same nonzero weight. Now consider the bases 29356 and 26354 . The first occurs exactly once as a transversal, while the second occurs exactly twice (as 26354 and as 42356). Since the transversals 29356 and 26354 get the same nonzero weight, the transversal 42356 must get weight 0. Therefore the edge joining the element 4 to the first set in the presentation has weight 0. So we may delete 4 from the first set and still have a presentation. This implies that $\{2, 7\}$ contains a cocircuit of $(\text{non-Pappus} \setminus 1)^*$ and hence contains a circuit of $\text{non-Pappus} \setminus 1$, which is false. So $(\text{non-Pappus} \setminus 1)^*$ is not nice. \square

We have written a MATHEMATICA program `nicetransversal.m` to test a presentation \mathcal{A} for niceness, using the MATHEMATICA function `Solve` to solve the niceness equations $c(S; \lambda) = 1$ for $S \in \mathcal{B}(M[\mathcal{A}])$. This program is available as part of the electronic version of this paper at arXiv.org. However, because the niceness equations are a polynomial system of degree r in a large number of variables, this program sometimes crashes for lack of memory or fails to complete even in several days of CPU time.

10.4 Rank-3 transversal matroids

Let us now make a systematic study of the niceness of rank-3 transversal matroids. We begin by recalling Brylawski's [13] algorithm for constructing transversal matroids. A *principal transversal matroid* (also called a *fundamental transversal matroid*) of rank r is obtained by beginning with a distinguished basis $B = \{1, 2, \dots, r\}$, which we view geometrically as a simplex with vertices $1, 2, \dots, r$, and adding elements as follows: First one can add elements freely to the flat of rank 0 (i.e. add loops). Then one can add elements freely to the flats of rank 1 (i.e. add elements parallel to the basis elements). Then one can add elements freely to the lines spanned by two basis elements: such elements pick up the dependencies that are forced by virtue of their lying on that line, but have no other dependencies. One continues this process by adding elements freely on the flats spanned by three basis elements, then by four basis elements, \dots and finally by r basis elements (the last are elements that are free in the matroid).

Theorem 10.5 (Brylawski [13]) *Every transversal matroid of rank r is a restriction of a principal transversal matroid of rank r , obtained by deleting some elements of the original basis B .*

Theorem 10.6 (Brylawski [13]) *A principal transversal matroid has a unique presentation, which consists of the r cocircuits that are the complements of the flats spanned by the sets $B \setminus i$ ($1 \leq i \leq r$).*

Applying this construction in the case $r = 3$, we conclude that the most general *simple* rank-3 principal transversal matroid is $C_{n_1, n_2, n_3; n'}$ with $n_1, n_2, n_3 \geq 2$ and $n' \geq 0$, defined as the rank-3 matroid on $n_1 + n_2 + n_3 + n' - 3$ elements consisting of consisting of three distinguished points (namely, the simplex vertices), three lines that join pairs of these vertices and contain n_1, n_2 and n_3 points, respectively, together with n' points freely added in the plane spanned by the vertices. All simple rank-3 transversal matroids can then be obtained by deleting zero or more simplex vertices of $C_{n_1, n_2, n_3; n'}$. By considering all the possibilities for simplex vertices to be deleted or not, and for the resulting lines to be nontrivial or trivial, we obtain the following classes of matroids:

- 1) $C_{n_1, n_2, n_3; n'}$ with $n_1, n_2, n_3 \geq 3$ and $n' \geq 0$.
- 2) $D_{n_1, n_2; n'}$ with $n_1, n_2 \geq 3$ and $n' \geq 0$, consisting of two intersecting lines containing n_1 and n_2 points, respectively, together with n' freely added points.
- 3) $E_{n_1, n_2, n_3; n'}$ with $n_1, n_2, n_3 \geq 3$ and $n' \geq 0$, consisting of an n_2 -point line that is met by nonintersecting lines with n_1 and n_3 points, respectively, together with n' freely added points.

- 4) $F_{n_1, n_2, n_3; n'}$ with $n_1, n_2, n_3 \geq 3$ and $n' \geq 0$, consisting of an intersecting pair of lines containing n_1 and n_2 points, respectively, and a third line containing n_3 points that does not meet the first two, together with n' freely added points.
- 5) $L_{n_1, n_2, n_3; n'}$ with $n_1, n_2, n_3 \geq 3$ and $n' \geq 0$, consisting of three nonintersecting lines having n_1, n_2 and n_3 points, respectively, together with n' freely added points.
- 6) $L_{n_1, n_2; n'}$ with $n_1, n_2 \geq 3$ and $n' \geq 0$, consisting of two nonintersecting lines having n_1 and n_2 points, respectively, together with n' freely added points.
- 7) $L_{n_1; n'}$ with $n_1 \geq 3$ and $n' \geq 1$, consisting of one n_1 -point line together with n' freely added points.
- 8) The uniform matroid $U_{3, n'}$ with $n' \geq 3$, consisting of n' points and no non-trivial lines.

We can now determine which of these matroids are nice:

Class 1. The smallest case $C_{3,3,3;0} \simeq \mathcal{W}^3$ is not nice (Example 10.3 above). And since, starting from any larger matroid $C_{n_1, n_2, n_3; n'}$ one can obtain $C_{3,3,3;0}$ by deleting elements, it follows that all such matroids are non-nice.

Class 2. The matroid $D_{n_1, n_2; n'}$ is nice when $n' = 0$ or 1 (Example 10.7). If $n' \geq 2$, the smallest case is $D_{3,3;2} \simeq F_7^{-5}$, which is not nice (Example 10.8). Therefore, all cases with $n' \geq 2$ are non-nice.

Class 3. The smallest case is $E_{3,3,3;0} \simeq P_7''$, which is not nice (Example 10.15).

Class 4. The smallest case is $F_{3,3,3;0}$, which is not nice; the argument is similar to that of Example 10.5 but is longer.

Class 5. The matroid $L_{n_1, n_2, n_3; 0}$ is nice (Example 10.4). All matroids $L_{n_1, n_2, n_3; n'}$ with $n' \geq 1$ are not nice (Example 10.5).

Class 6. By deletion from $L_{n_1, n_2, n_3; 0}$, we can conclude that $L_{n_1, n_2; n'}$ is nice when $n' \leq 2$. The smallest remaining case is $L_{3,3;3}$, which is not nice; the argument is similar to that of Example 10.5 but is longer.

Class 7. By deletion from $L_{n_1, n_2, n_3; 0}$, we can conclude that $L_{n_1; n'}$ is nice when $n' \leq 4$. The smallest remaining case is $L_{3;5}$, which is not nice (Example 10.6).

Class 8. All uniform matroids are nice (Example 10.1).

10.5 A wild speculation

Not all transversal matroids are nice; but this means only that our *method for proving* the half-plane property fails in these cases, not that the half-plane property itself fails. Indeed, we do not know a single example of a transversal matroid that fails the half-plane property. Moreover, we have conducted extensive numerical experiments (see Section 12) on the rank-3 transversal matroids discussed in the previous

subsection, and it seems plausible that all rank-3 transversal matroids have the half-plane property. Might *all* transversal matroids (and hence all gammoids) have the half-plane property? Unfortunately, we have no idea how to prove this.

Remark. The strongly base-orderable matroids [40, 11] form a minor-closed class that contains all the transversal matroids; so one might entertain the even stronger conjecture that “every strongly base-orderable matroid has the half-plane property”. However, this conjecture is false, since the matroid F_7^{-3} is strongly base-orderable but does not have the half-plane property (Example 11.7 below). To see that F_7^{-3} is strongly base-orderable, it suffices to note that a rank-3 matroid is strongly base-orderable \iff it is base-orderable \iff it has no restriction isomorphic to $M(K_4)$ [40].

11 Counterexamples

In this section we use Proposition 5.2(a) \implies (b) to show that certain polynomials do *not* have the half-plane property.

Example 11.1. Consider the Fano matroid F_7 with the ground set numbered as shown in Figure 6 (see Appendix A). Its basis set $\mathcal{B}(F_7)$ consists of all 3-element subsets of $[7]$ except $\{1, 2, 3\}$, $\{3, 4, 5\}$, $\{1, 5, 6\}$, $\{1, 4, 7\}$, $\{2, 5, 7\}$, $\{3, 6, 7\}$ and $\{2, 4, 6\}$. Let P be the basis generating polynomial $P_{\mathcal{B}(F_7)}$. If we take $x = \chi_{\{1,2,4,5\}}$ and $y = \chi_{\{3,6,7\}}$ (where χ_A denotes characteristic function of the set A), we obtain $p_{x,y}(\zeta) = 4\zeta^3 + 12\zeta^2 + 12\zeta$, whose roots are $\zeta = 0, (-3 \pm \sqrt{3}i)/2$. So the Fano matroid does not possess the half-plane property.

In fact, the Fano matroid is minor-minimal for failing to have the half-plane property. Indeed, it follows from Proposition 10.4 that *every* 7-element matroid lacking the half-plane property is minor-minimal. \square

Example 11.2. Consider next the non-Fano matroid F_7^- (Figure 6), which is obtained from F_7 by relaxing the circuit-hyperplane $\{2, 4, 6\}$, so that $\mathcal{B}(F_7^-) = \mathcal{B}(F_7) \cup \{\{2, 4, 6\}\}$. With the same choices of x and y , we obtain $p_{x,y}(\zeta) = 4\zeta^3 + 13\zeta^2 + 12\zeta$, whose roots are $\zeta = 0, (-13 \pm \sqrt{23}i)/8$. So the non-Fano matroid does not possess the half-plane property either; and, as noted above, it is minor-minimal. \square

Example 11.3. Consider next the matroid F_7^{--} obtained from F_7^- by relaxing $\{1, 4, 7\}$ (Figure 6). Choosing $x = \chi_{\{1,4,7\}}$ and $y = \chi_{\{2,3,5,6\}}$, we obtain $p_{x,y}(\zeta) = \zeta^3 + 12\zeta^2 + 13\zeta + 4$, whose roots are $\zeta \approx -10.834170$ and $\zeta \approx -0.582915 \pm 0.171501i$. So F_7^{--} does not possess the half-plane property either; and, as noted above, it is minor-minimal. \square

Example 11.4. A similar approach shows that the matroid $M(K_4) + e$ obtained from F_7^{--} by relaxing $\{3, 4, 5\}$ (Figure 6) also fails to possess the half-plane property.

But we can no longer choose x and y to be characteristic functions. Instead, let us take a unified approach to F_7 , F_7^- , F_7^{--} and $M(K_4) + e$ that gives additional insight into *why* the half-plane property fails.

Let us start with the graphic matroid $M(K_4)$, which of course *does* have the half-plane property (by Theorem 1.1). Let K_4 have vertex set $\{1, 2, 3, 4\}$. If we take $x = \chi_{\{12,13,14\}}$ and $y = \chi_{\{23,24,34\}}$, we find $p_{x,y}(\zeta) \equiv P_{\mathcal{B}(M(K_4))}(\zeta x + y) = \zeta(\zeta + 3)^2$, which has a double root at $\zeta = -3$. So $M(K_4)$ satisfies the condition of Proposition 5.2(b), but “just barely”: by suitable perturbations we may be able to split the double root into a pair of complex-conjugate roots, thereby proving the failure of the half-plane property for the perturbed matroid.

In the matroids F_7 *et al.*, let us consider 4 to be the “new element”, i.e. $F_7 \setminus 4 \simeq F_7^- \setminus 4 \simeq \dots \simeq M(K_4)$. Then the above choice of x and y corresponds to taking y to be the characteristic function of a 3-point line in $M(K_4)$ and x to be the characteristic function of the complementary set, e.g. $x = \chi_{\{1,2,5\}}$ and $y = \chi_{\{3,6,7\}}$. So let us perturb this slightly, by taking $x = \chi_{\{1,2,5\}} + \epsilon \chi_{\{4\}}$ and $y = \chi_{\{3,6,7\}} + a \epsilon \chi_{\{4\}}$ with $\epsilon, a \geq 0$. (Our previous choices for F_7 and F_7^- correspond to $\epsilon = 1$ and $a = 0$.) We then have:

$$\begin{aligned}
F_7: \quad p_{x,y}(\zeta) &= (1 + 3\epsilon)\zeta^3 + [6 + (6 + 3a)\epsilon]\zeta^2 + [9 + (3 + 6a)\epsilon]\zeta + 3a\epsilon \\
&\quad \text{Roots } \zeta = -(a/3)\epsilon + O(\epsilon^{3/2}) \text{ and } \zeta = -3 \pm 2\sqrt{a-3}\epsilon^{1/2} + O(\epsilon). \\
F_7^-: \quad p_{x,y}(\zeta) &= (1 + 3\epsilon)\zeta^3 + [6 + (7 + 3a)\epsilon]\zeta^2 + [9 + (3 + 7a)\epsilon]\zeta + 3a\epsilon \\
&\quad \text{Roots } \zeta = -(a/3)\epsilon + O(\epsilon^{3/2}) \text{ and } \zeta = -3 \pm \sqrt{3(a-3)}\epsilon^{1/2} + O(\epsilon). \\
F_7^{--}: \quad p_{x,y}(\zeta) &= (1 + 3\epsilon)\zeta^3 + [6 + (8 + 3a)\epsilon]\zeta^2 + [9 + (3 + 8a)\epsilon]\zeta + 3a\epsilon \\
&\quad \text{Roots } \zeta = -(a/3)\epsilon + O(\epsilon^{3/2}) \text{ and } \zeta = -3 \pm \sqrt{2(a-3)}\epsilon^{1/2} + O(\epsilon). \\
M(K_4) + e: \quad p_{x,y}(\zeta) &= (1 + 3\epsilon)\zeta^3 + [6 + (9 + 3a)\epsilon]\zeta^2 + [9 + (3 + 9a)\epsilon]\zeta + 3a\epsilon \\
&\quad \text{Roots } \zeta = -(a/3)\epsilon + O(\epsilon^{3/2}) \text{ and } \zeta = -3 \pm \sqrt{a-3}\epsilon^{1/2} + O(\epsilon).
\end{aligned}$$

So any choice of $a \in [0, 3)$ will yield nonreal roots for small $\epsilon > 0$ for all four matroids.

Let us conclude this example by remarking that a second (nonisomorphic) choice of x, y in $M(K_4)$ also yields a double root: for $x = \chi_{\{12,34\}}$ and $y = \chi_{\{13,14,23,24\}}$, we find $p_{x,y}(\zeta) = 4(\zeta + 1)^2$, which has a double root at $\zeta = -1$. An analogous perturbation of this choice also yields nonreal roots for F_7 *et al.*, provided that $a \in [0, 1)$. \square

Example 11.5. Consider the polynomial

$$P_\mu(z_1, \dots, z_7) = P_{\mathcal{B}(F_7)}(z) + \mu z_2 z_4 z_6. \quad (11.1)$$

For $\mu = 0$ (resp. $\mu = 1$) this is the basis generating polynomial of F_7 (resp. F_7^-). For $\mu = 4$ it is the polynomial $Q_A(z) = \det(AZ A^T)$ obtained from $Z = \text{diag}(z_1, \dots, z_7)$ and the matrix

$$A = \begin{bmatrix} 1 & 1 & 0 & 0 & 0 & 1 & 1 \\ 0 & 1 & 1 & 1 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 & 1 & 1 & 1 \end{bmatrix}, \quad (11.2)$$

which represents F_7^- over any field of characteristic $\neq 2$ (in particular, over \mathbb{C}); by Theorem 8.1, Q_A has the half-plane property. As in Example 11.4, let us choose $x = \chi_{\{1,2,5\}} + \epsilon\chi_{\{4\}}$ and $y = \chi_{\{3,6,7\}} + a\epsilon\chi_{\{4\}}$ with $\epsilon, a \geq 0$. We then obtain $p_{x,y}(\zeta) = (1 + 3\epsilon)\zeta^3 + [6 + (6 + \mu + 3a)\epsilon]\zeta^2 + [9 + (3 + 6a + \mu a)\epsilon]\zeta + 3a\epsilon$, whose roots are $\zeta = -(a/3)\epsilon + O(\epsilon^{3/2})$ and $\zeta = -3 \pm \sqrt{(4 - \mu)(a - 3)} \epsilon^{1/2} + O(\epsilon)$. So, if $-\infty < \mu < 4$ (resp. $\mu > 4$), then any choice $0 \leq a < 3$ (resp. $a > 3$) will yield nonreal roots for small $\epsilon > 0$. So P_μ has the half-plane property *only* for $\mu = 4$! \square

Example 11.6. Consider, more generally, the polynomial

$$P_{\mu,\nu,\rho}(z_1, \dots, z_7) = P_{\mathcal{B}(F_7)}(z) + \mu z_2 z_4 z_6 + \nu z_1 z_4 z_7 + \rho z_3 z_4 z_5. \quad (11.3)$$

For $(\mu, \nu, \rho) = (0, 0, 0)$ this is $P_{\mathcal{B}(F_7)}$; for $(\mu, \nu, \rho) = (1, 0, 0)$ it is $P_{\mathcal{B}(F_7^-)}$; for $(\mu, \nu, \rho) = (1, 1, 0)$ it is $P_{\mathcal{B}(F_7^{--})}$; for $(\mu, \nu, \rho) = (1, 1, 1)$ it is $P_{\mathcal{B}(M(K_4)+e)}$.

Some cases of the polynomial $P_{\mu,\nu,\rho}$ are of the form $Q_A(z) = \det(AZA^*)$ [where $Z = \text{diag}(z_1, \dots, z_7)$] for a suitable complex matrix A , and thus have the half-plane property by Theorem 8.1. Consider, for instance, the matrix

$$A = \begin{bmatrix} 1 & 1 & 0 & a & 0 & 1 & 1 \\ 0 & 1 & 1 & 1 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 & 1 & b & b \end{bmatrix} \quad (11.4)$$

For $a = 0$ and $b = e^{i\theta}$, we have $Q_A = P_{\mu,\nu,\rho}$ with $(\mu, \nu, \rho) = (2 + 2\cos\theta, 2 - 2\cos\theta, 0)$ [that is, $\mu, \nu \geq 0$ with $\mu + \nu = 4$ and $\rho = 0$].¹⁶ For $a = e^{\pm\pi i/3}$ and $b = 1$, we have $Q_A = P_{\mu,\nu,\rho}$ with $(\mu, \nu, \rho) = (3, 0, 1)$; and for $a = e^{\pm\pi i/3}$ and $b = e^{\mp 2\pi i/3}$, we have $Q_A = P_{\mu,\nu,\rho}$ with $(\mu, \nu, \rho) = (0, 3, 1)$. Moreover, by appropriately permuting columns of the matrix A (*without* permuting the column labels), the triplet (μ, ν, ρ) can be permuted at will.¹⁷ We do not know whether any additional cases of $P_{\mu,\nu,\rho}$ can be obtained from a determinant.

For $0 \leq \mu, \nu, \rho \leq 2$ with $\mu + \nu + \rho = 4$, the polynomial $P_{\mu,\nu,\rho}$ arises from the principal extension of $F_7 \setminus 4 \simeq M(K_4)$ by the new element 4 with weights

$$\lambda_2 = \lambda_6 = 1 - \mu/2 \quad (11.5a)$$

$$\lambda_1 = \lambda_7 = 1 - \nu/2 \quad (11.5b)$$

$$\lambda_3 = \lambda_5 = 1 - \rho/2 \quad (11.5c)$$

and thus has the half-plane property by Proposition 4.11.

Finally, as in Example 11.5, let us choose $x = \chi_{\{1,2,5\}} + \epsilon\chi_{\{4\}}$ and $y = \chi_{\{3,6,7\}} + a\epsilon\chi_{\{4\}}$ with $\epsilon, a \geq 0$. We then obtain the same $p_{x,y}$ as in Example 11.5, except that

¹⁶ We remark that the matrix A represents F_7^{--} over any field of at least four elements, provided that $a = 0$ and $b \notin \{0, 1, -1\}$.

¹⁷ This is because there are automorphisms of F_7 that realize an arbitrary permutation of the lines 246, 147 and 345.

μ is replaced by $\mu + \nu + \rho$. So we obtain nonreal roots for small $\epsilon > 0$ whenever $\mu + \nu + \rho \neq 4$. It follows (using also Theorem 6.1) that $P_{\mu,\nu,\rho}$ has the half-plane property *only* when $\mu, \nu, \rho \geq 0$ with $\mu + \nu + \rho = 4$. But we do not know whether these necessary conditions are sufficient. \square

Example 11.7. Consider the matroid F_7^{-3} obtained from F_7^- by relaxing the circuit-hyperplanes $\{2, 5, 7\}$ and $\{3, 6, 7\}$ (Figure 6). Let us take $x = \chi_{\{1,4,5\}} + \epsilon\chi_{\{2\}}$ and $y = \chi_{\{3,6,7\}}$. Then $p_{x,y}(\zeta) = (1 + 3\epsilon)\zeta^3 + (6 + 8\epsilon)\zeta^2 + (9 + 3\epsilon)\zeta + 1$, which has nonreal roots whenever $0.090685 \lesssim \epsilon \lesssim 0.494485$. So F_7^{-3} does not have the half-plane property; and, as noted above, it is minor-minimal.

Note, finally, that F_7^{-3} is a relaxation of P_7 , which is a $\sqrt[6]{1}$ -matroid and hence has the half-plane property (see Section 8). So relaxation does *not* in general preserve the half-plane property. \square

Example 11.8. Consider the rank-4 matroid P_8 represented over any field of characteristic $\neq 2$ by the matrix

$$\left[\begin{array}{cccc|cccc} 1 & 0 & 0 & 0 & 0 & 1 & 1 & 2 \\ 0 & 1 & 0 & 0 & 1 & 0 & 1 & 1 \\ 0 & 0 & 1 & 0 & 1 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & 2 & 1 & 1 & 0 \end{array} \right] \quad (11.6)$$

Its basis set $\mathcal{B}(P_8)$ consists of all 4-element subsets of $[8]$ except $\{1, 2, 3, 8\}$, $\{1, 2, 4, 7\}$, $\{1, 3, 4, 6\}$, $\{2, 3, 4, 5\}$, $\{1, 4, 5, 8\}$, $\{2, 3, 6, 7\}$, $\{1, 5, 6, 7\}$, $\{2, 5, 6, 8\}$, $\{3, 5, 7, 8\}$ and $\{4, 6, 7, 8\}$. If we take $x = \chi_{\{1,4,5,8\}}$ and $y = \chi_{\{2,3,6,7\}}$, we obtain $p_{x,y}(\zeta) = 16\zeta^3 + 28\zeta^2 + 16\zeta$, whose roots are $\zeta = 0, (-7 \pm \sqrt{15}i)/8$. So P_8 does not possess the half-plane property. Moreover, it is minor-minimal, since all the single-element contractions (resp. deletions) of P_8 are isomorphic to P_7 (resp. P_7^*), which are $\sqrt[6]{1}$ -matroids and hence have the half-plane property.

Observe now that $\{1, 4, 5, 8\}$ and $\{2, 3, 6, 7\}$ form the unique pair of disjoint circuit-hyperplanes in P_8 . Let us denote by P'_8 the matroid obtained from P_8 by relaxing one of these circuit-hyperplanes (say, $\{1, 4, 5, 8\}$), and by P''_8 the matroid obtained by relaxing both of them. Let us make the same choices of x and y as for P_8 . Then, for P'_8 one has $p_{x,y}(\zeta) = \zeta^4 + 16\zeta^3 + 28\zeta^2 + 16\zeta$, whose roots are $\zeta = 0$ and $\zeta \approx -14.093869, -0.953065 \pm 0.476353i$. And for P''_8 one has $p_{x,y}(\zeta) = \zeta^4 + 16\zeta^3 + 28\zeta^2 + 16\zeta + 1$, whose roots are $\zeta \approx -14.093459, -0.070955, -0.917793 \pm 0.397059i$. So neither P'_8 nor P''_8 possesses the half-plane property. We suspect (but are unable to prove) they are minor-minimal: all the single-element contractions (resp. deletions) of P'_8 are isomorphic to P_7 or P'_7 (resp. P_7^* or $(P'_7)^*$), and all the single-element contractions (resp. deletions) of P''_8 are isomorphic to P'_7 (resp. $(P'_7)^*$); and our numerical results (Section 12) suggest that P'_7 (and hence also its dual) has the half-plane property. \square

Example 11.9. Consider the Pappus and non-Pappus matroids (Figure 9 in Appendix A). Take $x = 2\chi_{\{1\}} + \chi_{\{3,4,6,7\}} + \epsilon\chi_{\{9\}}$ and $y = \chi_{\{2,5,8\}}$. We then have:

$$\begin{aligned}
\text{Pappus: } p_{x,y}(\zeta) &= (16 + 14\epsilon)\zeta^3 + (33 + 15\epsilon)\zeta^2 + (18 + 3\epsilon)\zeta + 1 \\
&\text{Roots } \zeta = -\frac{1}{16} + O(\epsilon) \text{ and } \zeta = -1 \pm (i\sqrt{2/15})\epsilon^{1/2} + O(\epsilon). \\
\text{non-Pappus: } p_{x,y}(\zeta) &= (16 + 14\epsilon)\zeta^3 + (33 + 16\epsilon)\zeta^2 + (18 + 3\epsilon)\zeta + 1 \\
&\text{Roots } \zeta = -\frac{1}{16} + O(\epsilon) \text{ and } \zeta = -1 \pm (i/\sqrt{15})\epsilon^{1/2} + O(\epsilon).
\end{aligned}$$

So the Pappus and non-Pappus matroids do not have the half-plane property. We do not know whether they are minor-minimal; but we suspect that they are, since our numerical experiments (Section 12) suggest that their two non-isomorphic deletions, $\text{non-Pappus}\backslash 1$ and $\text{Pappus}\backslash e \simeq \text{non-Pappus}\backslash 9$, do have the half-plane property. (Of course, all their contractions are rank-2 and hence have the half-plane property.) \square

It is worth noting that $\text{non-Pappus}\backslash 9$ is “borderline” for the half-plane property in the same sense that $M(K_4)$ is (Example 11.4), i.e. $p_{x,y}(\zeta)$ has a double root at $\zeta = -1$ for a suitable choice of x, y . This fact suggests that other extensions of $\text{non-Pappus}\backslash 9$ might also fail to have the half-plane property. This is indeed the case:

Example 11.10. Consider the matroid $(\text{non-Pappus}\backslash 9) + e$, obtained from $\text{non-Pappus}\backslash 9$ by adding a new element freely (let us call this new element 9). Then its bases are those of non-Pappus plus $\{2, 6, 9\}$ and $\{3, 5, 9\}$. Take $x = 2\chi_{\{1\}} + \chi_{\{3,4,6,7\}}$ and $y = \chi_{\{2,5,8\}} + \epsilon\chi_{\{9\}}$. We then have:

$$\begin{aligned}
(\text{non-Pappus}\backslash 9) + e: p_{x,y}(\zeta) &= 16\zeta^3 + (33 + 14\epsilon)\zeta^2 + (18 + 18\epsilon)\zeta + (1 + 3\epsilon) \\
&\text{Roots } \zeta = -\frac{1}{16} + O(\epsilon) \text{ and } \zeta = -1 \pm (i/\sqrt{15})\epsilon^{1/2} + O(\epsilon).
\end{aligned}$$

So $(\text{non-Pappus}\backslash 9) + e$ does not have the half-plane property. \square

12 Numerical experiments

Given a homogeneous multiaffine polynomial $P(x_1, \dots, x_n)$, we have searched numerically for counterexamples to the half-plane property using two methods:

Elementary method. Choose x_1, \dots, x_{n-1} uniformly at random in the rectangle $(0, 1) + (-1, 1)i$; then solve $P(x_1, \dots, x_n) = 0$ for x_n . If $\text{Re } x_n > 0$, we have found a counterexample to the half-plane property. (Thanks to homogeneity, there is no loss of generality in choosing x_1, \dots, x_{n-1} in the specified rectangular subset of \mathbb{C} .)

Method using Proposition 5.2. Choose vectors a, b uniformly at random in $[0, 1]^n$, and compute the roots of the univariate polynomial $p_{a,b}(\zeta) = P(\zeta a + b)$. If at least one of these roots has a nonzero imaginary part, we have found a counterexample to the half-plane property. (Thanks to homogeneity, there is no loss of generality in choosing a, b in $[0, 1]^n \subset \mathbb{R}^n$.)

We first applied these methods to some matroids for which the half-plane property is known to fail (F_7, F_7^-, \dots), in order to get a rough feeling for the rate of finding counterexamples. We then applied them to some matroids for which the half-plane

Matroid	Elementary Method		Method using Proposition 5.2	
	# tries	# counterexamples	# tries	# counterexamples
F_7	10^8	4152	10^6	81566
F_7^-	10^8	687	10^6	42620
F_7^{--}	10^8	34	10^6	12794
$M(K_4) + e$	10^8	0	10^6	1060
F_7^{-3}	10^8	0	10^6	695
F_7^{-4}	10^8	0	10^8	0
$\mathcal{W}^3 + e$	10^8	0	10^8	0
\mathcal{W}^{3+}	10^8	0	10^8	0
P'_7	10^8	0	10^8	0
$F_7^{-4} + e$	—	—	10^7	0
$\mathcal{W}^3 + e + f$	—	—	10^7	0
$\mathcal{W}^{3+} + e$	—	—	10^7	0
$P'_7 + e$	—	—	10^7	0
P_8	10^8	278	10^6	10930
P'_8	10^8	114	10^6	5590
P''_8	10^8	40	10^6	2723
V_8 (Vámos)	10^8	0	10^7	0
Pappus	10^8	0	10^6	544
non-Pappus	10^8	0	10^6	17
(non-Pappus $\setminus 9$) + e	10^8	0	10^7	6
non-Pappus $\setminus 1$	10^8	0	10^7	0
non-Pappus $\setminus 9$	10^8	0	10^8	0
$C_{3,3,3;3}$	10^7	0	10^7	0
$C_{4,4,4;4}$	10^7	0	10^6	0
$C_{5,5,5;5}$	10^7	0	10^6	0
$C_{6,6,6;6}$	10^6	0	10^5	0
$C_{7,7,7;7}$	10^6	0	10^5	0

Table 1: Results of numerical experiments to test the half-plane property.

property is an open question (F_7^{-4} , \mathcal{W}^{3+e} , \mathcal{W}^{3+} , P'_7 , \dots). The results of our numerical experiments are shown in Table 1. As can be seen, the method using Proposition 5.2 is vastly more powerful than the elementary method: in those cases where the half-plane property fails, the proportion of counterexamples is larger by a factor of ≈ 2000 – 20000 or even more. Roughly speaking, one pair a, b in Proposition 5.2 corresponds to a large set of counterexamples in the elementary method. Indeed, if we had used only the elementary method we would have failed to detect the failure of the half-plane property for $M(K_4) + e$, F_7^{-3} , Pappus, non-Pappus and $(\text{non-Pappus} \setminus 9) + e$, even with 10^8 tries.

Our numerical results suggest, first of all, that the matroids F_7^{-4} , $\mathcal{W}^3 + e$, \mathcal{W}^{3+} , P'_7 and their free extensions, as well as V_8 , $\text{non-Pappus} \setminus 1$ and $\text{non-Pappus} \setminus 9$, probably *do* all have the half-plane property. One urgent problem, therefore, is to try to prove (or disprove) these alleged facts; the proofs may well require new tools. We find it particularly surprising that the Vámos matroid V_8 has (or appears to have) the half-plane property: since this matroid is not representable over any field (much less over \mathbb{C}), neither the determinant condition nor the permanent condition applies to it — nor can the constructions in Section 4 lead to it, starting from a polynomial whose support is the collection of bases of a representable matroid — so we have no idea why it should have even the *weak* half-plane property.

Finally, we made a systematic study of the half-plane property for rank-3 transversal matroids, all of which are restrictions of $C_{n,n,n;n}$ for some $n \geq 3$ (see Section 10.4). The results in Table 1 strongly suggest that *all* rank-3 transversal matroids have the half-plane property. And they suggest the bold conjecture that perhaps all transversal matroids of *any* rank have the half-plane property.

13 Open questions

We conclude by presenting some open questions raised by, or related to, the results of this paper.

13.1 The strong half-plane property

Let us say that a polynomial P in n complex variables has the *strong half-plane property* if

$$\partial_x^{\mathbf{m}} |P(x + iy)|^2 \Big|_{x=0} \geq 0 \quad (13.1)$$

for all multi-indices \mathbf{m} and all $y \in \mathbb{R}^n$. It is easy to show that (13.1) implies the half-plane property: Fix $y \in \mathbb{R}^n$; then either $\partial_x^{\mathbf{m}} |P(x + iy)|^2 \Big|_{x=0} > 0$ for at least one multi-index \mathbf{m} , which implies $P(x + iy) \neq 0$ for all vectors $x > 0$, or else $\partial_x^{\mathbf{m}} |P(x + iy)|^2 \Big|_{x=0} = 0$ for all \mathbf{m} , which implies $P(x + iy) = 0$ for all vectors $x \geq 0$ and hence $P \equiv 0$ by analytic continuation from a real environment.

Question 13.1 *Does the half-plane property imply the strong half-plane property?*

It is easy to see that the answer is affirmative when $n = 1$ (just factor P); hence it is also true for general n if one considers only multi-indices \mathbf{m} with a single nonzero component. But we do not know whether it is true in general. This question was inspired by Newman’s “strong Lee–Yang theorem” [54, Section 3].

13.2 Same-phase theorem

Question 13.2 *Can the same-phase theorem (Theorems 6.1 and 6.2) be extended to a larger class of polynomials?*

The Fettweis–Basu lemma (Lemma 3.5), applied in many variables successively, may be a useful tool.

13.3 Support of polynomials with the half-plane property

In Theorems 7.1 and 7.2 we provided some *necessary* conditions for a subset $S \subseteq \mathbb{N}^E$ to be the support of a homogeneous polynomial with the half-plane property. It is natural to seek a generalization to the non-homogeneous case:

Problem 13.3 *Find necessary conditions for a subset $S \subseteq \mathbb{N}^E$ to be the support of a polynomial with the half-plane property.*

Two special cases of this problem were posed earlier:

Question 13.4 (= Question 7.4) *If P is multiaffine and has the half-plane property, is $\text{supp}(P)$ a delta-matroid? What if P also has the same-phase property?*

Question 13.5 (= Question 7.5) *Assume that P has the half-plane property and has definite parity (i.e. $\mathbf{m}, \mathbf{m}' \in \text{supp}(P)$ implies that $|\mathbf{m}| \equiv |\mathbf{m}'| \pmod{2}$). Is $\text{supp}(P)$ then a jump system? (Recall from Theorem 6.2 that all such polynomials have the same-phase property.)*

More generally, one can pose the probably-much-more-difficult problem:

Problem 13.6 *Find necessary and sufficient conditions for a subset $S \subseteq \mathbb{N}^E$ to be the support of a polynomial with the half-plane property.*

Already in the homogeneous multiaffine case this question looks difficult, and constitutes the converse of Theorem 7.1:

Question 13.7 (= Question 7.6) *Does every matroid M have the weak half-plane property? And if not, which ones do? (For instance, does the Fano matroid F_7 have the weak half-plane property?)*

By Corollary 8.2, every matroid representable over \mathbb{C} has the weak half-plane property. But we know *nothing* beyond this. Indeed, both of our *ab initio* methods for constructing polynomials with the half-plane property — the determinant method (Theorem 8.1) and the permanent method (Theorem 10.2) — lead always to polynomials P whose support matroid is \mathbb{C} -representable; and all the constructions of Section 4, applied to a polynomial P whose support matroid is \mathbb{C} -representable, lead to another polynomial with the same property.¹⁸ So even the following is an open problem:

Problem 13.8 *Construct a polynomial P with the half-plane property whose support matroid is not representable over \mathbb{C} . (Or prove that it is impossible.)*

Note that our numerical results (Section 12) suggest that the Vámos matroid V_8 , which is not representable over any field, does have the half-plane property.

Theorems 7.1 and 7.2 have the general form: if $P(x) = \sum_{\mathbf{m}} a_{\mathbf{m}} x^{\mathbf{m}}$ has the half-plane property and certain coefficients $a_{\mathbf{m}}$, $a_{\mathbf{m}'}$ are nonzero, then so are (one or more of) certain other coefficients $a_{\mathbf{m}''}$. It is natural to ask whether this qualitative result can be extended to a *quantitative* lower bound on some combination of the corresponding coefficients $a_{\mathbf{m}''}$.

Question 13.9 *Can Theorems 7.1 and 7.2 be extended to quantitative inequalities on the coefficients?*

This question is vaguely reminiscent of Kung’s discussion of the heuristic analogy between determinantal identities and basis-exchange properties in matroids [44, 45]. One approach might be to use Proposition 5.2(a) \implies (b); if successful, this would be a nice generalization of Theorem 5.3 to the higher-rank case.

13.4 (F, G) -representability of matroids

Question 13.10 (= Question 8.5) *When $-1 \notin G$, is the class of (F, G) -representable matroids always closed under contraction?*

Question 13.11 *If $-1 \notin G$, does weak (F, G) -representability necessarily imply (F, G) -representability?*

¹⁸ The operations of Sections 4.1–4.4, applied to a matroid that is representable over a field F , always produce another F -representable matroid: see [55, Proposition 3.2.4] for deletion and contraction, [55, Corollary 2.2.9] for duality, [55, Proposition 4.2.15] for direct sum, [55, Proposition 7.1.21] for parallel connection and series connection, and [55, Proposition 7.1.23] for 2-sum. The other operations of Section 4 (principal extension, truncation, coextension and cotruncation, and full-rank matroid union) can lead out of the class of F -representable matroids, but always produce a matroid that is representable over some finite extension field of F : see [14, Propositions 7.3.5 and 7.4.17] and [59]. In particular, applied to a \mathbb{C} -representable matroid, they always yield another \mathbb{C} -representable matroid.

Question 13.12 *Is Theorem 8.9 the prototype of a more general theorem asserting that, in certain cases, every (F, G) -representable matroid is in fact (F, G') -representable for some specified subgroup $G' \subsetneq G$?*

13.5 Determinant class, permanent class, and the half-plane property

In this paper we have given two distinct methods for constructing polynomials with the half-plane property: the determinant construction (Theorem 8.1) and the permanent construction (Theorem 10.2). It is natural to ask what is the relation (if any) between these two classes of polynomials, and between them and the class of all polynomials with the half-plane property. Let us pose this question as follows:

Fix integers r and n ($0 \leq r \leq n$), and let $\mathcal{P}_{r,n}$ be the set of degree- r homogeneous multiaffine polynomials in n variables with complex coefficients, $P(x) = \sum_{S \subseteq [n], |S|=r} a_S x^S$. Let $\mathcal{P}_{r,n}^+$ be the subset of $\mathcal{P}_{r,n}$ consisting of polynomials with non-negative coefficients. By identifying a polynomial with its coefficients, the spaces $\mathcal{P}_{r,n}$ and $\mathcal{P}_{r,n}^+$ can be thought of as $\mathbb{C}^{\binom{n}{r}}$ and $[0, \infty)^{\binom{n}{r}}$, respectively. Let us now define three subsets of $\mathcal{P}_{r,n}^+$ (thanks to Theorem 6.1, there is no loss of generality in restricting attention to $\mathcal{P}_{r,n}^+$):

- $\mathcal{H}_{r,n}$: the polynomials with the half-plane property.
- $\mathcal{D}_{r,n}$: the polynomials of “determinant class”, i.e. $P(x) = \det(AXA^*)$ [or equivalently $a_S = |\det(A \upharpoonright S)|^2$] for some $r \times n$ complex matrix A .
- $\mathcal{M}_{r,n}$: the polynomials of “permanent (or matching) class”, i.e. $P(x) = \text{per}(\Lambda X)$ [or equivalently $a_S = \text{per}(\Lambda \upharpoonright S)$] for some $r \times n$ nonnegative matrix Λ .

We have shown that $\mathcal{P}_{r,n}^+ \supseteq \mathcal{H}_{r,n} \supseteq \mathcal{D}_{r,n} \cup \mathcal{M}_{r,n}$, and these containments are in general strict (see Example 13.1 below).

Question 13.13 *What is the relation between the spaces $\mathcal{P}_{r,n}^+$, $\mathcal{H}_{r,n}$, $\mathcal{D}_{r,n}$ and $\mathcal{M}_{r,n}$?*

Most ambitiously, we can ask for a complete characterization of the determinant class $\mathcal{D}_{r,n}$ and the permanent class $\mathcal{P}_{r,n}$:

Problem 13.14 *Find necessary and sufficient conditions for a set of nonnegative real numbers $\{a_S\}_{S \subseteq [n], |S|=r}$ to be representable in the form $a_S = |\det(A \upharpoonright S)|^2$ for some $r \times n$ complex matrix A .*

Problem 13.15 *Find necessary and sufficient conditions for a set of nonnegative real numbers $\{a_S\}_{S \subseteq [n], |S|=r}$ to be representable in the form $a_S = \text{per}(\Lambda \upharpoonright S)$ for some $r \times n$ nonnegative matrix Λ .*

For the analogous problem with $a_S = \det(A \upharpoonright S)$, the necessary and sufficient condition is given by the Grassmann–Plücker syzygies [75, Proposition 1.6.1] [6, Section 1.2]. But the presence of the modulus-square seems to make Problem 13.14 quite difficult. Indeed, it seems nontrivial even in the rank-2 case. Here is an even more special case that illustrates some of the complexity:

Example 13.1. Consider the homogeneous degree-2 polynomial in $n \geq 3$ variables $P(z) = \frac{1}{2} \sum_{i,j=1}^n a_{ij} z_i z_j$ where

$$a_{ij} = \begin{cases} \mu & \text{if } (i, j) = (1, 2) \text{ or } (2, 1) \\ 1 & \text{in all other cases of } i \neq j \\ 0 & \text{if } i = j \end{cases} \quad (13.2)$$

One then finds, after some calculation, that:

- P has the half-plane property if and only if $0 \leq \mu \leq (2n - 4)/(n - 3)$.¹⁹
- P belongs to the determinant class if and only if:
 - For $n = 3$, $0 \leq \mu < \infty$.
 - For $n = 4$, $0 \leq \mu \leq 4$.
 - For $n = 5$, $\mu = 0$ or $\mu = 3$.
 - For $n \geq 6$, never.
- P belongs to the permanent class if and only if $0 \leq \mu \leq 2$.²⁰

□

¹⁹ SKETCH OF PROOF: Let A be the matrix defined in (13.2). Then it can be shown by induction on n that

$$\det(\lambda I - A) = (\lambda + 1)^{n-3}(\lambda + \mu)[\lambda^2 - (\mu + n - 3)\lambda + (n - 3)\mu - (2n - 4)].$$

So A has $n - 3$ eigenvalues -1 , one eigenvalue $-\mu$, and a pair of eigenvalues

$$\frac{\mu + n - 3 \pm \sqrt{\mu^2 - 2(n - 3)\mu + (n^2 + 2n - 7)}}{2}.$$

The claim then follows from Theorem 5.3.

²⁰ SKETCH OF PROOF: Suppose that there exist nonnegative numbers $\{\lambda_i^{(1)}\}_{i=1}^n$ and $\{\lambda_i^{(2)}\}_{i=1}^n$ such that

$$a_{ij} = \lambda_i^{(1)} \lambda_j^{(2)} + \lambda_j^{(1)} \lambda_i^{(2)} \quad \text{for all } i \neq j \quad (*)$$

By considering separately the cases $\lambda_1^{(1)} = 0$ and $\lambda_1^{(1)} \neq 0$ (and likewise for $\lambda_1^{(2)}$), one shows in each case that $\lambda_3^{(1)} = \dots = \lambda_n^{(1)} \equiv \alpha$ and $\lambda_3^{(2)} = \dots = \lambda_n^{(2)} \equiv \beta$. Using this fact together with (*), one shows (after some algebra) that $0 \leq \mu \leq 2$, and that every μ in this interval is attainable.

One can ask also about general properties of the spaces $\mathcal{H}_{r,n}$, $\mathcal{D}_{r,n}$ and $\mathcal{M}_{r,n}$. For example, are they convex? The answer is no in general; indeed, a convex combination of polynomials in $\mathcal{D}_{r,n} \cap \mathcal{M}_{r,n}$ need not even lie in $\mathcal{H}_{r,n}$. To see this, just let P and Q be polynomials depending on disjoint subsets of variables, e.g. $P(x_1, \dots, x_4) = x_1x_2$ and $Q(x_1, \dots, x_4) = x_3x_4$, both of which lie in $\mathcal{D}_{2,4} \cap \mathcal{M}_{2,4}$; but by Corollary 6.5, $(P + Q)/2$ does not even have the half-plane property.

On the other hand, by Theorem 7.1 we can “stratify” $\mathcal{H}_{r,n}$ as $\mathcal{H}_{r,n} = \bigcup_M \mathcal{H}_{r,n}(M)$ where

$$\mathcal{H}_{r,n}(M) = \{P \in \mathcal{H}_{r,n} : \text{supp}(P) = \mathcal{B}(M)\} \quad (13.3)$$

and the union runs over all rank- r matroids M on the ground set $[n]$; and likewise for $\mathcal{D}_{r,n}$ and $\mathcal{M}_{r,n}$. One might then ask whether the sets $\mathcal{H}_{r,n}(M)$, $\mathcal{D}_{r,n}(M)$ and $\mathcal{M}_{r,n}(M)$ are convex for fixed M . But this too is false [at least for $\mathcal{H}_{r,n}(M)$ and $\mathcal{D}_{r,n}(M)$]: in fact, a convex combination of polynomials in $\mathcal{D}_{r,n}(M)$ need not even lie in $\mathcal{H}_{r,n}(M)$. To see this, let P be the polynomial of Example 13.1, and let Q be the analogous polynomial with the indices $\{1, 2\}$ replaced by $\{3, 4\}$. Then, for $n = 4$, P and Q belong to $\mathcal{D}_{2,4}(U_{2,4})$ for $0 < \mu \leq 4$; but an easy calculation using Theorem 5.3 shows that $(P + Q)/2$ has the half-plane property only for $0 \leq \mu \leq 3$.

Finally, analogous examples show that the sets $\mathcal{H}_{r,n}(M)$ and $\mathcal{D}_{r,n}(M)$ need not be log-convex or harmonic-mean-convex. For example, let P be the polynomial of Example 13.1, and let Q be the analogous polynomial with the indices $\{1, 2\}$ replaced by $\{1, 3\}$. Then, for $n = 4$, P and Q belong to $\mathcal{D}_{2,4}(U_{2,4})$ for $0 < \mu \leq 4$; but a straightforward (though messy) calculation using Theorem 5.3 shows that the polynomial R obtained from P and Q by coefficientwise geometric mean (resp. coefficientwise harmonic mean²¹) has the half-plane property only for $\mu \geq 1/16$ (resp. $\mu \geq 1/7$).

13.6 Half-plane property for transversal matroids

As noted in Section 10.5, we have proven the half-plane property only for a subclass of transversal matroids (the “nice” ones), but we do not know a single example of a transversal matroid that fails the half-plane property. Our numerical experiments on rank-3 transversal matroids (Section 12) suggest the following conjecture:

Conjecture 13.16 *All rank-3 transversal matroids have the half-plane property.*

If this conjecture is indeed true, its proof will very likely require new techniques, which could potentially shed light also on other related problems. More ambitiously, we can raise the following question:

Question 13.17 *Might all transversal matroids (and hence all gammoids) have the half-plane property?*

Either a proof or a counterexample would be of considerable interest.

²¹ The harmonic mean of a and b is $2ab/(a + b)$.

Have HPP	Do not have HPP	HPP unknown
Q_7 (Example 10.7)	F_7 (Example 11.1)	F_7^{-4}
S_7 (Examples 10.4 and 10.7)	F_7^- (Example 11.2)	$\mathcal{W}^3 + e$
$M(K_4)^+$ (Example 4.1 and Corollary 8.2)	F_7^{--} (Example 11.3)	\mathcal{W}^{3+}
F_7^{-5} (Example 10.12)	$M(K_4) + e$ (Example 11.4)	P_7'
F_7^{-6} (Example 10.4)	F_7^{-3} (Example 11.7)	
P_7 (Corollary 8.2)		
P_7'' (Examples 10.17 and 10.18)		
P_7''' (Examples 10.4 and 10.7)		
$U_{3,7}$ (Theorem 9.1)		

Table 2: The 7-element rank-3 3-connected matroids, divided according to whether or not they have the half-plane property (HPP).

13.7 Half-plane property for 7-element rank-3 matroids

We have shown that all matroids of rank or corank at most 2 have the half-plane property (Corollary 5.5), as do all matroids on a ground set of at most 6 elements (Proposition 10.4). So the first nontrivial case arises with 7-element rank-3 matroids; we would like to know which ones have, and which ones do not have, the half-plane property. In Table 2 we divide the 7-element rank-3 3-connected matroids (see Appendix A.2) into three categories: those we have proven to have the half-plane property, those we have proven not to have the half-plane property, and those for which we have no proof either way. There are exactly four matroids in the latter category: F_7^{-4} , $\mathcal{W}^3 + e$, \mathcal{W}^{3+} and P_7' . Our numerical experiments (Section 12) suggest that these latter four matroids probably *do* have the half-plane property; but proving it may well require new techniques.

13.8 Algorithms

As noted in Sections 2.6, 5.1 and 7.2, both the half-plane property and the weak half-plane property are algorithmically testable, using quantifier-elimination methods. But, at least with existing algorithms and currently available computer hardware, these computations do not seem to be feasible in practice for any interesting matroids (e.g. 7-element rank-3 matroids).

Problem 13.18 *Find algorithms for testing the half-plane property (or, more ambitiously, the weak half-plane property) that are feasible in practice.*

More modestly, one can ask:

Problem 13.19 *Find heuristic numerical methods for testing the half-plane property that are “more powerful” than the method based on Proposition 5.2 (see Section 12).*

Acknowledgments

We wish to thank Sankar Basu, Jason Brown, Jim Geelen, Russ Lyons, Richard Stanley, Dirk Vertigan and Neil White for helpful comments and suggestions. We also wish to thank James Renegar and Adam Strzebonski for correspondence concerning quantifier-elimination algorithms; Jürgen Bokowski, Andreas Dress and Bernd Sturmfels for correspondence concerning determinants and Grassmann–Plücker identities; and Sankar Basu, Charles Desoer and Paul Penfield for giving us pointers to the engineering literature. Finally, we wish to thank Marc Noy and Dominic Welsh for organizing the Workshop on Tutte Polynomials and Related Topics (Barcelona, September 2001), at which this work was first presented.

This research was supported in part by NSA grants MDA904–99–1–0030 and MDA904–01–1–0026 (J.G.O.), NSF grants PHY–9900769 and PHY–0099393 (A.D.S.), and an operating grant from NSERC (D.G.W.).

A Matroids considered in this paper

The half-plane property is preserved by direct sums (Proposition 4.4) and 2-sums (Corollary 4.9); moreover, it is trivially preserved by adjoining loops or parallel elements. Therefore, we can restrict attention to simple 3-connected matroids. (In fact, every 3-connected matroid with at least 4 elements is automatically simple [55, Proposition 8.1.6].) Since all rank-1 and rank-2 matroids have the half-plane property (Corollary 5.5), we can restrict attention to matroids of rank ≥ 3 . Finally, since the half-plane property is invariant under duality (Proposition 4.2), we can restrict attention to matroids of rank $\leq \lfloor n/2 \rfloor$, where n is the number of elements.

In this appendix we list all 3-connected rank-3 matroids on 6 or 7 elements, as well as a few larger matroids that will play a role in this paper. We follow where possible the notation of [55, Appendix]; further information on many of these matroids can be found there.

A.1 Rank-3 matroids on 6 elements

The 3-connected rank-3 matroids on 6 elements are shown in Figure 4.

$M(K_4)$: Regular (in fact graphic and cographic). Self-dual. Not transversal or cotransversal. Has half-plane property (Theorem 1.1).

\mathcal{W}^3 : F -representable if and only if $|F| \geq 3$. Sixth-root-of-unity but not regular. Self-dual. Transversal but not nice; cotransversal but not co-nice (Example 10.3). Has half-plane property (Corollary 8.2).

Q_6 : F -representable if and only if $|F| \geq 4$. Not sixth-root-of-unity. Self-dual. Nice transversal and co-nice cotransversal (Example 10.7). Has half-plane property (Corollary 10.3).




Figure 4: The 3-connected rank-3 matroids on 6 elements. All of these are obtained from $M(K_4)$ by a sequence of relaxations.

P_6 : F -representable if and only if $|F| \geq 5$. Not sixth-root-of-unity. Self-dual. Nice transversal and co-nice cotransversal (Examples 10.4 and 10.7). Has half-plane property (Corollary 10.3).

$U_{3,6}$: F -representable if and only if $|F| \geq 4$. Not sixth-root-of-unity. Self-dual. Nice transversal and co-nice cotransversal (Example 10.1). Has half-plane property (Theorem 9.1).

A.2 Rank-3 matroids on 7 elements

Besides the uniform matroid $U_{3,7}$, we classify the 3-connected rank-3 matroids on 7 elements as follows [65]:

- (a) Those with a 4-point line (Figure 5).
- (b) Those with no 4-point line:
 - (b1) Those that are not ω -regular (Figure 6). All of these are obtained from the Fano matroid F_7 by a sequence of relaxations.
 - (b2) Those that are ω -regular (Figure 7). All of these are obtained from P_7 by a sequence of relaxations.




Figure 5: The 3-connected rank-3 matroids on 7 elements with a 4-point line.

A.2.1 Matroids with a 4-point line

Q_7 : F -representable if and only if $|F| \geq 5$. Not sixth-root-of-unity. Nice transversal (Example 10.7); co-transversal but not co-nice (Example 10.14). Non-nice principal extension of Q_6 ; non-nice free extension of $Q_7 \setminus 7$ (Example 4.4). Has half-plane property (Corollary 10.3).

S_7 : F -representable if and only if $|F| \geq 7$. Not sixth-root-of-unity. Nice transversal (Example 10.7); co-transversal (we don't know whether or not it is co-nice). Non-nice principal extension of P_6 ; non-nice free extension of $S_7 \setminus 7$. Has half-plane property (Corollary 10.3).

$M(K_4)^+$: F -representable if and only if $|F| \geq 3$. Sixth-root-of-unity but not regular. Not transversal or cotransversal. Nice principal extension of $M(K_4)$ with $\lambda_1 = \lambda_2 = \lambda_3 = 1/2$ (Example 4.1). This matroid is called O_7 in [29]. Has half-plane property (Example 4.1 or Corollary 8.2).

\mathcal{W}^{3+} : F -representable if and only if $|F| \geq 4$. Not sixth-root-of-unity. Transversal but not nice (Example 10.3); co-transversal but not co-nice (Example 10.13). Non-nice principal extension of \mathcal{W}^3 ; non-nice principal extension of $Q_7 \setminus 7$. Not known whether it has half-plane property.




Figure 6: The 3-connected rank-3 non-uniform matroids on 7 elements that have no 4-point line and are not ω -regular. All of these are obtained from the Fano matroid F_7 by a sequence of relaxations.




Figure 7: The 3-connected rank-3 non-uniform matroids on 7 elements that have no 4-point line and are ω -regular. All of these are obtained from P_7 by a sequence of relaxations.

A.2.2 The Fano matroid F_7 and its relaxations

F_7 : F -representable if and only if F has characteristic 2. Not sixth-root-of-unity. Not transversal or cotransversal. Doubly transitive automorphism group. Does not have half-plane property (Example 11.1).

F_7^- : F -representable if and only if F has characteristic $\neq 2$. Not sixth-root-of-unity. Not transversal or cotransversal. Does not have half-plane property (Example 11.2).

F_7^{--} : F -representable if and only if $|F| \geq 4$ (Example 11.6). Not sixth-root-of-unity. Not transversal or cotransversal. Non-nice principal extension of $M(K_4)$. Does not have half-plane property (Example 11.3).

$M(K_4) + e$: F -representable if and only if $|F| \geq 5$. Not sixth-root-of-unity. Not transversal or cotransversal. Non-nice free extension of $M(K_4)$ [Example 4.2]. Does not have half-plane property (Example 11.4).

F_7^{-3} : F -representable if and only if $|F| \geq 5$. Not sixth-root-of-unity. Not transversal; cotransversal but not co-nice (Example 10.9). Non-nice principal extension of \mathcal{W}^3 . Does not have half-plane property (Example 11.7).

- F_7^{-4} : F -representable if and only if $|F| \geq 4$. Not sixth-root-of-unity. Not transversal; cotransversal but not co-nice (Example 10.10). Non-nice principal extension of Q_6 . Not known whether it has half-plane property.
- $\mathcal{W}^3 + e$: F -representable if and only if $|F| \geq 5$. Not sixth-root-of-unity. Transversal but not nice (Example 10.3); cotransversal but not co-nice (Example 10.11). Non-nice free extension of \mathcal{W}^3 ; non-nice principal extension of Q_6 . Not known whether it has half-plane property.
- F_7^{-5} : F -representable if and only if $|F| \geq 5$. Not sixth-root-of-unity. Transversal but not nice (Example 10.8); co-nice cotransversal (Example 10.12). Non-nice free extension of Q_6 ; non-nice principal extension of P_6 . Has half-plane property (Corollary 10.3).
- F_7^{-6} : F -representable if and only if $|F| \geq 7$. Not sixth-root-of-unity. Nice transversal (Example 10.4); co-transversal (we don't know whether or not it is co-nice). Non-nice free extension of P_6 ; non-nice principal extension of $U_{3,6}$. Has half-plane property (Corollary 10.3).

A.2.3 P_7 and its relaxations

- P_7 : F -representable if and only if $|F| \geq 3$. Sixth-root-of-unity but not regular. k -regular for all $k \geq 1$ (see [66]). Not transversal or cotransversal. Has half-plane property (Corollary 8.2).
- P'_7 : F -representable if and only if $|F| \geq 4$. k -regular for all $k \geq 2$ (see [66]). Not transversal; cotransversal but not co-nice (Example 10.16). Non-nice principal extension of \mathcal{W}^3 . Not known whether it has half-plane property.
- P''_7 : F -representable if and only if $|F| \geq 5$. k -regular for all $k \geq 3$ (see [66]). Transversal but not nice (Example 10.15); co-nice cotransversal (Examples 10.17 and 10.18). Non-nice principal extension of Q_6 ; non-nice principal extension of R_6 . Has half-plane property (Corollary 10.3).
- P'''_7 : F -representable if and only if $|F| \geq 7$. k -regular for all $k \geq 4$ (see [66]). Nice transversal (Examples 10.4 and 10.7); co-transversal (we don't know whether or not it is co-nice). Non-nice free extension of R_6 ; non-nice principal extension of P_6 . Has half-plane property (Corollary 10.3).

A.3 Some rank-3 matroids on 8 or 9 elements

The matroids F_7^{-4} , $\mathcal{W}^3 + e$, \mathcal{W}^{3+} and P'_7 play a special role in this paper, as they are the only rank-3 7-element matroids for which we are unable to prove whether or




Figure 8: Some rank-3 8-element matroids that are free extensions of rank-3 7-element matroids.

not they have the half-plane property. Since our numerical experiments (Section 12) suggest that they probably *do* have the half-plane property, we have also investigated some single-element extensions of these matroids in an (unsuccessful) effort to find one that fails the half-plane property. For brevity we discuss here only the four matroids that are obtained as *free* extensions (see Figure 8):

$F_7^{-4} + e$: F -representable if and only if $|F| \geq 7$. Not sixth-root-of-unity. Not transversal; cotransversal but not co-nice (Example 10.10). Nice principal extension of F_7^{-4} with $\lambda_1 = 0, \lambda_2 = \dots = \lambda_7 = 1/4$ (Example 4.3). Every single-element deletion is isomorphic to F_7^{-4} , F_7^{-5} or $U_{3,7}$. Not known whether it has half-plane property.

$W^3 + e + f$: F -representable if and only if $|F| \geq 7$. Not sixth-root-of-unity. Transversal but not nice (Example 10.3); cotransversal but not co-nice (Example 10.11). Non-nice free extension of $W^3 + e$. Every single-element deletion is isomorphic to $W^3 + e$, F_7^{-5} or F_7^{-6} . Not known whether it has half-plane property.

$W^{3+} + e$: F -representable if and only if $|F| \geq 7$. Not sixth-root-of-unity. Transversal but not nice (Example 10.3); cotransversal but not co-nice (Example 10.13). Non-nice free extension of W^{3+} . Every single-element deletion is isomorphic to Q_7 , S_7 , W^{3+} , $W^3 + e$ or F_7^{-5} . Not known whether it has half-plane property.




Figure 9: The Pappus and non-Pappus matroids, along with the free extension $(\text{non-Pappus} \setminus 9) + e$.

$P'_7 + e$: F -representable if and only if $|F| \geq 7$. Not sixth-root-of-unity. Not transversal; cotransversal but not co-nice (Example 10.16). Non-nice free extension of P'_7 . Every single-element deletion is isomorphic to $\mathcal{W}^3 + e$, F_7^{-5} , P'_7 or P'''_7 . Not known whether it has half-plane property.

The Pappus and non-Pappus matroids have rank 3 and 9 elements, and are shown in Figure 9. We also consider some related matroids: the deletions $\text{non-Pappus} \setminus 1$ and $\text{non-Pappus} \setminus 9$, and the free extension $(\text{non-Pappus} \setminus 9) + e$. These matroids have the following properties:

Pappus: F -representable if and only if $|F| = 4$ or $|F| \geq 7$.²² Not sixth-root-of-unity. Not transversal or cotransversal. Transitive automorphism group. Every single-element deletion is isomorphic to $\text{non-Pappus} \setminus 9$. Does not have half-plane property (Example 11.9).

non-Pappus: Not representable over any field. Not sixth-root-of-unity. Not transversal or cotransversal. Every single-element deletion is isomorphic to

²² The assertion on [55, p. 516] is in error.

- non-Pappus\1 or non-Pappus\9. Does not have half-plane property (Example 11.9).
- non-Pappus\1: F -representable if and only if $|F| \geq 5$. Not sixth-root-of-unity. Not transversal; cotransversal but not co-nice (Example 10.19). Every single-element deletion is isomorphic to F_7^{-4} , $\mathcal{W}^3 + e$, F_7^{-5} , P'_7 or P''_7 . Not known whether it has half-plane property.
- non-Pappus\9: F -representable if and only if $|F| \geq 4$. Not sixth-root-of-unity. Not transversal or cotransversal. Every single-element deletion is isomorphic to F_7^{-4} or P'_7 . Not known whether it has half-plane property.
- (non-Pappus \ 9) + e : F -representable if $|F| \geq 7$. Not sixth-root-of-unity. Not transversal or cotransversal. Every single-element deletion is isomorphic to $F_7^{-4} + e$, $P'_7 + e$ or non-Pappus\9. Does not have half-plane property (Example 11.10).

A.4 Some rank-4 matroids on 8 elements

The rank-4 matroid P_8 is represented over any field of characteristic $\neq 2$ by the matrix²³

$$\left[\begin{array}{cccc|cccc} 1 & 0 & 0 & 0 & 0 & 1 & 1 & 2 \\ 0 & 1 & 0 & 0 & 1 & 0 & 1 & 1 \\ 0 & 0 & 1 & 0 & 1 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & 2 & 1 & 1 & 0 \end{array} \right] \quad (\text{A.1})$$

A geometric representation of P_8 over the reals can be obtained by starting from a 3-dimensional cube and then rotating one face of the cube by 45° in its plane [29, Figure 14]. There is a unique pair of disjoint circuit-hyperplanes in P_8 , namely $\{1, 4, 5, 8\}$ and $\{2, 3, 6, 7\}$. We denote by P'_8 (resp. P''_8) the matroid obtained from P_8 by relaxing one (resp. both) of these circuit-hyperplanes.

The Vámos matroid V_8 also has rank 4; it is pictured in [55, p. 76, Figure 2.4].

P_8 : F -representable if and only if F has characteristic $\neq 2$. Not sixth-root-of-unity. Self-dual. Not transversal or cotransversal. Transitive automorphism group. Every single-element contraction is isomorphic to P_7 ; every single-element deletion is isomorphic to $(P_7)^*$. See [29, 30] for more information. Does not have half-plane property (Example 11.8).

P'_8 : F -representable if and only if $|F| \geq 4$. Not sixth-root-of-unity. Self-dual. Not transversal or cotransversal. Every single-element contraction is isomorphic to P_7 or P'_7 ; every single-element deletion is isomorphic to $(P_7)^*$ or $(P'_7)^*$. Does not have half-plane property (Example 11.8).

²³ The $GF(3)$ -representation of P_8 on [55, p. 512] has a misprint: the bottom right element should be 0, not 1.

P_8'' : F -representable if and only if $|F| \geq 5$. Not sixth-root-of-unity. Self-dual. Not transversal or cotransversal. Transitive automorphism group. Every single-element contraction is isomorphic to P_7' ; every single-element deletion is isomorphic to $(P_7')^*$. See [29, 30] for more information. Does not have half-plane property (Example 11.8).

V_8 : Not representable over any field. Not sixth-root-of-unity. Self-dual. Not transversal or cotransversal. Every single-element contraction is isomorphic to F_7^{-4} or F_7^{-5} ; every single-element deletion is isomorphic to $(F_7^{-4})^*$ or $(F_7^{-5})^*$. Not known whether it has half-plane property.

A.5 Some transversal matroids

Some interesting transversal matroids are introduced in Sections 10.3 and 10.4. In particular, all rank-3 transversal matroids are classified in Section 10.4; and the rank- $(n+1)$ transversal matroid N_n is defined in Example 10.18.

B $(F, \{1\})$ -representability of matroids

We defined (F, G) -representations of matroids in Section 8.2 and noted that, when $-1 \in G$, such representations are covered under the theory of partial fields developed by Semple and Whittle [67]. When $-1 \notin G$, the class of (F, G) -representable matroids is less well-behaved. For example, neither reordering the columns, reordering the rows, nor pivoting is guaranteed to produce another (F, G) -representation. In this appendix, we consider (F, G) -representable matroids in the case that G is the trivial group $\{1\}$, and we characterize such matroids. If F has characteristic 2, then it is clear that a matroid M is $(F, \{1\})$ -representable if and only if M is binary. When the characteristic of F is not 2, the determination of all $(F, \{1\})$ -representable matroids, which is contained in Theorem B.1, is more difficult.

We now inductively define when a graph is a *chain of cycles*.

- (i) A single cycle C of length at least three is a chain of cycles, which we write as (C) .
- (ii) Suppose that (C_1, C_2, \dots, C_n) is a chain of cycles and that each of C_1, C_2, \dots, C_n is a cycle. Let e be an edge of C_n that is in none of C_1, C_2, \dots, C_{n-1} and let C_{n+1} be a cycle of length at least three that contains e but no other edge of (C_1, C_2, \dots, C_n) . Let $(C_1, C_2, \dots, C_{n+1})$ be the parallel connection of (C_1, C_2, \dots, C_n) and C_{n+1} across the edge e . Then $(C_1, C_2, \dots, C_{n+1})$ is a chain of cycles.

A graph G' is an *augmented chain of cycles* if G' can be obtained from a chain of cycles G by adding some (possibly empty) set of edges so that each added edge is parallel to some edge of G .

The next theorem is the main result of this appendix. The graph G_6 has 6 vertices and 9 edges and is obtained by attaching a triangle via parallel connection to each edge of a triangle.

Theorem B.1 *The following statements are equivalent for a matroid M and a field F of characteristic other than 2.*

- (i) M is $(F, \{1\})$ -representable.
- (ii) Each component of M is a loop, a rank-1 uniform matroid, or the cycle matroid of an augmented chain of cycles.
- (iii) M has no minor isomorphic to $U_{2,4}$, $M(K_4)$, $M(K_{2,3})$, or $M(G_6)$.

A consequence of Theorem B.1 is that, when the characteristic of F is not 2, the class of $(F, \{1\})$ -representable matroids is not closed under duality. For example, if G is the graph that is obtained from G_6 by contracting one edge incident with a degree-2 vertex, then G is an augmented chain of cycles, so $M(G)$ is $(F, \{1\})$ -representable. However, $M^*(G)$ has $M(K_{2,3})$ as a minor and so is not $(F, \{1\})$ -representable. We can also use $M(G)$ to show that, when the characteristic of F is not 2, the class of $(F, \{1\})$ -representable matroids is not closed under 2-sum, series connection, or parallel connection. To see this, we observe that $M(G_6)$ is the 2-sum of $M(G)$ and the graph obtained by adding an edge parallel to one edge of a triangle; $M(G_6)$ is also the series connection of $M(G)$ and $U_{1,2}$; and, finally, $M(G_6)$ is the parallel connection of the simplification of $M(G)$ and a triangle.

The proof of Theorem B.1 will use the following definition. Fix an integer $r \geq 1$; and, for $1 \leq i \leq r$, let \mathbf{e}_i be the column vector of length r with a 1 in the i th row and zeros elsewhere. An $r \times n$ matrix A will be called *special* if:

- (a) the first column is \mathbf{e}_1 , and the last column is \mathbf{e}_r ;
- (b) each column is either \mathbf{e}_i or $\mathbf{e}_i + \mathbf{e}_{i+1}$;
- (c) a column \mathbf{e}_i is always followed by either \mathbf{e}_i or $\mathbf{e}_i + \mathbf{e}_{i+1}$; and
- (d) a column $\mathbf{e}_i + \mathbf{e}_{i+1}$ is always followed by either $\mathbf{e}_i + \mathbf{e}_{i+1}$, \mathbf{e}_{i+1} , or $\mathbf{e}_{i+1} + \mathbf{e}_{i+2}$.

Note that a special matrix may have repeated columns; but if two columns are equal to c , say, then all intermediate columns must also be equal to c .

The next result is the main step in the proof of Theorem B.1.

Proposition B.2 *Let A be an $(F, \{1\})$ -representation of a connected loopless matroid M . If the number of rows of A equals $r(M)$, then A is special.*

PROOF. Let A be the $r \times n$ matrix $[a_{ij}]$. Assume that the columns of A are indexed, in order, by $1, 2, \dots, n$ where $E(M) = \{1, 2, \dots, n\}$. The proposition certainly holds if A has one row, so assume that $r \geq 2$. Note that A has no zero columns, because M is loopless; and A has no zero rows, because $r = r(M)$. Since every subdeterminant of A is in $\{0, 1\}$, it follows that A has no 2×2 submatrices of the form $\begin{bmatrix} * & 1 \\ 1 & 0 \end{bmatrix}$ or $\begin{bmatrix} 0 & 1 \\ 1 & * \end{bmatrix}$. Therefore:

B.3 *If $a_{ij} = 0$, then*

- (i) $a_{ij'} = 0$ for all j' with $1 \leq j' \leq j$, or $a_{i'j} = 0$ for all i' with $1 \leq i' \leq i$; and
- (ii) $a_{ij'} = 0$ for all j' with $j \leq j' \leq n$, or $a_{i'j} = 0$ for all i' with $i \leq i' \leq r$.

We rewrite these conditions as:

B.4 *If $a_{ij} = 0$, then*

- (i) *either all the entries north of a_{ij} are zero, or all the entries west of a_{ij} are zero; and*
- (ii) *either all the entries south of a_{ij} are zero, or all the entries east of a_{ij} are zero.*

Lemma B.5 *Let $a_{ij} = 0$. Then either*

- (i) $a_{i'j'} = 0$ for all i' and j' with $1 \leq i' \leq i$ and $j \leq j' \leq n$; or
- (ii) $a_{i'j'} = 0$ for all i' and j' with $i \leq i' \leq r$ and $1 \leq j' \leq j$.

PROOF. Since $a_{ij} = 0$, by (B.4)(i), either

- (a) all entries north of a_{ij} are 0; or
- (b) all entries west of a_{ij} are 0.

Suppose that (a) holds. Since column j is non-zero, if $1 \leq i' \leq i$, then, by (B.4)(ii), all entries east of $a_{i'j}$ are 0. Thus (i) holds. Now suppose that (b) holds. Then, since row i is non-zero, if $1 \leq j' \leq j$, then, by (B.4)(ii), all entries south of $a_{ij'}$ are zero, and (ii) holds. \square

Again it will be convenient to rewrite the last lemma as:

B.6 *If $a_{ij} = 0$, then either*

- (i) *all the entries north-east of a_{ij} are zero; or*
- (ii) *all the entries south-west of a_{ij} are zero.*

By combining (B.3) with the fact that A has no zero rows and no zero columns, we deduce that no column of A has $[1 \ 0 \ 1]^T$ as a submatrix and no row of A has $[1 \ 0 \ 1]$ as a submatrix. Thus every column of A and every row of A consists of a (possibly empty) sequence of zeros, followed by a non-empty sequence of ones, followed by a (possibly empty) sequence of zeros.

Let D_3 be the matrix $\begin{bmatrix} 1 & 1 & 0 \\ 1 & 1 & 1 \\ 0 & 1 & 1 \end{bmatrix}$. Then D_3 has determinant -1 and so the next lemma is immediate.

Lemma B.7 *The matrix A does not have D_3 as a submatrix.*

Lemma B.8 *If $a_{ij} = 1$ and $a_{(i+1)j} = 0$, then $j < n$ and $a_{i(j+1)} = 1$.*

PROOF. If $j = n$, then, since $a_{ij} = 1$, it follows by (B.4) that row $i + 1$ of A is zero; a contradiction. Thus $j < n$. Suppose that $a_{i(j+1)} = 0$. Then, by (B.6), all entries north-east of $a_{i(j+1)}$ are zero and all entries south-west of $a_{(i+1)j}$ are zero. Thus A has the block form $\begin{bmatrix} A_1 & 0 \\ 0 & A_2 \end{bmatrix}$ and so M is disconnected; a contradiction. \square

By a similar argument to that just given, we get:

Lemma B.9 *If $a_{ij} = 0$ and $a_{(i+1)j} = 1$, then $j \geq 2$ and $a_{(i+1)(j-1)} = 1$.*

Lemma B.10 $a_{r1} = 0 = a_{1n}$.

PROOF. Suppose that $a_{r1} = 1$. If $a_{i1} = 0$ for some $i < r$, then, by (B.5), row i is zero; a contradiction. Thus column 1 of A consists of all ones and, similarly, row r consists of all ones. Since $r \geq 2$, there is a column of A that does not consist of all ones. Let j be the first such column. Then $a_{j1} = 0$ and $\{1, 2, \dots, j-1\}$ is the ground set of a rank-1 component of M ; a contradiction. We conclude that $a_{r1} = 0$, and a similar argument shows that $a_{1n} = 0$. \square

Assume now that Proposition B.2 is false, and let A be a counterexample having the minimum number of columns. In particular, no two consecutive columns of A are equal. The following lemmas derive some properties of this minimal counterexample, culminating in a contradiction.

Lemma B.11 *Every column of A has at most two ones.*

PROOF. Suppose that, for some j with $1 < j < n$, column j has a unique one, which occurs in row i say. Then $1 < i < r$, and every entry north-east of $a_{(i-1)j}$ is zero, as is every entry south-west of $a_{(i+1)j}$. Thus M/j is disconnected, so M is a parallel connection with basepoint j . Indeed, M is the parallel connection of $M[A_1]$

and $M[A_2]$ where A_1 is the submatrix of A that consists of a_{ij} and all entries to its north-west, and A_2 is the submatrix of A that consists of a_{ij} and all entries to its south-east. Since M is connected, so are both $M[A_1]$ and $M[A_2]$. It follows that both A_1 and A_2 are special matrices and, therefore, so is A ; a contradiction. Thus every column of A except possibly the first or the last has at least two ones.

Now suppose that A has a single one in its first column. Then $a_{11} = 1$. Suppose that $M/1$ is connected. Let A_1 be the matrix obtained from A by deleting the first row and column. Then A_1 is special. By Lemma B.10, $a_{1n} = 0$. Let the first zero entry in row 1 of A be a_{1m} . Then $m \geq 3$ otherwise 1 is a coloop of M . Now $a_{11} = a_{12} = \dots = a_{1(m-1)} = 1$. Moreover, $a_{22} = a_{23} = \dots = a_{2(m-1)} = 1$ otherwise $M[A_1]$ has a loop. Since A has no consecutive equal columns and A_1 is special, $m \leq 4$. If $m = 3$, then columns 1 and 2 of A are $[1 \ 0 \ 0 \ \dots \ 0]^T$ and $[1 \ 1 \ 0 \ \dots \ 0]^T$, and it follows that A is special; a contradiction. We may now suppose that $m = 4$. Then $a_{24} = 0$ otherwise the submatrix of A determined by the first three rows and columns 2, 3, and 4 is D_3 ; a contradiction. Thus $a_{2j} = 0$ for all $j \geq 4$. Now the matrix A represents M over F . If we subtract row 2 from row 1 in A , we get another F -representation for M , which shows that 1 is a coloop of M ; a contradiction.

We may now assume that $M/1$ is disconnected. Then $M \setminus 1$ is connected. It follows that the matrix $A \setminus 1$, which is obtained from A by deleting the first column, is special. Thus column 2 of A equals e_1 . Therefore column 1 is parallel to column 2; a contradiction.

We conclude that the first column of A has at least two ones, and, by symmetry, the last column of A has at least two ones. \square

A submatrix C of a matrix D will be called *consecutive* if the rows of C are consecutive rows of D and the columns of C are consecutive columns of D .

Lemma B.12 *A has $\begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$ as a consecutive submatrix.*

PROOF. Proceed north from a_{r1} , which we know from Lemma B.10 is 0, until the first 1 is reached. Then we have $\begin{bmatrix} 1 \\ 0 \end{bmatrix}$ as a submatrix meeting consecutive rows. Then proceed east from the 0 in this submatrix until the first 1 is met. By Lemma B.8, we deduce that A has $\begin{bmatrix} 1 & 1 & \dots & 1 & 1 \\ 0 & 0 & \dots & 0 & 1 \end{bmatrix}$ as a consecutive submatrix, and the lemma follows. \square

Let J'_k be the $k \times k$ matrix all of whose entries are one except for that in the south-west corner, which is zero.

Lemma B.13 *Every consecutive J'_2 -submatrix of A is contained in a consecutive J'_3 -submatrix of A .*

PROOF. Suppose that $\begin{bmatrix} c_{11} & c_{12} & c_{13} \\ 1 & 1 & c_{23} \\ 0 & 1 & c_{33} \end{bmatrix}$ is a consecutive submatrix C of A containing $\begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$ as a submatrix. The additional row must exist and $c_{11} = 1$ otherwise A has a column with a single one. The additional column must exist and $c_{33} = 1$ otherwise A has a row with a single one and so has a coloop.

Suppose $c_{12} = 0$. Then adjoin an additional column x to A that is zero everywhere except in the row corresponding to the second row of C . This column should be added between the first and second columns of C with the resulting matrix being A' . The matroid $M[A']/x$ is disconnected. Thus $M[A]$ is the 2-sum of two matroids with basepoint x . Let A'_1 be the submatrix of A' consisting of the non-zero entry in column x and all entries north-west of it, and A'_2 be the submatrix of A' consisting of the non-zero entry in column x and all entries south-east of it. Then $M[A']$ is the parallel connection of $M[A'_1]$ and $M[A'_2]$ and so each of these is connected. Since A has a submatrix equal to A'_1 , the last matrix is an $(F, \{1\})$ -representation for a connected matroid. Since A'_2 is obtained by adjoining the column $[1 \ 0 \ 0 \ \dots \ 0]^T$ to the beginning of a submatrix of A , it too is an $(F, \{1\})$ -representation for a connected matroid. Thus each of A'_1 and A'_2 is special. It follows that A' is special and so too is A ; a contradiction. We conclude that $c_{12} = 1$.

Suppose $c_{23} = 0$. Let y be the column of A corresponding to the second column of C . Then $A \setminus y$ has the form $\begin{bmatrix} A_1 & 0 \\ 0 & A_2 \end{bmatrix}$ and so $M \setminus y$ is disconnected and the rank of A is the sum of the ranks, r_1 and r_2 , say, of A_1 and A_2 . We may apply row operations in A to transform A_2 into a matrix that has a permutation of I_{r_2} as a submatrix. These row operations will not affect the first r_1 rows of A . Then, by contracting the elements of M corresponding to the columns of I_{r_2} , we see that the submatrix A''_1 consisting of A_1 and the first r_1 rows of column y represents a connected matroid. This is because M is the series connection of this matroid and another with respect to the basepoint y . Thus A''_1 is special. But the last column of A''_1 has more than one one; a contradiction. We conclude that $c_{23} = 1$.

Finally, since D_3 is not a submatrix of A , we deduce that $c_{13} = 1$. \square

Lemma B.14 *For all $k \geq 2$, the matrix A has J'_k as a consecutive submatrix.*

PROOF. We shall argue by induction. By the last two lemmas, A has J'_2 and J'_3 as consecutive submatrices. Assume that A has J'_m as a consecutive submatrix, for some $m \geq 3$. The rows i and $i + 1$ of A that correspond to rows 1 and 2 of J'_m must differ. Therefore, in A , the submatrix induced by rows i and $i + 1$ has either

- (i) in the consecutive columns immediately after J'_m , a $2 \times t$ matrix H_1 of the form $\begin{bmatrix} 1 & 1 & \dots & 1 & 0 \\ 1 & 1 & \dots & 1 & 1 \end{bmatrix}$ for some $t \geq 1$;

(ii) in the consecutive columns immediately before J'_m , a $2 \times s$ matrix H_2 of the form $\begin{bmatrix} 1 & 1 & \dots & 1 & 1 \\ 0 & 1 & \dots & 1 & 1 \end{bmatrix}$ for some $s \geq 1$.

In case (i), the column of A corresponding to the last column of H_1 must have ones in the $m-1$ rows immediately below the unique zero otherwise (B.6) is contradicted. It follows that A has D_3 as a submatrix; a contradiction.

In case (ii), the unique zero in H_2 is in a consecutive submatrix of A equal to J'_2 . Thus, by Lemma B.13, the $(i-1)$ st row of A must have ones in all the columns corresponding to those of J'_m otherwise A has D_3 as a submatrix. We deduce that A has, as a consecutive submatrix, the matrix J''_m that is obtained by adjoining a row of ones to the beginning of J'_m .

The columns j and $j+1$ of A that correspond to the last two columns of J''_m must differ. If they differ in some row before row $i-1$ of A , let row u be the highest indexed such row where they differ. This row has $[1 \ 0]$ in columns j and $j+1$ of A . Then, by Lemma B.5, $a_{uj} = a_{u(j-1)} = \dots = a_{u(j-m+2)} = 1$ and so A has D_3 as a submatrix; a contradiction. We deduce that columns j and $j+1$ of A differ in some row after row $i+m-1$. Let row v be the smallest indexed such row where these columns differ. Then

$$\begin{bmatrix} a_{(v-1)j} & a_{(v-1)(j+1)} \\ a_{vj} & a_{v(j+1)} \end{bmatrix} = J'_2.$$

By Lemma B.13, this J'_2 is contained in a consecutive submatrix of A equal to J'_3 . To avoid having D_3 as a submatrix of A , it follows that $a_{(i-1)(j+2)} = a_{i(j+2)} = \dots = a_{(i+m-1)(j+2)} = 1$. Thus A has J'_{m+1} as a consecutive submatrix and the lemma follows by induction. \square

Proposition B.2 follows immediately from the last lemma since it implies that the matrix A is infinite. \square

Lemma B.15 *If A is a special matrix, then every square submatrix of A has determinant in $\{0, 1\}$.*

PROOF. We argue by induction on the number n of columns of A . If $n = 1$, then the result is immediate. Assume the result holds for $n = k$ and let $n = k + 1$. Now let A' be a square submatrix of A . We may assume that the columns of A are distinct since if A' has repeated columns, its determinant is 0. If A' meets the first column of A , then $\det A' = 0$ unless A' meets the first row of A . In the exceptional case, $\det A'$ equals the determinant of the submatrix A'_1 of A' that is obtained by deleting the first row and column. Now A'_1 is a submatrix of A_1 , the matrix obtained by deleting the first row and column of A . Moreover, it is not difficult to see that A_1 is special. Thus, by induction, $\det A' \in \{0, 1\}$. We may now assume that A' does not meet the first column of A . If A' does not meet the first row of A , then A' is a submatrix of

the special matrix A_1 and the result follows by the induction assumption. Thus we may assume that A' contains the entry a_{12} of A otherwise $\det A' = 0$. Since a_{12} is the only non-zero entry of the first row of A' , we have $\det A' = \det A'_1$. But A'_1 is a submatrix of the special matrix A_1 and again the result follows by the induction assumption. \square

We are now ready to prove Theorem B.1.

PROOF OF THEOREM B.1. We shall show first that (ii) implies (iii). Suppose that M satisfies (ii). Then every minor of M also satisfies (ii). The construction of M guarantees that M is the cycle matroid of a series-parallel network G . Thus M has no minor isomorphic to $U_{2,4}$ or $M(K_4)$. Moreover, G is an outerplanar graph, so M has no minor isomorphic to $M(K_{2,3})$. Finally, we observe that M does not have $M(G_6)$ as a minor since G_6 cannot be written as a chain of cycles. Thus (iii) holds.

Next we show that (iii) implies (ii). Suppose that M is a simple connected matroid that satisfies (iii) and has rank at least two, and assume that M is a minor-minimal matroid that does not satisfy (ii). The fact that M has none of $U_{2,4}$, $M(K_4)$, and $M(K_{2,3})$ as a minor ensures that $M \cong M(G)$ where G is an outerplanar graph with at least three vertices. Take an outerplanar embedding of G . The boundary of the infinite face is a Hamilton cycle C of G and all edges not in C are chords of C . Certainly no chords of C cross. Now take a chord e of C with endpoints x and y , say, such that there is an xy -path P_1 in C such that every vertex of $V(P_1) - \{x, y\}$ has degree 2 in G . Then $M \setminus P_1$ satisfies (iii) and hence also satisfies (ii). Thus $M \setminus P_1$ is the cycle matroid of a chain of cycles (C_1, C_2, \dots, C_n) , and M is obtained by taking the parallel connection of $M \setminus P_1$ and the circuit with ground set $P_1 \cup e$. If e is an edge of C_1 that is in no other C_i , then M is the cycle matroid of a chain of cycles. Therefore, by symmetry, we may now assume that either e is an edge that is common to C_i and C_{i+1} for some i , or e is in exactly one C_j for some j in $\{2, 3, \dots, n-1\}$. The first case cannot arise because G is outerplanar. The second case implies without difficulty that $M(G)$ has a minor isomorphic to $M(G_6)$. This contradiction implies that M satisfies (ii).

Now suppose that M satisfies (ii). We shall show that M satisfies (i). If the components M_1, M_2, \dots, M_m of M have $(F, \{1\})$ -representations A_1, A_2, \dots, A_m , respectively, then the matrix whose block form has A_1, A_2, \dots, A_m on the main diagonal and zeros elsewhere is an $(F, \{1\})$ -representation for M . Thus it suffices to prove that each component of M satisfies (i). Clearly a loop and a coloop are $(F, \{1\})$ -representable. Moreover, if the simple matroid associated with M is $(F, \{1\})$ -representable, then so too is M . Thus we may assume that M is simple. Let C be a k -cycle for some $k \geq 3$. Then the $(k-1) \times k$ matrix with columns $\mathbf{e}_1, \mathbf{e}_1 + \mathbf{e}_2, \mathbf{e}_2 + \mathbf{e}_3, \dots, \mathbf{e}_{k-2} + \mathbf{e}_{k-1}, \mathbf{e}_{k-1}$ is an F -representation for $M[C]$. The construction of a representation for a parallel connection of two matroids is straightforward (see, for example, [55, Proposition 7.1.21]). Using this, it is not difficult to show that the cycle matroid of a chain of cycles can be represented by a special matrix. It follows by Lemma B.15 that M

satisfies (i).

Finally, we show that (i) implies (ii). Assume that M is $(F, \{1\})$ -representable and consider a component M' of M of rank at least two. Let A be an $(F, \{1\})$ -representation for M' . We may assume that A has $r(M')$ rows otherwise we can delete some row from A and still retain an $(F, \{1\})$ -representation for M' . Then, by Proposition B.2, A is special. From the definition of a special matrix, it is not difficult to see that M' is the cycle matroid of an augmented chain of cycles. \square

References

- [1] L.V. Ahlfors, *Complex Analysis* (McGraw-Hill, New York, 1966).
- [2] N. Anderson, E.B. Saff and R.S. Varga, On the Eneström–Kakeya theorem and its sharpness, *Lin. Alg. Appl.* **28** (1979), 5–16.
- [3] T. Asano, Theorems on the partition functions of the Heisenberg ferromagnets, *J. Phys. Soc. Japan* **29** (1970), 350–359.
- [4] R.B.apat and T.E.S. Raghavan, *Nonnegative Matrices and Applications* (Cambridge University Press, Cambridge–New York, 1997).
- [5] S. Basu, R. Pollack and M.-F. Roy, On the combinatorial and algebraic complexity of quantifier elimination, *J. ACM* **43** (1996), 1002–1045.
- [6] J. Bokowski and B. Sturmfels, *Computational Synthetic Geometry*, Lecture Notes in Mathematics #1355 (Springer-Verlag, Berlin–Heidelberg, 1989).
- [7] N.K. Bose, A note on multivariable positive real functions, *IEEE Trans. Circuit Theory* **CT–19** (1972), 194–195.
- [8] A. Bouchet and W.H. Cunningham, Delta-matroids, jump systems, and bisubmodular polyhedra, *SIAM J. Discrete Math.* **8** (1995), 17–32.
- [9] R.L. Brooks, C.A.B. Smith, A.H. Stone and W.T. Tutte, The dissection of rectangles into squares, *Duke Math. J.* **7** (1940), 312–340.
- [10] J.I. Brown and C.J. Colbourn, Roots of the reliability polynomial, *SIAM J. Discrete Math.* **5** (1992), 571–585.
- [11] R.A. Brualdi, Transversal matroids, in *Combinatorial Geometries*, edited by Neil White, Encyclopedia of Mathematics and its Applications #29 (Cambridge University Press, Cambridge, 1987), Chapter 5, pp. 72–97.
- [12] O. Brune, Synthesis of a finite two-terminal network whose driving-point impedance is a prescribed function of frequency, *J. Math. and Phys.* **10** (1931), 191–236.

- [13] T.H. Brylawski, An affine representation for transversal geometries, *Stud. Appl. Math.* **54** (1975), 143–160.
- [14] T. Brylawski, Constructions, in *Theory of Matroids*, edited by Neil White, Encyclopedia of Mathematics and its Applications #26 (Cambridge University Press, Cambridge, 1986), Chapter 7, pp. 127–223.
- [15] B.F. Caviness and J.R. Johnson, editors, *Quantifier Elimination and Cylindrical Algebraic Decomposition* (Springer-Verlag, Wien–New York, 1998).
- [16] S. Chaiken, A combinatorial proof of the all minors matrix-tree theorem, *SIAM J. Alg. Disc. Meth.* **3** (1982), 319–329.
- [17] S. Chaiken and D.J. Kleitman, Matrix tree theorems, *J. Combin. Theory A* **24** (1978), 377–381.
- [18] W.-K. Chen, *Applied Graph Theory* (North-Holland, Amsterdam–London, 1971).
- [19] C.J. Colbourn, *The Combinatorics of Network Reliability* (Oxford University Press, New York–Oxford, 1987).
- [20] G.E. Collins, Quantifier elimination for real closed fields by cylindrical algebraic decomposition, in *Automata Theory and Formal Languages* (Lecture Notes in Computer Science #33), edited by H. Brakhage (Springer-Verlag, Berlin–Heidelberg–New York–Tokyo, 1975), pp. 134–183. Reprinted in: *Quantifier Elimination and Cylindrical Algebraic Decomposition*, edited by B.F. Caviness and J.R. Johnson (Springer-Verlag, Wien–New York, 1998), pp. 85–121.
- [21] W.H. Cunningham, A combinatorial decomposition theory, Ph.D. thesis, University of Waterloo, 1973.
- [22] D.M. Cvetković, M. Doob and H. Sachs, *Spectra of Graphs: Theory and Applications* (Academic Press, New York–San Francisco–London, 1980).
- [23] C.A. Desoer and E.S. Kuh, *Basic Circuit Theory* (McGraw-Hill, New York, 1969).
- [24] R.J. Duffin and T.D. Morley, Wang algebra and matroids, *IEEE Trans. Circuits and Systems* **25** (1978), 755–762.
- [25] A. Fettweis, Elementary symmetric functions and positive functions in several variables, in *Signal Processing, Scattering and Operator Theory, and Numerical Methods*, edited by M.A. Kaashoek, J.H. van Schuppen and A.C.M. Ran (Birkhäuser, Boston–Basel–Berlin, 1990), pp. 91–98.
- [26] A. Fettweis and S. Basu, New results on stable multidimensional polynomials — Part I: Continuous case, *IEEE Trans. Circuits and Systems* **34** (1987), 1221–1232.

- [27] F.R. Gantmacher, *The Theory of Matrices*, vol. 2 (Chelsea, New York, 1959).
- [28] J.F. Geelen, Lectures on jump systems, Center of Parallel Computing, University of Cologne, n.d.
- [29] J.F. Geelen, A.M.H. Gerards and A. Kapoor, The excluded minors for $GF(4)$ -representable matroids, *J. Combin. Theory B* **79** (2000), 247–299.
- [30] J.F. Geelen, J.G. Oxley, D.L. Vertigan and G.P. Whittle, On the excluded minors for quaternary matroids, *J. Combin. Theory B* **80** (2000), 57–68.
- [31] J.H. Grace, The zeros of a polynomial, *Proc. Cambridge Philos. Soc.* **11** (1902), 352–357.
- [32] D.H. Greene and D.E. Knuth, *Mathematics for the Analysis of Algorithms*, 3rd ed. (Birkhäuser, Boston, 1990).
- [33] J. Gregor, On quadratic Hurwitz forms I, *Aplikace Matematiky* **26**, 142–153 (1981).
- [34] D.Yu. Grigor’ev, Complexity of deciding Tarski algebra. *J. Symbolic Comput.* **5** (1988), 65–108.
- [35] O.J. Heilmann and E.H. Lieb, Theory of monomer-dimer systems, *Commun. Math. Phys.* **25** (1972), 190–232.
- [36] J. Heintz, M.F. Roy and P. Solernó, Sur la complexité du principe de Tarski-Seidenberg, *Bull. Soc. Math. France* **118** (1990), 101–126.
- [37] A. Hinkkanen, Schur products of certain polynomials, in *Lipa’s Legacy: Proceedings of the Bers Colloquium*, edited by Jozef Dodziuk and Linda Keen (American Mathematical Society, Providence, 1997), *Contemp. Math.* **211**, 285–295.
- [38] A.J. Hoffman, Eigenvalues and partitionings of the edges of a graph, *Linear Algebra Appl.* **5** (1972), 137–146.
- [39] T.W. Hungerford, *Algebra* (Springer-Verlag, New York, 1974, fifth printing 1989).
- [40] A.W. Ingleton, Transversal matroids and related structures, in *Higher Combinatorics*, edited by M. Aigner (Reidel, Dordrecht, 1977), pp. 117–131.
- [41] T. Koga, Synthesis of finite passive n -ports with prescribed positive real matrices of several variables, *IEEE Trans. Circuit Theory* **CT-15** (1968), 2–23.
- [42] S.G. Krantz, *Function Theory of Several Complex Variables*, 2nd ed. (Wadsworth & Brooks/Cole, Pacific Grove, California, 1992).

- [43] S. Krogdahl, The dependence graph for bases in matroids, *Discrete Math.* **19** (1977), 47–59.
- [44] J.P.S. Kung, Basis-exchange properties, in *Theory of Matroids*, edited by Neil White, Encyclopedia of Mathematics and its Applications #26 (Cambridge University Press, Cambridge, 1986), Chapter 4, pp. 62–75.
- [45] J.P.S. Kung, Twelve views of matroid theory, in *Combinatorial and Computational Mathematics* (Pohang, Korea, 15–17 February 2000), edited by Sungpyo Hong, Jin Ho Kwah, Ki Hang Kim and Fred W. Roush (World Scientific, Singapore, 2001), pp. 56–96.
- [46] T.D. Lee and C.N. Yang, Statistical theory of equations of state and phase transitions. II. Lattice gas and Ising model, *Phys. Rev.* **87** (1952), 410–419.
- [47] E.H. Lieb and A.D. Sokal, A general Lee-Yang theorem for one-component and multicomponent ferromagnets, *Commun. Math. Phys.* **80** (1981), 153–179.
- [48] L. Lovász and M.D. Plummer, *Matching Theory*, North-Holland Mathematics Studies #121, Annals of Discrete Mathematics #29 (North-Holland, Amsterdam–New York / Akadémiai Kiadó, Budapest, 1986).
- [49] L. Lovász, The membership problem in jump systems, *J. Combin. Theory B* **70** (1997), 45–66.
- [50] M. Marcus and H. Minc, *Introduction to Linear Algebra* (Macmillan, New York, 1965).
- [51] M. Marden, *Geometry of Polynomials*, 2nd ed., (American Mathematical Society, Providence, R.I., 1966).
- [52] J.H. Mason, A characterization of transversal independence spaces, in *Théorie des Matroïdes*, (Lecture Notes in Mathematics #211), edited by C.P. Bruter (Springer-Verlag, Berlin, 1971), pp. 86–94.
- [53] H. Minc, *Permanents* (Addison-Wesley, Reading MA, 1978).
- [54] C.M. Newman, Zeros of the partition function for generalized Ising systems, *Comm. Pure Appl. Math.* **27** (1974), 143–159.
- [55] J.G. Oxley, *Matroid Theory* (Oxford University Press, New York, 1992).
- [56] J. Oxley, C. Semple and D. Vertigan, Generalized $\Delta - Y$ exchange and k -regular matroids, *J. Combin. Theory B* **79** (2000), 1–65.
- [57] H. Ozaki and T. Kasami, Positive real functions of several variables and their applications to variable networks, *IEEE Trans. Circuit Theory* **CT-7** (1960), 251–260.

- [58] P. Penfield, R. Spence and S. Duinker, *Tellegen's Theorem and Electrical Networks* (MIT Press, Cambridge, MA, 1970).
- [59] M.J. Piff and D.J.A. Welsh, On the vector representation of matroids, *J. London Math. Soc.* (2) **2** (1970), 284–288.
- [60] J. Renegar, On the computational complexity and geometry of the first-order theory of the reals. I–III, *J. Symbolic Comput.* **13** (1992), 255–352.
- [61] G. Royle and A.D. Sokal, The Brown–Colbourn conjecture on zeros of reliability polynomials is false, in preparation.
- [62] W. Rudin, *Function Theory in Polydiscs* (Benjamin, New York, 1969).
- [63] D. Ruelle, Extension of the Lee–Yang circle theorem, *Phys. Rev. Lett.* **26** (1971), 303–304, 870 (erratum).
- [64] D. Ruelle, Some remarks on the location of zeroes of the partition function for lattice systems, *Commun. Math. Phys.* **31** (1973), 265–277.
- [65] C. Semple, k -regular matroids, Ph.D. thesis, Victoria University, Wellington, 1998.
- [66] C. Semple, On maximum-sized k -regular matroids, *Graphs Combin.* **15** (1999), 441–462.
- [67] C. Semple and G. Whittle, Partial fields and matroid representation, *Adv. Appl. Math.* **17** (1996), 184–208.
- [68] B. Simon, *The $P(\varphi)_2$ Euclidean (Quantum) Field Theory* (Princeton University Press, Princeton, 1974), Section IX.3.
- [69] J.H. Smith, Some properties of the spectrum of a graph, in *Combinatorial Structures and their Applications* (Proceedings of the Calgary International Conference, Calgary, Alberta, 1969), pp. 403–406 (Gordon and Breach, New York, 1970).
- [70] A.D. Sokal, Bounds on the complex zeros of (di)chromatic polynomials and Potts-model partition functions, *Combin. Probab. Comput.* **10** (2001), 41–77, cond-mat/9904146 at xxx.lanl.gov.
- [71] G. Szegő, Bemerkungen zu einem Satz von J. H. Grace über die Wurzeln algebraischer Gleichungen, *Math. Z.* **13** (1922), 28–55. Reprinted in: G. Szegő, *Collected Papers*, vol. 1, edited by Richard Askey (Birkhäuser, Boston–Basel–Stuttgart, 1982), pp. 506–534.

- [72] A. Tarski, *A Decision Method for Elementary Algebra and Geometry* (University of California Press, Berkeley–Los Angeles, 1951). Reprinted in: *Quantifier Elimination and Cylindrical Algebraic Decomposition*, edited by B.F. Caviness and J.R. Johnson (Springer-Verlag, Wien–New York, 1998), pp. 24–84.
- [73] D.G. Wagner, Zeros of reliability polynomials and f -vectors of matroids. *Combin. Probab. Comput.* **9** (2000), 167–190.
- [74] J.L. Walsh, On the location of the roots of certain types of polynomials, *Trans. Amer. Math. Soc.* **24** (1922), 163–180.
- [75] N. White, Coordinatizations, in *Combinatorial Geometries*, edited by Neil White, Encyclopedia of Mathematics and its Applications #29 (Cambridge University Press, Cambridge, 1987), Chapter 1, pp. 1–27.
- [76] N. White, Unimodular matroids, in *Combinatorial Geometries*, edited by Neil White, Encyclopedia of Mathematics and its Applications #29 (Cambridge University Press, Cambridge, 1987), Chapter 3, pp. 40–52.
- [77] G. Whittle, A characterisation of the matroids representable over $GF(3)$ and the rationals, *J. Combin. Theory B* **65** (1995), 222–261.
- [78] G. Whittle, On matroids representable over $GF(3)$ and other fields, *Trans. Amer. Math. Soc.* **349** (1997), 579–603.
- [79] H.S. Wilf, *Generatingfunctionology*, 2nd ed. (Academic Press, Boston, 1994).
- [80] D. Zeilberger, A combinatorial approach to matrix algebra, *Discrete Math.* **56** (1985), 61–72.