

Lower bounds for the energy in a crumpled elastic sheet { A minimal ridge

Shankar C Venkataramani

Department of Mathematics, University of Chicago, 5734 University Ave., Chicago, IL 60637, USA

E-mail: shankar@math.uchicago.edu

Abstract. We study the linearized Foppl { von Karm an theory of a long, thin rectangular elastic membrane that is bent through an angle 2θ . We prove rigorous bounds for the minimum energy of this configuration in terms of the plate thickness h and the bending angle. We show that the minimum energy scales as $h^{5/3} \theta^{7/3}$. This scaling is in sharp contrast with previously obtained results for the linearized theory of thin sheets with isotropic compression boundary conditions, where the energy scales as $h^2 \theta^2$.

AMS classification scheme numbers: 49J40, 74K15, 47J20, 46N10

Submitted to: Nonlinearity

1. Introduction

Everyday experience tells us that thin elastic sheets crumple, when con ned into a small volume, e.g. a sheet of paper "con ned" by one's hands. Crumpling also plays an important role in the mechanical behavior of packaging material and in the dissipation of the energy of collisions by the "crumple zones" of automobiles. Crumpling is therefore a problem of much intrinsic interest, but understanding this behavior is complicated by the complex morphology of a typical crumpled sheet.

Despite the complicated appearance of a crumpled sheet, the crumpling phenomenon is in itself very robust. It is easily observed in thin sheets made from a variety of materials, suggesting that it can be studied using simplified or idealized models that capture the essential features of thin elastic sheets. This approach leads one to consider a crumpled sheet as a minimum energy configuration for a simple elastic energy functional for thin sheets, viz. the Foppl { von Karm an (FvK) energy. Using this approach of elastic energy minimization, the crumpling response is now understood as a result of the elastic energy of the sheet concentrating on a small subset of the entire sheet [1, 2, 3]. The energy in a crumpled sheet is concentrated on a network of thin line-like creases (ridges) that meet in point-like vertices. Recent work has resulted in quantitative

understanding of both the vertices [4, 5, 6, 7, 8] and the ridges [9, 10]. Scaling laws governing the behavior of crumpled sheets have been obtained in the physics literature [1, 2, 9, 6].

Minimum energy configurations for the FvK energy have also been studied in the context of the blistering problem, viz. the buckling of membranes as a result of isotropic compression along the boundary. The blistering problem is relevant to the delamination of thin films that are chemically deposited at high temperatures, as well as the mechanical behavior of micro-fabricated thin-film diaphragms [11, 12].

There is a considerable body of mathematical work focused on the blistering problem [11, 13, 14, 15, 16, 17, 18]. Upper and lower bounds have been obtained for approximations to the elastic energy [11, 13, 16], for the FvK energy [15, 17] and for full three dimensional nonlinear elasticity [18]. The FvK energy and full three dimensional nonlinear elasticity give the same scaling for the upper and the lower bounds. This yields a rigorous scaling law for the energy of a blister. As we discuss below, the scaling laws for the blistering problem are different from those for the crumpling problem, even for the same energy functional. This indicates that the energy minimizing configurations for the FvK energy show an interesting dependence on the boundary conditions.

The minimization of the FvK elastic energy is an example of a non-convex variational problem that is regularized by a singular perturbation [19, 20]. It is well known that this can lead to a variety of multiple-scale behaviors including energy concentration and/or small scale oscillations in the minimizers [21].

Multiple scale behaviors, both microstructure and singularities, are ubiquitous in condensed matter systems. The crumpling phenomenon appears to be a particularly simple and tractable instance of multiple scale behavior. For this reason, there has been much recent interest in physics literature about the nature of the crumpling phenomenon [22, 23, 1, 2, 24]. A asymptotic analysis [9] and scaling arguments [2] show that the ridge energy scales as

$$E_r \propto L^{5/3};$$

where t is the thickness of the sheet, and L is the length of the ridge. Ben-Amar et al [1] and Mahadevan et al [4] show that the energy of a d-cone scales as

$$E_c \propto a^2 \log(L/a);$$

where a is the radius of the core associated with the vertex. It is clear that the ridge energy is asymptotically larger than the vertex energy as $L \rightarrow \infty$. However, which of these two energies is important for a given sheet depends on the relation between the nondimensional thickness $t = t/L$ of the sheet, and the "crossover thickness" $t_c = t_c(L)$ which is determined by setting $E_r = E_c$. t_c in turn depends on the values of the multiplicative constants for these scaling laws.

These multiplicative constants cannot be determined by scaling arguments. Mahadevan et al [6] estimate the constant for the vertex energy numerically using an ansatz for the shape of a sheet near the vertex, and they find that

$$E_c \approx 100 t^2 \log(L/a);$$

where θ is the complement of the tip angle of the cone [6, 7]. Based on Lobkovsky's work [9], Boudaoud et al [7] estimate the value of the constant in the ridge energy as 1. This implies, for the experiments in Refs. [6] and [7], the vertex energy dominates the ridge energy.

It would be useful to prove these scaling laws, and determine the multiplicative constants rigorously, that is, in an ansatz-free manner. In this paper, we propose a model problem that yields structures analogous to a single ridge in a crumpled sheet. We prove a rigorous lower bound (with a numerical value for the multiplicative constant) for the elastic energy in our model problem. This is a step toward rigorously proving the scaling law of Lobkovsky et al for the ridge energy [2]. We also discuss how the techniques in this paper can be extended to prove similar results for a "real" crumpled sheet, as opposed to our model problem.

This paper is organized as follows: In Sec. 2, we describe the problem of interest, set up the relevant energy functional and determine the appropriate boundary conditions. We also rescale the various quantities to a form that is suitable for further analysis, and recast the problem in terms of the rescaled quantities. In Sec. 3, we prove our main result, viz. a lower bound for the elastic energy for our boundary conditions. We present a concluding discussion in Sec. 4.

2. The Elastic energy

We are interested in a minimal ridge, i.e., the single crease that is formed when a long rectangular elastic strip is bent through an angle by clamping the lateral boundaries to a bent frame. This situation is depicted in Figure 1.

From the symmetry of the problem, it is clear that we only need to consider one half of the strip. This is represented schematically in Figure 2. We will use (the material) coordinates $(x; y)$ on the reference half strip $|x| \leq L, 0 \leq y \leq 1$. Also, we associate a $(u; v; w)$ coordinate system in space, so that as $y \rightarrow 1$, the half-strip is asymptotically in the $w = 0$ plane, as depicted in Figure 2. The grid in the figure is generated by the lines $x = \text{constant}$ and $y = \text{constant}$ that are straight in the reference (material) coordinates. w represents the out of plane deformation, and the in-plane coordinates are chosen so that the u and the v axes are asymptotically parallel to the x and y axes respectively as $y \rightarrow 1$. Since the sheet is bent through an angle 2θ , as $y \rightarrow 1$, the sheet will lie in the plane $w = v \tan(\theta)$. The symmetry of the two halves implies that the line $y = 0$ maps into the plane $w \tan(\theta) + v = 0$, which bisects the angle between the planes $w = 0$ (the asymptote as $y \rightarrow 1$) and $w = v \tan(\theta)$ (the asymptote as $y \rightarrow 1$). The symmetry between the two halves also implies that the tangent to the lines $x = \text{constant}$ at $y = 0$ should be perpendicular to the plane $w \tan(\theta) + v = 0$. Consequently $u_y = 0$ and $w_y = v_y \tan(\theta)$ at $y = 0$.

A mathematically justified way to obtain the elastic energy of the deformed sheet is to treat the sheet as a three dimensional (albeit thin) object and use a full nonlinear three dimensional elastic energy functional for the energy density. This approach however does




Figure 1. A minimal ridge. The boundary conditions are given by a frame (the thick solid lines) bent through an angle. The sheet is essentially flat outside the region bounded by the two dashed curves, and the bulk of the energy is concentrated in this region.




Figure 2. A schematic representation of our coordinate system and the boundary conditions imposed on the sheet. The grid is given by the lines $x = \text{constant}$ and $y = \text{constant}$. The two dashed lines are the tangent to the curve $x = 0$ and the line $w = w_0(0); u = 0$ respectively, and the angle between these lines is α .

not take advantage of the "thinness" of the sheet. In particular, we would like to treat the thin sheet as a two dimensional object. This will greatly reduce the complexity of the problem. The derivation of reduced dimensional descriptions of thin sheets has a long history. There is a classical theory for thin elastic sheets built on the work of Euler, Cauchy, Kirchhoff, Föppl and von Kármán [25, 26].

Many modern authors have considered the problem of deriving a reduced dimensional theory [27] from three dimensional elasticity in a mathematically rigorous fashion through Γ -convergence [28, 29]. Γ -convergence is the appropriate notion for convergence in variational problems. In the context of thin sheets, roughly speaking, finding the Γ -limit amounts to identifying an appropriate two dimensional energy functional whose minimizers give the limiting behavior of the minimizers of the full three dimensional energy in the limit the thickness $h \rightarrow 0$. This problem hasn't been solved in its entirety.

There are two situations for which reduced dimensional theories have been derived rigorously as Γ -limits of full three dimensional elasticity. Membrane theories [30, 31, 32] are applicable in situations where the stretching is essentially uniform through the thickness of the sheet and bending theories [33] are applicable in situations where the sheet is essentially unstretched, and all the elastic energy is due to strains that are first order in the thickness of the sheet.

Neither of these theories are applicable for them in a ridge. For them in a ridge, both the stretching (membrane) energy and the bending energy are important. In fact, a ridge is a result of the competition between these two energies. So we turn to the classical Föppl-von Kármán ansatz for an appropriate definition of the elastic energy [25, 26]. Although the FvK energy is not rigorously derived from three dimensional elasticity, it can be thought of as the sum of the membrane and the bending energies that have been derived rigorously in different limiting situations. We will further assume that, for the minimal ridge, the deflections w , u and v are small compared to the natural length scale L . The strains are of the order of the square of the maximum deflection divided by L and they are small. After some rescaling, the energy of the deformed sheet is given by the linearized FvK energy

$$I = \int_{\Omega} \left((u_x + \frac{w_x^2}{2} - 1)^2 + \frac{1}{2} (v_x + u_y + 2w_x w_y)^2 + (v_y + \frac{w_y^2}{2} - 1)^2 \right) dx dy + \frac{1}{2} \int_{\Omega} (w_{xx}^2 + 2w_{xy}^2 + w_{yy}^2) dx dy: \quad (1)$$

where x and y are reference coordinates on the sheet, u and v are in-plane coordinates, w is the out of plane displacement and h is the thickness of the sheet. The integrand for the first integral is given by the squares of the linearized strains,

$$\epsilon_{xx} = u_x + \frac{w_x^2}{2} - 1; \quad \epsilon_{xy} = \epsilon_{yx} = w_x w_y + \frac{1}{2} (v_x + u_y); \quad \epsilon_{yy} = v_y + \frac{w_y^2}{2} - 1:$$

The blistering of thin films is also described by the elastic energy in (1). A similar energy also describes multiple scale buckling in free elastic sheets (i.e. sheets that are not forced through the boundary conditions) that are not intrinsically flat [34].

The difference between the blistering problem and a minimal ridge is in the boundary conditions, which we describe below. Since the strains are assumed to be small, $v_y + w_y^2 \ll 1$. If the bending half-angle $\ll 1$, so that $\tan \theta \approx \theta$, we get $w_y = 0$ at $y = 0$. Since the deformation goes to zero far away from the bend, we have

$$u = x; v = w = 0; \text{ as } y \rightarrow \pm \infty;$$

Also, the lateral boundaries at $x = \pm L$ are clamped to the frame. Therefore,

$$u = x; v = w = 0; \text{ at } x = \pm L:$$

Note that the appropriate boundary condition for the minimal ridge at $y = 0$ is a free boundary condition, subject to the symmetry requirement $w + v \tan \theta = 0$. We are going to replace this free boundary condition with a Dirichlet boundary condition

$$v = v_0(x); w = w_0(x) \quad \text{at } y = 0;$$

where we will leave the functions $v_0(x)$ and $w_0(x)$ unspecified, except for a size condition. Defining

$$a = \max_{[-L, L]} (|v_0(x)| + |w_0(x)|);$$

we impose the size condition by requiring that a be "small".

2.1. Rescalings

The relevant small parameter in the problem is the non-dimensional thickness of the sheet, $\epsilon = h/L$. Following Lobkovsky [9], we introduce the rescaled coordinates and displacements by

$$x = LX; \quad y = \epsilon^{1/3} L^{2/3} Y;$$

and

$$w = \epsilon^{1/3} L^{2/3} W; \quad v = y + \epsilon^{1/3} L^{2/3} V; \quad u = x + \epsilon^{2/3} L^{1/3} U;$$

Since $\epsilon, L, x, y, u, v, w$ all have dimensions of a length, it is clear that the rescaled quantities X, Y, U, V, W are all dimensionless. With these rescalings, the dimensionless energy $I = \epsilon^{5/3} L^{1/3} \mathcal{I}$ is given by

$$\begin{aligned} \mathcal{I}(U; V; W) = & \int_{-L}^L \int_{-\infty}^{\infty} (U_X^2 + W_X^2)^2 + \frac{\epsilon^{2/3}}{2} (V_X + U_Y + 2W_X W_Y)^2 + \epsilon^{4/3} (V_Y + W_Y^2)^2 \\ & + W_{YY}^2 + 2\epsilon^{2/3} W_{XY}^2 + \epsilon^{4/3} W_{XX}^2 \, dX \, dY. \end{aligned} \quad (2)$$

Our quest for rigorous scaling results for the energy I reduces to the following { Show that the rescaled energy \mathcal{I} , of a minimizer $(U; V; W)$, is bounded above and below by positive constants uniform in the dimensionless thickness parameter ϵ , as $\epsilon \rightarrow 0$.

Setting

$$W_0(X) = \epsilon^{1/3} L^{2/3} w_0(LX); \quad V_0(X) = \epsilon^{1/3} L^{2/3} v_0(LX);$$

the rescaled quantities satisfy the boundary conditions

$$V = V_0(X); W = W_0(X); W_Y = 0 \quad \text{at } Y = 0;$$

and

$$U \rightarrow 0; V \rightarrow 0; W \rightarrow 0 \quad \text{as } Y \rightarrow 1;$$

We have the lateral boundary conditions

$$U = V = W = 0 \quad \text{at } X = \pm 1;$$

Also, the deformation at $Y = 0$ satisfies $|V_0(X)| \leq A$ and $|W_0(X)| \leq A$ where $A = L^{1/3} L^{2/3} a$.

3. Lower Bound

In this section, we prove a lower bound for the linearized Elastic energy in Eq. (1), provided that the length scale a associated with the boundary conditions satisfies a size condition.

Theorem 3.1. $I(u; v; w)$ is as defined in Eq. (1). There is a constant $b > 0$ such that, for all $u \in H^1$; $v \in H^1$ and $w \in H^2 \setminus W^{1,4}$ satisfying the boundary conditions

$$\begin{aligned} |u - x| &\leq 0; |v - y| \leq 0; w \rightarrow 0 \quad \text{as } y \rightarrow 1; \\ u = x; v = w = 0; &\text{ at } x = \pm L; \\ v = v_0(x); w = w_0(x) &\quad \text{at } y = 0; \end{aligned}$$

and the size condition

$$\max_{[L, 2L]} (|v_0(x)| + |w_0(x)|) \leq a \leq b L^{1/3} L^{2/3} L^{2/3};$$

we have the lower bound

$$I(u; v; w) \geq \frac{2}{5} L^{7/3} L^{5/3} L^{1/3};$$

Remark. We will prove the theorem by proving the scaled version of the statement, viz., the rescaled boundary conditions along with the rescaled size condition

$$\max (|V_0(X)| + |W_0(X)|) \leq A \leq b L^{2/3};$$

imply that

$$I \geq \frac{2}{5} L^{7/3};$$

Remark. The hypothesis for the theorem includes a size condition on the displacement at $y = 0$. This is somewhat unsatisfying, as it is not a priori obvious that a ridge in a "real" crumpled sheet will satisfy this condition.

In our search for a lower bound, we can assume $I(U;V;W) < 1$ w.l.o.g. From Eq. (2), it follows that for $\epsilon > 0$, $W \in H^2$. By the standard trace theorems, the boundary conditions $W(X;0) = W_0(X)$ and $W_Y(X;0) = 0$ are therefore assumed pointwise for almost every X .

Since $U = 0$ at $X = 1$, it follows that $\int_0^1 (U_X + W_X^2) dX = \int_0^1 W_X^2 dX$. Using this together with Jensen's inequality we get

$$I(U;V;W) = \int_0^1 \int_0^1 \frac{1}{2} (U_X + W_X^2)^2 + W_{YY}^2 dX dY \geq \int_0^1 \int_0^1 \frac{1}{2} W_X^2 dX + \int_0^1 \int_0^1 W_{YY}^2 dX dY \quad (3)$$

Thus the functional

$$E(W) = \int_0^1 \int_0^1 \frac{1}{2} W_X^2 dX + \int_0^1 \int_0^1 W_{YY}^2 dX dY$$

bounds $I(U;V;W)$ from below. Hence it suffices to obtain a lower bound for E with the given hypothesis to prove the theorem.

Let E_b and E_s denote the quantities

$$E_b = \int_0^1 \int_0^1 W_{YY}^2 dX dY;$$

and

$$E_s = \int_0^1 \int_0^1 \frac{1}{2} W_X^2 dX dY;$$

Although E_b and E_s are only lower bounds for the "true" bending and the stretching energies I_b and I_s , we will nonetheless call E_b and E_s the bending and the stretching energy for convenience.

For every X , we define

$$(X) = \int_0^1 W_{YY}^2(X;Y) dY;$$

and for every Y , we define

$$(Y) = \int_0^1 W_X^2 dX;$$

Since $E < 1$, $(X) > 0$ (a.e.) and $(Y) < 1$ (a.e.).

For any function f depending on X and Y , let $\|f\|_Y^2$ denote $\int_0^1 \int_0^1 f^2(X;Y) dX dY$, so that,

$$(Y) = \|W_X\|_Y^2;$$

(X) is a "local" (in X) measure of the bending energy, and $[(X)]^1$ can be thought of as the bending energy density in X that is obtained by integrating out the Y dependence. We can also think of (X) as the length scale associated with the bending energy as a

function of X , viz., we expect that the bending energy density in Y decays rapidly for $Y = (X)^{-1}$ (See Fig. 1). Likewise, $[(Y)]^2$ is a local (in Y) measure of the stretching energy. In terms of (X) and (Y) , the bending and the stretching energies are given by

$$E_b = \int_1^{Z_1} \frac{1}{(X)} dX \quad E_s = \int_0^{Z_1} \frac{1}{2} [(Y)]^2 dY:$$

We now begin our proof of the theorem. The idea behind the proof is to show that the stretching energy E_s can be bounded from below by a negative power of the bending energy E_b , so that the total energy $E_s + E_b$ tends to $+1$ as $E_b \rightarrow 0$ and $E_b \rightarrow 1$. This ensures the existence of a positive lower bound for E (and consequently also for I).

Lemma 3.2. For every Y , we have the inequality

$$kW - k_Y^2 \leq Y^2 - 2Y^3 E_b - 4A^2:$$

Proof. If $(X) > 0$, $W(X; \cdot)$ is a C^1 function by the Sobolev Embedding theorem and the boundary conditions imply that $W(X; 0) = W_0(X)$ and $W_Y(X; 0) = 0$. Consequently, for such an X ,

$$W(X; Y) = W_0(X) + \int_0^Y W_{YY}(X; \eta) (Y - \eta) d\eta:$$

We will estimate $W^2(X; Y)$ from this equation using the elementary inequalities

$$a - b^2 \leq (1 - \epsilon)a^2 + \frac{1}{\epsilon}b^2;$$

and

$$a + b^2 \leq (1 + \epsilon)a^2 + \frac{1 + \epsilon}{\epsilon}b^2;$$

for all $\epsilon > 0$. By our hypothesis on the boundary conditions,

$$W_0(X)^2 \leq A^2:$$

By Jensen's inequality

$$\begin{aligned} \int_0^Y W_{YY}(X; \eta) (Y - \eta) d\eta &\leq Y \int_0^Y W_{YY}^2(X; \eta) (Y - \eta)^2 d\eta \\ &\leq Y^3 \int_0^Y W_{YY}^2(X; \eta) d\eta \\ &\leq \frac{Y^3}{(X)} \end{aligned}$$

Combining these estimates, we get

$$W^2(X; Y) \leq (1 - \epsilon_1) Y^2 + \frac{1}{\epsilon_1} (1 + \epsilon_2) \frac{Y^3}{(X)} + \frac{1 + \epsilon_2}{\epsilon_2} A^2;$$

for all positive ϵ_1 and ϵ_2 . In particular, setting $\epsilon_1 = 1/2$ and $\epsilon_2 = 1$ yields

$$W^2(X; Y) \leq \frac{1}{2} Y^2 + \frac{2Y^3}{(X)} + 2A^2; \quad (4)$$

Integrating this inequality in X we obtain

$$kW \leq k_Y^2 Y^2 + 2Y^3 E_b + 4A^2; \quad (5)$$

proving the lemma. \square

Our proof is based on demonstrating that a small bending energy E_b will lead to a large stretching energy. This idea is quantified by the following lemma where we use Eq. (5) to estimate the stretching energy E_s from below in terms of the bending energy E_b .

Lemma 3.3. Let

$$= \frac{4E_b A^2}{3};$$

There is a constant $\epsilon > 0$ such that, if $\epsilon < \epsilon$, the stretching energy E_s satisfies

$$E_s \geq \frac{14}{5 \cdot 729} E_b^5;$$

Proof. We have,

$$(Y) = kW_X k_Y^2 - \frac{2}{4} kW k_Y^2;$$

by the Poincaré Inequality. Eq. (5) now yields

$$(Y) \leq \frac{2}{4} Y^2 + 2Y^3 E_b + 4A^2;$$

We will now extract the appropriate scaling dependence of (Y) and E_s on ϵ and E_b . Setting $Y^2 = 2Y^3 E_b$, we deduce that a characteristic scale Υ for Y is given by

$$\Upsilon = \frac{2}{2E_b};$$

Rescaling Y in terms of Υ , we obtain

$$(Y) = \frac{\epsilon^2}{16E_b^2} \frac{Y^2}{\Upsilon^2} + 1 + \frac{Y}{\Upsilon};$$

where ϵ is as defined above, i.e.

$$= \frac{4E_b A^2}{3};$$

Making the change of variables $Y = z\tilde{Y}$, and using $\tilde{Y}(0) = 0$, we see that

$$E_s = \int_0^1 \frac{1}{2} [\tilde{Y}'(z)]^2 dz = \frac{1}{512E_b^5} K(\tilde{Y}(1));$$

where

$$K(\tilde{Y}(1)) = \int_0^1 \max(z^2(1 - z), 0)^2 dz;$$

$K(\tilde{Y}(1))$ is clearly a continuous function of $\tilde{Y}(1)$ and $K(0) = 1/105$. Since $K(0) = 1/105 < 1/(5 \cdot 729)$, there is an $\epsilon > 0$ such that for all $\tilde{Y}(1) < \epsilon$, $K(\tilde{Y}(1)) < 1/(5 \cdot 729)$. The lemma follows. \square

We can now prove the theorem.

Proof. Let $b = \frac{5}{8} p^{1/3}$. The hypothesis imply

$$A < \frac{5}{8} p^{1/3} = 2^{2/3} b.$$

If $E_b = 2^{7/3} = 5$, there is nothing to prove. Therefore, we can assume that $E_b < 2^{7/3} = 5$. Then, it follows that

$$E_s = \frac{4E_b A^2}{3} < \frac{4}{3} b^2.$$

Consequently $E_s = 14/(5 \cdot 729 E_b^5)$. Minimizing $E_s + E_b$, we see that

$$I = E_s + E_b = \frac{14}{5 \cdot 729 E_b^5} + E_b = \frac{2^{7/3}}{5}.$$

The theorem follows by "undoing" our rescaling to express the results in terms of u, v, w, a and I . \square

Remark. In the appendix, we show that

$$K(\tilde{Y}(1)) = \frac{1}{105} - \frac{\tilde{Y}(1)^2}{6}.$$

Therefore, we can choose $\epsilon = 0.048$ and $b = 0.13$. We see that b is not exceedingly small. Rather it is $O(1)$. Also, it is not the best constant for this theorem, and we can get a better value by optimizing our choices for the constants ϵ_1 and ϵ_2 which we set to be $1/2$ and 1 respectively).

4. Discussion

We have proved a rigorous lower bound for the elastic energy of a ridge that is consistent with the results obtained by Lobkovsky et al [2] using scaling arguments, and by Lobkovsky [9] using matched asymptotics. In order to prove these scaling laws rigorously, we also need analogous upper bounds that are consistent with the same scaling. This approach has been used successfully for a variety of other variational

problems [35, 15, 17]. The upper bound is usually obtained by constructing a test solution that yields the bound. One is often guided by scaling arguments in the construction of the appropriate test solution. This is in contrast to the lower bounds, where one needs to obtain functional analysis type inequalities that captures the competition between distinct energies in the problem (e.g. lemma 3.3). It is this competition that determines the scaling behavior of the problem.

We have not constructed the upper bounds, because we believe that they will follow from the scaling ansatz in Lobkovsky's work [9]. We also believe that the upper bound will scale in the same manner as the lower bound, thereby giving us a rigorous scaling law for the energy of a single ridge.

As we remark before proving the theorem, our result is not directly applicable to the minimal ridge since we have the extra hypothesis

$$a \leq b^{2/3} \gamma^{1/3} L^{1/3};$$

We need to replace this restriction with a free boundary condition at $y = 0$ subject to the symmetry requirement $w \tan \theta + v = 0$, and the boundary condition $w_y = v_y \tan \theta$.

Despite this caveat, we claim that the analysis in this paper captures the essential features of the minimal ridge problem, namely the scaling in lemma 3.3, and consequently the scaling law for the energy of the ridge. We will show that this is indeed the case by investigating the problem with the "true" boundary conditions elsewhere.

A harder problem is to show that the scaling laws also hold for a real crumpled sheet, where the forcing is not through clamping the boundaries to a frame, but rather through the confinement in a small volume. In this case, there are interesting global geometric and topological considerations, some of which are explored in Refs. [36] and [37]. As Lobkovsky and Witten [10] argue, the boundary condition that the deformation goes to zero far away from the ridge implies that the ridges do not interact with each other significantly. The ridges can be considered the elementary excitations of a crumpled sheet. In our quest for upper and lower bounds, this translates to the statement that the test solutions for the upper bound can be constructed by piecing together local solutions near each ridge. Thus the upper bound is obtained relatively easily. The hard part is to show that the confinement actually leads to the formation of ridges, and that the competition between the bending and the stretching energy for this situation has the same form as in lemma 3.3. In this context, we expect that global topological considerations, as well as the non self-intersection of the sheet will play a key role in the analysis, as they do in the analysis of elastic rods (one dimensional objects) [38].

As we note above, the blistering problem is described by the same elastic energy (Eq. (1)), but with different boundary conditions. Our results show an interesting contrast with results for the blistering problem. Ben Belgacem et al. have shown that [17], for an isotropically compressed thin film, the energy of the minimizer satisfies

$$E \sim c^{3/2} L^{3/2} \gamma^{1/2} \epsilon^{1/2};$$

where L is a typical length scale of the domain, and ϵ is the compression factor. A construction for the upper bound strongly suggests that the minimizers develop an

initely branched network with oscillations on increasingly finer scales as $\epsilon \rightarrow 0$. In contrast, our results indicate that the energy of a minimal ridge satisfies

$$C \epsilon^{5/3} L^{1/3} \leq E \leq C \epsilon^{5/3} L^{1/3};$$

and the energy concentrates in a region of width $\epsilon^{1/3} L^{2/3}$. This shows that the nature of the solution of the variational problem for the elastic energy in (1) depends very strongly on the boundary conditions. In particular the very nature of the energy minimizers is different for the two problems. For the blistering problems, as $\epsilon \rightarrow 0$ the minimizers develop a branched network of folds refining towards the boundary. The minimizers therefore display the problem of small scale oscillations [39, 21]. The minimal ridge problem on the other hand displays the concentration phenomenon [39, 21] as $\epsilon \rightarrow 0$, with the energy concentrating on a region of width $\epsilon^{1/3} L^{2/3}$. It would be interesting to explore this issue further, and in particular, to understand the mechanisms by which the boundary conditions determine the nature of the minimizer.

Appendix

In this appendix, we prove the inequality

$$K(\epsilon) = \int_0^1 \max(z^2(1-z), \epsilon; 0)^2 dz \geq \frac{1}{105} - \frac{\epsilon}{6}.$$

Proof. Let $f \geq 0$. Then, for $f \geq \epsilon$, we have

$$[\max(f, \epsilon; 0)]^2 = f^2 - 2f\epsilon + \epsilon^2 = f^2 - 2f\epsilon.$$

For $0 \leq f \leq \epsilon$, we have

$$[\max(f, \epsilon; 0)]^2 = 0 = f^2 - 2f\epsilon.$$

Using these inequalities in the definition of $K(\epsilon)$ we obtain

$$K(\epsilon) = \int_0^1 z^2(1-z)^2 - 2\epsilon z^2(1-z) dz \geq \frac{1}{105} - \frac{\epsilon}{6}.$$

□

Acknowledgments

I wish to thank Brian D. Donna, Bob Kohn, L. Mahadevan, Stefan Müller and Tom Witten for helpful and enlightening conversations. This work was started during a visit to the Max-Planck Institute for Mathematics in the Sciences in Leipzig, and I would like to thank the MPI for their hospitality. This work was supported in part by the National Science Foundation through its MRSEC program under Award Number DMR-9808595, and by a NSF CAREER Award DMS-0135078. This work is also supported in part by a Research Fellowship from the Alfred P. Sloan Jr. Foundation.

References

- [1] M. Ben Amar and Y. Pomeau, "Crumpled paper," *Proc. Roy. Soc. London Ser. A*, vol. 453, p. 729, 1997.
- [2] A. Lobkovsky, S. Gentges, H. Li, D. Morse, and T. A. Witten, "Scaling properties of stretching ridges in a crumpled elastic sheet," *Science*, vol. 270, p. 1482, 1995.
- [3] E. M. Kramer and T. A. Witten, "Stress condensation in crushed elastic manifolds," *Phys. Rev. Lett.*, vol. 78, p. 1303, 1997.
- [4] E. Cerda and L. Mahadevan, "Conical surfaces and crescent singularities in crumpled sheets," *Phys. Rev. Lett.*, vol. 80, p. 2354, 1998.
- [5] S. Chab, F. Meb, and J.-C. Geminard, "Experimental study of developable cones," *Phys. Rev. Lett.*, vol. 80, p. 2358, 1998.
- [6] E. Cerda, S. Chab, F. Meb, and L. Mahadevan, "Conical dislocations in crumpling," *Nature*, vol. 401, p. 46, 1999.
- [7] A. Boudaoud, P. Patricio, Y. Couder, and M. Ben Amar, "Dynamics of singularities in a constrained elastic plate," *Nature*, vol. 407, no. 6805, pp. 718{720, 2000.
- [8] T. Mora and A. Boudaoud, "Thin elastic plates: On the core of developable cones," *Eurphys. Lett.*, vol. 59, no. 1, pp. 41{47, 2002.
- [9] A. E. Lobkovsky, "Boundary layer analysis of the ridge singularity in a thin plate," *Phys. Rev. E*, vol. 53, p. 3750, 1996.
- [10] A. E. Lobkovsky and T. A. Witten, "Properties of ridges in elastic membranes," *Phys. Rev. E*, vol. 55, p. 1577, 1997.
- [11] M. Ortiz and G. Giola, "Thermophology and folding patterns of buckling-driven thin-film blisters," *J. Mech. Phys. Solids*, vol. 42, no. 3, pp. 531{559, 1994.
- [12] G. Giola, A. Desimone, M. Ortiz, and A. M. Cuitino, "Folding energetics in thin-film diaphragms," *R. Soc. Lond. Proc. Ser. A Math. Phys. Eng. Sci.*, vol. 458, no. 2021, pp. 1223{1229, 2002.
- [13] W. Jin and R. V. Kohn, "Singular perturbation and the energy of folds," *J. Nonlinear Sci.*, vol. 10, no. 3, pp. 355{390, 2000.
- [14] B. Audoly, "Stability of straight delamination blisters," *Phys. Rev. Lett.*, vol. 83, p. 4124, 1999.
- [15] W. Jin and P. Stemberg, "Energy estimates for the von Kármán model of thin-film blistering," *J. Math. Phys.*, vol. 42, no. 1, pp. 192{199, 2001.
- [16] W. Jin and P. Stemberg, "In-plane displacements in thin-film blistering." preprint, 2001.
- [17] H. Ben Belgacem, S. Conti, A. Desimone, and S. Müller, "Rigorous bounds for the Föppl-von Kármán theory of isotropically compressed plates," *J. Nonlinear Sci.*, vol. 10, no. 6, pp. 661{683, 2000.
- [18] H. Ben Belgacem, S. Conti, A. Desimone, and S. Müller, "Energy scaling of compressed elastic films - three-dimensional elasticity and reduced theories," *Arch. Rat. Mech. Anal.*, vol. 164, pp. 1{37, 2002.
- [19] S. Müller, "Variational models for microstructure and phase transitions," in *Calculus of variations and geometric evolution problems* (Cetraro, 1996), pp. 85{210, Berlin: Springer, 1999.
- [20] P. Stemberg, "The effect of a singular perturbation on nonconvex variational problems," *Arch. Rational Mech. Anal.*, vol. 101, no. 3, pp. 209{260, 1988.
- [21] L. C. Evans, *Weak convergence methods for nonlinear partial differential equations*. Published for the Conference Board of the Mathematical Sciences, Washington, DC, 1990.
- [22] D. R. Nelson and L. Peliti, "Fluctuations in membranes with crystalline and hexatic order," *J. Phys. (Paris)*, vol. 48, p. 1085, 1987.
- [23] H. S. Seung and D. R. Nelson, "Defects in flexible membranes with crystalline order," *Phys. Rev. A*, vol. 38, p. 1005, 1988.
- [24] L. A. Cuccia and R. B. Lennox, "Molecular modeling of fullerenes with modular origami," *Abstr Pap Am Chem S*, vol. 208, p. 50, 1994.
- [25] A. E. H. Love, *A treatise on the Mathematical Theory of Elasticity*. Dover Publications, New

- York, 1944. Fourth Ed.
- [26] L.D. Landau and E.M. Lifshitz, *Theory of elasticity*. London: Pergamon Press, 1959.
 - [27] P.G. Ciarlet, *Mathematical elasticity*. Vol. II. Amsterdam: North-Holland Publishing Co., 1997. *Theory of plates*.
 - [28] E. De Giorgi and G. Dal Maso, *Γ -convergence and calculus of variations*, in *Mathematical theories of optimization* (Genova, 1981), pp. 121{143, Berlin: Springer, 1983.
 - [29] G. Dal Maso, *An introduction to Γ -convergence*. Boston, MA: Birkhäuser Boston Inc., 1993.
 - [30] H. Le Dret and A. Raoult, *Le modèle de membrane non linéaire comme limite variationnelle de l'élasticité non linéaire tridimensionnelle*, *C.R. Acad. Sci. Paris Ser. I Math.*, vol. 317, no. 2, pp. 221{226, 1993.
 - [31] H. Le Dret and A. Raoult, *The nonlinear membrane model as variational limit of nonlinear three-dimensional elasticity*, *J. Math. Pures Appl.* (9), vol. 74, no. 6, pp. 549{578, 1995.
 - [32] H. Le Dret and A. Raoult, *The membrane shell model in nonlinear elasticity: a variational asymptotic derivation*, *J. Nonlinear Sci.*, vol. 6, no. 1, pp. 59{84, 1996.
 - [33] G. Friesecke, S. Müller, and R.D. James, *Rigorous derivation of nonlinear plate theory and geometric rigidity*, *C.R. Math. Acad. Sci. Paris*, vol. 334, no. 2, pp. 173{178, 2002.
 - [34] E. Sharon, B. Roman, M. Marder, G.-S. Shin, and H.L. Swinney, *Buckling cascade in free thin sheets*, *Nature*, 419 (2002), p. 579.
 - [35] R.V. Kohn and S. Müller, *Surface energy and microstructure in coherent phase transitions*, *Comm. Pure Appl. Math.*, vol. 47, no. 4, pp. 405{435, 1994.
 - [36] S.C. Venkataramani, T.A. Witten, E.M. Kramer, and R.P. Gerch, *Limitations on the smooth continuum of an unstretchable manifold*, *J. Math. Phys.*, vol. 41, no. 7, pp. 5107{5128, 2000.
 - [37] B.A. D'Onofrio, T.A. Witten, S.C. Venkataramani, and E.M. Kramer, *Singularities, structures, and scaling in deformed m -dimensional elastic manifolds*, *Phys. Rev. E* (3), vol. 65, no. 1, part 2, pp. 016603, 25, 2002.
 - [38] O. Gonzalez, J.H. Maddocks, F. Schuricht, and H. von der Mosel, *Global curvature and self-contact of nonlinearly elastic curves and rods*, *Calc. Var. Partial Differential Equations*, vol. 14, no. 1, pp. 29{68, 2002.
 - [39] L. Tartar, *Compensated compactness and applications to partial differential equations*, in *Nonlinear analysis and mechanics: Heriot-Watt Symposium*, Vol. IV, pp. 136{212, Boston, Mass.: Pitman, 1979.