

The Cross Curvature Flow of 3-manifolds with Negative Sectional Curvature

Bennett Chow
UC San Diego

Richard S. Hamilton
Columbia University

1 The evolution equation

We introduce an evolution equation which deforms metrics on 3-manifolds with sectional curvature of one sign. Given a closed 3-manifold with an initial metric with negative sectional curvature, we conjecture that this flow will exist for all time and converge to a hyperbolic metric after a normalization. We shall establish a monotonicity formula in support of this conjecture. Note that in contrast to negative sectional curvature, every closed n -manifold admits a metric with negative Ricci curvature by the work of Gao and Yau [5] for $n = 3$ and Lohkamp [10] for all $n \geq 3$; When $n = 4$; Gromov and Thurston [6] have shown that there exist closed manifolds with arbitrarily pinched negative sectional curvature which do not admit metrics with constant negative sectional curvature. It is unknown whether such manifolds admit Einstein metrics. In particular, the stability result of Ye [13] assumes more than curvature pinching. When $n = 3$; it is an old conjecture, which is also a consequence of the Geometrization Conjecture, that any closed 3-manifold with negative sectional curvature admits a hyperbolic metric.

Let (M, g) be a 3-dimensional Riemannian manifold with negative sectional curvature. The Einstein tensor is $P_{ij} = R_{ij} - \frac{1}{2}Rg_{ij}$; We find it convenient to raise the indices: $P^{ij} = g^{ik}g^{jl}R_{kl}$, $\frac{1}{2}Rg^{ij}$; The cross curvature tensor is

$$h_{ij} = \frac{\det P^{k\ell}}{\det g^{k\ell}} V_{ij}$$

where V_{ij} is the inverse of P^{ij} : If the eigenvalues of P_{ij} are $a = R_{2323}$; $b = R_{1313}$; $c = R_{1212}$; then the eigenvalues of R_{ij} are $(b+c)$; $(a+c)$; $(a+b)$ and the eigenvalues of h_{ij} are bc ; ac ; ab : Hence if $M^3; g$ has negative sectional curvature, then both P_{ij} and h_{ij} are positive definite.

Lemma 1 We have the following identities

$$1. \operatorname{tr} P^{ij} = 0$$

Partially supported by NSF Grant DMS-0203926.

$$2. h^{ij} r_{ij} h_{jk} = \frac{1}{2} h^{ij} r_{jk} h_{ij}:$$

Note that the first identity is the contracted second Bianchi identity. The above identities imply that the cross curvature tensor is dual to the Ricci tensor in the following sense.

Lemma 2 Let $(M^n; g)$ be Riemannian manifold.

1. If the Ricci curvature is positive, then the identity map $\text{id} : (M; g_j) \rightarrow (M; R_{ij})$ is harmonic.
2. If $n = 3$ and the sectional curvature is negative (or positive), then $\text{id} : (M; h_{ij}) \rightarrow (M; g_{ij})$ is harmonic.

With this in mind we define the cross curvature flow $(XCF)^1$ by

$$\frac{\partial}{\partial t} g_{ij} = 2h_{ij}$$

if the sectional curvature is negative and by $\frac{\partial}{\partial t} g_{ij} = -2h_{ij}$ if the sectional curvature is positive. Note that the XCF is fully nonlinear whereas the Ricci flow is quasi-linear.

2 Short time existence

Let ϵ_{ijk} denote the volume form and raise indices by $\epsilon^{ijk} = g^{ip} g^{jq} g^{kr} \epsilon_{pqr}$. Our normalization is such that $\epsilon_{123} = \epsilon^{123} = 1$. We find it convenient to rewrite h_{ij} as

$$h_{ij} = \frac{1}{8} R_{i'pq}{}^{pqk} R_{kjrs}{}^{rs'}:$$

Lemma 3 If $(M; g)$ is a closed 3-manifold with negative (or positive) sectional curvature, then for any smooth initial metric a solution to the XCF exists for a short time.

Proof. We consider the case of negative sectional curvature since the case of positive sectional curvature is similar. Let $g_{i,j}$ denote a variation of the metric g_{ij} and let tildes also denote the variations of various curvature tensors. We have

$$\tilde{R}_{ijk}{}^{'} = \frac{1}{2} \left(\frac{\partial^2 g_{jk}}{\partial x^i \partial x^{'}} + \frac{\partial^2 g_{i'}}{\partial x^j \partial x^k} - \frac{\partial^2 g_{j'}}{\partial x^i \partial x^k} - \frac{\partial^2 g_{ik}}{\partial x^j \partial x^{'}} \right) +$$

where the dots denote terms with 1 or less derivatives of the metric. Applying the equality

$${}^{pqk} R_{kjrs}{}^{rs'} = 2P^{m'} \begin{matrix} p & q \\ j & m \end{matrix} \begin{matrix} p & q \\ m & j \end{matrix}$$

¹We owe this nice abbreviation to Ben Andrews.

yields

$$\begin{aligned} \tilde{R}_{ij} = & \frac{1}{8} \frac{\partial^2 g_{ip}}{\partial x^i \partial x^q} + \frac{\partial^2 g_{iq}}{\partial x^i \partial x^p} - \frac{\partial^2 g_{ip}}{\partial x^i \partial x^p} - \frac{\partial^2 g_{iq}}{\partial x^i \partial x^q} P^{m\prime} \left(\begin{matrix} p & q \\ j & m \end{matrix} \right. \\ & \frac{1}{8} \frac{\partial^2 g_{ip}}{\partial x^j \partial x^q} + \frac{\partial^2 g_{jq}}{\partial x^i \partial x^p} - \frac{\partial^2 g_{ip}}{\partial x^j \partial x^p} - \frac{\partial^2 g_{jq}}{\partial x^i \partial x^q} P^{m\prime} \left(\begin{matrix} p & q \\ i & m \end{matrix} \right. \\ & + \quad : \end{aligned}$$

The symbol of $E(g) = 2h$ is obtained from \tilde{R}_{ij} by replacing $\frac{\partial}{\partial x^i}$ by a cotangent vector ∂_i in the highest (second) order terms

$$DE(g)(\partial_i)\partial_j = P^{m\prime} \left(\partial_i g_{jm} + \partial_j g_{im} - \partial_i g_{jm} - \partial_j g_{im} \right) :$$

Since the sectional curvature is negative, $P^{m\prime}$ is positive and the eigenvalues of the symbol are nonnegative. One checks that the integrability condition is $L(h_{ij}) = 0$; where

$$L(T)_k + h^{ij} r_{ij} T_{jk} - \frac{1}{2} h^{ij} r_{ik} T_{ij} :$$

By Theorem 5.1 of [7], a solution to the XCF exists for short time. ■

3 Evolution of the Einstein tensor

We find it convenient to express the Einstein tensor as

$$P^{mn} = \frac{1}{4} g^{ijm} g^{kn} R_{ijk} , :$$

Lemma 4 The evolution of the Einstein tensor is given by

$$\frac{\partial}{\partial t} P^{ij} = r_k r_{ij} + P^{ik} P^{jl} - P^{ik} P^{jl} - \det P g^{ij} - H P^{ij}$$

where $H = g^{ij} h_{ij}$ and $\det P = \det P^{k\prime} = \det g^{k\prime}$:

Proof. We have

$$\begin{aligned} \frac{\partial}{\partial t} R_{ijk} = & r_{ij} r_{jk} + r_{jk} r_{ij} - r_{ij} r_{jk} h_j - r_{jk} r_{ij} h_k \\ & + g^{pq} (R_{ijkp} h_q + R_{ijp} h_{qk}) : \end{aligned}$$

Since the evolution of the volume form is given by $\frac{\partial}{\partial t} g^{ijk} = H g^{ijk}$ and $\frac{\partial}{\partial t} g^{ijk} = H g^{ijk}$; we have

$$\begin{aligned} \frac{\partial}{\partial t} P^{mn} = & \frac{1}{4} g^{ijm} g^{kn} (r_{ij} r_{jk} + r_{jk} r_{ij} - r_{ij} r_{jk} h_j - r_{jk} r_{ij} h_k) \\ & \frac{1}{4} g^{ijm} g^{kn} g^{pq} (R_{ijkp} h_q + R_{ijp} h_{qk}) - 2H P^{mn} \\ = & g^{ijm} g^{kn} r_{ij} r_{jk} h_j - \frac{1}{2} g^{ijm} g^{kn} g^{pq} R_{ijp} h_{qk} - 2H P^{mn} : \end{aligned}$$

The lemma follows from the identity

$$\frac{1}{2} g^{ijm} g^{kln} g^{pq} R_{ijp} h_{qk} + H P^{mn} = \det P g^{mn} :$$

■

4 Monotonicity of the volume of the Einstein tensor

Proposition 5 If $(M; g)$ is a 3-manifold with negative sectional curvature, then $\text{vol}(P_{ij})$ is nondecreasing under the XCF.

This follows from the more general computation

Lemma 6 For any $\alpha \in \mathbb{R}$

$$\begin{aligned} \frac{d}{dt} \int_M (\det P)^\alpha dV &= \int_M \left(\frac{1}{2} T^{ijk} T^{jik} - T^{i2} \right) (\det P)^\alpha dV \\ &\quad + (1 - 2\alpha) \int_M (\det P)^\alpha H dV \end{aligned}$$

where $T^{ijk} = P^i{}_r P^j{}_s P^k{}_t$; $T^i = V_{jk} T^{ijk} = P^{ij} r_j \log \det P$; and the norms are with respect to the metric V_{ij} :

Decomposing T^{ijk} into its irreducible components

$$T^{ijk} = E^{ijk} - \frac{1}{10} P^{ij} T^k + P^{ik} T^j + \frac{2}{5} P^{jk} T^i;$$

where the coefficients $\frac{1}{10}$ and $\frac{2}{5}$ are chosen so that $V_{ij} E^{ijk} = V_{ik} E^{ijk} = V_{jk} E^{ijk}$; we find that

$$T^{ijk} T^{jik} = E^{ijk} E^{jik} + T^{i2} :$$

Taking $\alpha = 1/2$ in the lemma, we have

$$\frac{d}{dt} \int_M (\det P)^{1/2} dV = \frac{1}{4} \int_M (E^{ijk} E^{jik} - T^{i2}) (\det P)^{1/2} dV \geq 0$$

and the proposition follows.

5 Approach to hyperbolic in an integral sense

We show that an integral measure of the difference of the metric from hyperbolic is monotone decreasing. Let

$$J = \int_M \frac{P}{3} (\det P)^{1/3} dV$$

where $P = g_{ij}P^{ij}$: By the arithmetic-geometric mean inequality, the integrand is nonnegative, and identically zero if and only if $P_{ij} = \frac{1}{3}P g_{ij}$; i.e., g_{ij} has constant curvature.

Theorem 7 Under the cross curvature flow $\frac{dJ}{dt} \leq 0$:

Proof. We compute $\frac{d}{dt} \int_M P dV = 3 \int_M \det P dV$: Combining this with the previous lemma with $n = 3$ and the decomposition of T^{ijk} into its irreducible components, we have

$$\frac{dJ}{dt} = \int_M \left[\frac{1}{6} \sum_{ijk} E^{ijk} E^{jik} + \frac{1}{3} T^i{}_i{}^2 - (\det P)^{1/3} dV \right]$$

which is nonpositive (0 if and only if g_{ij} has constant negative sectional curvature). ■

6 A maximum principle estimate

Let $\Delta = P^{ij} \nabla_i \nabla_j = \nabla_i \nabla_j P^{ij}$; which is an elliptic operator.

Proposition 8

$$\frac{\partial}{\partial t} \log \det P = \Delta \log \det P + \frac{1}{2} T^{ijk} T^{jik} - 2H :$$

By the maximum principle

$$\frac{d}{dt} \min_M \log \det P(t) \leq \max_M H(x;t) : \det P(x;t) = \min_M \det P(t) + 2X(t) :$$

Suppose for some $\epsilon < 1$ we have $\inf_M \int_{[0,T]} \det P = 0$: It is not difficult to show that there then exists a sequence of times $t_i \rightarrow T$ such that $\min_M \log \det P(t_i) \rightarrow -1$ and $X(t_i) \rightarrow 1$: Hence there exists $x_i \in M$ such that $\det P(x_i; t_i) \rightarrow 0$ and $H(x_i; t_i) \rightarrow 1$: Hence there exists a sequence of points and times where one of the sectional curvatures tends to zero and another tends to minus infinity.

7 Conclusion

In view of the well-developed theory of Ricci flow [9] and the ground breaking work of Perelman [11] on the second author's program for Ricci flow as an approach to the Thurston Geometrization and Poincare conjectures, it is hopeful that further progress can be made on the cross curvature flow. Recently Ben Andrews [1] has obtained new estimates for the cross curvature flow. In the case the universal cover of the initial 3-manifold is isometrically embedded as a hypersurface in Euclidean or Minkowski 4-space, the Gauss curvature flow

(see [4], [12], [3], [8], [2] for earlier works on the GCF) induces the XCF for the metric. In this case Andrews has proved convergence results. In general, he expects long time existence and convergence of the XCF to reduce to proving local in time regularity (higher derivative estimates).

References

- [1] Andrews, Ben, personal communication.
- [2] Andrews, Ben, Gauss curvature flow: the fate of the rolling stones, *Invent. Math.* 138 (1999), no. 1, 151{161; Motion of hypersurfaces by Gauss curvature, *Pacific J. Math.* 195 (2000), no. 1, 1{34.
- [3] Chow, Bennett, On Hamack's inequality and entropy for the Gaussian curvature flow, *Comm. Pure Appl. Math.* 44 (1991), no. 4, 469{483.
- [4] Firey, William J., Shapes of worn stones, *Mathematika* 21 (1974), 1{11.
- [5] Gao, L. Zhiyong; Yau, Shing-Tung, The existence of negatively Ricci curved metrics on three-manifolds, *Invent. Math.* 85 (1986), no. 3, 637{652.
- [6] Gromov, Mikhail; Thurston, William P., Pinching constants for hyperbolic manifolds, *Invent. Math.* 89 (1987), no. 1, 1{12.
- [7] Hamilton, Richard S., Three-manifolds with positive Ricci curvature, *J. Differential Geom.* 17 (1982), no. 2, 255{306.
- [8] Hamilton, Richard S., Remarks on the entropy and Hamack estimates for the Gauss curvature flow, *Comm. Anal. Geom.* 2 (1994), no. 1, 155{165.
- [9] Hamilton, Richard S., The formation of singularities in the Ricci flow, *Surveys in differential geometry, Vol. II* (Cambridge, MA, 1993), 7{136, *Internat. Press, Cambridge, MA*, 1995; Four-manifolds with positive isotropic curvature, *Comm. Anal. Geom.* 5 (1997), no. 1, 1{92; Non-singular solutions of the Ricci flow on three-manifolds, *Comm. Anal. Geom.* 7 (1999), no. 4, 695{729.
- [10] Lohkamp, Joachim, Metrics of negative Ricci curvature, *Ann. of Math.* (2) 140 (1994), no. 3, 655{683.
- [11] Perelman, Grisha, The entropy formula for the Ricci flow and its geometric applications, *math.DG/0211159*; Ricci flow with surgery on three-manifolds, *math.DG/0303109*; Finite extinction time for the solutions to the Ricci flow on certain three-manifolds, *math.DG/0307245*.
- [12] Tso, Kaising, Deforming a hypersurface by its Gauss-Kronecker curvature, *Comm. Pure Appl. Math.* 38 (1985), no. 6, 867{882.
- [13] Ye, Rugang, Ricci flow, Einstein metrics and space forms, *Trans. Amer. Math. Soc.* 338 (1993), no. 2, 871{896.