

THE SPECTRUM OF A RANDOM GEOMETRIC GRAPH IS CONCENTRATED

SANATAN RAI

ABSTRACT. Consider n points distributed uniformly in $[0, 1]^d$. Form a graph by connecting two points if their mutual distance is no greater than $r(n)$. This gives a random geometric graph, $G(\mathcal{X}_n; r(n))$, which is connected for appropriate $r(n)$. We show that the spectral measure of the transition matrix of the simple random walk (SRW) on $G(\mathcal{X}_n; r(n))$ is concentrated, and in fact converges to that of the graph on the deterministic grid.

1. INTRODUCTION

Let S be finite set contained in $[0, 1]^d$. Form a graph by connecting two points, $u, v \in S$ if $\|u - v\| \leq r$, obtaining a graph $G(S; r)$.

Let \mathcal{X}_n be a set on n points distributed iid $\text{Unif}[0, 1]^d$, we call $G \sim G(\mathcal{X}_n; r(n))$ a random geomtric graph. The function $r(n)$ is chosen to be such that $r(n) \downarrow 0$ as $n \uparrow \infty$, but such that G is a.s. and **whp** connected. Herein, **whp** denotes with high probability, ie with a probability greater than $1 - n^{-c}$ for some constant $c > 0$. We shall write \mathcal{D}_n for the set of n grid points that are the intersections of axes parallel lines with separation $n^{-1/d}$. Hence, $G(\mathcal{D}_n; r(n))$, is a deterministic graph.

For a graph G , $P(G)$ shall denote the transition probability matrix for the simple random walk (SRW) on G . That is:

$$P(G)_{uv} = \frac{[u \sim v]}{|\mathcal{N}(u)|},$$

where $\mathcal{N}(u)$ is the set of neighbours of vertex u , and $[u \sim v]$ is the indicator of the event $u \sim v$. Henceforth, we shall write $P(\mathcal{X}_n)$ and $P(\mathcal{D}_n)$ for $P(G(\mathcal{X}_n; r(n)))$ and $P(G(\mathcal{D}_n; r(n)))$, respectively, with $\text{Spec}(\mathcal{X}_n)$ and $\text{Spec}(\mathcal{D}_n)$ denoting their spectra. Furthermore, $\mu(\mathcal{X}_n)$ and $\mu(\mathcal{D}_n)$ shall stand for their spectral measures respectively.

Our main result is:

Theorem 1.1. *$\text{Spec}(\mathcal{X}_n)$ and $\text{Spec}(\mathcal{D}_n)$ are asymptotically equidistributed **whp**. Moreover, for $r(n)$ such that $M_n = o(r_n)$ a.s. and **whp**, the random spectral measure $\mu(\mathcal{X}_n)$:*

$$\mu(\mathcal{X}_n)(-\infty, x] := \frac{1}{n} |\{\lambda \in \text{Spec}(\mathcal{X}_n) : \lambda \leq x\}|$$

2000 *Mathematics Subject Classification.* **Primary** 60D05; **Secondary** 34L20.
Key words and phrases. Random geometric graphs; spectral measure.

is concentrated, ie, there exists a sequence of deterministic measures $\{\mu(\mathcal{D}_n)\}$, and $c_d > 0$ such that $(a(n) := n\pi_d r(n)^d)$:

$$\mathbb{P} \left\{ \|\mu(\mathcal{X}_n) - \mu(\mathcal{D}_n)\|_{\text{WS}} > \frac{t}{a(n)^{1/4}} \right\} \leq \frac{16na(n)^{1/4}}{t} \times \left[2 \exp \left(-\frac{1}{2} \left(\frac{t^4}{8t^4 + 4096} \right)^2 a(n) \right) + \exp \left(-c_d \frac{t^8}{512} a(n) \right) \right].$$

Here M_n is the minimum bottleneck matching distance between \mathcal{X}_n and \mathcal{D}_n in d -dimensions.

The norm $\|\cdot\|_{\text{WS}}$ is the **Wasserstein distance**:

$$\|\mu - \nu\|_{\text{WS}} := \sup_{f \text{ Lipschitz}} \left| \int f d\mu - \int f d\nu \right|.$$

1.1. Related work. Eigenvalues of random matrices with iid entries have been studied extensively, can be said to have begun with the work Wigner (1958), wherein he showed that the spectral measure of $N^{-1/2}X$ converges to the semicircle law, whenever X is Hermitian with iid complex entries. The rate of convergence for this class of random matrices was computed in Bai (1993). The work closest in spirit to ours is the paper by Guionnet and Zeituni (2000), wherein they prove rates of convergence using concentration of measure ideas. There is a vast body of research in this area, and rather than provide a comprehensive review, we only note here that the results and methods from this literature are not applicable to our case. In many, an exact expression of the limiting distributions of eigenvalues is used, and in some the Stiltjes transform is the main tool. Neither technique seems useful in this case. On the other hand, our proofs are elementary, and require only a knowledge of Chernoff-Höfdding bounds and basic real analysis.

Bounds on the second eigenvalue were obtained by Boyd et al. (2004), and used to establish the mixing times of the SRW and the fastest walk on $G(\mathcal{X}_n; r(n))$.

We also note that a Wilandt-Hoffman theorem for the eigenvalues of discrete matrix approximations to learning kernels were proved in Koltchinskii and Giné (2000); Koltchinskii (1998).

2. PROOF OF THE MAIN THEOREM

Two sequences of real numbers $\{x_n\}$ and $\{y_n\}$ are said to be **asymptotically equidistributed**, if for any L^1 function f (Pólya and Szégo (1998)):

$$\lim_{n \uparrow \infty} \left| \frac{1}{n} \sum_{i=1}^n f(x_i) - \frac{1}{n} \sum_{i=1}^n f(y_i) \right| = 0.$$

Suppose $\{A_n\}$ and $\{B_n\}$ are two sequences of matrices, such that $A_n, B_n \in \mathbb{M}_{n \times n}(\mathbb{R})$. We say that the two sequences are asymptotically equivalent iff

- (1) $\|A_n\|$ and $\|B_n\|$ are uniformly bounded, and
- (2) $\|A_n - B_n\|_{\text{HS}} \rightarrow 0$ as $n \uparrow \infty$.

Here $\|\cdot\|$ is the usual operator norm:

$$\|A\| := \sup_{\|x\|=1} |Ax|,$$

and $\|\cdot\|_{\text{HS}}$ is the **Hilbert-Schmidt norm**:

$$\|A\|_{\text{HS}} := \left(\frac{1}{n} \sum_{i,j} A_{ij}^2 \right)^{1/2} = \left(\frac{1}{n} \text{Tr}(A^T A) \right)^{1/2}.$$

With a few straightforward estimates, our main result shall follow easily from (Gray (1971)):

Theorem 2.1. *For any two matrices A and B :*

$$\left| \frac{1}{n} \sum_{i=1}^n \lambda_i(A) - \frac{1}{n} \sum_{i=1}^n \lambda_i(B) \right| \leq \|A - B\|_{\text{HS}}.$$

Therefore, if $\{A_n\}$ and $\{B_n\}$ are two asymptotically equivalent sequences of matrices, then their spectra are asymptotically equivalent.

To prove theorem 1.1 it suffices to show that $\{P(\mathcal{X}_n)\}$ and $\{P(\mathcal{D}_n)\}$ are asymptotically equivalent **whp**, and that $\|P(\mathcal{X}_n) - P(\mathcal{D}_n)\|_{\text{HS}}$ is concentrated.

Let M_n denote the length of the minimum bottleneck matching between \mathcal{X}_n and \mathcal{D}_n :

$$M_n \equiv M_n := \min_{\substack{\phi: \mathcal{X}_n \rightarrow \mathcal{D}_n \\ \phi \text{ matching}}} \max_{u \in \mathcal{X}_n} \|u - \phi(u)\|.$$

Then it is well known that:

Theorem 2.2.

$$M_n = \begin{cases} \mathcal{O}\left(\left(\frac{\log n}{n}\right)^{1/d}\right), & \text{when } d \geq 3, \text{ [Shor and Yukich (1991)]} \\ \mathcal{O}\left(\left(\frac{\log^{3/2} n}{n}\right)^{1/2}\right), & \text{when } d = 2, \text{ [Leighton and Shor (1989)]} \\ \mathcal{O}\left(\sqrt{\frac{\log \varepsilon^{-1}}{n}}\right), & \text{w.p. } 1 - \varepsilon, \text{ when } d = 1. \text{ [Goel et al. (2004)]} \end{cases}$$

Henceforth, for $u \in \mathcal{X}_n$ we shall write u' for its matched point in \mathcal{D}_n under a minimum bottleneck matching. We shall denote by $\mathcal{N}(u)$ the set of neighbours of u , and by $\mathcal{N}(u, u')$, the set of neighbours of u that are mapped to a neighbour of u' . We are now in a position to state a concentration result:

Lemma 2.3. *If $r(n)$ is such that $M_n = o(r(n))$ a.s. and **whp**, then there is a constant $c_d > 0$:*

$$\mathbb{P}\left\{\|P(\mathcal{X}_n) - P(\mathcal{D}_n)\|_{\text{HS}}^2 > \frac{t}{a(n)}\right\} \leq 2n \times \left[2 \exp\left(-\frac{1}{2} \left(\frac{t}{8t+16}\right)^2 a(n)\right) + \exp\left(-c_d \frac{t^2}{2} a(n)\right)\right]$$

where $a(n) := n\pi_d r(n)^d$.

Proof.

$$\begin{aligned}
\|P(\mathcal{X}_n) - P(\mathcal{D}_n)\|_{\text{HS}}^2 &= \frac{1}{n} \sum_{u \in \mathcal{X}_n} \sum_{v \in \mathcal{X}_n} \left(\frac{[u \sim v]}{|\mathcal{N}(u)|} - \frac{[u' \sim v']}{|\mathcal{N}(u')|} \right)^2 \\
&= \frac{1}{n} \sum_{u \in \mathcal{X}_n} \left(\frac{1}{|\mathcal{N}(u)|} + \frac{1}{|\mathcal{N}(u')|} - \frac{2|\mathcal{N}(u, u')|}{|\mathcal{N}(u)| |\mathcal{N}(u')|} \right) \\
(1) \quad &\leq \max_u \left(\frac{1}{|\mathcal{N}(u)|} + \frac{1}{|\mathcal{N}(u')|} - \frac{2|\mathcal{N}(u, u')|}{|\mathcal{N}(u)| |\mathcal{N}(u')|} \right).
\end{aligned}$$

So that:

$$\begin{aligned}
&\mathbb{P} \left\{ \|P(\mathcal{X}_n) - P(\mathcal{D}_n)\|_{\text{HS}}^2 > \frac{t}{a(n)} \right\} \leq \\
&n \max_u \mathbb{P} \left\{ \left(\frac{1}{|\mathcal{N}(u)|} + \frac{1}{|\mathcal{N}(u')|} - \frac{2|\mathcal{N}(u, u')|}{|\mathcal{N}(u)| |\mathcal{N}(u')|} \right) > \frac{t}{a(n)} \right\}
\end{aligned}$$

Now

$$\begin{aligned}
&\mathbb{P} \left\{ \left(\frac{1}{|\mathcal{N}(u)|} + \frac{1}{|\mathcal{N}(u')|} - \frac{2|\mathcal{N}(u, u')|}{|\mathcal{N}(u)| |\mathcal{N}(u')|} \right) > \frac{t}{a(n)} \right\} \\
&\leq \mathbb{P} \left\{ \frac{2}{|\mathcal{N}(u')|} + \left| \frac{1}{|\mathcal{N}(u)|} - \frac{1}{|\mathcal{N}(u')|} \right| - \frac{2|\mathcal{N}(u, u')|}{|\mathcal{N}(u)| |\mathcal{N}(u')|} > \frac{t}{a(n)} \right\} \\
&\leq \mathbb{P} \left\{ \left| \frac{1}{|\mathcal{N}(u)|} - \frac{1}{|\mathcal{N}(u')|} \right| > \frac{t}{8|\mathcal{N}(u')|} \right\} + \underbrace{\mathbb{P} \left\{ \frac{|\mathcal{N}(u)| - |\mathcal{N}(u, u')|}{|\mathcal{N}(u)|} > \frac{t}{8} \right\}}_{=:\ddagger},
\end{aligned}$$

since $n\pi_d r(n)^d/4 \leq |\mathcal{N}(u')| \leq n\pi_d r(n)^d$. To bound \ddagger , observe that if $\|u - v\| \leq r(n) - 2M_n$, then $\|u' - v'\| \leq r(n)$. Thus, all points within a radius of $r(n) - 2M_n$ map to neighbours of u' . Furthermore:

$$\begin{aligned}
\ddagger &\leq \mathbb{P} \left\{ \left| |\mathcal{N}(u)| - |\mathcal{N}(u')| \right| > \frac{t}{8t+16} |\mathcal{N}(u')| \right\} \\
&+ \mathbb{P} \left\{ \left| |\mathcal{N}(u')| - |\mathcal{N}(u, u')| \right| > \frac{t}{16} |\mathcal{N}(u')| \right\}
\end{aligned}$$

Now $|\mathcal{N}(u, u')|$ is stochastically greater than $\text{Bin}(n, \pi_d r(n)^d (1 - 2M_n/n)^d)$, so that the last term satisfies:

$$\mathbb{P} \left\{ \left| |\mathcal{N}(u')| - |\mathcal{N}(u, u')| \right| > \frac{t}{16} |\mathcal{N}(u')| \right\} \leq 2 \exp \left(-c_d \frac{t^2}{2} n\pi_d r(n)^d \right),$$

where $c_d > 0$ is such that:

$$c_d \frac{t}{8} \leq \min \left(\frac{1+t/8}{(1-2M_n^+/r(n))^d} - 1, \left| \frac{1-t/8}{(1-2M_n^+/r(n))^d} - 1 \right| \right)$$

for all large enough n . Here M_n^+ is a constant depending only on n such that $M_n \leq M_n^+$, a.s. and **whp**. Since we have chosen $r(n)$ such that $M_n/r(n) \rightarrow 0$, such a constant c_d always exists.

Putting all our estimates together, and using lemma A.1 we obtain:

$$\mathbb{P} \left\{ \|P(\mathcal{X}_n) - P(\mathcal{D}_n)\|_{\text{HS}}^2 > \frac{t}{a(n)} \right\} \leq 2n \times \left[\exp \left(-\frac{1}{2} \left(\frac{t}{t+8} \right)^2 a(n) \right) + \exp \left(-\frac{1}{2} \left(\frac{t}{8t+16} \right)^2 a(n) \right) + \exp \left(-c_d \frac{t^2}{2} a(n) \right) \right]$$

and the conclusion of the lemma follows. \blacksquare

Proposition 2.4. *When $r(n)$ is such that $M_n = r(n)$ a.s. and **whp**, for $g(n) \uparrow \infty$, the two sequences $\{P(\mathcal{X}_n)\}$ and $\{P(\mathcal{D}_n)\}$ are asymptotically equivalent **whp**.*

Proof. Since matrices are stochastic, $\|P(\mathcal{X}_n)\| = \|P(\mathcal{D}_n)\| = 1$, and hence they are of uniformly bounded norm. Lemma 2.3 implies that $\|P(\mathcal{X}_n) - P(\mathcal{D}_n)\|_{\text{HS}} \rightarrow 0$, **whp**. Hence $\{P(\mathcal{X}_n)\}$ and $\{P(\mathcal{D}_n)\}$ are asymptotically equivalent **whp**. \blacksquare

We now prove that the concentration of the Hilbert-Schmidt norm of the matrices $P(\mathcal{X}_n)$ implies concentration of the spectrum. We first need a lemma:

Lemma 2.5.

$$\mathbb{P} \left\{ \sup_{f \text{ Lipschitz}} \left| \frac{1}{n} \sum_{i=1}^n f(\lambda_i(\mathcal{X}_n)) - \frac{1}{n} \sum_{i=1}^n f(\lambda_i(\mathcal{D}_n)) \right| > 4\varepsilon \right\} \leq \frac{2}{\varepsilon} \mathbb{P} \{ \|P(\mathcal{X}_n) - P(\mathcal{D}_n)\|_{\text{HS}} > \varepsilon^2 \}.$$

Proof. When f is linear, the result follows directly from theorem 2.1 and lemma 2.4. To obtain a uniform bound over *all* Lipschitz f , we use the trick in Guionnet and Zeituni (2000) of approximating by a finite class of functions, for a given error bound.

To that end, let f be Lipschitz on $[-1, 1]$, and fix $\varepsilon > 0$. Define: $g_\varepsilon(x) := x[0 \leq x \leq \varepsilon] + \varepsilon[x > \varepsilon]$. Then for f_ε recursively defined by:

$$f_\varepsilon(x) = \sum_{i=0}^{\lceil x/\varepsilon \rceil} (2[f((i+1)\varepsilon) > f(i\varepsilon)] - 1) g_\varepsilon(x - i\varepsilon),$$

we must have $\|f - f_\varepsilon\| \leq \varepsilon$. Thus we can approximate any Lipschitz f to within ε by a weighted sum of at most $2/\varepsilon$ functions, each of which has two constant portions joined by a linear part. Since the weights are ± 1 , we have:

$$\begin{aligned} & \mathbb{P} \left\{ \sup_{f \text{ Lipschitz}} \left| \frac{1}{n} \sum_{i=1}^n f(\lambda_i(\mathcal{X}_n)) - \frac{1}{n} \sum_{i=1}^n f(\lambda_i(\mathcal{D}_n)) \right| > 4\varepsilon \right\} \\ & \leq \frac{2}{\varepsilon} \sup_k \mathbb{P} \left\{ \left| \frac{1}{n} \sum_{i=1}^n g_k(\lambda_i(\mathcal{X}_n)) - \frac{1}{n} \sum_{i=1}^n g_k(\lambda_i(\mathcal{D}_n)) \right| > \varepsilon^2 \right\} \\ & \leq \frac{2}{\varepsilon} \mathbb{P} \{ \|P(\mathcal{X}_n) - P(\mathcal{D}_n)\|_{\text{HS}} > \varepsilon^2 \} \end{aligned}$$

where $g_k(x) := g_\varepsilon(x - k\varepsilon)$, and the last inequality follows from theorem 2.1. \blacksquare

Proof of theorem 1.1. Proposition 2.4 and theorem 2.1 imply that the spectra are asymptotically equidistributed **whp**. By lemma 2.5, and the estimate in lemma

2.3:

$$\begin{aligned}
& \mathbb{P} \left\{ \|\mu(\mathcal{X}_n) - \mu(\mathcal{D}_n)\|_{\text{WS}} > \frac{t}{a(n)^{1/4}} \right\} \\
& \leq \frac{8a(n)^{1/4}}{t} \mathbb{P} \left\{ \|P(\mathcal{X}_n) - P(\mathcal{D}_n)\|_{\text{HS}} > \frac{t^4}{16\sqrt{a(n)}} \right\} \\
& \leq \frac{16na(n)^{1/4}}{t} \times \\
& \left[2 \exp \left(-\frac{1}{2} \left(\frac{t^4}{8t^4 + 4096} \right)^2 a(n) \right) + \exp \left(-c_d \frac{t^8}{512} a(n) \right) \right]
\end{aligned}$$

■

3. DISCUSSION AND OPEN PROBLEMS

Note that if we define $\mu_n^{\{x\}}$ to be the empirical measure

$$\mu_n^{\{x\}}(\alpha) = \frac{1}{n} \sum_{i=1}^n \delta(\alpha - x_i),$$

then $\{x_n\}$ and $\{y_n\}$ are asymptotically equidistributed if and only if $\|\mu_n^{\{x\}} - \mu_n^{\{y\}}\|_{\text{WS}} \rightarrow 0$, since we may approximate integrable functions by Lipschitz ones.

Simulations suggest that in fact, the eigenvalues of $P(\mathcal{X}_n)$ and $P(\mathcal{D}_n)$ are **asymptotically absolutely equally distributed** (Trench (2003)):

Conjecture 1. For any $f \in \mathcal{C}[-1, 1]$:

$$\lim_{n \uparrow \infty} \frac{1}{n} \sum_{i=1}^n |f(\lambda_i(\mathcal{X}_n)) - f(\lambda_i(\mathcal{D}_n))|^2 = 0,$$

where $\lambda_i(\mathcal{X}_n)$ and $\lambda_i(\mathcal{D}_n)$ are the i th largest eigenvalues of $P(\mathcal{X}_n)$ and $P(\mathcal{D}_n)$ respectively.

Such a result would have followed immediately, were it known that the matrices were symmetric. This is of course, not true. However, the matrices are *almost* symmetric in the asymptotic limit, so one would expect the matrices to be *almost* diagonalisable via unitary matrices. Then the conjecture would follow immediately from the theory of asymptotically equivalent matrices. As it stands, one needs some other manner of bounding the differences:

$$\frac{1}{n} \sum_{i=1}^n |\lambda_i(\mathcal{X}_n) - \lambda_i(\mathcal{D}_n)|^2,$$

to obtain say, a Wilandt-Hoffman type theorem. Then the conjecture would follow, the Stone-Weierstrass theorem.

It also seems reasonable to posit exponential tail bounds for the Wielandt-Hoffman type result.

Another natural question to ask is: what is the behaviour of the resolvent $R_n(x) := (I - xP(\mathcal{X}_n))^{-1}$, for $x \notin \text{Spec}(\mathcal{X}_n)$. Whereas, $(I - P(\mathcal{X}_n)) \rightarrow \Delta$, the asymptotic behaviour of the resolvent is not known. Convergence results for the resolvents of random walks in random environments, specifically, on \mathbb{Z}^d with ergodic (random) bond percolation were proved in Künnemann (1983).

APPENDIX A. AUXILLIARY LEMMATA

Let $X \sim \text{Bin}(n, p)$, then we have the following two tail bounds:

Lemma A.1 (Chernoff-Höfdding bounds for reciprocals). *For $t \geq 0$:*

$$\mathbb{P} \left\{ \left| \frac{1}{X} - \frac{1}{\mathbb{E}X} \right| > \frac{t}{\mathbb{E}X} \right\} \leq 2 \exp \left(-\frac{1}{2} \left(\frac{t}{1+t} \right)^2 \mathbb{E}X \right).$$

Proof. Note that:

$$\begin{aligned} \mathbb{P} \left\{ \left| \frac{1}{X} - \frac{1}{\mathbb{E}X} \right| > \frac{t}{\mathbb{E}X} \right\} &= \mathbb{P} \{ |X - \mathbb{E}X| > tX \} \\ &\leq \mathbb{P} \{ |X - \mathbb{E}X| > t(\mathbb{E}X - |X - \mathbb{E}X|) \} \\ &= \mathbb{P} \left\{ |X - \mathbb{E}X| > \frac{t}{1+t} \mathbb{E}X \right\} \\ &\leq 2 \exp \left(-\frac{1}{2} \left(\frac{t}{1+t} \right)^2 \mathbb{E}X \right). \end{aligned}$$

■

REFERENCES

- Zhi Dong Bai. Convergence rate of expected spectral distributions of large random matrices, part i: Wigner matrices, part ii: sample covariance matrices. *Annals of Probability*, 1993.
- Stephen Boyd, Arpita Ghosh, Balaji Prabhakar, and Devavrat Shah. Mixing times of random walk on geometric random graphs. *Submitted for publication*, 2004.
- Ashish Goel, Sanatan Rai, and Bhaskar Krishnamachari. Sharp thresholds for monotone properties in random geometric graphs. In *Proceedings of the Symposium on the Theory of Computing*, pages 13–23, 2004.
- Robert M. Gray. Toeplitz and circulant matrices: A review. *Revised 2002*, URL: <http://ee.stanford.edu/gray/toeplitz.pdf>, 1971.
- Alice Guionnet and Ofer Zeituni. Concentration of the spectral measure for large matrices. *Electronic Communications in Probability*, 9:119–36, 2000.
- Vladimir I. Koltchinskii. Asymptotics of spectral projections of some random matrices approximating integral operators. In Ernst Eberlein, Marjorie Hahn, and Michel Talagrand, editors, *High Dimensional Probability*, volume 43 of *Progress in Probability*, pages 191–227. Birkhäuser, 1998.
- Vladimir I. Koltchinskii and Evarist Giné. Random matrix approximation of spectra of integral operator. *Bernoulli*, 2000.
- Rolf Künnemann. The diffusion limit for reversible jump processes on \mathbb{Z}^d with ergodic random bond conductivities. *Communications in Mathematical Physics*, 90:27–68, 1983.
- Frank T. Leighton and Peter W. Shor. Tight bounds for minimax grid matching, with applications to the average case analysis of algorithms. *Combinatorica*, 9: 161–87, 1989.
- George Pólya and Gabor Szégo. *Problems and Theorems in Analysis I*. Springer-Verlag, reprint edition, January 1998.
- Peter W. Shor and Joseph E. Yukich. Minmax grid matching and empirical measures. *The Annals of Probability*, 19(3):1338–48, July 1991.

William F. Trench. Absolute equal distribution of the spectra of hermitian matrices.

Linear Algebra and Its Applications, 2003.

Eugene P. Wigner. On the distribution of the roots of certain symmetric matrices.

Annals of Mathematics, 67:325–27, 1958.

E-mail address: `sanat@stanford.edu`

DEPARTMENT OF MANAGEMENT SCIENCE AND ENGG., STANFORD UNIVERSITY, CA 94305.