

A q -ANALOG OF EULER'S DECOMPOSITION FORMULA FOR THE DOUBLE ZETA FUNCTION

DAVID M. BRADLEY

ABSTRACT. The double zeta function was first studied by Euler in response to a letter from Goldbach in 1742. One of Euler's results for this function is a decomposition formula, which expresses the product of two values of the Riemann zeta function as a finite sum of double zeta values involving binomial coefficients. In this note, we establish a q -analog of Euler's decomposition formula. More specifically, we show that Euler's decomposition formula can be extended to what might be referred to as a "double q -zeta function" in such a way that Euler's formula is recovered in the limit as q tends to 1.

1. INTRODUCTION

The Riemann zeta function is defined for $\Re(s) > 1$ by

$$\zeta(s) := \sum_{n=1}^{\infty} \frac{1}{n^s}. \quad (1)$$

Accordingly,

$$\zeta(s, t) := \sum_{n=1}^{\infty} \frac{1}{n^s} \sum_{k=1}^{n-1} \frac{1}{k^t}, \quad \Re(s) > 1, \quad \Re(s+t) > 2, \quad (2)$$

is known as the double zeta function. The sums (2), and more generally those of the form

$$\zeta(s_1, s_2, \dots, s_m) := \sum_{k_1 > k_2 > \dots > k_m > 0} \prod_{j=1}^m \frac{1}{k_j^{s_j}}, \quad \sum_{j=1}^m \Re(s_j) > n, \quad n = 1, 2, \dots, m, \quad (3)$$

have attracted increasing attention in recent years; see eg. [2, 4, 5, 6, 8, 9, 10, 12, 13, 18]. The survey articles [7, 14, 22, 23, 25] provide an extensive list of references. In (3) the sum is over all positive integers $k_1, \dots, k_m$ satisfying the indicated inequalities. Note that with positive integer arguments, $s_1 > 1$ is necessary and sufficient for convergence.

The problem of evaluating sums of the form (2) for integers $s > 1$, $t > 0$ seems to have been first proposed in a letter from Goldbach to Euler [16] in 1742. (See also [15, 17]

Date: August 13, 2021.

1991 Mathematics Subject Classification. Primary: 11M41; Secondary: 11M06, 05A30, 33E20, 30B50.

Key words and phrases. Euler sums, multiple harmonic series, q -analog, multiple zeta values, q -series, shuffle product, Lambert series.

and [1, p. 253].) Among other results for (2), Euler proved that if $s - 1$ and $t - 1$ are positive integers, then the decomposition formula

$$\zeta(s)\zeta(t) = \sum_{a=0}^{s-1} \binom{a+t-1}{t-1} \zeta(t+a, s-a) + \sum_{a=0}^{t-1} \binom{a+s-1}{s-1} \zeta(s+a, t-a) \quad (4)$$

holds. A combinatorial proof of Euler's decomposition formula (4) based on the Drinfel'd integral representations [2, 4, 5, 7, 8]

$$\begin{aligned} \zeta(s) &= \int_{1 > x_1 > \dots > x_s > 0} \left(\prod_{i=1}^{s-1} \frac{dx_i}{x_i} \right) \frac{dx_s}{1-x_s}, \\ \zeta(s, t) &= \int_{1 > x_1 > \dots > x_{s+t} > 0} \left(\prod_{i=1}^{s-1} \frac{dx_i}{x_i} \right) \frac{dx_s}{1-x_s} \left(\prod_{i=s+1}^{s+t-1} \frac{dx_i}{x_i} \right) \frac{dx_{s+t}}{1-x_{s+t}}, \end{aligned} \quad (5)$$

and the shuffle multiplication rule satisfied by such integrals is given in [5, eq. (10)]. It is of course well-known that (4) can also be proved algebraically by summing the partial fraction decomposition [20, p. 48] [19, Lemma 3.1]

$$\frac{1}{x^s(c-x)^t} = \sum_{a=0}^{s-1} \binom{a+t-1}{t-1} \frac{1}{x^{s-a}c^{t+a}} + \sum_{a=0}^{t-1} \binom{a+s-1}{s-1} \frac{1}{c^{s+a}(c-x)^{t-a}} \quad (6)$$

over appropriately chosen integers x and c . (See eg. [3].)

A q -analog of (3) was independently introduced in [11, 21, 24] as

$$\zeta[s_1, s_2, \dots, s_m] := \sum_{k_1 > k_2 > \dots > k_m > 0} \prod_{j=1}^m \frac{q^{(s_j-1)k_j}}{[k_j]_q^{s_j}}, \quad (7)$$

where

$$[k]_q := \sum_{j=0}^{k-1} q^j = \frac{1-q^k}{1-q}, \quad 0 < q < 1.$$

Observe that we now have

$$\zeta(s_1, \dots, s_m) = \lim_{q \rightarrow 1^-} \zeta[s_1, \dots, s_m],$$

so that (7) represents a generalization of (3). In this note, we establish a q -analog of Euler's decomposition formula (4).

2. MAIN RESULT

Our q -analog of Euler's decomposition formula naturally requires only the $m = 1$ and $m = 2$ cases of (7); specifically the q -analogs of (1) and (2) given by

$$\zeta[s] = \sum_{n>0} \frac{q^{(s-1)n}}{[n]_q^s} \quad \text{and} \quad \zeta[s, t] = \sum_{n>k>0} \frac{q^{(s-1)n} q^{(k-1)t}}{[n]_q^s [k]_q^t}. \quad (8)$$

We also define, for convenience, the sum

$$\varphi[s] := \sum_{n=1}^{\infty} \frac{(n-1)q^{(s-1)n}}{[n]_q^s} = \sum_{n=1}^{\infty} \frac{nq^{(s-1)n}}{[n]_q^s} - \zeta[s]. \quad (9)$$

We can now state our main result.

Theorem 1. *If $s-1$ and $t-1$ are positive integers, then*

$$\begin{aligned} \zeta[s]\zeta[t] &= \sum_{a=0}^{s-1} \sum_{b=0}^{s-1-a} \binom{a+t-1}{t-1} \binom{t-1}{b} (1-q)^b \zeta[t+a, s-a-b] \\ &\quad + \sum_{a=0}^{t-1} \sum_{b=0}^{t-1-a} \binom{a+s-1}{s-1} \binom{s-1}{b} (1-q)^b \zeta[s+a, t-a-b] \\ &\quad - \sum_{j=1}^{\min(s,t)} \frac{(s+t-j-1)!}{(s-j)!(t-j)!} \cdot \frac{(1-q)^j}{(j-1)!} \varphi[s+t-j]. \end{aligned}$$

Observe that the limiting case $q = 1$ of Theorem 1 reduces to Euler's decomposition formula (4).

3. A DIFFERENTIAL IDENTITY

Our proof of Theorem 1 relies on the following identity.

Lemma 1. *Let s and t be positive integers, and let x and y be non-zero real numbers. Then for all real q ,*

$$\begin{aligned} \frac{1}{x^s y^t} &= \sum_{a=0}^{s-1} \sum_{b=0}^{s-1-a} \binom{a+t-1}{t-1} \binom{t-1}{b} \frac{(1-q)^b (1+(q-1)y)^a (1+(q-1)x)^{t-1-b}}{x^{s-a-b} (x+y+(q-1)xy)^{t+a}} \\ &\quad + \sum_{a=0}^{t-1} \sum_{b=0}^{t-1-a} \binom{a+s-1}{s-1} \binom{s-1}{b} \frac{(1-q)^b (1+(q-1)x)^a (1+(q-1)y)^{s-1-b}}{y^{t-a-b} (x+y+(q-1)xy)^{s+a}} \\ &\quad - \sum_{j=1}^{\min(s,t)} \frac{(s+t-j-1)!}{(s-j)!(t-j)!} \cdot \frac{(1-q)^j}{(j-1)!} \cdot \frac{(1+(q-1)y)^{s-j} (1+(q-1)x)^{t-j}}{(x+y+(q-1)xy)^{s+t-j}}. \end{aligned}$$

Proof. Apply the partial differential operator

$$\frac{1}{(s-1)!} \left(-\frac{\partial}{\partial x} \right)^{s-1} \frac{1}{(t-1)!} \left(-\frac{\partial}{\partial y} \right)^{t-1}$$

to both sides of the identity

$$\frac{1}{xy} = \frac{1}{x+y+(q-1)xy} \left(\frac{1}{x} + \frac{1}{y} + q-1 \right).$$

□

Observe that when $q = 1$, Lemma 1 reduces to the identity

$$\frac{1}{x^s y^t} = \sum_{a=0}^{s-1} \binom{a+t-1}{t-1} \frac{1}{x^{s-a} (x+y)^{t+a}} + \sum_{a=0}^{t-1} \binom{a+s-1}{s-1} \frac{1}{(x+y)^{s+a} y^{t-a}},$$

from which the partial fraction identity (6) (proved by induction in [19]) trivially follows.

4. PROOF OF THEOREM 1

First, observe that if $s > 1$ and $t > 1$, then from (8),

$$\zeta[s]\zeta[t] = \sum_{n=1}^{\infty} \sum_{u+v=n} \frac{q^{(s-1)u}}{[u]_q^s} \cdot \frac{q^{(t-1)v}}{[v]_q^t},$$

where the inner sum is over all positive integers u and v such that $u+v=n$. Next, apply Lemma 1 with $x = [u]_q$, $y = [v]_q$, noting that then

$$1 + (q-1)x = q^u, \quad 1 + (q-1)y = q^v, \quad x + y + (q-1)xy = [u+v]_q.$$

After interchanging the order of summation, there comes

$$\begin{aligned} \zeta[s]\zeta[t] &= \sum_{a=0}^{s-1} \sum_{b=0}^{s-1-a} \binom{a+t-1}{t-1} \binom{t-1}{b} (1-q)^b S[s, t, a, b] \\ &\quad + \sum_{a=0}^{t-1} \sum_{b=0}^{t-1-a} \binom{a+s-1}{s-1} \binom{s-1}{b} (1-q)^b S[t, s, a, b] \\ &\quad - \sum_{j=1}^{\min(s,t)} \frac{(s+t-j-1)!}{(s-j)!(t-j)!} \cdot \frac{(1-q)^j}{(j-1)!} T[s, t, j], \end{aligned}$$

where

$$\begin{aligned}
S[s, t, a, b] &= \sum_{n=1}^{\infty} \sum_{u+v=n} \frac{q^{(s-1)u} q^{(t-1)v} q^{(t-1-b)u} q^{av}}{[u]_q^{s-a-b} [u+v]_q^{t+a}} = \sum_{n=1}^{\infty} \sum_{u+v=n} \frac{q^{(t+a-1)(u+v)} q^{(s-a-b-1)u}}{[u+v]_q^{t+a} [u]_q^{s-a-b}} \\
&= \sum_{n=1}^{\infty} \frac{q^{(t+a-1)n}}{[n]_q^{t+a}} \sum_{u=1}^{n-1} \frac{q^{(s-a-b-1)u}}{[u]_q^{s-a-b}} \\
&= \zeta[t+a, s-a-b]
\end{aligned}$$

and

$$\begin{aligned}
T[s, t, j] &= \sum_{n=1}^{\infty} \sum_{u+v=n} \frac{q^{(s-1)u} q^{(t-1)v} q^{(t-j)u} q^{(s-j)v}}{[u+v]_q^{s+t-j}} = \sum_{n=1}^{\infty} \sum_{u+v=n} \frac{q^{(s+t-j-1)(u+v)}}{[u+v]_q^{s+t-j}} \\
&= \varphi[s+t-j].
\end{aligned}$$

□

5. FINAL REMARKS

In [24], Zhao also gives a formula for the product $\zeta[s]\zeta[t]$. However, Zhao's formula is considerably more complicated than ours, as it is derived based on the q -shuffle rule [7, 11] satisfied by the Jackson q -integral analogs of the representations (5). Of course, we also have the very simple q -stuffle [11] formula $\zeta[s]\zeta[t] = \zeta[s, t] + \zeta[t, s] + \zeta[s+t] + (1-q)\zeta[s+t-1]$.

REFERENCES

- [1] B. Berndt, *Ramanujan's Notebooks Part I*, Springer, New York, 1985.
- [2] J. M. Borwein, D. J. Broadhurst, and D. M. Bradley, Evaluations of k -fold Euler/Zagier sums: a compendium of results for arbitrary k , *Electronic J. Combinatorics*, **4** (1997), no. 2, #R5. Wilf Festschrift.
- [3] D. Borwein, J. M. Borwein, and R. Girgensohn, Explicit evaluation of Euler sums, *Proc. Edinburgh Math. Soc.*, **38** (1995), 277–294.
- [4] J. M. Borwein, D. J. Broadhurst, D. M. Bradley, and P. Lisoněk, Special values of multiple polylogarithms, *Trans. Amer. Math. Soc.*, **353** (2001), no. 3, 907–941. <http://arXiv.org/abs/math.CA/9910045>
- [5] ———, J. M. Borwein, D. J. Broadhurst, D. M. Bradley, and P. Lisoněk, Combinatorial aspects of multiple zeta values, *Electronic J. Combinatorics*, **5** (1998), no. 1, #R38. <http://arXiv.org/abs/math.NT/9812020>
- [6] D. Bowman and D. M. Bradley, Resolution of some open problems concerning multiple zeta evaluations of arbitrary depth, *Compositio Mathematica*, **139** (2003), no. 1, 85–100. <http://arXiv.org/abs/math.CA/0310061>
- [7] ———, Multiple polylogarithms: A brief survey, *Proceedings of a Conference on q -Series with Applications to Combinatorics, Number Theory and Physics*, (B. C. Berndt and K. Ono eds.) Amer. Math. Soc., Contemporary Math., **291** (2001), 71–92. <http://arXiv.org/abs/math.CA/0310062>
- [8] ———, The algebra and combinatorics of shuffles and multiple zeta values, *J. Combinatorial Theory, Ser. A*, **97** (2002), no. 1, 43–61. <http://arXiv.org/abs/math.CO/0310082>

- [9] D. Bowman, D. M. Bradley, and J. Ryoo, Some multi-set inclusions associated with shuffle convolutions and multiple zeta values, *European J. Combinatorics*, **24** (2003), 121–127.
- [10] D. M. Bradley, Partition identities for the multiple zeta function, to appear in *Zeta Functions, Topology, and Physics*, Kinki University Mathematics Seminar Series, Developments in Mathematics. <http://arXiv.org/abs/math.CO/0402091>
- [11] ———, Multiple q -zeta values, *J. Algebra*, **283** (2005), no. 2, 752–798. doi: 10.1016/j.jalgebra.2004.09.017 <http://arXiv.org/abs/math.QA/0402093>
- [12] ———, Duality for finite multiple harmonic q -series, submitted for publication October 26, 2003. <http://arXiv.org/abs/math.CO/0402092>
- [13] D. J. Broadhurst and D. Kreimer, Association of multiple zeta values with positive knots via Feynman diagrams up to 9 loops, *Phys. Lett. B*, **393** (1997) no. 3-4, 403–412.
- [14] P. Cartier, Fonctions polylogarithmes, nombres polyzêtas et groupes pro-unipotents, *Séminaire Bourbaki*, 53^eème année, 2000–2001, N^o 885, 1–35.
- [15] L. Euler, *Meditationes Circa Singulare Serierum Genus*, Novi Comm. Acad. Sci. Petropol., **20** (1775), 140–186. Reprinted in “Opera Omnia,” ser. I, **15**, B. G. Teubner, Berlin (1927), 217–267.
- [16] ———, *Briefwechsel*, vol. 1, Birkhäuser, Basel, 1975.
- [17] L. Euler and C. Goldbach, *Briefwechsel 1729–1764*, Akademie-Verlag, Berlin, 1965.
- [18] Tu Quoc Thang Le and Jun Murakami, *Kontsevich’s integral for the Homfly polynomial and relations between values of multiple zeta functions*, *Topology and its Applications*, **62** (1995), no. 2, 193–206.
- [19] C. Markett, Triple sums and the Riemann zeta function, *J. Number Theory*, **48** (1994), 113–132.
- [20] N. Nielsen, *Die Gammafunktion*, Chelsea, New York, 1965, 47–59.
- [21] J. Okuda and Y. Takeyama, On relations for the q -multiple zeta values, preprint. <http://arXiv.org/abs/math.QA/0402152> v1, Feb. 10, 2004.
- [22] M. Waldschmidt, Multiple polylogarithms: an introduction, in *Number Theory and Discrete Mathematics*, Hindustan Book Agency and Birkhäuser Verlag, 2002, 1–12.
- [23] ———, Valeurs zêta multiples: une introduction, *Journal de Theorie des Nombres de Bordeaux*, **12** (2000), no. 2, 581–595.
- [24] J. Zhao, q -multiple zeta functions and q -multiple polylogarithms, <http://arXiv.org/abs/math.QA/0304448> v2, 23 May 2003.
- [25] W. Zudilin [V. V. Zudilin], Algebraic relations for multiple zeta values (Russian), *Uspekhi Mat. Nauk*, **58** (2003), no. 1, 3–32; translation in *Russian Math. Surveys*, **58** (2003), vol. 1, 1–29.

DEPARTMENT OF MATHEMATICS & STATISTICS, UNIVERSITY OF MAINE, 5752 NEVILLE HALL
ORONO, MAINE 04469-5752, U.S.A.

E-mail address: bradley@math.umaine.edu, dbradley@member.ams.org