

Refinements of Reverse Triangle Inequalities in Inner Product Spaces*

Arsalan Hojjat Ansari and Mohammad Sal Moslehian

Dept. of Math., Ferdowsi Univ.

P. O. Box 1159, Mashhad 91775

Iran

E-mail: msalm@math.um.ac.ir

URL: <http://www.um.ac.ir/~moslehian/>

Abstract

Refining some results of S. S. Dragomir, several new reverses of the triangle inequality in inner product spaces are obtained.

*2000 *Mathematics Subject Classification.* Primary 46C05; Secondary 26D15.

Key words and phrases. Triangle inequality, reverse inequality, Diaz-Metkalf inequality, inner product space.

1 Introduction.

It is interesting to know under which conditions the triangle inequality went the other way in a normed space X ; in other words, we would like to know if there is a constant c with the property that $c \sum_{k=1}^n \|x_k\| \leq \|\sum_{k=1}^n x_k\|$ for any finite set $x_1, \dots, x_n \in X$. M. Nakai and T. Tada [6] proved that the normed spaces with this property are precisely those of finite dimensional.

The first authors investigating reverse of the triangle inequality in inner product spaces were J. B. Diaz and F. T. Metcalf [1] by establishing the following result as an extension of an inequality given by M. Petrovich [7] for complex numbers:

Diaz-Metcalf Theorem. Let a be a unit vector in the inner product space $(H; \langle \cdot, \cdot \rangle)$. Suppose the vectors $x_k \in H, k \in \{1, \dots, n\}$ satisfy

$$0 \leq r \leq \frac{\operatorname{Re} \langle x_k, a \rangle}{\|x_k\|}, k \in \{1, \dots, n\}$$

Then

$$r \sum_{k=1}^n \|x_k\| \leq \left\| \sum_{k=1}^n x_k \right\|.$$

where equality holds if and only if

$$\sum_{k=1}^n x_k = r \sum_{k=1}^n \|x_k\| a.$$

Inequalities related to the triangle inequality are of special interest; cf. Chapter XVII of [5] and may be applied to get nice inequalities in complex numbers or to study vector-valued integral inequalities [2], [3].

Using several ideas and notation of [2], [3] we modify or refine results of S. Dragomir and get some new reverses of triangle inequality.

We use repeatedly the Cauchy-Schwarz inequality without mentioning it. The reader is referred to [8], [4] for the terminology on inner product spaces.

2 Main Results.

The following theorem is an strengthen of theorem 1 of [3] in which the real numbers r_1, r_2 are not neccesarily nonnegative. The proof seems to be different as well.

Theorem 1. Let a be a unit vector in the complex inner product space $(H; \langle \cdot, \cdot \rangle)$. Suppose the vectors $x_k \in H, k \in \{1, \dots, n\}$ satisfy

$$0 \leq r_1^2 \|x_k\| \leq \operatorname{Re} \langle x_k, r_1 a \rangle, 0 \leq r_2^2 \|x_k\| \leq \operatorname{Im} \langle x_k, r_2 a \rangle \quad (1)$$

for some $r_1, r_2 \in [-1, 1]$. Then we have the inequality

$$(r_1^2 + r_2^2)^{\frac{1}{2}} \sum_{k=1}^n \|x_k\| \leq \left\| \sum_{k=1}^n x_k \right\|. \quad (2)$$

The equality holds in (2) if and only if

$$\sum_{k=1}^n x_k = (r_1 + ir_2) \sum_{k=1}^n \|x_k\| a. \quad (3)$$

Proof. If $r_1^2 + r_2^2 = 0$, theorem is trivial. Assume that $r_1^2 + r_2^2 \neq 0$. Summing inequalities (1) over k from 1 to m , we have

$$\begin{aligned} (r_1^2 + r_2^2) \sum_{k=1}^n \|x_k\| &\leq \operatorname{Re} \left\langle \sum_{k=1}^n x_k, r_1 a \right\rangle + \operatorname{Im} \left\langle \sum_{k=1}^n x_k, r_2 a \right\rangle \\ &= \operatorname{Re} \left\langle \sum_{k=1}^n x_k, (r_1 + ir_2) a \right\rangle \\ &\leq \left| \left\langle \sum_{k=1}^n x_k, (r_1 + ir_2) a \right\rangle \right| \\ &\leq \left\| \sum_{k=1}^n x_k \right\| \|(r_1 + ir_2) a\| \\ &= (r_1^2 + r_2^2)^{\frac{1}{2}} \sum_{k=1}^n \|x_k\|. \end{aligned}$$

Hence 2 holds.

If (3) holds, then

$$\left\| \sum_{k=1}^n x_k \right\| = \|(r_1 + ir_2) \sum_{k=1}^n \|x_k\| a\| = (r_1^2 + r_2^2)^{\frac{1}{2}} \sum_{k=1}^n \|x_k\|.$$

Conversely, if the equality holds in (2), we have

$$\begin{aligned} (r_1^2 + r_2^2)^{\frac{1}{2}} \left\| \sum_{k=1}^n x_k \right\| &= (r_1^2 + r_2^2) \sum_{k=1}^n \|x_k\| \leq \operatorname{Re} \left\langle \sum_{k=1}^n x_k, (r_1 + ir_2)a \right\rangle \\ &\leq \left| \left\langle \sum_{k=1}^n x_k, (r_1 + ir_2)a \right\rangle \right| \leq (r_1^2 + r_2^2)^{\frac{1}{2}} \left\| \sum_{k=1}^n x_k \right\|. \end{aligned}$$

From this we deduce

$$\left| \left\langle \sum_{k=1}^n x_k, (r_1 + ir_2)a \right\rangle \right| = \left\| \sum_{k=1}^n x_k \right\| \|(r_1 + ir_2)a\|.$$

Consequently there exists $\eta \geq 0$ such that $\sum_{k=1}^n x_k = \eta(r_1 + ir_2)a$

From this we have

$$(r_1^2 + r_2^2)^{\frac{1}{2}} \eta = \|\eta(r_1 + ir_2)a\| = \left\| \sum_{k=1}^n x_k \right\| = (r_1^2 + r_2^2)^{\frac{1}{2}} \sum_{k=1}^n \|x_k\|.$$

Hence $\eta = \sum_{k=1}^n \|x_k\|$. \square

The next theorem is a refinement of Corollary 1 of [3] since, in the notation of the theorem, $\sqrt{2 - p_1^2 - p_2^2} \leq \sqrt{\alpha_1^2 + \alpha_2^2}$.

Theorem 2. Let a be a unit vector in the complex inner product space $(H; \langle \cdot, \cdot \rangle)$. Suppose the vectors $x_k \in H - \{0\}, k \in \{1, \dots, n\}$, such that

$$\|x_k - a\| \leq p_1, \|x_k - ia\| \leq p_2, p_1, p_2 \in (0, \sqrt{\alpha^2 + 1}) \quad (4)$$

where $\alpha = \min_{1 \leq k \leq n} \|x_k\|$. Let

$$\alpha_1 = \min \left\{ \frac{\|x_k\|^2 - p_1^2 + 1}{2\|x_k\|} : 1 \leq k \leq n \right\}, \alpha_2 = \min \left\{ \frac{\|x_k\|^2 - p_2^2 + 1}{2\|x_k\|} : 1 \leq k \leq n \right\},$$

Then we have the inequality

$$(\alpha_1^2 + \alpha_2^2)^{\frac{1}{2}} \sum_{k=1}^n \|x_k\| \leq \left\| \sum_{k=1}^n x_k \right\|$$

where the equality holds if and only if

$$\sum_{k=1}^n x_k = (\alpha_1 + i\alpha_2) \sum_{k=1}^n \|x_k\| a$$

Proof. From the first inequality in (4) we have

$$\langle x_k - a, x_k - a \rangle \leq p_1^2$$

$$\begin{aligned} \|x_k\|^2 + 1 - p_1^2 &\leq 2\operatorname{Re}\langle x_k, a \rangle, k = 1, \dots, n \\ \frac{\|x_k\|^2 - p_1^2 + 1}{2\|x_k\|} \|x_k\| &\leq \operatorname{Re}\langle x_k, a \rangle \end{aligned}$$

consequently

$$\alpha_1 \|x_k\| \leq \operatorname{Re}\langle x_k, a \rangle.$$

Similarly from the second inequality we obtain

$$\alpha_2 \|x_k\| \leq \operatorname{Re}\langle x_k, ia \rangle = \operatorname{Im}\langle x_k, a \rangle.$$

Now apply Theorem 1 for $r_1 = \alpha_1, r_2 = \alpha_2$. \square

Corollary 3. Let a be a unit vector in the complex inner product space $(H; \langle \cdot, \cdot \rangle)$. Suppose the vectors $x_k \in H - \{0\}, k \in \{1, \dots, n\}$ such that

$$\|x_k - a\| \leq 1, \|x_k - ia\| \leq 1.$$

Then

$$\frac{\alpha}{\sqrt{2}} \sum_{k=1}^n \|x_k\| \leq \left\| \sum_{k=1}^n x_k \right\|$$

in which $\alpha = \min_{1 \leq k \leq n} \|x_k\|$. The equality holds if and only if

$$\sum_{k=1}^n x_k = \alpha \frac{(1+i)}{2} \sum_{k=1}^n \|x_k\| a$$

Proof. Apply Theorem 2 for $\alpha_1 = \frac{\alpha}{2} = \alpha_2$. \square

Theorem 4. Let a be a unit vector in the inner product space $(H; \langle \cdot, \cdot \rangle)$ over the real or complex number field. Suppose that the vectors $x_k \in H - \{0\}, k \in \{1, \dots, n\}$ satisfy

$$\|x_k - a\| \leq p, p \in (0, \sqrt{\alpha^2 + 1}), \alpha = \min_{1 \leq k \leq n} \|x_k\|.$$

Then we have the inequality

$$\alpha_1 \sum_{k=1}^n \|x_k\| \leq \left\| \sum_{k=1}^n x_k \right\|$$

where

$$\alpha_1 = \min \left\{ \frac{\|x_k\|^2 - p^2 + 1}{2\|x_k\|} : 1 \leq k \leq n \right\}$$

The equality holds if and only if

$$\sum_{k=1}^n x_k = \alpha_1 \sum_{k=1}^n \|x_k\| a$$

Proof. The proof is similar to Theorem 2 in which we use Theorem 1 with $r_2 = 0$. \square

The next Theorem is a generalization of Theorem 1. It is a modification of Theorem 3 of [3], however our proof is apparently different.

Theorem 5. Let a_1, \dots, a_m be orthonormal vectors in the complex inner product space $(H; \langle \cdot, \cdot \rangle)$. Suppose that for $1 \leq t \leq m, r_t, \rho_t \in \mathbb{R}$ and that the vectors $x_k \in H, k \in \{1, \dots, n\}$ satisfy

$$0 \leq r_t^2 \|x_k\| \leq \operatorname{Re} \langle x_k, r_t a_t \rangle, 0 \leq \rho_t^2 \|x_k\| \leq \operatorname{Im} \langle x_k, \rho_t a_t \rangle, t \in \{1, \dots, m\} \quad (5)$$

Then we have the inequality

$$\left(\sum_{t=1}^m r_t^2 + \rho_t^2 \right)^{\frac{1}{2}} \sum_{k=1}^n \|x_k\| \leq \left\| \sum_{k=1}^n x_k \right\| \quad (6)$$

The equality holds in (7) if and only if

$$\sum_{k=1}^n x_k = \sum_{k=1}^n \|x_k\| \sum_{t=1}^m (r_t + i\rho_t) a_t. \quad (7)$$

Proof. If $\sum_{t=1}^m (r_t^2 + \rho_t^2) = 0$, theorem is trivial. Assume that $\sum_{t=1}^m (r_t^2 + \rho_t^2) \neq 0$. Summing inequalities (6) over k from 1 to n and again over t from 1 to m we get

$$\sum_{t=1}^m (r_t^2 + \rho_t^2) \sum_{k=1}^n \|x_k\| \leq \operatorname{Re} \left\langle \sum_{k=1}^n x_k, \sum_{t=1}^m r_t a_t \right\rangle + \operatorname{Im} \left\langle \sum_{k=1}^n x_k, \sum_{t=1}^m \rho_t a_t \right\rangle$$

$$\begin{aligned}
&= \operatorname{Re} \langle \sum_{k=1}^n x_k, \sum_{t=1}^m r_t a_t \rangle + \operatorname{Re} \langle \sum_{k=1}^n x_k, i \sum_{t=1}^m \rho_t a_t \rangle \\
&= \operatorname{Re} \langle \sum_{k=1}^n x_k, \sum_{t=1}^m (r_t + i \rho_t) a_t \rangle \\
&\leq |\langle \sum_{k=1}^n x_k, \sum_{t=1}^m (r_t + i \rho_t) a_t \rangle| \\
&\leq \left\| \sum_{k=1}^n x_k \right\| \left\| \sum_{t=1}^m (r_t + i \rho_t) a_t \right\| \\
&= \left\| \sum_{k=1}^n x_k \right\| \left(\sum_{t=1}^m (r_t^2 + \rho_t^2) \right)^{\frac{1}{2}}.
\end{aligned}$$

Then

$$\left(\sum_{t=1}^m (r_t^2 + \rho_t^2) \right)^{\frac{1}{2}} \sum_{k=1}^n \|x_k\| \leq \left\| \sum_{k=1}^n x_k \right\|.$$

If (8) holds, then

$$\left\| \sum_{k=1}^n x_k \right\| = \left\| \sum_{k=1}^n \|x_k\| \left\| \sum_{t=1}^m (r_t + i \rho_t) a_t \right\| \right\| = \sum_{k=1}^n \|x_k\| \left(\sum_{t=1}^m (r_t^2 + \rho_t^2) \right)^{1/2}.$$

Conversely, if the equality holds in (7) we obtain from (6) that

$$\begin{aligned}
&\left(\sum_{t=1}^m (r_t^2 + \rho_t^2) \right)^{1/2} \left\| \sum_{k=1}^n x_k \right\| = \\
&= \sum_{t=1}^m (r_t^2 + \rho_t^2) \sum_{k=1}^n \|x_k\| \leq \operatorname{Re} \langle \sum_{k=1}^n x_k, \sum_{t=1}^m (r_t + i \rho_t) a_t \rangle \leq \\
&\leq |\langle \sum_{k=1}^n x_k, \sum_{t=1}^m (r_t + i \rho_t) a_t \rangle| \leq \\
&\leq \left\| \sum_{k=1}^n x_k \right\| \left\| \sum_{t=1}^m (r_t + i \rho_t) a_t \right\| = \left\| \sum_{k=1}^n x_k \right\| \left(\sum_{t=1}^m (r_t^2 + \rho_t^2) \right)^{1/2}
\end{aligned}$$

Thus we have

$$|\langle \sum_{k=1}^n x_k, \sum_{t=1}^m (r_t + i \rho_t) a_t \rangle| = \left\| \sum_{k=1}^n x_k \right\| \left\| \sum_{t=1}^m (r_t + i \rho_t) a_t \right\|.$$

Consequently there exists $\eta \geq 0$ such that

$$\sum_{k=1}^n x_k = \eta \sum_{t=1}^m (r_t + i \rho_t) a_t$$

from which we have

$$\eta(\sum_{t=1}^m(r_t^2 + \rho_t^2))^{\frac{1}{2}} = \|\eta \sum_{t=1}^m(r_t + i\rho_t)a_t\| = \|\sum_{k=1}^n x_k\| = \sum_{k=1}^n \|x_k\|(\sum_{t=1}^m(r_t^2 + \rho_t^2))^{\frac{1}{2}}$$

Hence

$$\eta = \sum_{k=1}^n \|x_k\|. \square$$

Corollary 6. Let a_1, \dots, a_m be orthornormal vectors in the inner product space $(H; \langle \cdot, \cdot \rangle)$ over the real or complex number field. Suppose for $1 \leq t \leq m$ that the vectors $x_k \in H, k \in \{1, \dots, n\}$ satisfy

$$0 \leq r_t^2 \|x_k\| \leq Re \langle x_k, r_t a_t \rangle.$$

Then we have the inequality

$$(\sum_{t=1}^m r_t^2)^{\frac{1}{2}} \sum_{k=1}^n \|x_k\| \leq \|\sum_{k=1}^n x_k\|.$$

The equality holds if and only if

$$\sum_{k=1}^n x_k = \sum_{k=1}^n \|x_k\| \sum_{t=1}^m r_t a_t.$$

Proof. Apply Theorem 5 for $\rho_t = 0$. \square

Theorem 7. Let a_1, \dots, a_m be orthornormal vectors in the complex inner product space $(H; \langle \cdot, \cdot \rangle)$. Suppose that the vectors $x_k \in H - \{0\}, k \in \{1, \dots, n\}$ satisfy

$$\|x_k - a_t\| \leq p_t, \|x_k - ia_t\| \leq q_t, p_t, q_t \in (0, \sqrt{\alpha^2 + 1}), 1 \leq t \leq m$$

where $\alpha = \min_{1 \leq k \leq n} \|x_k\|$. Let

$$\alpha_t = \min\left\{\frac{\|x_k\|^2 - p_t^2 + 1}{2\|x_k\|} : 1 \leq k \leq n\right\}, \beta_t = \min\left\{\frac{\|x_k\|^2 - q_t^2 + 1}{2\|x_k\|} : 1 \leq k \leq n\right\}.$$

Then we have the inequality

$$(\sum_{t=1}^m \alpha_t^2 + \beta_t^2)^{\frac{1}{2}} \sum_{k=1}^n \|x_k\| \leq \|\sum_{k=1}^n x_k\|$$

where equality holds if and only if

$$\sum_{k=1}^n x_k = \sum_{k=1}^n \|x_k\| \sum_{t=1}^m (\alpha_t + i\beta_t) a_t.$$

Proof. For $1 \leq t \leq m, 1 \leq k \leq n$ it follows from $\|x_k - a_t\| \leq p_t$ that

$$\begin{aligned} \langle x_k - a_t, x_k - a_t \rangle &\leq p_t^2 \\ \frac{\|x_k\|^2 - p_t^2 + 1}{2\|x_k\|} \|x_k\| &\leq \operatorname{Re} \langle x_k, a_t \rangle \\ \alpha_t \|x_k\| &\leq \operatorname{Re} \langle x_k, a_t \rangle \end{aligned}$$

and similarly

$$\beta_t \|x_k\| \leq \operatorname{Re} \langle x_k, ia_t \rangle = \operatorname{Im} \langle x_k, a_t \rangle,$$

Now applying Theorem 4 with $r_t = \alpha_t, \rho_t = \beta_t$ we deduce the desired inequality. \square

Corollary 8. Let a_1, \dots, a_m be orthonormal vectors in the complex inner product space $(H; \langle \cdot, \cdot \rangle)$. Suppose that the vectors $x_k \in H, k \in \{1, \dots, n\}$ satisfy

$$\|x_k - a_t\| \leq 1, \|x_k - ia_t\| \leq 1, 1 \leq t \leq m$$

Then

$$\frac{\alpha}{\sqrt{2}} \sqrt{m} \sum_{k=1}^n \|x_k\| \leq \left\| \sum_{k=1}^n x_k \right\|.$$

The equality holds if and only if

$$\sum_{k=1}^n x_k = \alpha \frac{(1+i)}{2} \sum_{k=1}^n \|x_k\| \sum_{t=1}^m a_t$$

Proof. Applying Theorem 7 for $\alpha_t = \frac{\alpha}{2} = \beta_t$. \square

Remark. It is interesting to note that

$$\begin{aligned} \frac{\alpha}{\sqrt{2}} \sqrt{m} &\leq \frac{\left\| \sum_{k=1}^n x_k \right\|}{\sum_{k=1}^n \|x_k\|} \leq 1. \\ \alpha &\leq \sqrt{\frac{2}{m}} \end{aligned}$$

Corollary 9. Let a be a unit vector in the complex inner product space $(H; \langle \cdot, \cdot \rangle)$. Suppose that the vectors $x_k \in H - \{0\}, k \in \{1, \dots, n\}$ satisfy

$$\|x_k - a\| \leq p_1, \|x_k - ia\| \leq p_2, p_1, p_2 \in (0, 1].$$

Let

$$\alpha_1 = \min\left\{\frac{\|x_k\|^2 - p_1^2 + 1}{2\|x_k\|} : 1 \leq k \leq n\right\}, \alpha_2 = \min\left\{\frac{\|x_k\|^2 - p_2^2 + 1}{2\|x_k\|} : 1 \leq k \leq n\right\}.$$

If $\alpha_1 \neq (1 - p_1^2)^{\frac{1}{2}}$, or $\alpha_2 \neq (1 - p_2^2)^{\frac{1}{2}}$, then we have the following strictly inequality

$$(2 - p_1^2 - p_2^2)^{\frac{1}{2}} \sum_{k=1}^n \|x_k\| < \left\| \sum_{k=1}^n x_k \right\|$$

Proof. If equality holds, then by Theorem 2 we have

$$(\alpha_1^2 + \alpha_2^2)^{\frac{1}{2}} \sum_{k=1}^n \|x_k\| \leq \left\| \sum_{k=1}^n x_k \right\| = (2 - p_1^2 - p_2^2)^{\frac{1}{2}} \sum_{k=1}^n \|x_k\|$$

and so

$$(\alpha_1^2 + \alpha_2^2)^{\frac{1}{2}} \leq (2 - p_1^2 - p_2^2)^{\frac{1}{2}}.$$

On the other hand for $1 \leq k \leq n$,

$$\frac{\|x_k\|^2 - p_1^2 + 1}{2\|x_k\|} \geq (1 - p_1^2)^{\frac{1}{2}}$$

and so

$$\alpha_1 \geq (1 - p_1^2)^{\frac{1}{2}}.$$

Similarly

$$\alpha_2 \geq (1 - p_2^2)^{\frac{1}{2}}.$$

Hence

$$(2 - p_1^2 - p_2^2)^{\frac{1}{2}} \leq (\alpha_1^2 + \alpha_2^2)^{\frac{1}{2}}$$

Thus

$$\sqrt{\alpha_1^2 + \alpha_2^2} = (2 - p_1^2 - p_2^2)^{\frac{1}{2}}.$$

Therefore

$$\alpha_1 = (1 - p_1^2)^{\frac{1}{2}} \text{ and } \alpha_2 = (1 - p_2^2)^{\frac{1}{2}}$$

a contradiction. \square

The following result looks like Corollary 2 of [3].

Theorem 10. Let a be a unit vector in the complex inner product space $(H; \langle \cdot, \cdot \rangle)$, $M \geq m > 0$, $L \geq \ell > 0$ and $x_k \in H - \{0\}$, $k \in \{1, \dots, n\}$ such that

$$Re\langle Ma - x_k, x_k - ma \rangle \geq 0, Re\langle Lia - x_k, x_k - lia \rangle \geq 0$$

or equivalently,

$$\|x_k - \frac{m+M}{2}a\| \leq \frac{M-m}{2}, \|x_k - \frac{L+\ell}{2}ia\| \leq \frac{L-\ell}{2}.$$

Let

$$\alpha_{m,M} = \min\left\{\frac{\|x_k\|^2 + mM}{(m+M)\|x_k\|} : 1 \leq k \leq n\right\}$$

and

$$\alpha_{\ell,L} = \min\left\{\frac{\|x_k\|^2 + \ell L}{(\ell+L)\|x_k\|} : 1 \leq k \leq n\right\}$$

Then we have the inequality

$$(\alpha_{m,M}^2 + \alpha_{\ell,L}^2)^{\frac{1}{2}} \sum_{k=1}^n \|x_k\| \leq \left\| \sum_{k=1}^n x_k \right\|.$$

The equality holds if and only if

$$\sum_{k=1}^n x_k = (\alpha_{m,M} + i\alpha_{\ell,L}) \sum_{k=1}^n \|x_k\| a.$$

Proof. For each $1 \leq k \leq n$, it follows from

$$\|x_k - \frac{m+M}{2}a\| \leq \frac{M-m}{2}$$

that

$$\left\langle x_k - \frac{m+M}{2}a, x_k - \frac{m+M}{2}a \right\rangle \leq \left(\frac{M-m}{2}\right)^2.$$

Hence

$$\|x_k\|^2 + mM \leq (m+M)Re\langle x_k, a \rangle.$$

Then

$$\frac{\|x_k\|^2 + mM}{(m+M)\|x_k\|} \|x_k\| \leq \operatorname{Re}\langle x_k, a \rangle$$

consequently

$$\alpha_{m,M} \|x_k\| \leq \operatorname{Re}\langle x_k, a \rangle.$$

Similarly from the second inequality we deduce

$$\alpha_{\ell,L} \|x_k\| \leq \operatorname{Im}\langle x_k, a \rangle.$$

Applying Theorem 1 for $r_1 = \alpha_{m,M}$, $r_2 = \alpha_{\ell,L}$, we infer the desired inequality. \square

Theorem 11. Let a be a unit vector in the complex inner product space $(H; \langle \cdot, \cdot \rangle)$, $M \geq m > 0$, $L \geq \ell > 0$ and $x_k \in H - \{0\}$, $k \in \{1, \dots, n\}$ such that

$$\operatorname{Re}\langle Ma - x_k, x_k - ma \rangle \geq 0, \operatorname{Re}\langle Lia - x_k, x_k - \ell ia \rangle \geq 0$$

or equivalently

$$\|x_k - \frac{m+M}{2}a\| \leq \frac{M-m}{2}, \|x_k - \frac{L+\ell}{2}ia\| \leq \frac{L-\ell}{2}.$$

Let

$$\alpha_{m,M} = \min\left\{\frac{\|x_k\|^2 + mM}{(m+M)\|x_k\|} : 1 \leq k \leq n\right\}$$

and

$$\alpha_{\ell,L} = \min\left\{\frac{\|x_k\|^2 + \ell L}{(\ell+L)\|x_k\|} : 1 \leq k \leq n\right\}.$$

If $\alpha_{m,M} \neq 2\frac{\sqrt{mM}}{m+M}$, or $\alpha_{\ell,L} \neq 2\frac{\sqrt{\ell L}}{\ell+L}$, then we have

$$2\left(\frac{mM}{(m+M)^2} + \frac{\ell L}{(\ell+L)^2}\right)^{\frac{1}{2}} \sum_{k=1}^n \|x_k\| < \left\|\sum_{k=1}^n x_k\right\|.$$

Proof. If $2\left(\frac{mM}{(m+M)^2} + \frac{\ell L}{(\ell+L)^2}\right)^{\frac{1}{2}} \sum_{k=1}^n \|x_k\| = \left\|\sum_{k=1}^n x_k\right\|$ then by theorem 10 we have

$$(\alpha_{m,M}^2 + \alpha_{\ell,L}^2)^{\frac{1}{2}} \sum_{k=1}^n \|x_k\| \leq \left\|\sum_{k=1}^n x_k\right\| = 2\left(\frac{mM}{(m+M)^2} + \frac{\ell L}{(\ell+L)^2}\right)^{\frac{1}{2}} \sum_{k=1}^n \|x_k\|.$$

Consequently

$$(\alpha_{m,M}^2 + \alpha_{\ell,L}^2)^{\frac{1}{2}} \leq 2\left(\frac{mM}{(m+M)^2} + \frac{\ell L}{(\ell+L)^2}\right)^{\frac{1}{2}}.$$

On the other hand for $1 \leq k \leq n$, $\frac{\|x_k\|^2 + mM}{(m+M)\|x_k\|} \geq 2\frac{\sqrt{mM}}{m+M}$, and $\frac{\|x_k\|^2 + \ell L}{(\ell+L)\|x_k\|} \geq 2\frac{\sqrt{\ell L}}{\ell+L}$,
so

$$(\alpha_{m,M}^2 + \alpha_{\ell,L}^2)^{\frac{1}{2}} \geq 2\left(\frac{mM}{(m+M)^2} + \frac{\ell L}{(\ell+L)^2}\right)^{\frac{1}{2}}.$$

Then

$$(\alpha_{m,M}^2 + \alpha_{\ell,L}^2)^{\frac{1}{2}} = 2\left(\frac{mM}{(m+M)^2} + \frac{\ell L}{(\ell+L)^2}\right)^{\frac{1}{2}}.$$

Hence

$$\alpha_{m,M} = 2\frac{\sqrt{mM}}{m+M}$$

and

$$\alpha_{\ell,L} = 2\frac{\sqrt{\ell L}}{\ell+L}$$

a contradiction. \square

Finally we mention two applications of our results to the complex numbers.

Corollary 12. Let $a \in C$ with $|a| = 1$. Suppose that $z_k \in C, k \in \{1, \dots, n\}$ such that

$$|z_k - a| \leq p_1, |z_k - ia| \leq p_2, p_1, p_2 \in (0, \sqrt{\alpha^2 + 1})$$

where

$$\alpha = \min\{|z_k| : 1 \leq k \leq n\}.$$

Let

$$\alpha_1 = \min\left\{\frac{|z_k|^2 - p_1^2 + 1}{2|z_k|} : 1 \leq k \leq n\right\}, \alpha_2 = \min\left\{\frac{|z_k|^2 - p_2^2 + 1}{2|z_k|} : 1 \leq k \leq n\right\}.$$

Then we have the inequality

$$\sqrt{\alpha_1^2 + \alpha_2^2} \sum_{k=1}^n |z_k| \leq \left| \sum_{k=1}^n z_k \right|.$$

The equality holds if and only if

$$\sum_{k=1}^n z_k = (\alpha_1 + i\alpha_2) \left(\sum_{k=1}^n |z_k| \right) a.$$

Proof. Apply Theorem 2 for $H = C$. \square

Corollary 13. Let $a \in C$ with $|a| = 1$. Suppose that $z_k \in C, k \in \{1, \dots, n\}$ such that

$$|z_k - a| \leq 1, |z_k - ia| \leq 1.$$

If $\alpha = \min\{|z_k| : 1 \leq k \leq n\}$. Then we have the inequality

$$\frac{\alpha}{\sqrt{2}} \sum_{k=1}^n |z_k| \leq \left| \sum_{k=1}^n z_k \right|$$

the equality holds if and only if

$$\sum_{k=1}^n z_k = \alpha \frac{(1+i)}{2} \left(\sum_{k=1}^n |z_k| \right) a.$$

Proof. Apply Corollary 3 for $H = C$. \square

References

- [1] J.B. Diaz and F.T. Metcalf, A complementary triangle inequality in Hilbert and Banach spaces, Proc. Amer. Math. Soc. 17(1) (1966), 88-97.
- [2] S.S. Dragomir, Reverses of the triangle inequality in inner product spaces, arXiv:math.FA/0405495.
- [3] S.S. Dragomir, Some reverses of the generalized triangle inequality in complex inner product spaces, arXiv:math.FA/0405497.
- [4] S.S. Dragomir, *Discrete inequalities of the Cauchy-Bunyakovsky-Schwarz type*. Nova Science Publishers, Inc., Hauppauge, NY, 2004.
- [5] D.S. Mitrinović, J.E. Pečarić and A.M. Fink, Classical and New Inequalities in Analysis, Kluwer Academic Publishers, Dordrecht/Boston/London, 1993.

- [6] M. Nakai and T. Tada, The reverse triangle inequality in normed spaces, New Zealand J. Math. 25 (1996), no. 2, 181-193.
- [7] M. Petrovich, Module d'une somme, L'Enseignement Mathématique, 19 (1917), 53-56.
- [8] Th.M. Rassias, *Inner Product Spaces and Applications*, Chapman-Hall, 1997.