

A homotopy aspect of representation theory

Haibao Duan

Institute of Mathematics, Chinese Academy of Sciences

Beijing 100080, dhb@math.ac

Abstract

We bring a linkage from representation theory of Lie groups to homotopy theory for maps between flag manifolds. As applications we derive from representation theory abundant families of distinguishable homotopy classes of maps between flag manifolds.

2000 Mathematical Subject Classification: 57R22 (22E46).

1. Introduction

A current topic in algebraic geometry is the relationship between representation theory of Lie groups and Schubert calculus in flag manifolds (cf. [LG, Chapter 4], [K, VIII], [T]). It seems not aware in the existing literature that there is a direct connection between representation theory and homotopy theory for maps between flag manifolds, implicitly contained in the classical works of Borel and Hirzebruch [BH]. This connection may be summarized as “*character factors through homotopy*” and is presented in the Theorem in §5. Applying this idea we obtain abundant families of maps between flag manifolds whose homotopy classes are distinguishable by the characters of representations (Corollary 1–3 in §5).

We begin with a problem concerning homotopy classification for maps between flag manifolds (§2), followed by a brief introduction to what we need from representation theory (§3, §4). The main results are stated and established respectively in §5 and §6. The final section §7 is devoted to further comments.

2. A realization problem in homotopy theory

For a compact connected semi-simple Lie group G with a maximal torus $T \subset G$, write G/T for the flag manifold of G modulo T . If G is the unitary group $U(n)$ of order n and if $T^n \subset U(n)$ is the diagonal subgroup, we write $F(n)$ instead of $U(n)/T^n$.

The primary objective of our study is the set $[G/T, F(n)]$ of all homotopy classes of maps from G/T to $F(n)$ (cf. §7.1 for references on the topic). For this purpose one may examine the correspondence that sends a map f to the induced cohomology homomorphism f^* .

$$(A) \quad r : [G/T, F(n)] \rightarrow \text{Hom}[H^2(F(n)), H^2(G/T)]$$

$$(\text{resp. } r_1 : [G/T, F(n)] \rightarrow \text{Hom}[H^*(F(n)), H^*(G/T)]),$$

where $\text{Hom}[H^*(F(n)), H^*(G/T)]$ is the set of all ring maps from $H^*(F(n))$ to $H^*(G/T)$. It follows from rational homotopy theory [GH] that

the r_1 is finite to one;

We recall also from Bott and Samelson [BS] that

the ring $H^(G/T)$ is torsion free and over rationals generated multiplicatively by elements from $H^2(G/T)$.*

For these reasons a solution to the following problem is an important step towards understanding the set $[G/T, F(n)]$.

Problem. Which homomorphisms $h : H^2(F(n)) \rightarrow H^2(G/T)$ are induced by maps $G/T \rightarrow F(n)$?

3. Notions

For a compact connected Lie group G with a fixed maximal torus T , consider the transgression

$$(B) \quad \tau_G : H^1(T) = \text{Hom}(T, S^1) \rightarrow H^2(G/T)$$

in the fibration $T \hookrightarrow G \rightarrow G/T$ [BH, §10.1], where S^1 is the circle group, and where as indicated the 1-dimensional cohomology $H^1(T)$ is canonically identified with $\text{Hom}(T, S^1)$, the set of 1-dimensional unitary representation of T (cf. Lemma 6 in §6).

If $G = U(n)$ is the unitary group, the diagonal subgroup $T^n \subset U(n)$ has the factorization $T^n = \text{diag}\{e^{i\theta_1}, \dots, e^{i\theta_n}\}$. Let $t'_k \in H^1(T^n) = \text{Hom}(T^n, S^1)$ be the projection onto the k^{th} factor, $1 \leq k \leq n$, and put $t_k = \tau_{U(n)}(t'_k) \in H^2(F(n))$.

Lemma 1. In $H^2(F(n))$ one has $t_1 + \dots + t_n = 0$. Further, the classes t_1, \dots, t_{n-1} form a basis of $H^2(F(n))$.

Proof. This follows from the description of the ring $H^*(F(n))$ due to Borel [B]: $H^*(F(n)) = \mathbb{Z}[t_1, \dots, t_n] / \langle e_i \mid 1 \leq i \leq n \rangle$, where the $e_i \in \mathbb{Z}[t_1, \dots, t_n]$ is the i^{th} elementary symmetric function in the t_1, \dots, t_n . \square

Consider next a compact connected semi-simple G with a fixed maximal torus T . Let $p : \tilde{G} \rightarrow G$ be the universal cover of G and let $\tilde{T} \subset \tilde{G}$ be the maximal torus that corresponds to T under p . Then p induces an isomorphism $\tilde{G}/\tilde{T} \cong G/T$ between flag varieties. It is known that ([BH, §10.1], [DZZ, Theorem 2])

Lemma 2. *A set $\{\omega_i \in \text{Hom}(\tilde{T}, S^1) \mid 1 \leq i \leq m = \dim \tilde{T}\}$ of fundamental dominant weights of \tilde{G} relative to \tilde{T} constitutes a basis of $H^1(\tilde{T}) = \text{Hom}(\tilde{T}, S^1)$ (cf. Lemma 6 in §6), and the transgression $\tau_{\tilde{G}} : H^1(\tilde{T}) \rightarrow H^2(G/T)$ is an isomorphism.*

Consequently, if we make no difference in notation between elements in $H^1(\tilde{T})$ and their images under $\tau_{\tilde{G}}$ in $H^2(G/T)$ (as in [BH]), we have

Lemma 3. *The set $\{\omega_i \in H^2(G/T) \mid 1 \leq i \leq m\}$ of fundamental dominant weights of \tilde{G} is a basis of $H^2(G/T)$.*

Remark 1. The classes $\omega_i \in H^2(G/T)$, $1 \leq i \leq m$, are of particular interests in the algebraic intersection theory on G/T . With respect to the classical Schubert cell decomposition of G/T [BGG], they are precisely the *special Schubert classes* on G/T (cf. [DZZ, §5.2]).

In view of Lemma 1 and 3, every homomorphism $h : H^2(F(n)) \rightarrow H^2(G/T)$ has a canonical numerical characterization as

$$(C) \quad h(t_k) = a_{k,1}\omega_1 + \cdots + a_{k,m}\omega_m, \quad 1 \leq k \leq n-1,$$

where $a_{k,i} \in \mathbb{Z}$. As a result, if we let \mathbb{Z}^+ be the set of non-negative integers, and let $\mathbb{Z}^+[\omega_1, \dots, \omega_m, \rho]$ be the semiring of polynomials in the variables $\omega_1, \dots, \omega_m, \rho$ with coefficients in \mathbb{Z}^+ that is subject to the relation

$$(D) \quad \omega_1 \cdots \omega_m \rho = 1,$$

we may introduce a map

$$s : \text{Hom}(H^2(F(n)), H^2(G/T)) \rightarrow \mathbb{Z}^+[\omega_1, \dots, \omega_m, \rho]$$

by the simple algorithm: if $h : H^2(F(n)) \rightarrow H^2(G/T)$ is given as that in (C), we put

$$s(h) = \sum_{1 \leq k \leq n-1} \omega_1^{a_{k,1}} \cdots \omega_m^{a_{k,m}} + \omega_1^{b_1} \cdots \omega_m^{b_m},$$

where $b_i = -(a_{1,i} + \cdots + a_{n-1,i})$. We note that although the $s(h)$ may appear as an element in the semiring $\mathbb{Z}^+[\omega_1^\pm, \dots, \omega_m^\pm]$ of Laurent polynomials, the obvious relations $\omega_k^{-1} = \rho \prod_{j \neq k} \omega_j$, $1 \leq k \leq m$, from (D) are sufficient to convert it to an element in $\mathbb{Z}^+[\omega_1, \dots, \omega_m, \rho]$.

4. Preliminaries from representation theory

Denote by $R^+(G)$ the semiring of isomorphism classes $\{V\}$ of G -Modules V over \mathbb{C} . As in §3 let \tilde{G} be the universal cover of a semi-simple G with $\tilde{T} \subset \tilde{G}$ the maximal torus that corresponds to T in G . The *character homomorphism* of \tilde{G} is the semiring map

$$\chi : R^+(\tilde{G}) \rightarrow R^+(\tilde{T}) = \mathbb{Z}^+[\omega_1, \dots, \omega_m, \rho],$$

given by restriction $\{V\} \rightarrow \{V \mid \tilde{T}\}$, where the identification

$$R^+(\tilde{T}) = \mathbb{Z}^+[\omega_1, \dots, \omega_m, \rho]$$

follows from Lemma 2.

For a $\{V\} \in R^+(\tilde{G})$, the polynomial $\chi(V) \in \mathbb{Z}^+[\omega_1, \dots, \omega_m, \rho]$ is called *the character of the \tilde{G} -module V* . We will be in particular interested in the subset $\Omega(G) = \text{Im}\chi$ of $R^+(\tilde{T})$. It admits the partition

$$\Omega(G) = \coprod_{1 \leq n} \Omega_n(G)$$

with $\Omega_n(G) = \{g \in \Omega(G) \mid g(1, \dots, 1) = n\}$. Alternatively, the $\Omega_n(G)$ consists of all characters of n -dimensional representations of \tilde{G} .

Indeed, elements in $\Omega(G)$ (resp. in $\Omega_n(G)$) are abundant. To explain this we consider the set $\Lambda(G) = \{a_1\omega_1 + \dots + a_m\omega_m \in H^2(G/T) \mid a_m \in \mathbb{Z}^+\}$ of *dominant weights* of G . For a $\lambda \in \Lambda(G)$ let L_λ be the complex line bundle over G/T with first Chern class $c_1(L_\lambda) = \lambda$, and let V_λ be the \tilde{G} -module of holomorphic sections of L_λ . Since the set $\{V_\lambda \mid \lambda \in \Lambda(G)\}$ presents all finite dimensional complex representations of \tilde{G} (cf. [LG, p. 69]) and since the χ is injective, we have

Lemma 4. *The set of infinitely many polynomials*

$$\{\chi(V_\lambda) \in \mathbb{Z}^+[\omega_1, \dots, \omega_m, \rho] \mid \lambda \in \Lambda(G)\}$$

is a basis for the semiabelian group $\Omega(G)$ (over \mathbb{Z}^+).

We emphasize at this stage that there have been several methods computing the character $\chi(V_\lambda) \in \mathbb{Z}^+[\omega_1, \dots, \omega_m, \rho]$ in terms of the $\lambda \in \Lambda(G)$. These are the Weyl character formula [Hu], Demazure character formula [D] and the Lakshmibai-Seshadri character formula ([LS], [Li]). Consequently, the $\Omega(G)$ admits concrete presentation as a set of polynomials. We present such an example for further use.

If $G = SU(m)$ is the special unitary group of order m , then $(\tilde{G}, \tilde{T}) = (G, T)$, $G/T = F(m)$. Consider the semiring map

$$\alpha : R^+(T) = \mathbb{Z}^+[\omega_1, \dots, \omega_{m-1}, \rho] \rightarrow \mathbb{Z}^+[y_1, \dots, y_m]$$

by $\omega_k \rightarrow y_1 \cdots y_k$, $1 \leq k \leq m-1$; $\rho \rightarrow y_2 y_3^2 \cdots y_m^{m-1}$, where $\mathbb{Z}^+[y_1, \dots, y_m]$ is the semiring of polynomials in the y_1, \dots, y_m with coefficients in \mathbb{Z}^+ that is subject to the relation $y_1 \cdots y_m = 1$.

For a partition $\mu = (\mu_1 \geq \mu_2 \geq \cdots \geq \mu_m \geq 0)$ let $s_\mu(y_1, \dots, y_m) \in \mathbb{Z}^+[y_1, \dots, y_m]$ be *Schur symmetric function* associated to μ [M, p.40], and let $S[y_1, \dots, y_m] \subset \mathbb{Z}^+[y_1, \dots, y_m]$ be the semiabelian group generated over \mathbb{Z}^+ by all Schur functions $s_\mu(y_1, \dots, y_m)$ associated to the μ with last part $\mu_m = 0$. Then (cf. [FH, Chapter 15], [LG; Chapter 5])

Lemma 5. α maps $\Omega(SU(m))$ isomorphically onto $S[y_1, \dots, y_m]$.

For the geometries underlying the transformations α , we refer to [DZZ; Example 3] or [DZ₂; (4.1)].

5. Character factors through homotopy

Let $Hom^0(\tilde{G}, U(n))$ be the set of all homomorphisms $g : \tilde{G} \rightarrow U(n)$ that satisfy $g(\tilde{T}) \subset T^n$. The obvious inclusion $i_n : Hom^0(\tilde{G}, U(n)) \rightarrow R^+(\tilde{G})$ is surjective onto the subset $R_n^+(\tilde{G})$ of isomorphism classes of n -dimensional complex representations of \tilde{G} . On the other hand, since every $g \in Hom^0(\tilde{G}, U(n))$ preserves the maximal torus, there is a ready-made map

$$\rho : Hom^0(\tilde{G}, U(n)) \rightarrow [G/T, F(n)]$$

defined by the commutivity of the diagram

$$\begin{array}{ccc} \tilde{G} & \xrightarrow{g} & U(n) \\ \downarrow & & \downarrow \\ G/T = \tilde{G}/\tilde{T} & \xrightarrow{\rho(g)} & F(n) \end{array},$$

where the vertical maps are the obvious quotients. We may now organize the relevant correspondences in the diagram below.

$$\begin{array}{ccccc} Hom^0(\tilde{G}, U(n)) & & \xrightarrow{i_n} & & R^+(\tilde{G}) \\ \rho \downarrow & & & & \downarrow \chi \\ [G/T, F(n)] & \xrightarrow{r} & Hom[H^2(F(n)), H^2(G/T)] & \xrightarrow{s} & \mathbb{Z}^+[\omega_1, \dots, \omega_m, \rho] \end{array}.$$

Theorem (*Character factors through homotopy*). $\chi \circ i_n = s \circ r \circ \rho$.

The proof of the Theorem will be postponed until the next section and at this moment, we show how it leads to a partial solution to our problem. From the Theorem we obtain

Corollary 1. *For any polynomial $g \in \Omega_n(G)$, there exists a map $f : G/T \rightarrow F(n)$ such that $s(f^*) = g$. Furthermore, one may read the induced map $f^* : H^2(F(n)) \rightarrow H^2(G/T)$ from the polynomial g .*

The first part of Corollary 1 may be rephrased as

Corollary 2. *For a $h \in \text{Hom}(H^2(F(n)), H^2(G/T))$ with $s(h) \in \Omega(G)$, there exists a map $f : G/T \rightarrow F(n)$ such that $f^* = h$.*

In algebraic geometry, Schur functions appear as the polynomial representatives of Schubert classes in complex Grassmanian manifolds (cf. [L], [St]). In representation theory, Schur functions occur as the characters of irreducible representations of unitary groups [T], [FH]. Combining Lemma 5 with Corollary 1 we get the following result illustrating the fashion by which homotopy theory interacts with combinatorics of symmetric functions.

Corollary 3. *For any symmetric function $g \in S[y_1, \dots, y_m]$ with $g(1, \dots, 1) = n$, there exists a map $f : F(m) \rightarrow F(n)$ such that $\alpha \circ s(f^*) = g$.*

Remark 2. According to Corollary 3, every Schur function $s_\mu(y_1, \dots, y_m)$ with $s_\mu(1, \dots, 1) = n$ can arise as the $\alpha \circ s$ -image of a linear map

$$f_\mu^* : H^2(F(n)) \rightarrow H^2(F(m)).$$

We refer to [P] for three definitions of Schur functions in combinatorics. It would be natural to expect that useful properties of Schur functions can be derived from the linear maps f_μ^* .

6. Proof of the Theorem

The binary operation

$$\text{Hom}(T, S^1) \times \text{Hom}(T, S^1) \rightarrow \text{Hom}(T, S^1)$$

by $\alpha\beta = \mu \circ (\alpha \times \beta) \circ \Delta$ furnishes the set $\text{Hom}(T, S^1)$ with the structure of an abelian group, where $\Delta : T \rightarrow T \times T$ is the diagonal embedding and where $\mu : S^1 \times S^1 \rightarrow S^1$ is the product in S^1 . Consider the map

$$\iota : \text{Hom}(T, S^1) \rightarrow H^1(T)$$

by $\iota(\alpha) = \alpha^*[S^1]$, where $\alpha^* : H^1(S^1) \rightarrow H^1(T)$ is induced by α and where $[S^1] \in H^1(S^1) = \mathbb{Z}$ is the fundamental class of the oriented circle S^1 . The following standard fact clarifies the identification $H^1(T) = \text{Hom}(T, S^1)$ that we have adopted in (B) and in Lemma 2.

Lemma 6. *The correspondence ι is an isomorphism of abelian groups. In particular, one has*

$$\iota(\omega_1^{a_1} \cdots \omega_n^{a_n}) = a_1 \omega_1 + \cdots + a_m \omega_m \text{ (in } H^1(T)),$$

where we use $\omega_i \in H^1(T)$ instead of $\iota(\omega_i)$ (as in Lemma 2 for the sake of simplicity).

We are now ready to establish the Theorem. Each representation $g \in \text{Hom}^0(\tilde{G}, U(n))$ induces a bundle map

$$\begin{array}{ccccc} \tilde{T} & \hookrightarrow & \tilde{G} & \rightarrow & G/T \\ g' \downarrow & & g \downarrow & & \rho(g) \downarrow \\ T^n & \hookrightarrow & U(n) & \rightarrow & F(n) \end{array},$$

where the g' is the restriction of g to \tilde{T} . From the naturality of the transgression (B) we have the commutative diagram

$$\begin{array}{ccc} H^1(T^n) & \xrightarrow{\tau_{U(n)}} & H^2(F(n)) \\ g'^* \downarrow & & \rho(g)^* \downarrow \\ H^1(\tilde{T}) & \xrightarrow[\cong]{\tau_{\tilde{G}}} & H^2(G/T) \end{array}.$$

With the $\tau_{U(n)}$ and $\tau_{\tilde{G}}$ being specified respectively in Lemma 1 and 2, we find that if the $\rho(g)^*$ is given by

$$\rho(g)^*(t_k) = a_{k,1}\omega_1 + \cdots + a_{k,m}\omega_m, \quad 1 \leq k \leq n-1 \text{ (cf. (C))},$$

in $H^2(G/T)$, we have

$$\begin{aligned} \text{(E)} \quad g'^*(t'_k) &= a_{k,1}\omega_1 + \cdots + a_{k,m}\omega_m, \quad 1 \leq k \leq n-1, \text{ and} \\ g'^*(t'_n) &= b_1\omega_1 + \cdots + b_m\omega_m \end{aligned}$$

in $H^1(\tilde{T})$, where $b_i = -(a_{1,i} + \cdots + a_{n-1,i})$. However, as is standard, the character $\chi(g)$ can be computed in terms of (E) as

$$\chi(g) = \iota^{-1}[g'^*(t'_1)] + \cdots + \iota^{-1}[g'^*(t'_n)]$$

This agrees with $s(\rho(g)^*)$ by Lemma 6.

7. Endnotes

This paper is by no means a final exposition in the topic, but is part of a larger effort to understand maps between flag manifolds by means of cohomology.

7.1. During the past two decades many works were devoted to the study of self-maps of flag manifolds, cf. [GH], [H₁], [H₂], [HH], [P₁], [P₂], [DZ₁], [Du].

However, the corresponding investigation into maps between different flag manifolds has not yet received as much attention as it deserves. The aim of present work is to demonstrate concrete examples (from representation theory) indicating the richness of the latter topic.

7.2. Our method can be extended to the study of homotopy classes of maps $G/T \rightarrow G'/T'$ between two arbitrary flag manifolds. For instance in the Problem in §2, one may replace the flag manifold $F(n)$ by

$$D(n) = SO(n)/T \text{ (resp. } T(n) = Sp(n)/T \text{)},$$

where the $SO(n)$ (resp. $Sp(n)$) is the special orthogonal group (resp. symplectic group) of order n and where $T \subset SO(n)$ (resp. $T \subset Sp(n)$) is a maximal torus. In these cases results analogous to the Theorem hold, only the character homomorphisms for the real and quaternionic representations will be involved and the correspondence s in §3 would be modified accordingly.

7.3. Let $Map(G/T, F(n))$ be the space of all continuous maps $G/T \rightarrow F(n)$. For a $f \in Map(G/T, F(n))$, let $M_f \subset Map(G/T, F(n))$ be the path-connected component containing f . In [Me] and [S], W. Meier and S. Smith initiated the project to determine the rational homotopy type of the space M_f .

Conjecture. The rational homotopy type of M_f is determined by the polynomial $s \circ r(f) \in \mathbb{Z}^+[\omega_1, \dots, \omega_m, \rho]$.

More precisely, if $g \in Hom^0(U(m), U(n))$ is such that

$$\alpha \circ \chi(g) = l(t_1 + \dots + t_m) + (n - lm) \text{ (cf. Corollary 3),}$$

then, modulo the products of odd dimensional spheres, the rational homotopy type of the space $M_{\rho(g)}$ has been determined by S. Smith [S]. We observe that the sum $t_1 + \dots + t_m$ is the Schur function associated to the partition $\mu = (1, 0, \dots, 0)$ ([M, P.42]). We note also from Lemma 5 that corresponding to any Schur function $s_\mu(y_1, \dots, y_m)$ with $s_\mu(1, \dots, 1) = n$, there is a representation $g_\mu \in Hom^0(U(m), U(n))$ with $\alpha \circ \chi(g_\mu) = s_\mu(y_1, \dots, y_m)$.

Problem. For a partition $\mu = (\mu_1, \mu_2, \dots, \mu_m)$, determine the rational homotopy type of the space $M_{\rho(g_\mu)}$.

Acknowledgements. This paper is based on a talk given at the International conference on *homotopy theory and related topics* held in Seoul, February 1-4, 2005. The author is grateful to S. Smith for suggestive discussion during the conference.

References

- [B] A. Borel, Sur la cohomologie des espaces fibres principaux et des homogenes de groupes de Lie compacts, *Ann. Math.*, 57(1953), 115-207.
- [BH] A. Borel and F. Hirzebruch, Characteristic classes and homogeneous space, I., *Amer. J. Math.*, 80 (1958), 458-538.
- [BS] R. Bott and H. Samelson, The cohomology ring of G/T , *Nat. Acad. Sci.* 41 (7) (1955), 490-492.
- [BGG] N. Bernstein, I.M. Gelfand and S. I. Gelfand, Schubert cells and cohomology of the spaces G/P , *Russian Math. Surv.* 28(1973), 1-26.
- [D] M. Demazure, Désingularization des variétés de Schubert généralisées, *Ann. Sci. École. Norm. Sup.* (4) 7(1974), 53-88.
- [Du] H. Duan, Self-maps of the Grassmannian of complex structures. *Compositio Math.* 132 (2002), no. 2, 159–175.
- [DZ₁] H. Duan and Xu-an Zhao, The classification of cohomology endomorphisms of certain flag manifolds. *Pacific J. Math.* 192 (2000), no. 1, 93–102.
- [DZ₂] H. Duan, Xu-an Zhao, The height function on the 2-dimensional cohomology of a flag manifold, *J. Lie Theory*, 15(1)(2005), 219-226.
- [DZZ] H. Duan, Xu-an Zhao and Xuezhi Zhao, The Cartan matrix and enumerative calculus. *J. Symbolic Comput.* 38 (2004), no. 3, 1119–1144.
- [FH] W. Fulton and J. Harris, Representation theory. A first course. *Graduate Texts in Mathematics*, 129. Springer-Verlag, New York, 1991.
- [GH] H. Glover, and W. Homer, Self-maps of flag manifolds. *Trans. Amer. Math. Soc.* 267 (1981), no. 2, 423–434.
- [H₁] M. Hoffman, On fixed point free maps of the complex flag manifold, *Indiana Univ. Math. J.* 33 (1984), no. 2, 249–255.
- [H₂] M. Hoffman, Endomorphisms of the cohomology of complex Grassmannians. *Trans. Amer. Math. Soc.* 281 (1984), no. 2, 745–760.
- [HH] M. Hoffman and W. Homer, On cohomology automorphisms of complex flag manifolds, *Proc. AMS.* 91(4)(1984), 643-648.
- [Hu] J. E. Humphreys, Introduction to Lie algebras and representation theory, *Graduated Texts in Math.* 9, Springer-Verlag New York, 1972.

- [K] S. Kumar, Kac-Moody groups, their flag varieties and representation theory. Progress in Mathematics, 204. Birkhäuser Boston, Inc., Boston, MA, 2002. xvi+606 pp.
- [L] L. Lesieur, Les problemes d'intersections sur une variete de Grassmann, C. R. Acad. Sci. Paris, 225 (1947), 916-917.
- [Li] P. Littelmann, A Littelwood–Richardson rule for symmetrizable Kac–Moody algebra, Invent. Math. 116(1994), 329-346.
- [LG] V. Lakshmibai and N. Gonciulea, Flag varieties, Hermann-Acutalities Mathematiques, Hermann Editeurs Des Sciences et Des Arts, 2001.
- [LS] V. Lakshmibai and C. S. Seshadri, Standard monomial theory and Schubert varieties—a survey, Proceedings of the conference on "Algebraic groups and Applications", 1991, 279-322, Manoj Prakashan.
- [M] I. G. Macdonald, Symmetric functions and Hall polynomials, Oxford Mathematical Monographs, Oxford University Press, Oxford, second ed., 1995.
- [Me] W. Meier, Rational universal fibrations and flag manifolds, Math. Ann. 258(1982), 329–340.
- [P] P. Pragacz, Algebro-geometric applications of Schur S- and Q-polynomials, Topics in invariant Theory (M.-P. Malliavin, ed.), Lecture Notes in Math., Vol. 1478, Springer-Verlag, Berlin and New York, 1991, 130-191.
- [P₁] S. Papadima, Rigidity properties of compact Lie group module maximal tori, Math. Ann. 275(1986) 637-652.
- [P₂] S. Papadima, Rational homotopy equivalences of Lie type. Math. Proc. Cambridge Philos. Soc. 104 (1988), no. 1, 65–80.
- [S] S. Smith, Rational classification of simple function space components for flag manifolds. Canad. J. Math. 49 (1997), no. 4, 855–864.
- [St] R. Stanley, Some combinatorial aspects of the Schubert calculus, Combinatoire et représentation du groupe symétrique, Strasbourg (1976), 217-251.
- [T] H. Tamvakis, The connection between representation theory and Schubert calculus, *preprint available on arXiv: math.AG/0306414*.