

Coordination of Multiple Dynamic Agents with Asymmetric Interactions

Hong Shi, Long Wang, Tianguang Chu

Intelligent Control Laboratory, Center for Systems and Control,
Department of Mechanics and Engineering Science,
Peking University, Beijing 100871, P. R. China

Abstract In this paper, we consider multiple mobile agents moving in Euclidean space with point mass dynamics. Using a coordination control scheme, we can make the group generate stable flocking motion. The control laws are a combination of attractive/repulsive and alignment forces, and the control law acting on each agent relies on the position information of all agents in the group and the velocity information of its neighbors. By using the control laws, all agent velocities become asymptotically the same, collisions can be avoided between all agents, and the final tight formation minimizes all agent global potentials. Moreover, we show that the velocity of the center of mass is invariant and is equal to the final common velocity. Furthermore, we study the motion of the group when the velocity damping is taken into account. We prove that the common velocity asymptotically approaches zero, and the final configuration minimizes the global potential of all agents. In this case, we can properly modify the control scheme to generate the same stable flocking. Finally, we provide some numerical simulations to further illustrate our results.

Keywords Collective behavior, swarm, robot teams, coordination, flocking, asymmetric interactions, multi-agent systems, collision avoidance, stability.

1 Introduction

In nature, flocking can be found everywhere and it can be regarded as a typical behavior of large number of interacting dynamic agents. This exists in the form of flocking of birds, schooling of fish, and swarming of bacteria. Understanding the mechanisms and operational principles in them can provide useful ideas for developing distributed cooperative control and coordination of multiple mobile autonomous agents/robots. In recent years, distributed control/coordination of the motion of multiple dynamic agents/robots has emerged as a topic of major interest [1]–[4]. This is partly due to recent technological advances in communication and computation, and wide applications of multi-agent systems in many engineering areas including cooperative control of unmanned aerial vehicles (UAVs), scheduling of automated highway systems, schooling for underwater vehicles, attitude alignment for satellite clusters and congestion control in communication networks

This work was supported by the National Natural Science Foundation of China (No. 10372002 and No. 60274001) and the National Key Basic Research and Development Program (No. 2002CB312200).

[5][8]. Correspondingly, there has been considerable effort in modelling and exploring the collective dynamics in physics, biology, and control engineering, and trying to understand how a group of autonomous creatures or man-made mobile autonomous agents/robots can cluster in formations without centralized coordination and control [10][23].

In order to generate computer animation of the motion of flocks, Reynolds [9] modelled the boid as an object moving in a three dimensional environment based on the positions and velocities of its nearby flockmates and introduced the following three rules (named steering forces) [9]:

- 1) Collision Avoidance: avoid collisions with nearby flockmates,
- 2) Velocity Matching: attempt to match velocity with nearby flockmates, and
- 3) Flock Centering: attempt to stay close to nearby flockmates.

Subsequently, Vicsek et al. [10] proposed a simple model of autonomous agents (i.e., points or particles). In the model, all agents move at a constant identical speed and each agent updates its heading as the average of the heading of agent itself with its nearest neighbors plus some additive noise. They demonstrated numerically that all agents will eventually move in the same direction, despite the absence of centralized coordination and control. In fact, Vicsek's model can be seen as a special case of Reynolds's model, and it only considers the velocity matching between agents. Jadbabie et al. [11] and Savkin [12] used two kinds of completely different methods to provide the theoretical explanation for the observed behaviors in Vicsek's model, respectively. According to the results in [9], Tanner et al. [13] studied a swarm model that consists of multiple mobile agents moving on the plane with double integrator dynamics. They introduced a set of control laws that enabled the group to generate stable flocking motion and provided strictly theoretical justification. However, it is perhaps more reasonable to take the agents' masses into account and consider the point mass model in which each agent moves in n -dimensional space based on the Newton's law. In this paper, we investigate the collective behavior of multi-agent systems in n -dimensional space with point mass dynamics.

In [13], the authors used an undirected graph to describe the neighboring relations between agents, which means that the neighboring relations are mutual. In other words, they only considered the case with bidirectional information exchange between agents. However, under some circumstances, the information exchange is not mutual. In fact, due to the agent differences, they maybe have different action forces on different agents and even have different sense ranges, hence, the influence intensities between two agents might be different with each other and even their information can not be exchanged with each other at all. For example, in a group of agents with spherical sense neighborhoods but with different radii of the neighborhoods or a group of agents with conic sense neighborhoods, the information exchange among them might be unidirectional. A group of mobile robots with conic vision range is just an example. In this paper, the results in [13] are extended to a directed graph. We consider the stability properties of the group in the case of directed information exchange. In order to generate stable flocking, we introduce a set of control laws so that each agent regulates its velocity based on a fixed set of "neighbors" and regulates its position such that its global potential become minimum. Note that, in this paper, we only consider the fixed topology of the neighboring relations, and the case that the information topology is dynamic will be discussed in another paper. Here, the control laws are a combination of attractive/repulsive and alignment forces. By using the control laws, all agent velocities become asymptotically the same, collisions can be avoided between all agents, and the final tight formation minimizes all agent potentials.

This paper is organized as follows: In Section 2, we formulate the problem to be investigated. Some basic concepts and results in graph theory are provided in Section 3. We analyze the system stability with some specific control laws in Section 4. Some numerical simulations are presented to further illustrate our results in Section 5. Finally, we briefly summarize our results in Section 6.

2 Problem Formulation

We consider a group of N agents moving in an n -dimensional Euclidean space, each has point mass dynamics described by

$$\begin{aligned}\dot{\underline{x}}^i &= \underline{v}^i; \\ m_i \dot{\underline{v}}^i &= \underline{u}^i; \quad i = 1; \dots; N;\end{aligned}\quad (1)$$

where $\underline{x}^i = (\underline{x}_1^i; \dots; \underline{x}_n^i)^T \in \mathbb{R}^n$ is the position vector of agent i ; $\underline{v}^i = (\underline{v}_1^i; \dots; \underline{v}_n^i)^T \in \mathbb{R}^n$ is its velocity vector, $m_i > 0$ is its mass, and $\underline{u}^i = (\underline{u}_1^i; \dots; \underline{u}_n^i)^T \in \mathbb{R}^n$ is the (force) control input acting on agent i . $\underline{x}^{ij} = \underline{x}^i - \underline{x}^j$ denotes the relative position vector between agents i and j .

Our aim is to make the whole group move at a common velocity and maintain constant distances between all agents. We first consider the ideal case, that is, we ignore the velocity damping. In order to achieve our objective, we try to decrease the velocity differences between agents, and at the same time, regulate their distances such that their global potentials become minimum. Hence, we choose the control law for each agent to be a combination of two components. The control input \underline{u}^i for agent i is

$$\underline{u}^i = \dot{\underline{x}}^i + \underline{v}^i; \quad (2)$$

where $\dot{\underline{x}}^i$ is used to regulate the potentials among agents and \underline{v}^i is used to regulate the velocity of agent i to the weighted average of its "neighbors". $\dot{\underline{x}}^i$ is derived from the social potential fields which is described by artificial social potential function, V^i , which is a function of the relative distances between agent i and its neighbors. Collision-free and cohesion in the group can be guaranteed by this term. Note that $\dot{\underline{x}}^i$ indicates the tendency of collision avoidance and cohesion of the flocks, whereas \underline{v}^i indicates the tendency of agent velocity matching.

Certainly, in some cases, the velocity damping can not be ignored. For example, the objects moving in viscous environment and the mobile objects with high speeds, such as air vehicles, are subject to the influence of velocity damping. Then, under these circumstances, the model in (1) should be the following form

$$\begin{aligned}\dot{\underline{x}}^i &= \underline{v}^i; \\ m_i \dot{\underline{v}}^i &= \underline{u}^i - k_i \underline{v}^i;\end{aligned}\quad (3)$$

where $k_i > 0$ is the "velocity damping gain", $k_i \underline{v}^i$ is the velocity damping term, and \underline{u}^i is the control input for agent i . Note that we assume the damping force is in proportion to the magnitude of velocity. And, because the "velocity damping gain" is determined by the shape and size of the object, the property of medium, and some other factors, we assume that the damping gains k_i , $i = 1; \dots; N$ are not equal to each other. Certainly, in some cases, the assumption of the same gain is enough. In order to achieve our aim,

the velocity damping should be cancelled by some terms in the control laws. Thus, we modify the control scheme to be

$$\mathbf{u}^i = \mathbf{v}^i + \mathbf{v}^i + k_i \mathbf{v}^i: \quad (4)$$

3 Main Results

In this section, we investigate the stability properties of multiple mobile agents with point mass dynamics described in (1). We present explicit control input in (2) for the terms \mathbf{v}^i and \mathbf{v}^i . In this paper, the control law acting on each agent is based on two kinds of information topologies that is the position information topology and the velocity information topology. We will employ algebraic graph theory as basic tools to study the properties of the group. Some concepts and results in graph theory are given in the Appendix.

In this paper, we assume that each agent is equipped with two onboard sensors: the position sensor which is used to sense the position information of the neighbors and the velocity sensor which is used to sense the velocity information of its neighbors, and assume that all the sensors can sense instantaneously. Correspondingly, we define two kinds of structure topologies to describe the neighboring relations between the agents. We will use an undirected graph G to describe the position sensor information flow and use a weighted directed graph D to describe the velocity sensor information flow.

First, we make the following definitions and assumptions.

Definition 1: (Position neighboring graph) The position neighboring graph, $G = (V; E)$, is an undirected graph consisting of a set of vertices, $V = \{n_1; \dots; n_N\}$, indexed by the agents in the group, and a set of edges, $E = \{(n_i; n_j) \mid n_i, n_j \in V, i \neq j\}$, which contain unordered pairs of vertices that represent the position neighboring relations.

Definition 2: (Velocity neighboring graph) The velocity neighboring graph, $D = (V; E)$, is a directed graph consisting of a set of vertices, $V = \{n_1; \dots; n_N\}$, indexed by the agents in the group, and a set of arcs, $E = \{(n_i; n_j) \mid n_i, n_j \in V, i \neq j\}$, which contain ordered pairs of vertices that represent the velocity neighboring relations.

Note that, in E , an arc $(n_i; n_j)$ represents a unidirectional velocity information exchange link from n_i to n_j , which means that agent i can sense the velocity of agent j .

Assumption 1: The position neighboring graph G is complete.

In order to make the total potential of each agent be global minimum and at the same time, ensure collision-free in the group, we assume that the position neighboring graph is complete. This means that, each agent can always obtain the position information of all the other agents in the group. Certainly, in the case that the position neighboring relation is determined by a certain neighborhood around the agent and consequently cause the topology of the neighboring graph G to be dynamic, we can also guarantee collision avoidance in the group.

Assumption 2: The velocity neighboring graph D is weakly connected.

In this paper, we consider a group of mobile agents with fixed topology, so D is weakly connected and does not change with time. Denote the set $N_i = \{j \mid a_{ij} > 0\}$, $i \in \{1, \dots, N\}$ which contains all neighbors of agent i . If agent j is a neighbor of agent i , we denote $j \in N_i$, and otherwise we denote $j \notin N_i$.

Definition 3 [13]: (Potential function) Potential V^{ij} is a differentiable, nonnegative, radially unbounded function of the distance $\|x^i - x^j\|$ between agents i and j , such that

i) $V^{ij}(kx^{ij}) \leq 1$ as $kx^{ij} \leq 0$,

ii) V^{ij} attains its unique minimum when agents i and j are located at a desired distance.

Functions $V^{ij}, i, j = 1; \dots; N$ are the artificial social potential functions that govern the interindividual interactions. Cohesion and separation can be achieved by artificial potentials [6]. One example of such potential function is the following

$$V(x) = a \ln x^2 + \frac{b}{x^2};$$

where $x \in \mathbb{R}_+ = (0; 1)$ is variable, $a > 0$ and $b > 0$ are some constants. It is easy to see that V attains its unique minimum when $x = \sqrt{\frac{b}{a}}$. Hence, when the distance $kx^i - x^j k$ between agents i and j is $\sqrt{\frac{b}{a}}$, the potential function V^{ij} attains its unique minimum.

By the definition of V^{ij} , the total potential of agent i can be expressed as

$$V^i = \sum_{j=1; j \neq i}^N V^{ij}(kx^{ij}k); \quad (5)$$

Agent dynamics are different in ideal case (i.e., velocity damping is ignored) and nonideal case. This means that the agent has different motion equations in the two cases. Hence, in what follows, we will discuss the motion of the group in the two different cases, respectively.

3.1 Ideal Case

In this case, in order to achieve our control aim, we take the control law u^i to be

$$u^i = \sum_{j=1; j \neq i}^N w_{ij} (v^i - v^j) + \sum_{j=1; j \neq i}^N r_{x^i} V^{ij}; \quad (6)$$

Note that, $w_{ij} \geq 0$, and $w_{ii} = 0, i, j = 1; \dots; N$ represent the interaction coefficients. And $w_{ij} > 0$ if agent j is a neighbor of agent i , and is 0 otherwise. We denote $W = [w_{ij}]$. Thus, by the weakly connectivity of the velocity neighboring graph, $W + W^T$ is irreducible. The control law in (6) implies that we adopt the local velocity regulation and the global potential regulation to achieve our aim.

In the discussion to follow, we will need the concept of weight balance condition defined below:

Weight Balance Condition [20]: consider the weight matrix $W = [w_{ij}] \in \mathbb{R}^{N \times N}$, for all $i = 1; \dots; N$, we assume that $\sum_{j=1}^N w_{ij} = \sum_{j=1}^N w_{ji}$.

The weight balance condition has a graphical interpretation: consider the directed graph associated with a matrix, weight balance means that, for any node in this graph, the weight sum of all incoming edges equals the weight sum of all outgoing edges [25]. The weight balance condition can find physical interpretations in engineering systems such as water flow, electrical current, and traffic systems.

Proposition 1: Let D be a weighted directed graph such that the weight balance condition is satisfied. Then D is strongly connected if and only if it is weakly connected.

Proof: It is obvious that if D is strongly connected, then it is weakly connected. Hence, we only need to prove that if D is weakly connected, then it is strongly connected.

In the following, we will use the way of contradiction to prove it. Assume that D is weakly connected, but not strongly connected, then we denote all strongly connected components of D as D_1, \dots, D_m where m is an integer and $m > 1$. If there is an arc starting in D_i and ending in D_j , then any arc joining D_i to D_j must start in D_i . Hence we can define a directed graph \tilde{D} with the strongly connected components of D as its vertices, and such that there is an arc from D_i to D_j in \tilde{D} if and only if there is an arc in D starting in D_i and ending in D_j . Obviously that the directed graph \tilde{D} can not contain any cycles since otherwise the number of strongly connected components of D will be equal to or less than $m - 1$. It follows that there is a strongly connected component, D_1 say, such that any arc that ends on a vertex in it must start at a vertex in it. Since D is weakly connected, there is at least one arc that starts in D_1 and ends on a vertex not in D_1 . Consequently, in D_1 , the sum of in-degree of all vertices is less than the sum of out-degree of all vertices. This means that there must be a vertex in D such that the weight balance condition can not be satisfied. Thus we have the contradiction.

Hence, if a weighted directed graph is weakly connected and the weights of each agent satisfy the weight balance condition, then the directed graph must be strongly connected.

3.1.1 Stability Analysis

Before presenting the main results of this paper, we first prove the following important lemma.

Lemma 1: Let $A \in \mathbb{R}^{n \times n}$ be any diagonal matrix with positive diagonal entries. Then

$$A \text{spanflg}^\perp \setminus \text{spanflg} = 0;$$

where $\mathbf{1} = (1, \dots, 1)^T \in \mathbb{R}^n$, spanflg is the space spanned by vector $\mathbf{1}$, and spanflg^\perp is the orthogonal complement space of spanflg .

Proof: Let $p \in A \text{spanflg}^\perp \setminus \text{spanflg}$: Then $p \in \text{spanflg}^\perp$ and there is some $q \in \text{spanflg}^\perp$ such that $p = Aq$. It follows that $q^T A q = q^T p = 0$: Since A is positive definite by assumption, we have $q = 0$ and hence $p = 0$.

Theorem 1: By taking the control law in (6), under Assumption 2 and the weight balance condition, all agent velocities in the group described in (1) become asymptotically the same, collision avoidance can be ensured between all agents and the group final configuration minimizes all agent global potentials.

Proof: Choose the following positive semidefinite function

$$J = \frac{1}{2} \sum_{i=1}^N V^i + m_i v^{iT} v^i :$$

It is easy to see that J is the sum of the total artificial potential energy and the total kinetic energy of all agents in the group. Define the level sets of J in the space of agent velocities and relative distances

$$= \{v^i; x^{ij} \mid J \leq c\} : \quad (7)$$

In what follows, we will prove that the set is compact. In fact, the set $\{v^i; x^{ij} \mid J \leq c\}$ such that $J \leq c$ ($c > 0$) is closed by continuity. Moreover, boundedness can be proved under Assumption 1, namely, from $J \leq c$, we have that $V^{ij} \leq c$. Potential V^{ij} is radially

unbounded, so there must be a positive constant d such that $kx^{ij}k \leq d$, for all $i, j = 1, \dots, N$. In the same way, $w^i \leq 2cm_i$, thus $kv^i k \leq \frac{2cm_i}{2cm_i}$.

By the symmetry of V^{ij} with respect to x^{ij} and $x^{ij} = x^{ji}$, it follows that

$$\frac{\partial V^{ij}}{\partial x^{ij}} = \frac{\partial V^{ij}}{\partial x^i} = \frac{\partial V^{ij}}{\partial x^j}; \quad (8)$$

and therefore

$$\frac{d}{dt} \sum_{i=1}^N \frac{1}{2} V^i = \sum_{i=1}^N r_{x^i} V^i = \dot{J}.$$

Calculating the time derivative of J along the solution of system (1), we have

$$\begin{aligned} \dot{J} &= \sum_{i=1}^N v^{iT} \sum_{j \in \mathcal{N}_i} w_{ij} (v^i - v^j) = v^T (L - I_n) v \\ &= \frac{1}{2} v^T (L + L^T - I_n) v; \end{aligned} \quad (9)$$

where $v = (v^1; \dots; v^N)^T$ is the stack vector of all agent velocity vectors, $L = [l_{ij}]$ with

$$l_{ij} = \begin{cases} \sum_{k=1, k \neq i}^N w_{ik} & i = j; \\ -w_{ij} & i \neq j; \end{cases} \quad (10)$$

is the Laplacian matrix of the weighted velocity neighboring graph, and $(L + L^T) - I_n$ is the Kronecker product of $L + L^T$ and I_n , with I_n the identity matrix of order n .

From the definition of matrix L , under the weight balance condition, it is easy to see that $L + L^T$ is symmetric and has the properties that every row sum is equal to 0, the diagonal elements are positive, and all the other elements are nonpositive. By matrix theory [25], all eigenvalues of $L + L^T$ are nonnegative. Hence, matrix $L + L^T$ is positive semidefinite. By the connectivity of graph D , we know that $L + L^T$ is irreducible and the eigenvector associated with the single zero eigenvalue is 1_N . On the other hand, it is known that the identity matrix I_n has an eigenvalue $= 1$ of n multiplicity and n linearly independent eigenvectors

$$p^1 = [1; 0; \dots; 0]^T; p^2 = [0; 1; 0; \dots; 0]^T; \dots; p^n = [0; \dots; 0; 1]^T.$$

By matrix theory [25], the eigenvalues of $(L + L^T) - I_n$ are nonnegative, $= 0$ is an eigenvalue of multiplicity n and the associated eigenvectors are

$$q^1 = [p^1; \dots; p^1]^T; \dots; q^n = [p^n; \dots; p^n]^T.$$

Thus $\dot{J} = 0$, and $\dot{J} = 0$ implies that all agents have the same velocity vector, that is, the vector $v_k = (v_k^1; \dots; v_k^N)^T$ ($k = 1, \dots, n$), which is composed of every corresponding k th component $v_k^1; \dots; v_k^N$ of $v^1; \dots; v^N$, is contained in $\text{span}\{1\}$, where $1 = (1; \dots; 1)^T \in \mathbb{R}^N$. It follows that $\dot{x}^{ij} = 0$, $(i, j) \in \mathcal{E}$.

We use LaSalle's invariance principle [26] to establish convergence of system trajectories to the largest positively invariant subset of the set defined by $E = \{v | \dot{J} = 0\}$. In E , the agent velocity dynamics are

$$\dot{v}^i = \frac{1}{m_i} u^i = \frac{1}{m_i} \sum_{j=1, j \neq i}^N r_{x^i} V^{ij} = \frac{1}{m_i} r_{x^i} V^i$$

and therefore it follows that

$$\underline{v} = (M - I_n) \begin{bmatrix} r_{x^1} V^1 \\ \vdots \\ r_{x^N} V^N \end{bmatrix} = (M - B) \begin{bmatrix} r_{x^{i_1}} V^{i_1} \\ \vdots \\ r_{x^{i_N}} V^{i_N} \end{bmatrix}; \quad (11)$$

where $M = \text{diag}(\frac{1}{m_1}, \dots, \frac{1}{m_N})$, and the matrix B is the incidence matrix of the position neighboring graph. Hence

$$\underline{v}_k = (M - B) [r_{x^{i_j}} V^{i_j}]_k; \quad k = 1, \dots, n;$$

Thus, $\underline{v}_k \in \text{range}(M - B)$, $k = 1, \dots, n$. By matrix theory, we have

$$\text{range}(M - B) = M \text{range}(B) = M \text{range}(B B^T) = M \text{spanflg},$$

and therefore

$$\underline{v}_k \in M \text{spanflg}; \quad k = 1, \dots, n; \quad (12)$$

In any invariant set of E , by $\underline{v}_k \in \text{spanflg}$, we have

$$\underline{v}_k \in \text{spanflg}; \quad (13)$$

By Lemma 1, we get from (12) and (13)

$$\underline{v}_k \in (M \text{spanflg}) \setminus \text{spanflg} = \{0\}; \quad k = 1, \dots, n;$$

Thus, in steady state, all agent velocities no longer change and from (11), the potential V^i of each agent is globally minimized. Collision-free can be ensured between the agents since otherwise it will result in $V^i \rightarrow \infty$.

Remark 1: If we take the control law for agent i to be

$$u^i = \sum_{j \in N_i} (v^i - v^j) \sum_{j=1, j \neq i}^N r_{x^i} V^{ij}; \quad (14)$$

then the weight balance condition implies that, in the velocity neighboring graph, for each vertex, the number of arcs starting at it is equal to the number of arcs ending on it. When we take the control law in (14), by using the same analysis method as in Theorem 1, we can also obtain the same conclusion.

Note that, from (9), we see that the interaction coefficients in control law (6) can influence the decaying rate of the total energy J . Hence, we conclude that the convergence rate of the system will be influenced by the interaction coefficients. Explicit analysis on this topic will be presented in Section 4.1.3.

3.1.2 Common Velocity

In this section, we will show that the common velocity can be obtained by the initial velocities of all agents.

The position vector of the center of mass in system (1) is defined as

$$\mathbf{x} = \frac{\sum_{i=1}^N m_i \mathbf{x}^i}{\sum_{i=1}^N m_i};$$

Thus, the velocity vector of the center of mass is

$$\underline{v} = \frac{\sum_{i=1}^N m_i \underline{v}^i}{\sum_{i=1}^N m_i}.$$

By using control law (6), we obtain

$$\underline{\dot{v}} = -\frac{1}{\sum_{i=1}^N m_i} \sum_{i=1}^N \sum_{j \in \mathcal{N}_i} w_{ij} (\underline{v}^i - \underline{v}^j) + \sum_{j=1; j \neq i}^N r_{xi} \underline{v}^{ij}.$$

By the symmetry of function V^{ij} with respect to x^{ij} , under the weight balance condition, we get $\underline{\dot{v}} = 0$. This means that, by using control law (6), the velocity of the center of mass is invariant.

Therefore, combining Theorem 1 and the analysis above, we have the following theorem.

Theorem 2: By taking the control law in (6), under Assumption 2 and the weight balance condition, the final common velocity is equal to the initial velocity of the center of mass, that is, the final velocity \underline{v}_f is

$$\underline{v}_f = \frac{\sum_{i=1}^N m_i \underline{v}^i(0)}{\sum_{i=1}^N m_i};$$

where $\underline{v}^i(0)$ is the velocity value of agent i at initial time $t = 0$, $i = 1; \dots; N$.

Remark 2: Note that, by the calculation above, we can see that the final common velocity is determined by the masses and the initial velocities of all agents, and does not rely on the neighboring relations and the magnitudes of the interaction coefficients under Assumption 2 and the weight balance condition.

Remark 3: Even if the velocity neighboring graph is not connected, under the weight balance condition, the velocity of the center of mass is still invariant by using control law (6). However, in this case, the final velocities of all agents might be different. In fact, when the velocity neighboring graph is not connected, under the weight balance condition, control law (6) only ensures that all agents from the same connected group will have the same final velocity, and the final velocities of any two different connected groups might not be equal to each other.

Remark 4: Using the control law in (6), from Theorems 1 and 2, we know that if the initial velocity of the center of mass is zero, the center of mass will not drift. All agents adjust their positions and velocities to minimize the total potential, and the final common velocity of all agents is zero.

Hence, by using control law (6), under Assumption 2, the whole group can move ahead at a common nonzero velocity if and only if the initial velocity of the center of mass is not zero.

Definition 4: The average velocity of all agents is defined as $\bar{\underline{v}} = \frac{1}{N} \sum_{i=1}^N \underline{v}^i$.

Remark 5: If we modify the control law \underline{u}^i to be

$$\underline{u}^i = \sum_{j \in \mathcal{N}_i} m_j w_{ij} (\underline{v}^j - \underline{v}^i) + \sum_{j=1; j \neq i}^N m_j r_{xi} \underline{v}^{ij}; \quad (15)$$

where m_i and w_{ij} are defined as before, by choosing the Lyapunov function

$$J = \frac{1}{2} \sum_{i=1}^N (V^i + v^{iT} v^i);$$

under Assumption 2 and the weight balance condition, we can still get the results as in Theorem 1. Since the proof is similar to the proof of Theorem 1, we omit the details.

Moreover, by using the control law in (15), under Assumption 2 and the weight balance condition, we can obtain that the average velocity of all agents in group (1) is invariant and therefore the final velocity of the group is the average of the initial velocities of all agents, that is,

$$v_f = \frac{\sum_{i=1}^N v^i(0)}{N};$$

where $v^i(0)$ is the velocity value of agent i at initial time $t = 0$, $i = 1; \dots; N$. The final common velocity does not rely on the agents' masses, the neighboring relations, or the magnitudes of the interaction coefficients under Assumption 2 and the weight balance condition.

3.1.3 Convergence Rate Analysis

From the discussion above, we know that the coupling coefficients can influence the decay-ing rate of the energy function J , hence, we guess that the coupling coefficients can also influence the convergence rate of system (1). In the following, we will present qualitative analysis of the influence of the weights w_{ij} on the convergence rate of the system.

We consider the dynamics of the error system. From the discussion in 4.1.2, the velocity of the center of mass in system (1) is invariant. Thus, we define the following error vectors:

$$\begin{aligned} e^i &= x^i - x; \\ e_v^i &= v^i - v; \end{aligned}$$

where x and v are the position vector and the velocity vector of the center of mass, respectively. Hence, the error dynamics is given by

$$\begin{aligned} \dot{e}^i &= e_v^i; \\ \dot{e}_v^i &= \frac{1}{m_i} u^i; \quad i = 1; \dots; N; \end{aligned} \quad (16)$$

By the definition of V^{ij} and $e^i = x^i - x$, we get

$$r_{x^i} V^{ij}(kx^{ij}k) = r_{e^i} V^{ij}(ke^{ij}k);$$

By using the control law in (6), we obtain

$$\dot{e}_v^i = \frac{1}{m_i} \sum_{j \in N_i} w_{ij} (e_v^i - e_v^j) - \sum_{j=1; j \neq i}^N r_{e^i} V^{ij}(ke^{ij}k); \quad (17)$$

We choose the following positive semi-definite function

$$J = \frac{1}{2} \sum_{i=1}^N (V^i)^2 + m_i e_v^{iT} e_v^i$$

which is the energy function of the error system (16). $(V^i)^T$ is the potential of agent i in (16) and it equals V^{ij} by the definition of potential function V^{ij} .

Calculating the time derivative of J , we have

$$\begin{aligned} \dot{J} &= \sum_{i=1}^N \sum_{j \neq i} w_{ij} e_v^{iT} (e_v^i - e_v^j) = e_v^T (L - I_n) e_v \\ &= -\frac{1}{2} e_v^T (L + L^T - I_n) e_v; \end{aligned} \quad (18)$$

where $e_v = (e_v^{1T}; \dots; e_v^{NT})^T$, and L and I_n are defined as before.

Using the same analysis method as in Theorem 1, we have $\dot{J} \leq 0$, and $\dot{J} = 0$ implies that $e_v^1 = e_v^2 = \dots = e_v^N$. This occurs only when $e_v^1 = e_v^2 = \dots = e_v^N = 0$, that is, this occurs only when all agents have the same velocity. In other words, if there exist two agents with different velocities, the energy function J is strictly monotone decreasing with time. Certainly, before the group forms the final tight configuration, there might be the case that all agents have the same velocity, but due to the regulation of the potentials among agents, it instantly changes into the case that not all agents have the same velocity except when the group has achieved the final stable state. Hence, the decaying rate of energy is equivalent to the convergence rate of the system. It is easy to see that when all agents have not achieved the common velocity, for any solution of the error system (16), e_v must be in the subspace spanned by eigenvectors of $(L + L^T - I_n)$ corresponding to the nonzero eigenvalues. Thus, from (18), we have $\dot{J} = -\lambda_2 e_v^T e_v$, where λ_2 denotes the second smallest real eigenvalue of matrix $L + L^T$. Therefore, we have the following conclusion: The convergence rate of the system relies on the second smallest real eigenvalue of matrix $L + L^T$ with L defined as in (10).

3.2 Nonideal case

We know that, in some cases, the velocity damping should not be ignored. Then, if we still take control law (6), what will be the motion of the group? In fact, in this case, the total force acting on the i th agent is

$$u^i = \sum_{j \neq i} w_{ij} (v^i - v^j) + \sum_{j=1; j \neq i}^N r_{xi} V^{ij} - k_i v^i; \quad (19)$$

where w_{ij} and k_i are defined as before.

The following theorem shows the motion and the final configuration of the group.

Theorem 3: By taking the control law in (6), under Assumption 2 and the weight balance condition, all agent velocities in the group described in (3) become asymptotically the same, all agents finally stop moving, collision avoidance can be ensured between all agents, and the group finally converges to minimize all agent global potentials.

Proof: Taking the Lyapunov function J defined as in Theorem 1, that is,

$$J = \frac{1}{2} \sum_{i=1}^N (V^i + m_i v^{iT} v^i);$$

We can show analogously that the set $\Omega = \{(v^i; x^{ij}) \mid \dot{J} \leq -c \text{ for } c > 0\}$ is compact.

Calculating the time derivative of J , we have

$$\dot{J} = -\frac{1}{2} \mathbf{v}^T (\mathbf{L} + \mathbf{L}^T) \mathbf{v} - \mathbf{v}^T (\mathbf{H} - \mathbf{I}_n) \mathbf{v};$$

where \mathbf{v} and \mathbf{L} are defined as in Theorem 1, and $\mathbf{H} = \text{diag}(k_1; \dots; k_N)$ with $k_i > 0$ is the velocity damping gain. It is easy to see that \mathbf{H} is positive definite. Using the same analysis method as in Theorem 1, we know that $\dot{J} = 0$, and $J = 0$ implies that $\mathbf{v}^1 = \dots = \mathbf{v}^N$ and they all must equal zero. We denote $E = \{\mathbf{v} | J = 0\}$. In E , the agent velocity dynamics become

$$\dot{\mathbf{v}}^i = -\frac{1}{m_i} \sum_{j=1, j \neq i}^N \mathbf{r}_{xi} V^{ij} = -\frac{1}{m_i} \mathbf{r}_{xi} \mathbf{V}^i;$$

Following the proof of Theorem 1, we can conclude that $\mathbf{v}_k = 0$, hence $\dot{\mathbf{v}}^i = 0$, $i = 1; \dots; N$, which means that the agent velocity no longer changes in steady state. All agents will finally stop moving, and the final configuration minimizes all agent global potentials. Furthermore, during the course of motion, collisions can be avoided between the agents.

Remark 6: It can be shown that if we use control law (15), we can still obtain all results in Theorem 3.

Remark 7: From Theorem 3, we know that due to damping, all agents eventually stop moving. This is because when all agents eventually move ahead at a common velocity, control input (6) equals zero.

In order to make the group have the same properties as in ideal case, the control law should contain the velocity damping term. Hence, we modify the control scheme to be (4), where \mathbf{v}^i and $\dot{\mathbf{v}}^i$ are defined as in (6). Then, the actual total force acting on agent i is

$$\mathbf{u}^i = \sum_{j=1, j \neq i}^N \mathbf{w}_{ij} (\mathbf{v}^i - \mathbf{v}^j) - \sum_{j=1, j \neq i}^N \mathbf{r}_{xi} V^{ij};$$

Following Theorems 1 and 2, we can easily obtain the same stable flocking motion and the final common velocity, that is, when the velocity damping is taken into account, by using control scheme (4), under Assumption 2 and the weight balance condition, all agent velocities in the group described in (3) become asymptotically the same, collision-free can be ensured between all agents, the group final configuration minimizes all agent global potentials, and the final common velocity is equal to the initial velocity of the center of mass.

4 Simulations

In this section, we will present some numerical simulations for the system described in (1) in order to illustrate the results obtained in the previous sections.

These simulations all are performed with ten agents moving on the plane whose initial positions, velocities and the velocity neighboring relations are selected randomly, but they satisfy: 1) all initial positions are chosen within a ball of radius $R = 15$ [m] centered at the origin, 2) all initial velocities are selected with arbitrary directions and magnitudes in the range of $(0, 10)$ [m/s], and 3) the velocity neighboring graph is connected. All agents

have different masses to each other and they are randomly selected in the range of $(0, 1)$ [kg].

Note that, because the position neighboring graph is complete, we will not describe it. In the following figures, we only present the velocity neighboring relations.

Figs. 1{6 show the results in one of our simulations, where the control laws are taken in the form of (6) with the explicit potential function

$$V^{ij} = \frac{1}{2} \ln kx^{ij}k^2 + \frac{5}{2kx^{ij}k^2}; \quad i, j = 1; \dots; 10;$$

The interaction coefficient matrix W is generated randomly such that $\sum_{j=1}^{p_0} w_{ij} = \sum_{j=1}^{p_0} w_{ji}$, $w_{ii} = 0$, and the nonzero w_{ij} satisfy $0 < w_{ij} < 1$ for all $i, j = 1; \dots; 10$. We run the simulation for 200 seconds.

In Figs. 1{4, the blue lines all represent the bidirectional neighboring relations and the red lines with arrows represent the unidirectional neighboring relations. Fig. 1 shows the group initial state which includes the initial positions, velocities and the velocity neighboring relations. Figs. 2 and 3 depict the motion trajectories of all agents and the configurations of the group, respectively, where the black solid arrow direction represent the motion direction of the agents, and the dotted lines represent the agent trajectories. In order to indicate the influence of potential function on the group cohesion and configuration, we present the group configuration in Fig. 2 at time $t = 60$ s. It can be seen from Figs. 2 and 3 that, during the course of motion, all agents regulate their positions to minimize their potentials and regulate their velocities to become the same. Fig. 4 shows the final steady state configuration and the common velocity at $t = 200$ s. By numerical calculation, we can obtain that all agents achieve the same velocity approximately at $t = 128.92$ s and the final common velocity equals the initial velocity of the center of mass. In Fig. 5, the star represents the initial position of the center of mass, and it can be seen from it that the velocity of the center of mass is invariant. Fig. 6 is the velocity curves. The solid arrow indicates the tendency of velocity variation. Fig. 6 distinctly demonstrates that all agent velocities asymptotically approach the same.

Hence, numerical simulation also indicates that, by using the control law in (6), under the assumption of the connectivity of the velocity neighboring graph and the weight balance condition, stable flocking motion can be achieved.

For the case that the initial velocity of the center of mass is zero, we also perform some simulations. Fig. 7 is one of them and we run its associated simulation for 3000 seconds. In Fig. 7, the star represents the position of the center of mass. In the simulation, the center of mass is always stationary, the final configuration no longer changes, the whole group does not drift, and all agents finally stop moving.

5 Conclusions

In this paper, we have investigated the collective behavior of multiple mobile agents moving in n -dimensional space with point mass dynamics and introduced a set of control laws which enable the group to generate stable flocking motion. We analyzed the group properties in two different cases, respectively. When we ignored the velocity damping, using a coordination control scheme, we can make the group generate stable flocking

motion. The control laws are a combination of attractive/repulsive and alignment forces and the control law acting on each agent relies on the position information of all agents in the group and the velocity information of all its neighboring agents. The control laws ensure that all agent velocities become asymptotically the same, collisions can be avoided between all agents, and the final tight formation minimizes all agent global potentials. Moreover, we analyzed the magnitude and direction of the final velocity and showed that the final common velocity is equal to the initial velocity of the center of mass of the system. When the velocity damping is taken into account, in order to generate stable flocking, we properly modified the control scheme such that the velocity damping was cancelled by some terms in the control laws. Finally, numerical simulations were worked out to further verify our theoretical results.

6 Appendix: Graph Theory Preliminaries

In this section, we briefly summarize some basic concepts and results in graph theory that have been used in this paper. More comprehensive discussions can be found in [27].

A undirected graph G consists of a vertex set $V = \{n_1; n_2; \dots; n_m\}$ and an edge set $E = \{(n_i; n_j) : n_i, n_j \in V\}$, where an edge is an unordered pair of distinct vertices of V . If $n_i, n_j \in V$, and $(n_i; n_j) \in E$, then we say that n_i and n_j are adjacent or neighbors, and denote this by writing $n_j \sim n_i$. A graph is called complete if every pair of vertices are adjacent. A path of length r from n_i to n_j in a undirected graph is a sequence of $r + 1$ distinct vertices starting with n_i and ending with n_j such that consecutive vertices are adjacent. If there is a path between any two vertices of G , then G is connected. In this paper, we always assume that the graph is simple graph, which means that there is no self-loops and each element of E is unique. An oriented graph is a graph together with a particular orientation, where the orientation of a graph G is the assignment of a direction to each edge, so edge $(n_i; n_j)$ is an directed edge (arc) from n_i to n_j . The incidence matrix B of an oriented graph G is the $m \times n$ matrix with rows and columns indexed by the vertices and edges of G , respectively, such that the ij -entry is equal to 1 if edge j is ending on vertex n_i , -1 if edge j is beginning with vertex n_i , and 0 otherwise. Define the Laplacian matrix of G as $L(G) = BB^T$: $L(G)$ is always positive semidefinite. Moreover, for a connected graph, $L(G)$ has a single zero eigenvalue, and the associated right eigenvector is $\mathbf{1}_m$.

A directed graph D consists of a vertex set $V = \{n_1; n_2; \dots; n_m\}$ and an arc set $E = \{(n_i; n_j) : n_i, n_j \in V\}$, where an arc, or directed edge, is an ordered pair of distinct vertices of V . In this paper, we always assume that $n_i \neq n_j$, meaning that there is no self-loops, and assume that each element of E is unique. Let $D = (V; E; A)$ be a weighted directed graph. $A = [a_{ij}]$ is the weighted adjacency matrix, where a_{ij} is the weight of arc $(n_i; n_j)$, $a_{ij} \geq 0$ for all $i, j \in I = \{1; 2; \dots; m\}$; $a_{ii} = 0$ for all $i \in I$. The set of neighbors of vertex n_i is defined as $N_i = \{j \in I : a_{ij} > 0\}$. The in-degree and out-degree of vertex n_i are, respectively, defined as

$$\deg_{\text{in}}(n_i) = \sum_{j=1}^m a_{ji}; \quad \deg_{\text{out}}(n_i) = \sum_{j=1}^m a_{ij}.$$

The weighted graph D Laplacian matrix is defined as $L(D) = D - A$, where D is the degree matrix of D which is a diagonal matrix and its i th diagonal element is $d_{ii} = \deg_{\text{out}}(n_i)$.

By definition, $\lambda = 0$ is an eigenvalue of the Laplacian matrix $L(D)$ and $\mathbf{1}_m$ is its associated right eigenvector. A path of length r from n_0 to n_r in a directed graph is a sequence of $r + 1$ distinct vertices starting with n_0 and ending with n_r such that (n_{k-1}, n_k) is an arc of D for $k = 1; \dots, r$. A weak path is a sequence of $r + 1$ distinct vertices such that for $k = 1; \dots, r$, either (n_{k-1}, n_k) or (n_k, n_{k-1}) is an arc. A directed graph is strongly connected if any two vertices can be joined by a path and is weakly connected if any two vertices can be joined by a weak path.

References

- [1] N.E. Leonard and E. Fierelli, "Virtual leaders, artificial potentials and coordinated control of groups," in Proc. IEEE Conference on Decision and Control, Orlando, Florida USA, vol. 3, pp. 2968-2973, December 2001.
- [2] K. W. Arbuthnott and J. Lazarus, "Tendency-distance models of social cohesion in animal groups," J. Theoretical Biology, vol. 150, pp. 473-488, 1991.
- [3] I. Suzuki and M. Yamashita, "Distributed anonymous mobile robots: Formation of geometric patterns," SIAM J. Computing, vol. 28, no. 4, pp. 1347-1363, 1999.
- [4] J.H. Reif and H. Wang, "Social potential fields: A distributed behavioral control for autonomous robots," Robotics and Autonomous Systems, vol. 27, no. 3, pp. 171-194, May 1999.
- [5] F. Giulietti, L. Pollini, and M. Innocenti, "Autonomous formation flight," IEEE Control Systems Magazine, vol. 20, no. 6, pp. 34-44, December 2000.
- [6] E. Rimon and D.E. Koditschek, "Exact robot navigation using artificial potential functions," IEEE Transactions on Robotics and Automation, vol. 8, no. 5, pp. 501-518, October 1992.
- [7] R. Bachmayer and N.E. Leonard, "Vehicle networks for gradient descent in a sampled environment," in Proc. IEEE Conference on Decision and Control, Las Vegas, Nevada USA, vol. 1, pp. 112-117, December 2002.
- [8] R. Olfati-Saber and R.M. Murray, "Consensus problems in networks of agents with switching topology and time-delays," IEEE Transactions on Automatic Control, vol. 49, no. 9, pp. 1520-1533, September 2004.
- [9] C.W. Reynolds, "Flocks, herds, and schools: A distributed behavioral model," Computer Graphics (ACM SIGGRAPH '87 Conference Proceedings), vol. 21, no. 4, pp. 25-34, July 1987.
- [10] T. Vicsek, A. Czirak, E. Ben-Jacob, I. Cohen, and O. Shochet, "Novel type of phase transition in a system of self-driven particles," Physical Review Letters, vol. 75, no. 6, pp. 1226-1229, August 1995.
- [11] A. Jadbabaie, J. Lin, and A.S. Morse, "Coordination of groups of mobile autonomous agents using nearest neighbor rules," IEEE Transactions on Automatic Control, vol. 48, no. 6, pp. 988-1001, June 2003.

- [12] A . V . Savkin, \Coordinated collective motion of autonomous mobile robots: Analysis of Vicsek's model," IEEE Transactions on Automatic Control, vol. 49, no. 6, pp. 981{983, June 2004.
- [13] H . G . Tanner, A . Jadbabaie, and G . J. Pappas, \Stable flocking of mobile agents, Part I: Fixed topology," in Proc. IEEE Conference on Decision and Control, Maui, Hawaii USA, vol. 2, pp. 2010{2015, December 2003.
- [14] H . G . Tanner, A . Jadbabaie, and G . J. Pappas, \Stable flocking of mobile agents, Part II: Dynamic topology," in Proc. IEEE Conference on Decision and Control, Maui, Hawaii USA, vol. 2, pp. 2016{2021, December 2003.
- [15] H . Shi, L. Wang, T . Chu, and W . Zhang, \Coordination of a group of mobile autonomous agents," International Conference on Advances in Intelligent Systems| Theory and Applications, Luxembourg, November 2004.
- [16] V . Gazi and K . M . Passino, \Stability analysis of swarms," IEEE Transactions on Automatic Control, vol. 48, no. 4, pp. 692{697, April 2003.
- [17] V . Gazi and K . M . Passino, \Stability analysis of social foraging swarms," IEEE Transactions on Systems, Man and Cybernetics| Part B : Cybernetics, vol. 34, no. 1, pp. 539{557, February 2004.
- [18] Y . Liu, K . M . Passino, and M . Polycarpou, \Stability analysis of m -dimensional asynchronous swarms with a fixed communication topology," IEEE Transactions on Automatic Control, vol. 48, no. 1, pp. 76{95, January 2003.
- [19] Y . F . Liu and K . M . Passino, \Stable social foraging swarms in a noisy environment," IEEE Transactions on Automatic Control, vol. 49, no. 1, pp. 30{44, January 2004.
- [20] L. Wang, H . Shi, T . Chu, W . Zhang and L . Zhang, \Aggregation of foraging swarms," Lecture Notes in Artificial Intelligence, vol. 3339, pp. 766{777, Springer-Verlag, 2004.
- [21] T . Chu, L. Wang, and S. Mu, \Collective behavior analysis of an anisotropic swarm model," in Proc. of the 16th International Symposium on Mathematical Theory of Networks and Systems, Leuven, Belgium, pp. 1{14, July 2004.
- [22] H . Shi, L. Wang, and T . Chu, \Swarming behavior of multi-agent systems," in Proc. of the 23rd Chinese Control Conference, Wuxi, China, pp. 1027{1031, August, 2004.
- [23] B . Liu, T . Chu, L. Wang, and F . Hao, \Self-organization in a group of mobile autonomous agents," in Proc. of the 23rd Chinese Control Conference, Wuxi, China, pp. 45{49, August, 2004.
- [24] B . Liu, T . Chu, L. Wang., and Z . Wang, \Swarm dynamics of a group of mobile autonomous agents," Chinese Physics Letters, vol. 22, no. 1, pp. 254-257, 2005.
- [25] R . A . Horn and C . R . Johnson, Matrix Analysis. New York: Cambridge Univ. Press, 1985.
- [26] H . K . Khalil, Nonlinear Systems, Upper Saddle River, NJ: PrenticeHall, 1996.
- [27] C . Godsil and G . Royle, Algebraic Graph Theory. New York: SpringerVerlag, 2001.