

Fisher Information Matrix of General Stable Distributions Close to the Normal Distribution

By Muneya MATSUI*

Graduate School of Economics, University of Tokyo,
Hongo 7-3-1, Bunkyo-Ku, Tokyo 113-0033, Japan

Abstract

We investigate behavior of the Fisher information matrix of general stable distributions. Du-Mouchel (1975, 1983) proved that the Fisher information $I_{\alpha\alpha}$ of characteristic exponent α diverges to infinity as α approaches 2. Nagaev and Shkol'nik (1988) made more detailed analysis of $I_{\alpha\alpha}$ and derived asymptotic behavior of $I_{\alpha\alpha}$ diverging to infinity as α approaches 2 in the symmetric case. Extending their work in this paper we have obtained behavior of the Fisher information matrix of general stable distributions as α approaches 2 by detailed study of behavior of the corresponding density and its score functions. We clarify the limiting values of the 4×4 Fisher Information matrix with respect to the location μ , the scale σ , the characteristic exponent α and the skewness parameter β .

Keywords

General stable distributions, information matrix, normal distribution, score functions.

1 Notations and preliminary results

The family of stable distributions has enjoyed great interest of researchers in many fields e.g., mathematics, physics, cosmology and even economics. These applications are summarized in Uchaikin and Zolotarev (1999). In statistical inference estimation of stable parameters has been of great interest. In recent years, maximum likelihood estimation of stable distributions has become feasible (see Brorsen and Yang (1990), Nolan (2001) or Matsui and Takemura (2004)). Even in time series models like GARCH using general stable distributions, maximum likelihood estimation is possible owing to recent development of global algorithm (see Liu and Brorsen (1995)). Since the Fisher information matrix gives useful criteria for the accuracy of estimation, it is indispensable to analyze that of general stable distributions. Near Gaussian distribution ($\alpha = 2$), the information of α , $I_{\alpha\alpha}$ diverges to ∞ and asymptotic behavior of $I_{\alpha\alpha}$ as $\alpha \uparrow 2$ is of great interest. Nagaev and Shkol'nik (1988) have solved this problem excellently for symmetric stable distributions. However for general stable distributions Nagaev and Shkol'nik (1988) stated "We note the problems under study are as yet unresolved for non-symmetric stable distributions." In this paper we investigate and obtain asymptotic behavior of $I_{\theta\theta}$, $\theta = \mu, \sigma, \alpha, \beta$, as $\alpha \uparrow 2$ under general stable distributions.

Let

$$\Phi(t) = \Phi(t; \mu, \sigma, \alpha, \beta) = \exp \left(-|\sigma t|^\alpha \left\{ 1 + i\beta(\operatorname{sgn} t) \tan \left(\frac{\pi\alpha}{2} \right) (|\sigma t|^{1-\alpha} - 1) \right\} + i\mu t \right)$$

*mmatsui@grad.e.u-tokyo.ac.jp

denote the characteristic function of general stable distribution ($\alpha \neq 1$) with parameters

$$\theta = (\theta_1, \theta_2, \theta_3, \theta_4) = (\mu, \sigma, \alpha, \beta),$$

$$\mu \in \mathbf{R}, \sigma > 0, 0 < \alpha \leq 2, |\beta| \leq 1.$$

For the standard case $(\mu, \sigma) = (0, 1)$ we simply write the characteristic function as

$$(1.1) \quad \Phi(t; \alpha, \beta) = \exp \left(-|t|^\alpha \left\{ 1 + i\beta(\operatorname{sgn} t) \tan \left(\frac{\pi\alpha}{2} \right) (|t|^{1-\alpha} - 1) \right\} \right).$$

This is Zolotarev's (M) parameterization (see p.11 of Zolotarev (1986)). The corresponding density is written as $f(x; \mu, \sigma, \alpha, \beta)$ and $f(x; \alpha, \beta)$ in the standard case. Then

$$f(x; \mu, \sigma, \alpha, \beta) = \frac{1}{\sigma} f \left(\frac{x - \mu}{\sigma}; \alpha, \beta \right).$$

We also write the density of $N(0, 2)$ as

$$f(x; 2) = \frac{1}{2\sqrt{\pi}} \exp \left(-\frac{x^2}{4} \right).$$

For the rest of this paper, without loss of generality we consider the Fisher information only at the standard case $(\mu, \sigma) = (0, 1)$.

The first derivative of $f(x; \alpha, \beta)$ with respect to x is denoted by $f'(x; \alpha, \beta)$ and the partial derivative with respect to θ_i , i.e.,

$$\left. \frac{\partial f(x; \mu, \sigma, \alpha, \beta)}{\partial \theta_i} \right|_{(\mu, \sigma) = (0, 1)}$$

is denoted by $f_{\theta_i}(x; \alpha, \beta)$. Note that

$$(1.2) \quad f_\mu(x; \alpha, \beta) = -f'(x; \alpha, \beta),$$

$$(1.3) \quad f_\sigma(x; \alpha, \beta) = -f(x; \alpha, \beta) - x f'(x; \alpha, \beta).$$

The components of Fisher information matrix I are defined as follows.

$$I_{ij} = I_{\theta_i \theta_j} = \int_{-\infty}^{\infty} \frac{\partial f}{\partial \theta_i} \frac{\partial f}{\partial \theta_j} \frac{1}{f} dx.$$

Let

$$\zeta = -\beta \tan \left(\frac{\pi\alpha}{2} \right),$$

$$\varrho = \frac{2}{\pi\alpha} \arctan \left(\beta \tan \left(\frac{\pi\alpha}{2} \right) \right).$$

From Theorem 1 of Nolan (1997), which is a modified version of the formula (2.2.18) of Zolotarev (1986), the density $f(x; \alpha, \beta)$ for the case $\alpha \neq 1$ and $x \neq \zeta$ is written as

$$(1.4) \quad f(x; \alpha, \beta) = \frac{\alpha |x - \zeta|^{1/(\alpha-1)}}{2|\alpha - 1|} \int_{-\varrho^*}^1 A(\varphi; \alpha, \beta) \exp \left(-|x - \zeta|^{\alpha/(\alpha-1)} A(\varphi; \alpha, \beta) \right) d\varphi,$$

where $\varrho^* = \varrho \operatorname{sgn}(x - \zeta)$ and

$$(1.5) \quad A(\varphi; \alpha, \beta) = \left(\cos \left(\frac{\pi}{2} \alpha \varrho \right) \right)^{\frac{1}{\alpha-1}} \left(\frac{\cos \left(\frac{\pi}{2} \varphi \right)}{\sin \left\{ \frac{\pi}{2} \alpha (\varphi + \varrho) \right\}} \right)^{\frac{\alpha}{\alpha-1}} \frac{\cos \left[\frac{\pi}{2} \{ \alpha \varrho + (\alpha - 1) \varphi \} \right]}{\cos \left(\frac{\pi}{2} \varphi \right)}$$

is a positive function. Here we should note the relation $f(x; \alpha, \beta) = f(-x; \alpha, -\beta)$ for $x - \zeta < 0$, which will be used many times. As $\alpha \uparrow 2$, ζ and ϱ converge to 0 and at $\alpha = 2$ we obtain an unusual representation of Gaussian distribution $N(0, 2)$,

$$f(x; 2) = x \int_0^1 1 / \left(2 \sin \left(\frac{\pi}{2} \varphi \right) \right)^2 \exp \left(-x^2 / \left(2 \sin \left(\frac{\pi}{2} \varphi \right) \right)^2 \right) d\varphi, \quad x > 0.$$

The remainder of this paper consists of three sections. In Section 2 we derive behavior of $f'(x; \alpha, \beta)$, $f_\mu(x; \alpha, \beta)$ and $f_\sigma(x; \alpha)$ as $\alpha \uparrow 2$ by careful analysis of the formula (1.4). In Section 3 behavior of $f_\alpha(x; \alpha, \beta)$ and $f_\beta(x; \alpha, \beta)$ are obtained by utilizing the inversion formula. The information matrix of general stable distributions is given in Section 4.

2 Density and derivatives of general stable distributions close to the normal distribution

In this Section the density and derivatives of general stable distributions as $\alpha \uparrow 2$ and $x \rightarrow \infty$ are obtained. In the following we write

$$\Delta = 2 - \alpha.$$

As $\Delta \rightarrow 0$ we can show

$$(2.1) \quad f(x; \alpha, \beta) = f(x; 2) + O(\Delta)$$

uniformly in x by the finiteness of $f_\alpha(x; \alpha, \beta)$, which can be easily verified by the formula (3.2) in the proof of Lemma 3.1. However if we consider the ratio $f(x; 2)/f(x; \alpha, \beta)$ rather than the difference $f(x; 2) - f(x; \alpha, \beta)$ of the densities as $x \rightarrow \infty$, $f(x; 2)$ is much smaller than $f(x; \alpha, \beta)$ and we can not obtain an accurate approximation of $f(x; \alpha, \beta)$ without closer investigation of the remainder $O(\Delta)$. In the case of $x \rightarrow \infty$ we can utilize the asymptotic expansion

$$(2.2) \quad f(x; \alpha, \beta) = \Delta(1 + \beta)(x - \zeta)^{\Delta-3} + o(\Delta x^{-4})$$

from the expansion (2.5.4) on p.94 of Zolotarev (1986), Bergström (1953) or Section XVII.6 of Feller (1971). (2.5.4) of Zolotarev (1986) is in terms of (B) representation with characteristic function defined on p.12 of Zolotarev (1986). Hence as $x \rightarrow \infty$

$$f_B(x; \alpha, \beta) = \Delta(1 + \beta_B)x^{\Delta-3} + o(\Delta x^{-4}),$$

where f_B and β_B means the density or β of (B) representation. We obtain (2.2) by using the relation

$$f(x; \alpha, \beta) = \left(\cos \left(\frac{\pi}{2} \alpha \varrho \right) \right)^{\frac{1}{\alpha}} f_B \left(\left(\cos \left(\frac{\pi}{2} \alpha \varrho \right) \right)^{\frac{1}{\alpha}} (x - \zeta); \alpha, \beta_B \right)$$

by noting that $\cos(\frac{\pi}{2} \alpha \varrho) \rightarrow 1$ and $\beta_B = \beta + O(\Delta^2)$ as $\Delta \rightarrow 0$. Then the problem is which of the two approximations (2.1) and (2.2) is dominant for large x . We give an answer to this problem in the following two theorems, which are based on Theorem 1 of Nagaev and Shkol'nik (1988). We can also see the precise statement on p.129 of Uchaikin and Zolotarev (2003) concerning this problem.

Theorem 2.1 Let β , $-1 < \beta < 1$, be fixed and let $\beta^* = \beta \operatorname{sgn}(x - \zeta)$. Define

$$\begin{aligned} F_1(x; \alpha, \beta) &= f(x - \zeta; 2), \\ F_2(x; \alpha, \beta) &= (1 + \beta^*)\Delta(x - \zeta)^{\Delta-3}, \\ g(x; \alpha, \beta) &= F_1(x; \alpha, \beta) + F_2(x; \alpha, \beta). \end{aligned}$$

For an arbitrarily small $\epsilon > 0$ there exist Δ_0 and x_0 such that for all $\Delta < \Delta_0$ and $|x| > x_0$,

$$|f(x; \alpha, \beta)/g(x; \alpha, \beta) - 1| < \epsilon.$$

Furthermore, for an arbitrarily small constant $\delta > 0$,

$$g(x; \alpha, \beta) = \begin{cases} F_1(x; \alpha, \beta) (1 + o(\Delta^{\delta/2})) & \text{if } |x - \zeta| \leq (2 - \delta)(\log 1/\Delta)^{1/2}, \\ F_2(x; \alpha, \beta) (1 + o(\Delta^{\delta/2})) & \text{if } |x - \zeta| \geq (2 + \delta)(\log 1/\Delta)^{1/2}, \end{cases}$$

uniformly in $|x| > x_0$.

Theorem 2.2 Define

$$\begin{aligned} F'_1(x; \alpha, \beta) &= -\frac{x - \zeta}{2} f(x - \zeta; 2), \\ F'_2(x; \alpha, \beta) &= -3(1 + \beta^*)\Delta(x - \zeta)^{\Delta-4}, \\ g'(x; \alpha, \beta) &= F'_1(x; \alpha, \beta) + F'_2(x; \alpha, \beta). \end{aligned}$$

Under the same conditions and notations of Theorem 2.1, for an arbitrarily small $\epsilon > 0$ there exist Δ_0 and x_0 such that for all $\Delta < \Delta_0$ and $|x| > x_0$,

$$|f'(x; \alpha, \beta)/g'(x; \alpha, \beta) - 1| < \epsilon.$$

Furthermore, for an arbitrarily small constant $\delta > 0$,

$$g'(x; \alpha, \beta) = \begin{cases} F'_1(x; \alpha, \beta) (1 + o(\Delta^{\delta/2})) & \text{if } |x - \zeta| \leq (2 - \delta)(\log 1/\Delta)^{1/2}, \\ F'_2(x; \alpha, \beta) (1 + o(\Delta^{\delta/2})) & \text{if } |x - \zeta| \geq (2 + \delta)(\log 1/\Delta)^{1/2}, \end{cases}$$

uniformly in $|x| > x_0$.

We need the following 4 lemmas concerning behavior of $A(\varphi; \alpha, \beta)$ in (1.5) for $\varphi \doteq 1$ to prove Theorem 2.1 and Theorem 2.2, which correspond to the lemmas of Theorem 1 of Nagaev and Shkol'nik (1988). In the lemmas and the proofs of Theorems 2.1 and 2.2 ϵ and ϵ' denote arbitrarily small positive constants because in the end of the proofs of Theorems 2.1 and 2.2 we let ϵ and $\epsilon' \downarrow 0$ separately. From now on the Fisher information is evaluated at a fixed value of β , $|\beta| < 1$. For notational abbreviation we sometimes write $A(\varphi; \alpha, \beta)$ as $A(\varphi)$. We also write

$$\lambda = 1 - \varphi, \quad \varphi_\Delta = 1 - \Delta^{1/2-\epsilon}.$$

Lemma 2.1 As $\Delta \rightarrow 0$ and for $0 \leq \lambda \leq \Delta/\epsilon'$,

$$A(1 - \lambda) = \frac{(\lambda/\Delta)^{\frac{1}{1-\Delta}} (1 + \beta + \lambda/\Delta)}{(1 + \beta + 2\lambda/\Delta)^2} (1 + o(\Delta \log(1/\Delta)))$$

uniformly in λ .

Lemma 2.2 As $\Delta \rightarrow 0$

$$A(\varphi_\Delta) = \frac{1}{4} + \frac{\pi^2}{16} \Delta^{1-2\epsilon} + o(\Delta), \quad A'(\varphi_\Delta) = -\frac{\pi^2}{8} \Delta^{1/2-\epsilon} - \frac{(1+\beta)^2}{8} \Delta^{1/2+3\epsilon} + O(\Delta).$$

Lemma 2.3 As $\Delta \rightarrow 0$ and $\lambda \rightarrow 0$ such that $\lambda \geq \Delta^{1/2-\epsilon}$,

$$A(1-\lambda) = \frac{1}{4} + \frac{\pi^2}{16} \lambda^2 + o(\lambda^2), \quad A'(1-\lambda) = -\frac{\pi^2}{8} \lambda - \frac{\Delta^2(1-\lambda+\beta)^2}{8\lambda^3} + o(\lambda^2),$$

$$A''(1-\lambda) = \frac{\pi^2}{8} + \frac{\pi^2}{8} \lambda^2 - \frac{3}{8} \frac{\Delta^2(1-\lambda+\beta)^2}{\lambda^4} - \frac{3}{4} \frac{\Delta^2(1-\lambda+\beta)}{\lambda^3} \left(1 - \frac{\Delta(1-\lambda+\beta)^2}{\lambda^2}\right) + o(\lambda^2).$$

Note that $\Delta/\epsilon' \leq \Delta^{1/2-\epsilon} \leq \lambda$ implies $\Delta = o(\lambda^2)$ and we have to consider terms like Δ^2/λ^4 or Δ^2/λ^3 in Lemma 2.3.

Lemma 2.4 If Δ is sufficiently small, $A(\varphi)$ is a monotonically decreasing function on $(-\varrho^*, 1)$.

The proofs of the lemmas are given in Appendix.

In the following $\bar{\epsilon} > 0$ is an appropriate positive constant.

Proof of Theorem 2.1 We prove the assertion in the case $x - \zeta > 0$ first. As in the proof of Theorem 1 of Nagaev and Shkol'nik (1988), let $\tau, \eta > 0$ be arbitrarily small numbers. We denote $z = (x - \zeta)^{\alpha/(\alpha-1)}$ and $\varphi_0 = \varphi_\Delta - z^{-1/2+\tau}$. Then we divide the integral

$$H = \int_{-\varrho^*}^1 A(\varphi; \alpha, \beta) \exp(-zA(\varphi; \alpha, \beta)) d\varphi$$

into six subintegrals,

$$H = \sum_{k=1}^6 H_k,$$

where each H_k corresponds to the integration of H for the k -th interval of $[-\varrho^*, 1-\eta)$, $[1-\eta, \varphi_0)$, $[\varphi_0, \varphi_\Delta)$, $[\varphi_\Delta, 1-\Delta/\epsilon')$, $[1-\Delta/\epsilon', 1-\Delta\epsilon')$ and $[1-\Delta\epsilon', 1]$. First we calculate H_i 's utilizing Lemmas 2.1-2.4 for fixed ϵ and ϵ' . Later we see that H_3 and H_6 dominate the others terms. We also let ϵ and $\epsilon' \downarrow 0$ in the end.

Calculation of H_1 :

From Lemma 2.2, $A(\varphi_\Delta) > \frac{1}{4}$. By Lemma 2.4 and $1-\eta < \varphi_\Delta$, for sufficiently small Δ there exists a constant $\rho_1 \in (0, 1)$ such that $\rho_1 A(1-\eta) > \frac{1}{4}$. Then for a constant $\gamma > \frac{1}{4}$ we easily find

$$(2.3) \quad \begin{aligned} H_1 &= \frac{1}{z(1-\rho_1)} \int_{-\varrho^*}^{1-\eta} z(1-\rho_1) A(\varphi) \exp\{-z\rho_1 A(\varphi) - z(1-\rho_1) A(\varphi)\} d\varphi \\ &\leq \frac{1}{z(1-\rho_1)} \exp(-z\rho_1 A(1-\eta)) \\ &= O(\exp(-\gamma z)/z). \end{aligned}$$

Calculation of H_2 :

From Lemma 2.4

$$\begin{aligned} H_2 &= \int_{1-\eta}^{\varphi_0} A(\varphi) \exp(-zA(\varphi)) d\varphi \\ &\leq \eta A(1-\eta) \exp(-zA(\varphi_0)) \\ &= O(\exp(-zA(\varphi_0))). \end{aligned}$$

Then we can write

$$A(\varphi_0) = A(\varphi_\Delta) + A'(\varphi_\Delta)(\varphi_0 - \varphi_\Delta) + \frac{1}{2}A''(\xi)(\varphi_0 - \varphi_\Delta)^2,$$

where $\varphi_0 \leq \xi \leq \varphi_\Delta$. By Lemma 2.2 and Lemma 2.3 we can write

$$\begin{aligned} zA(\varphi_\Delta) &= \frac{z}{4} + R_1(\Delta, z, \epsilon), \\ zA'(\varphi_\Delta)(\varphi_0 - \varphi_\Delta) &= R_2(\Delta, z, \epsilon), \\ zA''(\xi)(\varphi_0 - \varphi_\Delta)^2 &\leq \bar{c}z^{2\tau}, \end{aligned}$$

where

$$\begin{aligned} R_1(\Delta, z, \epsilon) &= O(z\Delta^{1-2\epsilon}) > 0, \\ R_2(\Delta, z, \epsilon) &= O(\Delta^{1/2-\epsilon}z^{1/2+\tau}) > 0. \end{aligned}$$

Here we use functions $R_i(\Delta, z, \epsilon)$, $i = 1, 2$, for convenience in comparing H_2 with H_3 . Then

$$(2.4) \quad H_2 = O\left(\exp\left(-\frac{z}{4} - R_1(\Delta, z, \epsilon) - R_2(\Delta, z, \epsilon) - \bar{c}z^{2\tau}\right)\right).$$

Calculation of H_3 :

We show the detailed calculation of H_3 because this is the dominant term in H as $\Delta \rightarrow 0$ and $x \rightarrow \infty$.

We have for $\xi \in [\varphi_0, \varphi_\Delta]$,

$$H_3 = \exp(-zA(\varphi_\Delta)) \int_{\varphi_0}^{\varphi_\Delta} A(\varphi) \exp\left(-zA'(\varphi_\Delta)(\varphi - \varphi_\Delta) - \frac{z}{2}A''(\xi)(\varphi - \varphi_\Delta)^2\right) d\varphi.$$

Utilizing Lemmas 2.2 and 2.3, for some constants $\rho_1, \rho_2 \in (0, 1)$ we obtain

$$\begin{aligned} A(\varphi) &= \frac{1}{4} + \frac{\pi^2}{16}\lambda^2 + o(\lambda^2), \\ -zA(\varphi_\Delta) &= -\frac{z}{4} - \frac{\pi^2}{16}\Delta^{1-2\epsilon}z + o(\Delta z), \\ -zA'(\varphi_\Delta)(\varphi - \varphi_\Delta) &= -\left(\frac{\pi^2}{8}\Delta^{1/2-\epsilon} + \frac{1}{8}(1+\beta)^2\Delta^{1/2+3\epsilon}\right)\rho_1 z^{1/2+\tau} + O(\Delta z^{1/2+\tau}), \\ -\frac{z}{2}A''(\xi)(\varphi - \varphi_\Delta)^2 &= -\frac{z}{2}\frac{\pi^2}{8}(\varphi - \varphi_0)^2 \\ &\quad -\left(\frac{\pi^2}{8}\lambda^2 - \frac{3}{8}\frac{\Delta^2(1-\lambda+\beta)^2}{\lambda^4} - \frac{3}{4}\frac{\Delta^2(1-\beta+\lambda)}{\lambda^3}\left(1 - \frac{\Delta(1-\lambda+\beta)^2}{\lambda^2}\right) + o(\lambda^2)\right)\frac{\rho_2}{2}z^{2\tau}. \end{aligned}$$

Summarizing the main terms, we obtain

$$\begin{aligned}
A(\varphi) &= \frac{1}{4} + O(\lambda^2), \\
-zA(\varphi_\Delta) &= -\frac{z}{4} - R_1(\Delta, z, \epsilon), \\
-zA'(\varphi_\Delta)(\varphi - \varphi_\Delta) &= -R_3(\Delta, z, \epsilon), \\
-\frac{z}{2}A''(\xi)(\varphi - \varphi_\Delta)^2 &= -\frac{z}{2}\frac{\pi^2}{8}(\varphi - \varphi_0)^2 + O(\lambda^2 z^{2\tau}) + O(\Delta^2/\lambda^4 z^{2\tau}) \\
&= -\frac{z}{2}\frac{\pi^2}{8}(\varphi - \varphi_0)^2 + O(z^{-1+4\tau}) + O(\Delta^{4\epsilon} z^{2\tau}),
\end{aligned}$$

where

$$0 < R_3(\Delta, z, \epsilon) = O(\Delta^{1/2-\epsilon} z^{1/2+\tau}) \leq R_2(\Delta, z, \epsilon).$$

Note $\varphi_0 \leq \varphi \leq \varphi_\Delta \Leftrightarrow \Delta^{1/2-\epsilon} \leq \lambda \leq \Delta^{1/2-\epsilon} + z^{-1/2+\tau}$. Then we obtain

$$H_3 = R(\lambda, \Delta, z, \epsilon) \frac{1}{4} \exp\left(-\frac{z}{4}\right) \int_{\varphi_0}^{\varphi_\Delta} \exp\left(-\frac{z}{2}\frac{\pi^2}{8}(\varphi - \varphi_\Delta)^2\right) d\varphi,$$

where

$$R(\lambda, \Delta, z, \epsilon) = (1 + O(\lambda^2)) \exp(-R_1(\Delta, z, \epsilon) - R_3(\Delta, z, \epsilon) + O(z^{-1+4\tau}) + O(\Delta^{4\epsilon} z^{2\tau})).$$

Here the inequality

$$\int_x^\infty e^{-y^2/2} dy \leq \frac{1}{x} e^{-x^2/2}, \quad x > 0$$

gives

$$\begin{aligned}
\int_{\varphi_0}^{\varphi_\Delta} \exp\left(-\frac{z}{2}\frac{\pi^2}{8}(\varphi - \varphi_0)^2\right) d\varphi &= \frac{1}{\sqrt{z}} \int_0^\infty \exp\left(-\frac{z}{2}\frac{\pi^2}{8}\varphi^2\right) d\varphi - \frac{1}{\sqrt{z}} \int_{z^\tau}^\infty \exp\left(-\frac{z}{2}\frac{\pi^2}{8}\varphi^2\right) d\varphi \\
&= \frac{2}{\sqrt{\pi z}} + O\left(\frac{1}{z^{1/2+\tau}} \exp\left(\frac{-z^{2\tau}}{2}\right)\right).
\end{aligned}$$

Then we obtain

$$(2.5) \quad H_3 = \exp\left(-\frac{z}{4} - R_1(\Delta, z, \epsilon) - R_3(\Delta, z, \epsilon)\right) \frac{1}{2\sqrt{\pi z}} \exp(O(z^{-1+4\tau}) + O(\Delta^{4\epsilon} z^{2\tau})) (1 + o(1)).$$

Written in this form, H_3 can be easily compared with H_2 , i.e., the formula (2.4). Further we will see later that H_3 will be dominant if $z \leq (4 - \delta) \log 1/\Delta$ for an arbitrarily small constant $\delta > 0$. Then we only have to consider the convergence of H_3 for $z \leq O(\log 1/\Delta)$. For $z \leq O(\log 1/\Delta)$, $R_1(\Delta, z, \epsilon) \downarrow 0$, $R_2(\Delta, z, \epsilon) \downarrow 0$ and $R_3(\Delta, z, \epsilon) \downarrow 0$ as $\Delta \rightarrow 0$. Finally we obtain

$$(2.6) \quad H_3 = \frac{1}{2\sqrt{\pi z}} \exp\left(-\frac{z}{4}\right) (1 + o(1)) \quad \text{if } z \leq O(\log 1/\Delta).$$

Calculation of H_4 :

From Lemma 2.4

$$\begin{aligned}
H_4 &\leq \int_{\varphi_\Delta}^{1-\Delta/\epsilon'} A(\varphi_\Delta) \exp(-zA(1 - \Delta/\epsilon')) d\varphi \\
&\leq \Delta^{1/2-\epsilon} A(\varphi_\Delta) \exp(-zA(1 - \Delta/\epsilon')).
\end{aligned}$$

Here $\lambda/\Delta \leq 1/\epsilon'$. Then by Lemma 2.1 and 2.4 as $\Delta \rightarrow 0$

$$A(1 - \Delta/\epsilon') = \frac{\epsilon'^{\frac{\Delta}{\Delta-1}} \{\epsilon'(1 + \beta) + 1\}}{\{\epsilon'(1 + \beta) + 2\}^2} = \frac{1}{4}(1 - w(\epsilon')),$$

where $w(\cdot)$ is a nonnegative function such as

$$\lim_{t \rightarrow 0} w(t) = 0.$$

Finally we obtain

$$(2.7) \quad H_4 \leq O\left(\Delta^{1/2-\epsilon} \exp\left(-\frac{z}{4}(1 - w(\epsilon'))\right)\right).$$

For the comparison of H_4 with H_3 and H_6 , which is needed in later argument, we state some properties of H_4 . For a given small number $\delta > 0$, there exists ϵ' such that $\delta > w(\epsilon')$. Then for $z \leq 4 \log 1/\Delta$,

$$\Delta^\delta = \exp\left(-\frac{\delta}{4} \times 4 \log 1/\Delta\right) \leq \exp\left(-\frac{\delta}{4}z\right).$$

Consequently

$$(2.8) \quad H_4 \leq O\left(\Delta^{1/2-\delta-\epsilon} \exp\left(-\frac{z}{4}(1 + \delta - w(\epsilon'))\right)\right).$$

In this representation H_4 can be easily compared with H_3 . For $z \geq 4 \log 1/\Delta$

$$(2.9) \quad H_4 \leq O(\Delta \exp(-\bar{c}z)).$$

In this representation H_4 can be easily compared with H_6 .

Calculation of H_5 :

From Lemma 2.1 for $\epsilon' \leq \lambda/\Delta \leq 1/\epsilon'$, $A(\varphi)$ is bounded. Then

$$(2.10) \quad \begin{aligned} H_5 &= \int_{1-\Delta/\epsilon'}^{1-\Delta\epsilon'} A(\varphi) \exp(-zA(\varphi)) d\varphi \\ &\leq (\Delta/\epsilon' - \Delta\epsilon') \sup_{\varphi \in (1-\Delta/\epsilon', 1-\Delta\epsilon')} A(\varphi) \exp(-zA(\varphi)) \\ &\leq O(\Delta \exp(-\bar{c}z)). \end{aligned}$$

Calculation of H_6 :

From Lemma 2.1 for $\lambda/\Delta \leq \epsilon'$

$$\begin{aligned} A(1 - \lambda; \alpha, \beta) &= \frac{(\lambda/\Delta)^{1/(1-\Delta)}}{1 + \beta} (1 + O(\epsilon') + O(\Delta \log(1/\Delta))) \\ &= \frac{(\lambda/\Delta)^{1/(1-\Delta)}}{1 + \beta} (1 + R_4(\Delta, \epsilon')), \end{aligned}$$

where $R_4(\Delta, \epsilon') = O(\epsilon') + O(\Delta \log(1/\Delta))$. Substituting the above results into H_6 and replacing $\varphi = 1 - \lambda$ by λ , we obtain

$$H_6 = \left(\frac{1 + R_4}{1 + \beta}\right) \int_0^{\Delta\epsilon'} (\lambda/\Delta)^{1/(1-\Delta)} \exp\left(-z \left(\frac{1 + R_4}{1 + \beta}\right) (\lambda/\Delta)^{1/(1-\Delta)}\right) d\lambda.$$

Furthermore, replacing

$$\lambda = \Delta \left(\frac{1 + \beta}{1 + R_4} \frac{x}{z} \right)^{1-\Delta}$$

by x and defining

$$g(z) = \epsilon'^{1/(1-\Delta)} \left(\frac{1 + R_4}{1 + \beta} \right) z,$$

H_6 is written as

$$H_6 = \frac{\Delta(1 + \beta)^{1-\Delta}}{z^{2-\Delta}} \int_0^{g(z)} x \exp(-x) x^{-\Delta} dx \times \{(1 + R_4)^{(\Delta-1)}(1 - \Delta)\}.$$

Then the integration of the right side of the equation is evaluated as

$$\begin{aligned} \int_0^{g(z)} x \exp(-x) dx + O(\Delta) &= [-xe^{-x} - e^{-x}]_0^{g(z)} + O(\Delta) \\ &= 1 + O(ge^{-g}) + O(\Delta). \end{aligned}$$

Here for arbitrarily small constant ϵ' and any $\Delta \rightarrow 0$, $g(z) \rightarrow \infty$ as $z \rightarrow \infty$. Therefore

$$(2.11) \quad H_6 = \Delta(1 + \beta)z^{\Delta-2}(1 + o(1)).$$

For fixed ϵ and ϵ' the formulas (2.3), (2.4), (2.5), (2.7), (2.10) and (2.11) are proved.

Comparing (2.3), (2.4), (2.5), (2.7), (2.10) and (2.11), the order of $H_i, i = 1, \dots, 6$, is summarized as follows:

$$\max(H_1, H_2, H_4, H_5) = o(\max(H_3, H_6)).$$

The relative dominance of H_3 (2.6) and H_6 (2.11) depends on the value of z . For an arbitrarily small $\delta > 0$,

$$\begin{aligned} H_6 &= o(H_3) \quad \text{if } z \leq (4 - \delta) \log 1/\Delta, \\ H_3 &= o(H_6) \quad \text{if } z \geq (4 + \delta) \log 1/\Delta. \end{aligned}$$

These are easily confirmed if we substitute $(4 \pm \delta) \log 1/\Delta$ into z .

Now for proving the uniformity in Δ and z we let $\epsilon, \epsilon' \downarrow 0$ depending on Δ and z . We first let $\epsilon \downarrow 0$. We can take $\epsilon = (\log 1/\Delta)^{\delta-1}$ for a constant $\delta \in (0, 1)$. Concerning Lemmas 2.1-2.4, we must consider the relations of $\Delta^{2\epsilon} \geq \Delta/\lambda^2$ in Lemma 2.3. In the other lemmas the results do not change. Substituting $\epsilon = (\log 1/\Delta)^{\delta-1}$ into $\Delta^{2\epsilon}$ and taking logarithm of $\Delta^{2\epsilon}$

$$\log \Delta^{2\epsilon} = -2(\log 1/\Delta)^\delta \rightarrow -\infty.$$

Hence $\Delta = o(\lambda^2)$ and the conclusion of Lemma 2.3 is not changed. Concerning the calculations of H_i , the order of dominance is not affected by the limiting operation $\epsilon \downarrow 0$. For the comparison of H_3 and H_6 we may substitute $\epsilon = (\log 1/\Delta)^{\delta-1}$ and $z = (4 \pm \delta) \log 1/\Delta$ into (2.5) and (2.11). The other comparisons are easy. Then we have only to confirm that H_3 satisfies (2.6) because the calculation of H_6 is not affected by ϵ . If we substitute $\epsilon = (\log 1/\Delta)^{\delta-1}$ into (2.5), we obtain (2.6) since

$$z\Delta^{1-2\epsilon} \rightarrow 0, \quad \Delta^{1/2-\epsilon} z^{1/2+\tau} \rightarrow 0, \quad \Delta^{4\epsilon} z^{2\tau} \rightarrow 0, \quad \text{for } z \leq 4 \log 1/\Delta$$

imply

$$R_1(\Delta, z, \epsilon) \downarrow 0, \quad R_2(\Delta, z, \epsilon) \downarrow 0, \quad R_3(\Delta, z, \epsilon) \downarrow 0, \quad O(\Delta^{4\epsilon} z^{2\tau}) \rightarrow 0, \quad \text{for } z \leq 4 \log 1/\Delta.$$

Next we consider $\epsilon' \downarrow 0$ depending on Δ . We may consider $\epsilon' = (\log 1/\Delta)^{\delta'-1}$ for an arbitrarily small $\delta' \in (0, 1)$. Concerning Lemma 2.1 ϵ' must satisfy $\Delta/\epsilon' \leq \Delta^{1/2-\epsilon}$. This is easily confirmed. ϵ' does not affect the other lemmas. Concerning the calculations H'_i s, we must consider H_4 , H_5 and H_6 . The other H'_i s are not affected by ϵ' . From (2.7) and the definition of $w(\epsilon')$, the calculation of H_4 concerning the order, i.e., formulas (2.8) and (2.9) are not changed. We investigate H_5 in detail. If $[1 - \Delta/\epsilon, 1 - \Delta\epsilon']$, then $\epsilon' \leq \lambda/\Delta \leq 1/\epsilon'$ and $A(\varphi)$ of Lemma 2.1 satisfies

$$\frac{1 + o(1)}{1 + \beta} \epsilon'^{1/(1-\Delta)} \leq A(\varphi).$$

Hence

$$\begin{aligned} H_5 &\leq (\Delta/\epsilon' - \Delta\epsilon') \times \sup_{\varphi \in (1-\Delta/\epsilon, 1-\Delta\epsilon')} A(\varphi) \exp(-zA(\varphi)) \\ &\leq O\left((\Delta/\epsilon' - \Delta\epsilon') \times \epsilon'^{\frac{1}{1-\Delta}} \exp\left(-\bar{c}z\epsilon'^{\frac{1}{1-\Delta}}\right)\right) \\ &\leq O\left(\Delta \exp\left(-\bar{c}z\epsilon'^{1/(1-\Delta)}\right)\right). \end{aligned}$$

Then for $z \leq (4 - 4\delta'') \log 1/\Delta$ and for an arbitrarily small positive constant δ'' ,

$$H_5 \leq O(\Delta) \leq O\left(\Delta^{\delta''} \exp\left(-\frac{z}{4}\right)\right) = o(H_3).$$

For $z \geq (4 - 4\delta'') \log 1/\Delta$

$$H_5 = O(\Delta \exp(-\bar{c}z^{\delta'})) = o(H_6).$$

Hence the order and dominant terms of H_i 's are not changed. For the calculation of H_6 , $g(z)$ must go to ∞ as $z \rightarrow \infty$. Considering $z \geq (4 + \delta) \log 1/\Delta$, we can easily confirm this condition. Thus we can let $\epsilon' \downarrow 0$ depending on Δ . Therefore convergence of $H_i, i = 1, \dots, 6$, only depends on Δ and z and is faster as $\Delta \rightarrow 0$ and $z \rightarrow \infty$.

Finally, substituting $z = (x - \zeta)^{\alpha/(\alpha-1)}$ into H_3 and H_6 and multiplying these by $\alpha/(2(\alpha - 1))(x - \zeta)^{1/(\alpha-1)}$, we obtain the result for $x - \zeta > 0$. For $x - \zeta < 0$ we utilize the relations $f(x; \alpha, \beta) = f(-x; \alpha, -\beta)$ of general stable distributions. \square

Proof of Theorem 2.2 From (1.4) the derivative of the density is calculated as follows.

$$(2.12) \quad f'(x; \alpha, \beta) = \frac{1}{\alpha - 1} (x - \zeta)^{-1} f(x; \alpha, \beta) - \frac{\alpha^2}{2(\alpha - 1)^2} (x - \zeta)^{2/(\alpha-1)} \times H',$$

where

$$H' = \int_{-\varrho^*}^1 A^2(\varphi; \alpha, \beta) \exp\left(-(x - \zeta)^{\alpha/(\alpha-1)} A(\varphi; \alpha, \beta)\right) d\varphi.$$

Since behavior of $f(x; \alpha, \beta)$ was obtained in Theorem 2.1, we evaluate the integration of the second term of (2.12). As in Theorem 2.1 we divide integral H' into subintegrals,

$$H' = \sum_{k=1}^6 H'_k,$$

where each H'_k corresponds to the k -th interval of $[0, 1 - \mu)$, $[1 - \mu, \varphi_0)$, $[\varphi_0, \varphi_\Delta)$, $[\varphi_\Delta, 1 - \Delta/\epsilon')$, $[1 - \Delta/\epsilon', 1 - \Delta\epsilon')$ and $[1 - \Delta\epsilon', 1]$. From Lemmas 2.1, 2.2 and 2.3, for $\lambda \leq \Delta/\epsilon'$

$$(2.13) \quad A^2(1 - \lambda) = \frac{(\lambda/\Delta)^{\frac{2}{1-\Delta}}(1 + \beta + \lambda/\Delta)^2}{(1 + \beta + 2\lambda/\Delta)^4}(1 + O(\Delta \log(1/\Delta))),$$

$$A^2(\varphi_\Delta) = \frac{1}{16} + \frac{\pi^2}{32}\Delta^{1-2\epsilon} + o(\Delta),$$

and for $\varphi \leq \varphi_\Delta \Leftrightarrow \Delta^{1/2-\epsilon} \leq \lambda$

$$A^2(1 - \lambda) = \frac{1}{16} + \frac{\pi^2}{32}\lambda^2 + o(\lambda^2).$$

Considering $A^2(\varphi, \Delta)$ term of the integrand, we easily find the following results on H'_i similar to $H_i, i = 1, \dots, 5$, in the proof of Theorem 2.1.

$$(2.14) \quad H'_1 \leq O(\exp(-\gamma z)/z^2), \quad \text{for } \gamma > \frac{1}{4},$$

$$(2.15) \quad H'_2 \leq O\left(\exp\left(-\frac{z}{4} + O(z\Delta^{1-2\epsilon}) + O(z^{1/2+\tau}\Delta^{1/2-\epsilon}) - \bar{c}z^{2\tau}\right)\right),$$

$$(2.16) \quad H'_3 = \frac{1}{8\sqrt{z\pi}} \exp\left(-\frac{z}{4}\right)(1 + o(1)),$$

$$(2.17) \quad H'_4 \leq O\left(\Delta^{1/2-\epsilon} \exp\left(-\frac{z}{4}(1 - w(\epsilon'))\right)\right),$$

$$(2.18) \quad H'_5 \leq O(\Delta \exp(-\bar{c}z)).$$

Later we see H'_3 and H'_6 dominate the others terms as H_3 and H_6 dominate the other H_i 's in Theorem 2.1. Since the calculation H'_6 is somewhat different, we treat H'_6 separately. From (2.13), for $\lambda/\Delta \leq \epsilon'$

$$A^2(1 - \lambda; \alpha, \beta) = \frac{(\lambda/\Delta)^{\frac{2}{1-\Delta}}}{(1 + \beta)^2} (1 + R_4(\Delta, \epsilon'))^2,$$

where $R_4(\Delta, \epsilon')$ is defined in the proof of Theorem 2.1. Substituting the above result into H'_6 and replacing $\varphi = 1 - \lambda$ by λ , we obtain

$$H'_6 = \left(\frac{1 + R_4}{1 + \beta}\right)^2 \int_0^{\Delta\epsilon'} (\lambda/\Delta)^{2/(1-\Delta)} \exp\left(-z \left(\frac{1 + R_4}{1 + \beta}\right) (\lambda/\Delta)^{1/(1-\Delta)}\right) d\lambda.$$

Furthermore, replacing

$$\lambda = \Delta \left(\frac{1 + \beta}{1 + R_4} \frac{x}{z}\right)^{1-\Delta}$$

by x and defining

$$g(z) = \epsilon'^{1/(1-\Delta)} \left(\frac{1 + R_4}{1 + \beta}\right) z,$$

H'_6 is written as

$$H'_6 = \frac{\Delta(1 + \beta)^{1-\Delta}}{z^{3-\Delta}} \int_0^{g(z)} x^2 \exp(-x) x^{-\Delta} dx \times \{(1 + R_4)^{(\Delta-1)}(1 - \Delta)\}.$$

Then the integration of the right side of the equation is evaluated as

$$\begin{aligned} \int_0^{g(z)} x^2 \exp(-x) dx + O(\Delta) &= [-x^2 e^{-x} - 2x e^{-x} - 2e^{-x}]_0^{g(z)} + O(\Delta) \\ &= 2 + O(g^2 e^{-g}) + O(\Delta). \end{aligned}$$

Here for arbitrarily small constant ϵ' and any $\Delta \rightarrow 0$, $g(z) \rightarrow \infty$ as $z \rightarrow \infty$. Therefore

$$(2.19) \quad H'_6 = 2(1 + \beta)\Delta z^{\Delta-3}(1 + o(1)).$$

Hence for fixed ϵ and ϵ' formulas (2.14)-(2.19) are proved.

From (2.14)-(2.19)

$$\max(H'_1, H'_2, H'_4, H'_5) = o(\max(H'_3, H'_6)).$$

The relative dominance H'_3 and H'_6 depend on the value of z . For an arbitrarily small $\delta > 0$

$$\begin{aligned} H'_6 &= o(H'_3) \quad \text{if } z \leq (4 - \delta) \log 1/\Delta, \\ H'_3 &= o(H'_6) \quad \text{if } z \geq (4 + \delta) \log 1/\Delta. \end{aligned}$$

These are easily confirmed if we substitute $(4 \pm \delta) \log 1/\Delta$ into z .

Substituting $z = (x - \zeta)^{\alpha/(\alpha-1)}$ into H'_3 and H'_6 and multiplying each of them by $\alpha^2/(2(\alpha - 1)^2)(x - \zeta)^{2/(\alpha-1)}$ we obtain the second integral of the equation (2.12). Combining this with Theorem 2.1, we obtain the desired result. For $x - \zeta < 0$ we use the relation $f(x; \alpha, \beta) = f(-x; \alpha, -\beta)$. For proving uniformity in Δ and z we need to let $\epsilon, \epsilon' \downarrow 0$. We omit the derivation because it is quite similar to that in the proof of Theorem 2.1.

□

From Theorem 2.1, Theorem 2.2 and formulas (1.2), (1.3), we obtain behavior of the score functions as $\alpha \uparrow 2$. These score functions in the corollaries below are needed for obtaining the Fisher information matrix of the general stable distributions.

Corollary 2.1 *Under the same conditions and notations of Theorem 2.1 and 2.2, define*

$$g_\mu(x; \alpha, \beta) = -g'(x; \alpha, \beta)/g(x; \alpha, \beta).$$

Then for an arbitrarily small $\epsilon > 0$, there exist Δ_0 and x_0 such that for all $\Delta < \Delta_0$ and $|x| > x_0$,

$$|f_\mu(x; \alpha, \beta)/g_\mu(x; \alpha, \beta) - 1| < \epsilon.$$

Furthermore define

$$\begin{aligned} F_\mu^1(x; \alpha, \beta) &= \frac{x - \zeta}{2}, \\ F_\mu^2(x; \alpha, \beta) &= \frac{3}{x - \zeta}. \end{aligned}$$

Then

$$\frac{g_\mu(x; \alpha, \beta)}{g(x; \alpha, \beta)} = \begin{cases} F_\mu^1(x; \alpha, \beta)(1 + o(\Delta^{\delta/2})) & \text{if } |x - \zeta| \leq (2 - \delta)(\log 1/\Delta)^{1/2} \\ F_\mu^2(x; \alpha, \beta)(1 + o(\Delta^{\delta/2})) & \text{if } |x - \zeta| \geq (2 + \delta)(\log 1/\Delta)^{1/2}, \end{cases}$$

uniformly in $|x| > x_0$. For the remaining interval $(2 - \delta)(\log 1/\Delta)^{1/2} \leq x \leq (2 + \delta)(\log 1/\Delta)^{1/2}$,

$$\left| \frac{g_\mu(x; \alpha, \beta)}{g(x; \alpha, \beta)} \right| \leq \bar{c}x$$

uniformly in $|x| > x_0$.

Corollary 2.2 *Under the same conditions and notations of Theorems 2.1 and 2.2, define*

$$g_\sigma(x; \alpha, \beta) = -g(x; \alpha, \beta) - (x - \zeta)g'(x; \alpha, \beta).$$

Then for an arbitrarily small $\delta > 0$, there exist Δ_0 and x_0 such that for all $\Delta < \Delta_0$ and $|x| > x_0$

$$|f_\sigma(x; \alpha, \beta)/g_\sigma(x; \alpha, \beta) - 1| < \epsilon.$$

Furthermore define

$$\begin{aligned} F_\sigma^1(x; \alpha, \beta) &= \frac{(x - \zeta)^2}{2}, \\ F_\sigma^2(x; \alpha, \beta) &= 2. \end{aligned}$$

Then

$$\frac{g_\sigma(x; \alpha, \beta)}{g(x; \alpha, \beta)} = \begin{cases} F_\sigma^1(x; \alpha, \beta)(1 + o(\Delta^{1/2})) & \text{if } |x - \zeta| \leq (2 - \delta)(\log 1/\Delta)^{1/2} \\ F_\sigma^2(x; \alpha, \beta)(1 + o(\Delta^{1/2})) & \text{if } |x - \zeta| \geq (2 + \delta)(\log 1/\Delta)^{1/2}. \end{cases}$$

For the remaining interval $(2 - \delta)(\log 1/\Delta)^{1/2} \leq x \leq (2 + \delta)(\log 1/\Delta)^{1/2}$,

$$\left| \frac{g_\sigma(x; \alpha, \beta)}{g(x; \alpha, \beta)} \right| \leq \bar{c}x^2$$

uniformly in $|x| > x_0$.

3 Derivatives of density with respect to the parameters α and β

In this section we obtain the derivatives of the density with respect to the parameters α and β by analyzing the inversion formula. The density of general stable distributions can be written as

$$(3.1) \quad f(x; \alpha, \beta) = \frac{1}{\pi} \operatorname{Re} \int_0^\infty e^{-itx} \Phi(t; \alpha, \beta) dt,$$

where $\Phi(t; \alpha, \beta)$ is defined in (1.1). Utilizing (3.1), we derive the following two lemmas concerning derivatives with respect to α and β , which are extensions of Lemma 5 of Nagaev and Shkol'nik (1988) to non-symmetric case.

Lemma 3.1 *As $\Delta = 2 - \alpha \rightarrow 0$, there exists x_0 and for all $|x| \geq x_0$,*

$$f_\alpha(x; \alpha, \beta) = -\frac{1}{|y|^{1+\alpha}} \left\{ 1 + \beta^* + \Delta(M_1 + M_2 \log |y|) + \frac{M_3}{|y|} + \frac{M_4 + M_5 \log |y|}{|y|^\alpha} \right\},$$

where $y = x - \zeta$ and M_1, \dots, M_5 are some constants.

Lemma 3.2 *Under the same conditions and notations of Lemma 3.1,*

$$f_\beta(x; \alpha, \beta) = \frac{\Delta \operatorname{sgn} y}{|y|^{1+\alpha}} \left(1 + \Delta M_6 + \frac{M_7}{|y|} + \frac{M_8}{|y|^\alpha} \right),$$

where M_6 , M_7 and M_8 are some constants. Furthermore

$$|f_\beta(x; \alpha, \beta)| = O(\Delta),$$

uniformly $x \in \mathbf{R}$.

Proof of Lemma 3.1 First we consider the case of $y = x - \zeta > 0$. Differentiating f by α , we get

$$(3.2) \quad f_\alpha(x; \alpha, \beta) = \frac{1}{\pi} \operatorname{Re} \int_0^\infty e^{-ity} \exp \left(-t^\alpha \left(1 - i\beta \tan \left(\frac{\pi\alpha}{2} \right) \right) \right) \\ \times \left\{ -t^\alpha \log t \left(1 - i\beta \tan \left(\frac{\pi\alpha}{2} \right) \right) + (t^\alpha - t) \frac{i\beta\pi}{2} / \left(\cos \left(\frac{\pi\alpha}{2} \right) \right)^2 \right\} dt.$$

Transforming $f_\alpha(x; \alpha, \beta)$ by $t \rightarrow s/y$, we obtain

$$f_\alpha(x; \alpha, \beta) = \frac{1}{\pi y} \operatorname{Re} \int_0^\infty e^{-is} \exp \left(-s^\alpha y^{-\alpha} \left(1 - i\beta \tan \left(\frac{\pi\alpha}{2} \right) \right)^2 \right) \\ \times \left\{ -s^\alpha y^{-\alpha} (\log s + \log y) \left(1 - i\beta \tan \left(\frac{\pi\alpha}{2} \right) \right) + (s^\alpha y^{-\alpha} - sy^{-1}) \frac{i\beta\pi}{2} / \left(\cos \left(\frac{\pi\alpha}{2} \right) \right)^2 \right\} ds$$

Further transforming $f_\alpha(x; \alpha, \beta)$ by $s \rightarrow te^{i\varsigma}$, $\varsigma = -\frac{\pi}{2\alpha}$ as in Lemma 5 of Nagaev and Shkol'nik (1988), we find

$$f_\alpha(x; \alpha, \beta) = \frac{1}{\pi y} \operatorname{Re} e^{i\varsigma} \int_0^\infty \exp \left(-ite^{i\varsigma} + it^\alpha y^{-\alpha} \left(1 - i\beta \tan \left(\frac{\pi\alpha}{2} \right) \right) \right) \\ \times \left\{ it^\alpha y^{-\alpha} (\log t + i\varsigma + \log y) \left(1 - i\beta \tan \left(\frac{\pi\alpha}{2} \right) \right) - (it^\alpha y^{-\alpha} + ty^{-1} e^{i\varsigma}) \frac{i\beta\pi}{2} / \left(\cos \frac{\pi\alpha}{2} \right)^2 \right\} dt \\ = K_1 + K_2,$$

where K_1 and K_2 are obtained in formulas (E.2) and (E.4) of appendix E. Then the lemma is proved for $y = x - \zeta > 0$. For $y = x - \zeta < 0$, we use the relation $f_\alpha(x; \alpha, \beta) = f_\alpha(-x; \alpha, -\beta)$. \square

Proof of Lemma 3.2 First we assume $y = x - \zeta > 0$. Differentiating the density (3.1) by β and doing the same transform as in Lemma 3.1, we obtain

$$f_\beta(x; \alpha, \beta) = \frac{\tan(\frac{\pi\alpha}{2})}{\pi y} \left(\frac{1}{y^\alpha} \operatorname{Re} \int_0^\infty e^{i\varsigma} e^{-ite^{i\varsigma}} t^\alpha dt - \frac{1}{y} \operatorname{Re} \int_0^\infty i e^{2i\varsigma} e^{-ite^{i\varsigma}} t dt + \frac{c}{y^{2\alpha}} + \frac{c}{y^{1+\alpha}} \right).$$

The calculation of $f_\beta(x; \alpha, \beta)$ is easy by the equations (E.1) and (E.3) in appendix E. As $\Delta \rightarrow 0$, $\tan \frac{\pi\alpha}{2} = -\frac{\pi}{2}\Delta + o(\Delta^2)$. Finally we utilize the relation $f_\beta(x; \alpha, \beta) = -f_\beta(x; \alpha, -\beta)$ and the first part of the lemma is proved. The second part of the lemma is obvious from

$$|f_\beta(x; \alpha, \beta)| \leq \frac{|\tan \frac{\alpha\pi}{2}|}{\pi} \int_0^\infty e^{-t^\alpha} |t^\alpha - t| dt.$$

\square

Table 1: Limit of information matrix at $\alpha = 2$

$I_{\theta\theta}$	μ	σ	α	β
μ	0.5	0	0	0
σ	0	2.0	$-\infty$	0
α	0	$-\infty$	∞	0
β	0	0	0	0

4 Information matrix of general stable distributions

The Fisher information matrix of general stable distributions are derived in this section. Table 1 gives limiting values of the information matrix at $\alpha = 2$. To the author's knowledge other than the diagonal element and $I_{\mu\sigma}$, these limiting results have not been given in literature. In following theorem we obtain asymptotic behavior of information matrix in more detail for the important diverging cases.

Theorem 4.1 *As $\Delta = 2 - \alpha \rightarrow 0$, behavior of the Fisher information matrix of general stable distributions at $\mu = 0$, $\sigma = 1$, $\beta \neq \pm 1$, is given as follows.*

$$\begin{bmatrix} I_{\mu\mu} & I_{\mu\sigma} & I_{\mu\alpha} & I_{\mu\beta} \\ * & I_{\sigma\sigma} & I_{\sigma\alpha} & I_{\sigma\beta} \\ * & * & I_{\alpha\alpha} & I_{\alpha\beta} \\ * & * & * & I_{\beta\beta} \end{bmatrix} = \begin{bmatrix} 0.5 + o(1) & o(1) & o(1) & O(\Delta) \\ * & 2.0 + o(1) & -\frac{1}{2} \log \log 1/\Delta \times (1 + o(1)) & o(\Delta \log \log 1/\Delta) \\ * & * & \frac{1}{4\Delta \log 1/\Delta} \times (1 + o(1)) & o\left(\frac{1}{\log 1/\Delta}\right) \\ * & * & * & \frac{\Delta}{4(1-\beta^2) \log 1/\Delta} \times (1 + o(1)) \end{bmatrix}.$$

Note that for $I_{\mu\sigma}$ and $I_{\mu\alpha}$ the rates of convergences to 0 are not obtained and only the limiting values are obtained. For $I_{\sigma\alpha}$, $I_{\alpha\alpha}$ and $I_{\alpha\beta}$ the exact rates of convergences are obtained. However for symmetric stable distributions since $I_{\mu\sigma} = I_{\mu\alpha} = 0$, the above 3×3 matrix presents exact limiting behavior of the Fisher information matrix as $\Delta \rightarrow 0$. The idea of the proof is based on the proof of Theorem 2 of Nagaev and Shkol'nik (1988).

Proof of Theorem 4.1 In our proof we use the following notations. T is a sufficiently large constant such that Theorem 2.1 and Corollaries 2.1 and 2.2 are applicable. We also denote

$$x_1(\Delta) = (2 - \delta)(\log 1/\Delta)^{1/2}, \quad x_2(\Delta) = (2 + \delta)(\log 1/\Delta)^{1/2}, \quad x_3(\Delta) = \exp(\Delta^{-1/2}),$$

where $\delta > 0$ is an arbitrarily small constant. Further we use the notation “const” for some proper constants. We prove $I_{\alpha\alpha}$ in some detail and the proof of the other $I_{\theta\theta}$ are given in Appendix F .

Proof of $I_{\alpha\alpha}$:

In our proof we utilize Theorem 2.1 and Lemma 3.1. We divide integral of Fisher information matrix

into two parts,

$$\begin{aligned} I_{\alpha\alpha} &= \int_0^\infty \frac{\{f_\alpha(x + \zeta; \alpha, \beta)\}^2}{f(x + \zeta; \alpha, \beta)} dx + \int_0^\infty \frac{\{f_\alpha(x - \zeta; \alpha, -\beta)\}^2}{f(x - \zeta; \alpha, -\beta)} dx \\ &= I_{\alpha\alpha}^1 + I_{\alpha\alpha}^2. \end{aligned}$$

This is obtained by the relation $f(x; \alpha, \beta) = f(-x; \alpha, -\beta)$ for $x - \zeta < 0$. $I_{\alpha\alpha}^1$ is calculated first. Further we divide integration $I_{\alpha\alpha}^1$ into five subintegrals,

$$I_{\alpha\alpha}^1 = \sum_{k=1}^5 I_{\alpha\alpha}(k),$$

where each $I_{\alpha\alpha}(k)$ corresponds to the k -th interval of $[0, T)$, $[T, x_1(\Delta))$, $[x_1(\Delta), x_2(\Delta))$, $[x_2(\Delta), x_3(\Delta))$ and $[x_3(\Delta), \infty)$.

Clearly

$$(4.1) \quad I_{\alpha\alpha}(1) < \infty.$$

For $I_{\alpha\alpha}(2)$

$$f(x + \zeta; \alpha, \beta) = f(x; 2)(1 + o(1)), \quad f_\alpha(x + \zeta; \alpha, \beta) = \text{const} \times x^{\Delta-3}.$$

Then

$$(4.2) \quad I_{\alpha\alpha}(2) = \text{const} \times \int_T^{x_1(\Delta)} x^{2\Delta-6} \exp\left(\frac{x^2}{4}\right) dx \leq \text{const} \times \frac{1}{\Delta^{1-\delta}(\log 1/\Delta)^{5/2-\Delta}}.$$

For $I_{\alpha\alpha}(3)$

$$f(x + \zeta; \beta, \alpha) = \{(1 + \beta)\Delta x^{\Delta-3} + f(x; 2)\}(1 + o(1)), \quad f_\alpha(x + \zeta; \alpha, \beta) = \text{const} \times x^{\Delta-3}.$$

Then

$$(4.3) \quad I_{\alpha\alpha}(3) \leq \text{const} \times \frac{1}{\Delta} \int_{x_1(\Delta)}^{x_2(\Delta)} x^{\Delta-3} dx = \text{const} \times \frac{\delta}{\Delta \log 1/\Delta}.$$

For $I_{\alpha\alpha}(4)$

$$f(x + \zeta; \alpha, \beta) = (1 + \beta)\Delta x^{\Delta-3}(1 + o(1)), \quad f_\alpha(x + \zeta; \alpha, \beta) = -(1 + \beta)x^{\Delta-3}(1 + o(1)).$$

Then

$$(4.4) \quad I_{\alpha\alpha}(4) = \frac{1 + \beta}{\Delta} \int_{x_2(\Delta)}^{x_3(\Delta)} x^{\Delta-3} dx (1 + o(1)) = \frac{1 + \beta}{8\Delta \log 1/\Delta} (1 + o(1)).$$

For $I_{\alpha\alpha}(5)$

$$f(x + \zeta; \alpha, \beta) = (1 + \beta)\Delta x^{\Delta-3}(1 + o(1)), \quad f_\alpha(x + \zeta; \alpha, \beta) = -(1 + \beta + \Delta \log x)x^{\Delta-3}(1 + o(1)).$$

Then

$$\begin{aligned}
(4.5) \quad I_{\alpha\alpha}(5) &= \text{const} \times \frac{1}{\Delta} \int_{x_3(\Delta)}^{\infty} x^{\Delta-3} \{\max(1+\beta, \Delta \log x)\}^2 dx \\
&\leq \text{const} \times \Delta \int_{x_3(\Delta)}^{e^{(1+\beta)/\Delta}} x^{\Delta-3} (\log x)^2 dx + \text{const} \times \frac{1}{\Delta} \int_{e^{(1+\beta)/\Delta}}^{\infty} x^{\Delta-3} dx \\
&\leq \text{const} \times \frac{1}{\Delta} \int_{x_3(\Delta)}^{\infty} x^{\Delta-3} dx \\
&= O(e^{-2/\Delta^{1/2}}/\Delta) \rightarrow 0, \text{ as } \Delta \rightarrow 0.
\end{aligned}$$

From the formulas (4.1)-(4.5) we obtain

$$I_{\alpha\alpha}^1 = \frac{1+\beta}{8\Delta \log 1/\Delta} (1 + o(1)).$$

Setting $\beta \rightarrow -\beta$ in $I_{\alpha\alpha}^1$, we obtain $I_{\alpha\alpha}^2$. Adding $I_{\alpha\alpha}^1$ and $I_{\alpha\alpha}^2$, we prove the assertion. \square

A Proof of Lemma 2.1

As $\Delta \rightarrow 0$, $\alpha\varrho = \frac{2}{\pi} \arctan(\beta \tan(\frac{\pi\alpha}{2}))$ in $A(\varphi, \alpha, \beta)$ is expanded as

$$\alpha\varrho = \frac{2}{\pi} \arctan\left(\beta \tan\left(\frac{\pi\alpha}{2}\right)\right) = -\beta\Delta - \frac{\beta(1-\beta^2)}{3} \left(\frac{\pi}{2}\right)^2 \Delta^3 + o(\Delta^4).$$

Since the smaller remainder terms other than $-\beta\Delta$ are not needed in the following argument we substitute $-\beta\Delta$ into $A(\varphi, \alpha, \beta)$ for $\alpha\varrho$ in advance. Then for convenience we write $A(\varphi, \alpha, \beta)$ of (1.5) as

$$\begin{aligned}
(A.1) \quad A(\varphi, \alpha, \beta) &= \frac{\cos(\frac{\pi}{2}\{(1-\Delta)\varphi - \beta\Delta\})}{\sin(\frac{\pi}{2}\{(2-\Delta)\varphi - \beta\Delta\})} \left(\frac{\cos(\frac{\pi}{2}\varphi) \cos(\frac{\pi}{2}\beta\Delta)}{\sin(\frac{\pi}{2}\{(2-\Delta)\varphi - \beta\Delta\})} \right)^{1/(1-\Delta)} \\
&= \frac{C(\varphi, \Delta, \beta)}{B(\varphi, \Delta, \beta)} \left(\frac{D(\varphi)E(\Delta, \beta)}{B(\varphi, \Delta, \beta)} \right)^{1/(1-\Delta)},
\end{aligned}$$

where

$$\begin{aligned}
B(\varphi, \Delta, \beta) &= \sin\left(\frac{\pi}{2}\{(2-\Delta)\varphi - \beta\Delta\}\right), \\
C(\varphi, \Delta, \beta) &= \cos\left(\frac{\pi}{2}\{(1-\Delta)\varphi - \beta\Delta\}\right), \\
D(\varphi) &= \cos\left(\frac{\pi}{2}\varphi\right), \\
E(\Delta, \beta) &= \cos\left(\frac{\pi}{2}\beta\Delta\right).
\end{aligned}$$

Then substituting $1 - \lambda$ for φ we obtain

$$(A.2) \quad B(1 - \lambda, \Delta, \beta) = \sin\left(\frac{\pi}{2}(\Delta + 2\lambda - \Delta\lambda + \beta\Delta)\right),$$

$$(A.3) \quad C(1 - \lambda, \Delta, \beta) = \sin\left(\frac{\pi}{2}(\Delta + \lambda - \Delta\lambda + \beta\Delta)\right),$$

$$(A.4) \quad D(1 - \lambda) = \sin\left(\frac{\pi}{2}\lambda\right),$$

$$(A.5) \quad E(\Delta, \beta) = \cos\left(\frac{\pi}{2}\beta\Delta\right).$$

Note that if $\Delta \rightarrow 0$, then $\lambda \rightarrow 0$ since $0 \leq \lambda \leq \Delta/\epsilon'$ and $\epsilon' > 0$ is an arbitrarily small fixed number (or even when ϵ' converges to 0 slower than Δ). Expanding in terms of $\Delta \rightarrow 0$, we have

$$\begin{aligned} B(1 - \lambda, \Delta, \beta) &= \frac{\pi}{2}\Delta \left(1 + 2\frac{\lambda}{\Delta} - \lambda + \beta\right) + O(\Delta^3), \\ C(1 - \lambda, \Delta, \beta) &= \frac{\pi}{2}\Delta \left(1 + \frac{\lambda}{\Delta} - \lambda + \beta\right) + O(\Delta^3), \\ D(1 - \lambda) &= \frac{\pi}{2}\Delta \times \frac{\lambda}{\Delta} + O(\Delta^3), \\ E(\Delta, \beta) &= 1 - \frac{1}{2}\left(\frac{\pi}{2}\beta\Delta\right)^2 + o(\Delta^3). \end{aligned}$$

Substituting these expansions to $A(\varphi, \Delta, \beta)$ of (A.1) and utilizing $x^{1/(\Delta-1)} = x^{-1} - x^{-1} \log x \Delta + o(\Delta)$, we prove the desired result.

□

B Proof of Lemma 2.2

We prove $A(\varphi_\Delta, \Delta, \beta)$ first. When $\varphi = \varphi_\Delta = 1 - \Delta^{1/2-\epsilon}$, we replace λ by $\Delta^{1/2-\epsilon}$ in the formulas (A.2)-(A.4). Then

$$\begin{aligned} B(\varphi_\Delta, \Delta, \beta) &= \sin\left(\frac{\pi}{2}\{2\Delta^{1/2-\epsilon} + (1 + \beta)\Delta - \Delta^{3/2-\epsilon}\}\right), \\ C(\varphi_\Delta, \Delta, \beta) &= \sin\left(\frac{\pi}{2}\{\Delta^{1/2-\epsilon} + (1 + \beta)\Delta - \Delta^{3/2-\epsilon}\}\right), \\ D(\varphi_\Delta) &= \sin\left(\frac{\pi}{2}\Delta^{1/2-\epsilon}\right). \end{aligned}$$

Note that $E(\Delta, \beta)$ is the same as in Proof of Lemma 2.1 and we omit it. As $\Delta \rightarrow 0$ we have

$$(B.1) \quad \begin{aligned} B(\varphi_\Delta, \Delta, \beta) &= \pi\Delta^{1/2-\epsilon} \left\{1 + \frac{1}{2}(1 + \beta)\Delta^{\frac{1}{2}+\epsilon} - \frac{\pi^2}{6}\Delta^{1-2\epsilon} - \frac{1}{2}\Delta - \frac{\pi^2}{4}(1 + \beta)\Delta^{3/2-\epsilon} + o(\Delta^{3/2})\right\}, \end{aligned}$$

$$(B.2) \quad \begin{aligned} C(\varphi_\Delta, \Delta, \beta) &= \frac{\pi}{2}\Delta^{1/2-\epsilon} \left\{1 + (1 + \beta)\Delta^{1/2+\epsilon} - \frac{1}{6}\frac{\pi^2}{4}\Delta^{1-2\epsilon} - \Delta - \frac{\pi^2}{8}(1 + \beta)\Delta^{3/2-\epsilon} + o(\Delta^{3/2})\right\}, \end{aligned}$$

$$(B.3) \quad D(\varphi_\Delta) = \frac{\pi}{2}\Delta^{1/2-\epsilon} \left(1 - \frac{1}{6}\frac{\pi^2}{4}\Delta^{1-2\epsilon} + o(\Delta^{3/2})\right).$$

Substituting these expansions into $A(\varphi, \Delta, \beta)$ of (A.1), we obtain the desired result.

Next we prove assertion concerning $A'(\varphi_\Delta, \Delta, \beta)$. From the equation (1.5), we can write

$$\begin{aligned}
\text{(B.4)} \quad A'(\varphi, \Delta, \beta) &= A(\varphi, \Delta, \beta) \\
&\times \frac{\pi}{2} \left\{ \frac{(2-\Delta)^2 \cos(\frac{\pi}{2}\{(2-\Delta)\varphi - \beta\Delta\})}{\Delta-1 \sin(\frac{\pi}{2}\{(2-\Delta)\varphi - \beta\Delta\})} - (1-\Delta) \frac{\sin(\frac{\pi}{2}\{(1-\Delta)\varphi - \beta\Delta\})}{\cos(\frac{\pi}{2}\{(1-\Delta)\varphi - \beta\Delta\})} - \frac{1}{1-\Delta} \frac{\sin(\frac{\pi}{2}\varphi)}{\cos(\frac{\pi}{2}\varphi)} \right\} \\
&= A(\varphi, \Delta, \beta) \times \frac{\pi}{2} \left\{ \frac{(2-\Delta)^2}{\Delta-1} \frac{F(\varphi, \Delta, \beta)}{B(\varphi, \Delta, \beta)} - (1-\Delta) \frac{G(\varphi, \Delta, \beta)}{C(\varphi, \Delta, \beta)} - \frac{1}{1-\Delta} \frac{H(\varphi)}{D(\varphi)} \right\},
\end{aligned}$$

where

$$\begin{aligned}
F(\varphi, \Delta, \beta) &= \cos\left(\frac{\pi}{2}\{(2-\Delta)\varphi - \beta\Delta\}\right), \\
G(\varphi, \Delta, \beta) &= \sin\left(\frac{\pi}{2}\{(1-\Delta)\varphi - \beta\Delta\}\right), \\
H(\varphi) &= \sin\left(\frac{\pi}{2}\varphi\right).
\end{aligned}$$

Then substituting φ_Δ for φ , we obtain

$$\begin{aligned}
F(\varphi_\Delta, \Delta, \beta) &= -\cos\left(\frac{\pi}{2}\{2\Delta^{1/2-\epsilon} + (1+\beta)\Delta - \Delta^{3/2-\epsilon}\}\right), \\
G(\varphi_\Delta, \Delta, \beta) &= \cos\left(\frac{\pi}{2}\{\Delta^{1/2-\epsilon} + (1+\beta)\Delta - \Delta^{3/2-\epsilon}\}\right), \\
H(\varphi_\Delta) &= \cos\left(\frac{\pi}{2}\Delta^{1/2-\epsilon}\right).
\end{aligned}$$

As $\Delta \rightarrow 0$,

$$\text{(B.5)} \quad F(\varphi_\Delta, \Delta, \beta) = -1 + \frac{1}{2}\pi^2\Delta^{1-2\epsilon} + \frac{1}{2}\pi^2(1+\beta)\Delta^{3/2-\epsilon} + o(\Delta^{3/2}),$$

$$\text{(B.6)} \quad G(\varphi_\Delta, \Delta, \beta) = 1 - \frac{\pi^2}{8}\Delta^{1-2\epsilon} - \frac{\pi^2}{4}(1+\beta)\Delta^{3/2-\epsilon} + o(\Delta^{3/2}),$$

$$\text{(B.7)} \quad H(\varphi_\Delta) = 1 - \frac{\pi^2}{8}\Delta^{1-2\epsilon} + o(\Delta^{3/2}).$$

Combining (B.1)-(B.3), (B.5)-(B.7), $A(\varphi, \Delta, \beta)$ and $A'(\varphi, \alpha, \beta)$ of (B.4), we prove the lemma.

□

C Proof of Lemma 2.3

We further expand $B(\varphi, \Delta, \beta)$, $C(\varphi, \Delta, \beta)$ and $D(\varphi, \Delta)$ in (A.2)-(A.4) with respect to λ . As $\Delta \rightarrow 0$ and $\lambda \rightarrow 0$ such that $\Delta^{1/2-\epsilon} \leq \lambda$,

$$\begin{aligned}
\text{(C.1)} \quad B(1-\lambda, \Delta, \beta) &= \pi\lambda \left\{ 1 - \frac{1}{6}(\pi\lambda)^2 + \frac{1}{120}(\pi\lambda)^4 + \frac{\Delta(1-\lambda+\beta)}{2\lambda} - \frac{\pi^2\lambda\Delta}{4}(1-\lambda+\beta) + o(\lambda^4) \right\},
\end{aligned}$$

$$\begin{aligned}
\text{(C.2)} \quad C(1-\lambda, \Delta, \beta) &= \frac{\pi}{2}\lambda \left\{ 1 - \frac{1}{6}\left(\frac{\pi}{2}\lambda\right)^2 + \frac{1}{120}\left(\frac{\pi}{2}\lambda\right)^4 + \frac{\Delta(1-\lambda+\beta)}{\lambda} - \frac{\pi^2\lambda\Delta}{8}(1-\lambda+\beta) + o(\lambda^4) \right\},
\end{aligned}$$

$$\begin{aligned}
\text{(C.3)} \quad D(1-\lambda, \Delta) &= \frac{\pi}{2}\lambda \left\{ 1 - \frac{1}{6}\left(\frac{\pi}{2}\lambda\right)^2 + \frac{1}{120}\left(\frac{\pi}{2}\lambda\right)^4 + o(\lambda^5) \right\}.
\end{aligned}$$

Substituting these equations to $A(\varphi, \Delta, \beta)$, we obtain the result.

Concerning $A'(\varphi, \alpha, \beta)$ we use formulas

$$\begin{aligned} F(1 - \lambda, \Delta, \beta) &= -\cos\left(\frac{\pi}{2}\{2\lambda + (1 - \lambda + \beta)\Delta\}\right), \\ G(1 - \lambda, \Delta, \beta) &= \cos\left(\frac{\pi}{2}\{\lambda + (1 - \lambda + \beta)\Delta\}\right), \\ H(1 - \lambda) &= \cos\left(\frac{\pi}{2}\lambda\right). \end{aligned}$$

As $\Delta \rightarrow 0$ and $\lambda \rightarrow 0$ under $\Delta^{1/2-\epsilon} \leq \lambda$,

$$(C.4) \quad F(1 - \lambda, \Delta, \beta) = -1 + \frac{1}{2}(\pi\lambda)^2 + o(\lambda^3),$$

$$(C.5) \quad G(1 - \lambda, \Delta, \beta) = 1 - \frac{1}{2}\left(\frac{\pi}{2}\lambda\right)^2 + o(\lambda^3),$$

$$(C.6) \quad H(1 - \lambda) = 1 - \frac{1}{2}\left(\frac{\pi}{2}\lambda\right)^2 + o(\lambda^3).$$

Substituting (C.1)-(C.3) and (C.4)-(C.6) to $A'(\varphi, \alpha, \beta)$ of (B.4), we obtain the desired assertion. From the equation (1.5) or $A'(\varphi, \alpha, \beta)$ of (B.4), we can write

$$\begin{aligned} (C.7) \quad A''(\varphi, \Delta, \beta) &= A'(\varphi, \Delta, \beta) \times \frac{\pi}{2} \left\{ \frac{(2 - \Delta)^2}{\Delta - 1} \frac{F(\varphi, \Delta, \beta)}{B(\varphi, \Delta, \beta)} - (1 - \Delta) \frac{G(\varphi, \Delta, \beta)}{C(\varphi, \Delta, \beta)} - \frac{1}{1 - \Delta} \frac{H(\varphi)}{D(\varphi)} \right\} \\ &+ A(\varphi) \frac{\pi^2}{4(1 - \Delta)} \left\{ \frac{(2 - \Delta)^3}{(\sin(\frac{\pi}{2}\{(2 - \Delta)\varphi - \beta\Delta\}))^2} - \frac{(1 - \Delta)^3}{(\cos(\frac{\pi}{2}\{(1 - \Delta)\varphi - \beta\Delta\}))^2} - \frac{1}{(\cos(\frac{\pi}{2}\varphi))^2} \right\} \\ &= A'(\varphi, \Delta, \beta) \times \frac{\pi}{2} \left\{ \frac{(2 - \Delta)^2}{\Delta - 1} \frac{F(\varphi, \Delta, \beta)}{B(\varphi, \Delta, \beta)} - (1 - \Delta) \frac{G(\varphi, \Delta, \beta)}{C(\varphi, \Delta, \beta)} - \frac{1}{1 - \Delta} \frac{H(\varphi)}{D(\varphi)} \right\} \\ &+ A(\varphi) \frac{\pi^2}{4(1 - \Delta)} \left\{ \frac{(2 - \Delta)^3}{B^2(\varphi, \Delta, \beta)} - \frac{(1 - \Delta)^3}{C^2(\varphi, \Delta, \beta)} - \frac{1}{D^2(\varphi)} \right\}. \end{aligned}$$

Combining (C.1)-(C.3), (C.4)-(C.6), $A'(\varphi, \alpha, \beta)$ of (B.4) and $A''(\varphi, \alpha, \beta)$ of (C.7), we obtain the desired assertion.

□

D Proof of Lemma 2.4

We only need to prove for $x - \zeta > 0$, because for $x - \zeta < 0$ $f(x; \alpha, \beta) = f(-x; \alpha, -\beta)$ and ϱ^* in (1.5) does not change, namely

$$\varrho^* = \text{sgn}(x - \zeta) \frac{\pi\alpha}{2} \arctan\left(\beta \tan\left(\frac{\pi\alpha}{2}\right)\right) = \text{sgn}(-x + \zeta) \frac{\pi\alpha}{2} \arctan\left(-\beta \tan\left(\frac{\pi\alpha}{2}\right)\right).$$

We use notation $\beta_B \in (-1, 1)$ which corresponds to the parameter $\beta \in (-1, 1)$ for (B) representation of characteristic functions. The reason is that the proof in terms of (B) representation is simpler since $\alpha\varrho = -\beta_B\Delta$ (see p.74 of Zolotarev 1986 ; the notation ϱ in our paper corresponds to θ in Zolotarev) and we do not need to expand $\alpha\varrho$ with respect to β as in the proof of Lemma 2.3. Moreover our purpose is only to determine the sign of $A'(\varphi)$. We divide the integral interval as $(-\varrho, 1 - \Delta/\epsilon']$,

$[1 - \Delta/\epsilon', 1)$ and prove the lemma for each interval. Except for the interval $(-\varrho, 0)$ we follow the line of the proof of Lemma 4 of Nagaev and Shkol'nik (1988). For $(1 - \Delta/\epsilon', 1)$ the proof is obvious from Lemma 2.1. For $(-\varrho, 1 - \Delta/\epsilon']$ we have to consider the sign of ϱ . Note that

$$\text{sgn}(\varrho) = -\text{sgn}(\beta_B).$$

First for $0 \leq -\varrho \leq \varphi \leq 1 - \Delta/\epsilon' \Leftrightarrow \Delta/\epsilon' \leq \lambda \leq 1 + \varrho$, we get

$$A'(\varphi) = \frac{-1}{1 - \Delta} \frac{\pi}{2} A(\varphi) A_1(\varphi),$$

where

$$\begin{aligned} A_1(1 - \lambda) &= -(2 - \Delta)^2 \cot\left(\frac{\pi}{2}(\Delta + 2\lambda - \Delta\lambda + \beta_B\Delta)\right) \\ &\quad + (1 - \Delta)^2 \cot\left(\frac{\pi}{2}(\Delta + \lambda - \Delta\lambda + \beta_B\Delta)\right) + \cot\left(\frac{\pi}{2}\lambda\right). \end{aligned}$$

From the formula (4.3.70) in [4] we have

$$A_1(1 - \lambda) = \frac{2(1 + \beta_B)^2 \Delta^2}{\pi \lambda (\Delta + 2\lambda - \Delta\lambda + \beta_B\Delta)(\Delta + \lambda - \Delta\lambda + \beta_B\Delta)} + \sum_{k=0}^{\infty} d_k \gamma_k(\lambda),$$

where d_k 's are positive numbers and

$$\gamma_k(\lambda) = \left(\frac{\pi}{2}\lambda\right)^{2k+1} [(2 - \Delta)^2 \{2 - \Delta + (1 + \beta_B)\Delta/\lambda\} - (1 - \Delta)^2 \{1 - \Delta + (1 + \beta_B)\Delta/\lambda\}].$$

Since $\Delta/\lambda \leq \epsilon'$ for arbitrarily small $\epsilon' > 0$, $A_1(\varphi) > 0$ is easily confirmed.

Secondly we investigate the case $-\varrho < \varphi \leq 0 \Leftrightarrow \alpha\varphi = \gamma\Delta$ where $\gamma \in (\beta_B, 0)$ for $\beta_B < 0$. Then

$$\begin{aligned} A_1(\varphi) &= (2 - \Delta)^2 \cot\left(\frac{\pi}{2}\alpha(\varphi + \varrho)\right) + (1 - \Delta)^2 \tan\left(\frac{\pi}{2}\{\alpha\varrho + (\alpha - 1)\varphi\}\right) + \tan\left(\frac{\pi}{2}\varrho\right) \\ &= (2 - \Delta)^2 \cot\left(\frac{\pi}{2}(\gamma - \beta_B)\Delta\right) + (1 - \Delta)^2 \tan\left(\frac{\pi}{2}\left(\gamma - \beta_B - \frac{\gamma}{2 - \Delta}\right)\Delta\right) + \tan\left(\frac{\pi}{2}\frac{\gamma\Delta}{2 - \Delta}\right). \end{aligned}$$

As $\Delta \rightarrow 0$,

$$A_1(\varphi) = \frac{(2 - \Delta)^2}{\frac{\pi}{2}(\gamma - \beta_B)\Delta} + O(\Delta).$$

We easily find that $A_1(\varphi) > 0$ for sufficiently small Δ .

□

E Calculations of f_α

For all $y \geq y_0 = x_0 - \zeta$, we have

$$\exp\left(it^\alpha y^{-\alpha} \left(1 - i\beta \tan\left(\frac{\pi\alpha}{2}\right)\right)\right) = 1 + c_0 t^\alpha y^{-\alpha} \left(1 - i\beta \tan\left(\frac{\alpha\pi}{2}\right)\right),$$

where $c_0 \in \mathbf{C}$ is some complex constant. Utilizing this expansion in $f_\alpha(x; \alpha, \beta)$ we obtain

$$\begin{aligned}
K_1 &= \frac{1}{\pi y} \operatorname{Re} e^{i\varsigma} \int_0^\infty \exp \left(-ite^{i\varsigma} + it^\alpha y^{-\alpha} \left(1 - i\beta \tan \left(\frac{\pi\alpha}{2} \right) \right) \right) \\
&\quad \times it^\alpha y^{-\alpha} (\log t + i\varsigma + \log y) \left(1 - i\beta \tan \left(\frac{\pi\alpha}{2} \right) \right) dt \\
&= \frac{1}{\pi y^{1+\alpha}} \operatorname{Re} e^{i\varsigma} \int_0^\infty e^{-ite^{i\varsigma}} \left\{ 1 + c_0 t^\alpha y^{-\alpha} \left(1 - i\beta \tan \left(\frac{\alpha\pi}{2} \right) \right) \right\} \\
&\quad \times it^\alpha (\log t + i\varsigma + \log y) \left(1 - i\beta \tan \left(\frac{\pi\alpha}{2} \right) \right) dt \\
&= \frac{1}{\pi y^{1+\alpha}} \operatorname{Re} e^{i\varsigma} \int_0^\infty e^{-ite^{i\varsigma}} it^\alpha (\log t + i\varsigma + \log y) \left(1 - i\beta \tan \left(\frac{\alpha\pi}{2} \right) \right) dt \\
&\quad + \frac{1}{\pi y^{1+2\alpha}} \operatorname{Re} e^{i\varsigma} \int_0^\infty e^{-ite^{i\varsigma}} i c_0 t^{2\alpha} \left(1 - \beta \tan \left(\frac{\alpha\pi}{2} \right) \right)^2 (\log t + i\varsigma + \log y) dt \\
&= \frac{1}{\pi y^{1+\alpha}} \operatorname{Re} e^{i\varsigma} \int_0^\infty e^{-ite^{i\varsigma}} it^\alpha (\log t + i\varsigma + \log z) dt + \frac{\Delta(c + c \log y)}{y^{1+\alpha}} + \frac{c + c \log y}{y^{1+2\alpha}}.
\end{aligned}$$

Note that As $\Delta \rightarrow 0$, $\tan \frac{\pi\alpha}{2} = -\frac{\pi}{2} \Delta + o(\Delta^2)$. Then we calculate the integrations in K_1 above. Write $\lambda_1 = \cos \varsigma$ and $\lambda_2 = \sin \varsigma$. Here at $\alpha = 2$, $\lambda_1 = 1/\sqrt{2}$ and $\lambda_2 = -1/\sqrt{2}$. From the formulas (3.5), (4.40) in part I and (3.7), (4.17) in part II of Oberhettinger (1990), we have

$$\begin{aligned}
\operatorname{Re} e^{i\varsigma} \int_0^\infty e^{-ite^{i\varsigma}} it^\alpha \log t dt &= \int_0^\infty e^{t\lambda_2} t^2 \log t \{ \lambda_1 \sin(t\lambda_1) - \lambda_2 \cos(t\lambda_1) \} dt + \Delta c \\
&= 4\lambda_1 \lambda_2 \arctan \left(-\frac{\lambda_1}{\lambda_2} \right) + 8\lambda_1 \lambda_2 (\lambda_2^2 - \lambda_1^2) \left(\frac{3}{2} - \text{EU} \right) + \Delta c \\
&= -\frac{\pi}{2} + \Delta c,
\end{aligned}$$

$$\begin{aligned}
\text{(E.1)} \quad -\operatorname{Re} e^{i\varsigma} \int_0^\infty e^{-ite^{i\varsigma}} t^\alpha dt &= -\int_0^\infty e^{t\lambda_2} t^2 \{ \lambda_2 \sin(t\lambda_1) + \lambda_1 \cos(t\lambda_1) \} dt + \Delta c \\
&= -4\varsigma \lambda_1 \lambda_2 + \Delta c \\
&= -\frac{\pi}{2} + \Delta c,
\end{aligned}$$

$$\begin{aligned}
\operatorname{Re} e^{i\varsigma} \int_0^\infty e^{-ite^{i\varsigma}} it^\alpha dt &= \int_0^\infty e^{t\lambda_2} t^2 \{ \lambda_1 \sin(t\lambda_1) - \lambda_2 \cos(t\lambda_1) \} dt + \Delta c \\
&= -2\lambda_1^4 + 2\lambda_2^4 + \Delta c \\
&= \Delta c,
\end{aligned}$$

where EU means Euler's constant. Substituting these equations into K_1 , we obtain

$$\text{(E.2)} \quad K_1 = -\frac{1}{y^{1+\alpha}} \left\{ 1 + \Delta(c + c \log y) + \frac{c + c \log y}{y^\alpha} \right\}.$$

For K_2 we do the similar calculations.

$$\begin{aligned}
K_2 &= -\frac{1}{\pi y} \operatorname{Re} e^{i\varsigma} \int_0^\infty \exp\left(-ite^{i\varsigma} + it^\alpha y^{-\alpha} \left(1 - i\beta \tan\left(\frac{\pi\alpha}{2}\right)\right)\right) \\
&\quad \times (it^\alpha y^{-\alpha} + ty^{-1}e^{i\varsigma}) \frac{i\beta\pi}{2} / \left(\cos\frac{\pi\alpha}{2}\right)^2 dt \\
&= \frac{\beta}{2y(\cos(\frac{\pi\alpha}{2}))^2} \operatorname{Re} e^{i\varsigma} \int_0^\infty e^{-ite^{i\varsigma}} \left\{1 + ct^\alpha y^{-\alpha} \left(1 - i\beta \tan\left(\frac{\pi\alpha}{2}\right)\right)\right\} (t^\alpha y^{-\alpha} - ity^{-1}e^{i\varsigma}) dt \\
&= \frac{\beta}{2y(\cos(\frac{\pi\alpha}{2}))^2} \left\{ \frac{1}{y^\alpha} \operatorname{Re} e^{i\varsigma} \int_0^\infty e^{-ite^{i\varsigma}} t^\alpha dt - \frac{1}{y} \operatorname{Re} e^{2i\varsigma} \int_0^\infty e^{-ite^{i\varsigma}} it dt + \frac{c}{y^{2\alpha}} + \frac{c}{y^{1+\alpha}} \right\}.
\end{aligned}$$

Substituting the following two equations into K_2 , we obtain K_2 . The first equation is obtained from formula (E.1) in the calculation of K_1 . The second equation is obtained from the formulas (3.5) in part I and (3.7) in part II of Oberhettinger (1990).

$$\begin{aligned}
\operatorname{Re} e^{i\varsigma} \int_0^\infty e^{-ite^{i\varsigma}} t^\alpha dt &= -2 + \Delta c, \\
(E.3) \quad -\operatorname{Re} ie^{2i\varsigma} \int_0^\infty e^{-ite^{i\varsigma}} t dt &= \int_0^\infty e^{t\lambda_2} t \{2\lambda_1 \lambda_2 \cos(t\lambda_1) + (\lambda_2^2 - \lambda_1^2) \sin(t\lambda_1)\} dt \\
&= 2\lambda_1 \lambda_2 (\lambda_2^2 - \lambda_1^2) + (\lambda_2^2 - \lambda_1^2) (-2\lambda_1 \lambda_2) \\
&= 0.
\end{aligned}$$

Hence

$$(E.4) \quad K_2 = \frac{\beta}{y^{1+\alpha}} \left\{ -1 + \Delta c + \frac{c}{y^\alpha} + \frac{c}{y} \right\}.$$

F Proof of $I_{\theta\theta}$ other than $I_{\alpha\alpha}$

Here we give proofs for $I_{\beta\beta}, I_{\alpha\beta}, I_{\sigma\alpha}, I_{\sigma\beta}, I_{\mu\beta}$ and $I_{\mu\alpha}$. In the following we assume $T > x_0$ where x_0 is defined in Theorem 2.1 and Corollaries 2.1 and 2.2.

Proof of $I_{\beta\beta}$:

Here we utilize Theorem 2.1 and Lemma 3.2. However the proof of $I_{\beta\beta}$ is essentially the same as the proof of $I_{\alpha\alpha}$ because

$$f_\beta(x; \alpha, \beta) = -\frac{\Delta}{1+\beta} \operatorname{sgn}(x - \zeta) f_\alpha(x; \alpha, \beta) (1 + o(1))$$

for sufficiently large $|x - \zeta|$. Therefore we omit the detailed derivation of $I_{\beta\beta}$.

Proof of $I_{\alpha\beta}$:

Here we utilize Theorem 2.1, Lemma 3.1 and Lemma 3.2. From the same reason as $I_{\beta\beta}$ we omit the proof.

Proof of $I_{\sigma\alpha}$:

We make use of Lemma 3.1 and Corollary 2.2. We divide integration $I_{\sigma\alpha}$ into two subintegrals,

$$\begin{aligned}
I_{\sigma\alpha} &= \int_0^\infty \frac{f_\sigma(x + \zeta; \alpha, \beta) f_\alpha(x + \zeta; \alpha, \beta)}{f(x + \zeta; \alpha, \beta)} dx + \int_0^\infty \frac{f_\sigma(x - \zeta; \alpha, -\beta) f_\alpha(x - \zeta; \alpha, -\beta)}{f(x - \zeta; \alpha, -\beta)} dx \\
&= I_{\sigma\alpha}^1 + I_{\sigma\alpha}^2.
\end{aligned}$$

$I_{\sigma\alpha}^1$ is calculated first. Further we divide integration $I_{\sigma\alpha}^1$ into four subintegrals,

$$I_{\sigma\alpha}^1 = \sum_{k=1}^4 I_{\sigma\alpha}(k),$$

where each $I_{\sigma\alpha}(k)$ corresponds to the integration of $I_{\sigma\alpha}^1$ for the k -th interval of $[0, T)$, $[T, x_1(\Delta))$, $[x_1(\Delta), x_2(\Delta))$ and $[x_2(\Delta), \infty)$. For $I_{\sigma\alpha}(1)$ clearly

$$(F.1) \quad I_{\sigma\alpha}(1) < \infty.$$

For $I_{\sigma\alpha}(2)$

$$\frac{f_{\sigma}(x + \zeta, \alpha, \beta)}{f(x + \zeta, \alpha, \beta)} = \frac{x^2}{2}(1 + o(1)), \quad f(x + \zeta, \alpha, \beta) = -(1 + \beta)x^{\Delta-3}(1 + o(1)).$$

Then

$$(F.2) \quad \begin{aligned} I_{\sigma\alpha}(2) &= -(1 + \beta) \int_T^{x(\Delta)} \frac{x^{\Delta-1}}{2} dx (1 + o(1)) \\ &= -\frac{1 + \beta}{4} \log \log 1/\Delta (1 + o(1)). \end{aligned}$$

For $I_{\sigma\alpha}(3)$

$$\left| \frac{f_{\sigma}(x + \zeta, \alpha, \beta)}{f(x + \zeta, \alpha, \beta)} \right| \leq \text{const} \times x^2, \quad |f_{\alpha}(x + \zeta; \alpha, \beta)| = (1 + \beta)x^{\Delta-3}(1 + o(1)),$$

and

$$\left| \frac{f_{\sigma}(x + \zeta; \alpha, \beta)f_{\alpha}(x + \zeta; \alpha, \beta)}{f(x + \zeta; \alpha, \beta)} \right| = \text{const} \times x^{\Delta-1}(1 + o(1)).$$

It is easy to see

$$(F.3) \quad I_{\sigma\alpha}(3) = \text{const} \times \int_{x_1(\Delta)}^{x_2(\Delta)} x^{\Delta-1} dx = O(\delta).$$

For $I_{\sigma\alpha}(4)$

$$\left| \frac{f_{\sigma}(x + \zeta; \alpha, \beta)}{f(x + \zeta; \alpha, \beta)} \right| \leq 2(1 + o(1)), \quad |f_{\alpha}(x + \zeta; \alpha, \beta)| = (1 + \beta)x^{\Delta-3}(1 + o(1)).$$

Thus we easily show

$$(F.4) \quad I_{\sigma\alpha}(4) \rightarrow 0, \quad \text{as } \Delta \rightarrow 0.$$

From (F.1), (F.2), (F.3) and (F.4), we obtain

$$I_{\sigma\alpha}^1 = -\frac{1}{4}(1 + \beta) \log \log 1/\Delta (1 + o(1)).$$

Setting $\beta \rightarrow -\beta$ in $I_{\sigma\alpha}^1$, we obtain $I_{\sigma\alpha}^2$. Adding $I_{\sigma\alpha}^1$ and $I_{\sigma\alpha}^2$, we prove the assertion.

Proof of $I_{\sigma\beta}$:

Here we utilize Lemma 3.2 and Corollary 2.2. From the same reason as $I_{\beta\beta}$ we omit the proof. Note that the result can be expected from behavior of $\Delta \times I_{\sigma\alpha}$ and the asymmetry of $f_{\beta}(x; \alpha, \beta)$ around ζ , which we see in Lemma 3.2.

Proof of $I_{\mu\beta}$:

Here we utilize Lemma 3.2 and Corollary 2.1. We divide integration $I_{\mu\beta}$ into two subintegrals.

$$\begin{aligned} I_{\mu\beta} &= \int_0^{\infty} \frac{f_{\mu}(x + \zeta; \alpha, \beta) f_{\beta}(x + \zeta; \alpha, \beta)}{f(x + \zeta; \alpha, \beta)} dx + \int_0^{\infty} \frac{f_{\mu}(x - \zeta; \alpha, -\beta) f_{\beta}(x - \zeta; \alpha, -\beta)}{f(x - \zeta; \alpha, -\beta)} dx \\ &= I_{\mu\beta}^1 + I_{\mu\beta}^2. \end{aligned}$$

$I_{\mu\beta}^1$ is calculated first. Further we divide integration $I_{\mu\beta}^1$ into two subintegrals,

$$I_{\mu\beta}^1 = \sum_{k=1}^2 I_{\mu\beta}(k),$$

where each $I_{\mu\beta}(k)$ corresponds to the integration of $I_{\mu\beta}^1$ for the k -th interval of $[0, T)$ and $[T, \infty)$. For $I_{\mu\beta}(1)$

$$f_{\beta}(x + \zeta; \alpha, \beta) = O(\Delta), \quad \left| \frac{f_{\mu}(x + \zeta; \alpha, \beta)}{f(x + \zeta; \alpha, \beta)} \right| \leq \text{const}.$$

Then $I_{\mu\beta}(1) = O(\Delta)$. For $I_{\mu\beta}(2)$ it follows that

$$\left| \frac{f_{\mu}(x + \zeta; \alpha, \beta)}{f(x + \zeta; \alpha, \beta)} \right| \leq \text{const} \times x, \quad f_{\beta}(x + \zeta; \alpha, \beta) = \text{const} \times \Delta x^{\Delta-3},$$

uniformly in $x \in [T, \infty)$. Thus we find $I_{\mu\beta}(2) = O(\Delta)$ easily. Setting $\beta \rightarrow -\beta$ in $I_{\mu\beta}^1$, we obtain $I_{\mu\beta}^2$. Adding $I_{\mu\beta}^1$ and $I_{\mu\beta}^2$, we obtain the desired result.

Proof of $I_{\mu\alpha}$:

Here we utilize Lemma 3.1 and Corollary 2.1. We divide integration $I_{\sigma\alpha}$ into two subintegrals,

$$\begin{aligned} x I_{\mu\alpha} &= \int_0^{\infty} \frac{f_{\mu}(x + \zeta; \alpha, \beta) f_{\alpha}(x + \zeta; \alpha, \beta)}{f(x + \zeta; \alpha, \beta)} dx - \int_0^{\infty} \frac{f_{\mu}(x - \zeta; \alpha, -\beta) f_{\alpha}(x - \zeta; \alpha, -\beta)}{f(x - \zeta; \alpha, -\beta)} dx \\ &= I_{\mu\alpha}^1 + I_{\mu\alpha}^2. \end{aligned}$$

$I_{\mu\alpha}^1$ is calculated first. From Corollary 2.1 and Lemma 3.2 we can find an integrable function $g_1(x)$ such that for $x \in (0, \infty)$ and all sufficiently small Δ

$$\left| \frac{f_{\mu}(x + \zeta; \alpha, \beta) f_{\alpha}(x + \zeta; \alpha, \beta)}{f(x; \alpha, \beta)} \right| \leq g_1(x).$$

Hence we can apply the dominated convergence theorem to $I_{\mu\alpha}^1$. Then

$$\lim_{\Delta \rightarrow 0} I_{\mu\alpha}^1 = \int_0^{\infty} \frac{f_{\mu}(x; 2) f_{\alpha}(x; 2)}{f(x; 2)} dx.$$

We apply the same arguments as $I_{\mu\alpha}^1$ to $I_{\mu\alpha}^2$ and obtain

$$\lim_{\Delta \rightarrow 0} I_{\mu\alpha}^2 = - \int_0^{\infty} \frac{f_{\mu}(x; 2) f_{\alpha}(x; 2)}{f(x; 2)} dx.$$

Adding $I_{\mu\alpha}^1$ and $I_{\mu\alpha}^2$, we prove our assertion.

□

References

- [1] H. Bergström, *On some expansions of stable distribution functions*, Ark. Mat., **2** (1953), pp. 375–378.
- [2] B. W. Brorsen and S. R. Yang, *Maximum likelihood estimates of symmetric stable distribution parameters*. Comm. Statist. Simul., **19** (1990), pp. 1459–1464.
- [3] W. H. DuMouchel, *Stable distributions in statistical inference 2: Information from stably distributed samples*, J. Amer. Statist. Assoc., **70** (1975), pp. 386–393.
- [4] W. H. DuMouchel, *Estimating the stable index α in order to measure tail thickness: a critique*, Ann. Statist., **11** (1983), pp. 1019–1031.
- [5] W. Feller, *An Introduction to Probability Theory and Its Applications*, Vol. 2, 2nd ed., Wiley, New York, 1971.
- [6] S. Liu and B. W. Brorsen, *Maximum likelihood estimation of a GARCH-stable model*, Journal of Applied Econometrics, **10** (1995), pp. 273–285.
- [7] M. Matsui and A. Takemura, *Some improvements in numerical evaluation of symmetric stable density and its derivatives*, Discussion Paper CIRJE-F-292, Faculty of Economics, University of Tokyo, 2004. Also available at <http://arxiv.org/abs/math.ST/0408321>.
- [8] A. V. Nagaev and S. M. Shkol'nik, *Some properties of symmetric stable distributions close to the normal distribution*, Theory of Probability and its Applications, **33** (1988), pp. 139–144.
- [9] J. P. Nolan, *Numerical computation of stable densities and distributions*, Comm. Statist. Stochastic models, **13** (1997), pp. 759–774.
- [10] J. P. Nolan, *Maximum likelihood estimation and diagnostics for stable distributions*. Lévy Processes: Theory and Applications, (O. E. Barndorff-Nielsen et al. eds.), Birkhauser, Boston, (2001), pp. 379–400.
- [11] F. Oberhettinger, *Tables of Fourier Transforms and Fourier Transforms of Distributions*, Springer-Verlag, Berlin, 1990.
- [12] V. V. Uchikin and V. M. Zolotarev, *Chance and Stability, Stable Distributions and their Applications*, Modern Probability and Statistics, Monographs, VSP BV, Utrecht, The Netherland, 1999.
- [13] V. M. Zolotarev, *One-Dimensional Stable Distributions*, Transl. of Math. Monographs, **65**, Amer. Math. Soc., Providence, RI, 1986. (Transl. of the original 1983 Russian)