

AN ADAPTATION THEORY FOR NONPARAMETRIC CONFIDENCE INTERVALS¹

By T. Tony Cai and Mark G. Low

University of Pennsylvania

A nonparametric adaptation theory is developed for the construction of confidence intervals for linear functionals. A between class modulus of continuity captures the expected length of adaptive confidence intervals. Sharp lower bounds are given for the expected length and an ordered modulus of continuity is used to construct adaptive confidence procedures which are within a constant factor of the lower bounds. In addition, minimax theory over nonconvex parameter spaces is developed.

1. Introduction. The problem of estimating a linear functional occupies a central position in nonparametric function estimation. It is most complete in the Gaussian settings:

$$(1) \quad dY(t) = f(t) dt + n^{-1/2} dW(t); \quad \frac{1}{2} \leq t \leq \frac{1}{2};$$

where $W(t)$ is standard Brownian motion and

$$(2) \quad Y(i) = f(i) + n^{-1/2} z_i; \quad i \in M;$$

where z_i are i.i.d. standard normal random variables and M is a finite or countably infinite index set. In particular, minimax estimation theory has been well developed in Ibragimov and Hasminskii (1984), Donoho and Liu (1991) and Donoho (1994).

Confidence sets also play a fundamental role in statistical inference. In the context of nonparametric function estimation variable size confidence intervals, bands and balls have received particular attention recently. For any confidence set there are two main interrelated issues which need to

Received February 2003; revised December 2003.

¹Supported in part by NSF Grant DMS-03-06576.

AMS 2000 subject classifications. Primary 62G99; secondary 62F12, 62F35, 62M99.

Key words and phrases. Adaptation, between class modulus, confidence intervals, coverage, expected length, linear functionals, minimax estimation, modulus of continuity, white noise model.

This is an electronic reprint of the original article published by the Institute of Mathematical Statistics in The Annals of Statistics, 2004, Vol. 32, No. 5, 1805-1840. This reprint differs from the original in pagination and typographic detail.

be considered together, coverage probability and the expected size of the confidence set.

One common technique for constructing confidence bands and intervals is through the bootstrap. In this context it has been noted that intervals based on the bootstrap often have poor coverage probability. See, for example, Hall (1992) and Hardle and Marron (1991). Picard and Tribouley (2000) construct adaptive confidence intervals for functions at a point using a wavelet method which achieve optimal coverage accuracy up to a logarithmic factor although in this case the issue of optimal expected length is not addressed. On the other hand Li (1989), Beran and Dumbgen (1998) and Genovese and Wasserman (2002) have constructed confidence balls which guarantee coverage probability. Closer to the present work, adaptive confidence bands have been constructed in the special case of shape restricted functions. In this context Hengartner and Stark (1995) and Dumbgen (1998) give a variable width confidence band which adapts to local smoothness while maintaining a given level of coverage probability.

In this paper we focus on the construction of confidence intervals for linear functionals which adapt to the unknown function. This adaptation problem can be made precise by considering collections of parameter spaces $\{F_j; j \in J\}$, where J is some index set. For such a collection of parameter spaces the confidence interval should have a given coverage probability over the union of the parameter spaces. Subject to this constraint the goal is to minimize the maximum expected length simultaneously over each of the parameter spaces.

For example, consider the simple and most easily explained case of two nested spaces, $F_1 \subset F$. An adaptive confidence interval must attain optimal expected length performance over both F_1 and F while satisfying a given coverage probability over F . More specifically write $\mathcal{I}_{F_1, F}$ for the collection of all confidence intervals which cover the linear functional Tf with minimum coverage probability of at least $1 - \alpha$ over the parameter space F . Denote by $L(CI; G) = \sup_{f \in G} E_f(L(CI))$ the maximum expected length of a confidence interval CI over G where $L(CI)$ is the length of the CI . Then a benchmark for the evaluation of the maximum expected length over F_1 for any $CI \in \mathcal{I}_{F_1, F}$ is given by

$$(3) \quad L(F_1; F) = \inf_{CI \in \mathcal{I}_{F_1, F}} L(CI; F_1):$$

In particular, when $F_1 = F$ set $L(F) = L(F; F)$, which gives the minimum expected length of confidence intervals of level $1 - \alpha$ over F . For convex F , Donoho (1994) constructed fixed length intervals centered at a new estimator which have length within a small constant factor of $L(F)$.

The major result in the present paper is the construction of confidence intervals which have expected length within a constant factor of $L(F_j; F)$

simultaneously over a collection of convex parameter spaces F_j where $F = [F_j]$. The construction of such intervals is general and is applicable to collections of arbitrary convex parameter spaces. It is shown in Cai and Low (2003) that in particular cases, such as collections of convex functions, the general procedure can be modified to yield simple and easily implementable procedures.

The main technical tools used in the derivation of the general adaptive confidence intervals are geometric quantities, the ordered and between class moduli of continuity which are defined as follows. For a linear functional T and parameter spaces F and G there are ordered moduli of continuity $\omega(\cdot; F; G)$ associated with the Gaussian models (1) and (2) defined by

$$(4) \quad \omega(\cdot; F; G) = \sup \{ |Tf - Tg| : \|f - g\|_k \leq \eta; f \in F; g \in G \}$$

where $\|\cdot\|_k$ is the $L_2(\frac{1}{2}; \frac{1}{2})$ function norm in the white noise model (1) and the $\|\cdot\|_2$ sequence norm over the index set M in the Gaussian model (2). As we shall give a unified treatment of both models it is convenient in the notation used throughout the paper not to distinguish the function norm and the sequence norm. It is implicit that for results concerning the white noise model (1) the notation $\|\cdot\|_k$ always refers to the L_2 function norm whereas for the sequence model (2) it always refers to the $\|\cdot\|_2$ sequence norm. When $G = F$, $\omega(\cdot; F; F)$ is the modulus of continuity over F introduced by Donoho and Liu (1991) and will be denoted by $\omega(\cdot; F)$.

For two parameter spaces F and G and a given linear functional T , the between class modulus of continuity is defined as $\omega_+(\cdot; F; G) = \max \{ \omega(\cdot; F; G); \omega(\cdot; G; F) \}$, or equivalently

$$(5) \quad \omega_+(\cdot; F; G) = \sup \{ |Tf - Tg| : \|f - g\|_k \leq \eta; f \in F; g \in G \}$$

The between class and ordered moduli were first introduced in Cai and Low (2002) in the context of adaptive estimation under mean squared error where they were shown to be instrumental in characterizing the possible degree of adaptability over two convex classes F and G in the same way that the modulus of continuity $\omega(\cdot; F)$ used by Donoho and Liu (1991) and Donoho (1994) captures the minimax difficulty of estimation over a single convex parameter space F .

The paper is organized as follows. Section 2 covers adaptation over two convex parameter spaces F_1 and F_2 where the theory is most easily understood. A lower bound based on the between class modulus as defined in (5) is given for $L(F_1; F)$ where $F = F_1 \cup F_2$. An adaptive confidence interval attaining this bound is also constructed by using the ordered modulus as given in (4). Various examples are used to illustrate the adaptation theory.

More generally let $\{F_j; j \in J\}$ be a collection of convex parameter spaces with nonempty intersections and let $F = [F_j]$. The goal is then to simultaneously minimize $L(CI; F_j)$ for confidence intervals $CI \in \mathcal{I}_F$. For each

parameter space F_j , $L(F_j; F)$ provides a lower bound on the maximum expected length over F_j for any $CI \in \mathcal{I}_F$. In Section 3 a complete treatment is given for nested F_j , possibly infinite in number. For any collection of nested convex parameter spaces a variable length confidence interval is constructed which for a given level of coverage has expected length within a constant factor of the minimum expected length simultaneously over all parameter spaces in the collection.

Section 4 treats the case of a general finite collection of convex parameter spaces. A more complicated procedure results in an interval which also has expected length within a constant factor of the minimum expected length although the constant factor now depends on the number of parameter spaces in the collection. Finally in Section 5 it is shown, by example, that the rate of growth in this constant factor as a function of the number of parameter spaces cannot in general be avoided. In addition, the adaptation theory developed in this paper is used to extend the minimax theory to a finite union of convex parameter spaces. This extension is given in Section 5.

2. Adaptation over two parameter spaces. In this section we consider adaptation over two parameter spaces. For the development of this theory, it is convenient for a given α to provide a benchmark for the maximum expected length over F_1 of confidence intervals with a given coverage probability of $1 - \alpha$ over $F = F_1 \cup F_2$, namely to provide a lower bound for $L(F_1; F)$ as defined in (3). This benchmark is given in Section 2.1 for arbitrary parameter spaces.

We give a complete treatment of adaptation when the two parameter spaces are convex. In this case adaptive intervals attaining the lower bound given in Section 2.1 are constructed. The adaptive procedure is given in Section 2.2. Examples illustrating the theory are given in Section 2.3.

It is convenient to write $a_1 \rightarrow b_1$ whenever $0 < \liminf a_1 = b_1 = \limsup a_1 < 1$, where l ranges over either a continuous or discrete index set.

2.1. Lower bound on the length of confidence intervals. The following simple two-point Normal mean problem is the basis for a surprisingly useful general lower bound on the expected length of $1 - \alpha$ level confidence intervals. We shall see later that the two-point bound is easy to apply for adaptation theory because each point can be chosen to lie in different parameter spaces. Previous work on confidence intervals for bounded Normal means as in Pratt (1961), Zeytinoglu and Mintz (1984) and Stark (1992) is useful for minimax theory but it is not applicable for general adaptation problems.

Let $X \sim N(\theta; \sigma^2)$ and suppose that $\theta^2 = f_0; 1g$ where $0 < \theta_1$. Consider the following simple statistical decision theory problem: construct confidence intervals $CI(X)$ for θ which have smallest expected length under θ_0

subject to the coverage constraint

$$P(CI(X)) \geq 1 - \alpha \quad \text{for } \alpha = 0, 1.$$

Throughout the paper set $z = \Phi^{-1}(1 - \alpha/2)$ where Φ is the cumulative density function of a standard Normal distribution. In addition write $L(CI)$ for the length of a confidence interval CI .

Proposition 1. Let $X \sim N(\mu, \sigma^2)$ and suppose that $\mu = f_0; 1g$ where $0 < 1$. Let $CI(X)$ be a $1 - \alpha$ level confidence interval for μ . Then

$$(6) \quad E_1 L(CI(X)) \geq (1 - \alpha) \left(1 + \frac{1 - \alpha}{2} z \right) +$$

for $i = 0, 1$. Moreover there exists a confidence interval which attains the lower bounds simultaneously for both $i = 0$ and $i = 1$.

Proof. It is clear that it suffices to consider confidence intervals $CI(X)$ of three possible forms: $[0; 1]$, f_0g and f_1g . The problem is then to minimize $P_0(CI(X) = [0; 1])$ subject to the constraints $P_0(CI(X) = f_1g) = 1 - \alpha$ and $P_1(CI(X) = f_0g) = 1 - \alpha$.

It follows from the Neyman-Pearson lemma that, subject to the constraint that $P_1(CI(X) = f_0g) = 1 - \alpha$,

$$P_0(CI(X) = f_0g) \leq \frac{1 - \alpha}{2} z.$$

Hence

$$\begin{aligned} E_0 L(CI(X)) &= (1 - \alpha) P_0(CI(X) = [0; 1]) \\ &= (1 - \alpha) (1 - P_0(CI(X) = f_1g) - P_0(CI(X) = f_0g)) \\ &\geq (1 - \alpha) \left(1 - \frac{1 - \alpha}{2} z \right). \end{aligned}$$

The bound for 1 follows similarly.

It is easy to see that an interval attaining the lower bound for 0 and 1 is given by

$$CI(X) = \begin{cases} f_0g; & \text{if } X \leq -z, \\ [0; 1]; & \text{if } -z < X < 0 + z, \\ f_1g; & \text{if } X \geq 0 + z, \end{cases}$$

when $-z < 0 + z$. Otherwise set

$$CI(X) = \begin{cases} f_0g; & \text{if } X \leq \frac{0 + 1}{2}, \\ f_1g; & \text{if } X > \frac{0 + 1}{2}. \end{cases}$$

In this case the confidence interval always has zero length and coverage of at least 1.

Based on the two-point bound given in Proposition 1 the following theorem gives a lower bound for infinite-dimensional Gaussian models.

Theorem 1. Let $0 < \epsilon < \frac{1}{2}$ and let F_1, F be two parameter spaces. Then

$$(7) \quad L(F_1; F) \geq \frac{1}{2} \left(1 + \frac{\epsilon}{P_n} \right) \mathbb{E}_{P_n} \left[\int F_1; F \right]$$

where $L(F_1; F)$ is defined in (3) and $\mathbb{E}_{P_n} \left[\int F_1; F \right]$ is the between class modulus as given in (5).

Proof. We shall focus on the proof for the white noise with drift model (1). The proof for the sequence model (2) is analogous. Fix $\epsilon > 0$. For any $\epsilon > 0$ there are functions $f_1 \in F_1$ and $f_2 \in F$ such that

$$\left| \int f_2 - \int f_1 \right| \geq \frac{\epsilon}{P_n} \mathbb{E}_{P_n} \left[\int F_1; F \right]$$

and such that

$$\|f_2 - f_1\|_{L_2} \leq \frac{\epsilon}{P_n}.$$

Denote by P_i the probability measure associated with the white noise process

$$dY(t) = f_i(t) dt + \frac{1}{P_n} dW(t); \quad \frac{1}{2} \leq t \leq \frac{1}{2}; i = 1, 2.$$

Let $\sigma_n = n\|f_1 - f_2\|_{L_2}^2$. Then a sufficient statistic for the family of measures $\{P_i; i = 1, 2\}$ is given by the log-likelihood ratio $S_n = \log(dP_2/dP_1)$ with

$$S_n \begin{cases} \geq N \frac{\sigma_n}{2} & \text{under } P_1, \\ \leq -N \frac{\sigma_n}{2} & \text{under } P_2. \end{cases}$$

An equivalent sufficient statistic is thus given by

$$Q_n = \frac{\int f_1 + \int f_2}{2} + \frac{\int f_2 - \int f_1}{\sigma_n} S$$

where

$$Q_n \begin{cases} \geq N \frac{(\int f_2 - \int f_1)^2}{\sigma_n} & \text{under } P_1, \\ \leq -N \frac{(\int f_2 - \int f_1)^2}{\sigma_n} & \text{under } P_2. \end{cases}$$

It follows from Proposition 1 that for any confidence interval $CI(Q_n)$ based on Q_n ,

$$E_{f_1} L(CI(Q_n)) \leq \int_{f_2}^{\int_{f_1}} \frac{\int_{f_2}^{\int_{f_1}}}{z} +$$

where $\int_{f_1}^{\int_{f_2}} = \frac{\int_{f_2}^{\int_{f_1}}}{n}$. Hence

$$E_{f_1} L(CI(Q_n)) \leq \int_{f_2}^{\int_{f_1}} (1 - \frac{p}{n} k f_2 - f_1 k_2 - z)_+ \\ + \frac{1}{n} \int_{f_1}^{\int_{f_2}} ; F_1 ; F (1 - (\frac{1}{n} - z))_+ :$$

Letting $\epsilon \rightarrow 0$, it follows that for any $\epsilon > 0$,

$$L(CI(Q_n); F_1) \leq \frac{1}{n} \int_{f_1}^{\int_{f_2}} ; F_1 ; F (1 - (\frac{1}{n} - z))_+ :$$

By the sufficiency of Q_n , it follows that for any confidence interval $CI \in \mathcal{I}_F$

$$(8) \quad L(CI; F_1) \leq \sup_{\epsilon > 0} \frac{1}{n} \int_{f_1}^{\int_{f_2}} ; F_1 ; F (1 - (\frac{1}{n} - z))_+ :$$

The theorem follows on taking $\epsilon = z$.

Remark 1. Although the primary use of this theorem is for adaptive confidence intervals, it can also be used to show that from a minimax point of view there is relatively little to gain by using variable length intervals. In the minimax setting Donoho (1994) showed that over a given convex parameter space F , fixed length confidence intervals for a linear functional Tf with coverage of at least $1 - \alpha$ must have maximum length at least $2! \left(\frac{2z}{n} ; F \right)$ and that fixed length confidence intervals can be centered on a new estimator with maximum length at most $2! \left(\frac{2z - \alpha}{n} ; F \right)$. By taking $F_1 = F$, Theorem 1 yields that the minimax expected length of a $1 - \alpha$ level confidence interval over any parameter space F satisfies

$$(9) \quad L(F) \leq \frac{1}{2} \int_{f_1}^{\int_{f_2}} ; F : :$$

This shows that for any given $\alpha < 1/2$ the optimal variable length confidence intervals must have maximum expected length at least a fixed constant factor of the length of the shortest fixed length confidence interval when the parameter space F is convex.

2.2. Adaptive confidence interval. There are at least two natural ways to define adaptive confidence intervals over a collection of convex parameter spaces $\{F_i; i = 1, \dots, k\}$. Let $F = \bigcup_{i=1}^k F_i$. Call a confidence interval $CI_2 I_{\mathcal{F}}$ adaptive over the collection $\{F_i; i = 1, \dots, k\}$ if, for all $1 \leq i \leq k$,

$$(10) \quad L(CI; F_i) \leq C_i(\alpha) + \frac{Z}{\sqrt{n}}; F_i; F;$$

where $C_i(\alpha)$ are constants depending on α only. In other words a confidence interval which adapts over the parameter spaces F_i attains the lower bound given in Theorem 1 for each i while maintaining coverage over F . We shall show that such adaptive confidence intervals can always be constructed when k is finite.

It is also reasonable, in light of the minimax discussion given above, to term a confidence interval $CI_2 I_{\mathcal{F}}$ adaptive over the collection of parameter spaces F_i if, for all $1 \leq i \leq k$,

$$(11) \quad L(CI; F_i) \leq C_i(\alpha) + \frac{Z}{\sqrt{n}}; F_i$$

where $C_i(\alpha)$ are constants depending on α only. We shall call such a confidence interval strongly adaptive. It is clear that a confidence interval which is strongly adaptive is also adaptive. However strongly adaptive confidence intervals do not always exist. Low (1997) has given examples where $L(F_1; F) = L(F_1)$, in which case strongly adaptive estimators do not exist. Other examples are given in Section 2.3 and throughout the paper. On the other hand, when $L(F_1; F) = L(F_1)$ strongly adaptive estimators do exist and any estimator which is adaptive is also strongly adaptive.

In this section the focus is on adaptation over two parameter spaces where the theory is most easily understood. For two parameter spaces $F_1 \subseteq F$, Theorem 1 gives a lower bound for the maximum expected length over F_1 of confidence intervals with guaranteed coverage over F . We now show that the lower bound can in fact be attained within a constant factor not depending on n when F_1 is convex and F is the union of F_1 and another convex set F_2 .

Let $\{F_1, F_2\}$ be a pair of convex parameter spaces with nonempty intersection and let $F = F_1 \cup F_2$. Our first objective is to construct a confidence interval for a linear functional Tf which has guaranteed coverage probability of $1 - \alpha$ over F and has maximum expected length over F_1 within a constant factor of the lower bound given in Theorem 1, namely, for any $CI_2 I_{\mathcal{F}}$,

$$(12) \quad L(CI; F_1) \leq \frac{1}{2} + \frac{Z}{\sqrt{n}}; F_1; F;$$

The construction of the adaptive confidence interval relies on the ordered modulus $!_{(n)}(F_i; F_j)$ as given in (4). For $1 \leq i, j \leq 2$, set

$$!_{i,j} = !_{\frac{Z_{=2}}{n}}(F_i; F_j) :$$

Cai and Low (2004) give an algorithm for the construction of a linear estimator $\hat{T}_{i,j}$ which has variance bounded by

$$(13) \quad \text{Var}(\hat{T}_{i,j}) \leq \frac{1}{Z_{=2}^2} !_{i,j}^2$$

and bias which satisfies

$$(14) \quad \inf_{f \in F_j} (E(\hat{T}_{i,j}) - T(f)) \leq \frac{1}{2} !_{i,j}$$

and

$$(15) \quad \sup_{f \in F_i} (E(\hat{T}_{i,j}) - T(f)) \leq \frac{1}{2} !_{i,j} :$$

We shall use the linear estimators $\hat{T}_{i,j}$ to construct a confidence interval which has guaranteed coverage probability over F and which also has expected length over F_1 within a constant factor of the lower bound given by (26). For $j = 1$ and 2 define the confidence intervals $CI_{j,}$ by

$$(16) \quad CI_{j,} = \left[\min_{i=1,2} \hat{T}_{i,j} - \frac{3}{2} !_{i,j}; \max_{i=1,2} \hat{T}_{i,j} + \frac{3}{2} !_{j,i} \right] :$$

The following result shows that the confidence interval $CI_{1,}$ attains the lower bound on the maximum expected length over F_1 given in (7) within a constant factor not depending on n and satisfies the constraint that it has the minimum coverage of 1 for all $f \in F$.

Lemma 1. Let F_1 and F_2 be convex parameter spaces with $F_1 \setminus F_2 \neq \emptyset$ and let $F = F_1 \cup F_2$. Let the interval $CI_{j,}$ be defined as in (16) for $j = 1$ and 2 . Then $CI_{j,} \subset I_{\mathbb{R}}$ and $CI_{j,}$ has expected length over F_j which satisfies

$$(17) \quad L(CI_{j,}; F_j) \leq \frac{9}{Z_{=2}} + 4 !_{+} + \frac{Z_{=2}}{n} !_{F_j; F} :$$

Lemma 1 follows from the proof of Proposition 4 given in Section 4.1.

Remark 2. Theorem 1 and Lemma 1 together show that under the conditions of Lemma 1,

$$(18) \quad L(F_1; F) \leq !_{+} + \frac{Z_{=2}}{n} !_{F_1; F} :$$

Although the interval $CI_{1;}$ has guaranteed coverage probability over F and optimal expected length over F_1 , it may not have optimal expected length over F because the expected length over F_2 is not controlled. On the other hand, by symmetry $CI_{2;}$ has guaranteed coverage probability over F and optimal expected length over F_2 . By Bonferroni, the confidence interval $CI = CI_{1;=2} \setminus CI_{2;=2}$ also has coverage probability of at least 1 and so $CI \subset I_{\mathbb{F}}$. Furthermore, it is easy to see that it has optimal expected length over both F_1 and F_2 and hence also over F . In other words the confidence interval CI is a 1 level adaptive confidence interval over F_1 and F_2 .

Proposition 2. Let F_1 and F_2 be convex parameter spaces with $F_1 \setminus F_2 \neq \emptyset$ and let $F = F_1 \cup F_2$. Let the interval $CI_{j;}$ be defined as in (16) and let $CI = CI_{1;=2} \setminus CI_{2;=2}$. Then CI is a 1 level adaptive confidence interval over F_1 and F_2 . That is, $CI \subset I_{\mathbb{F}}$ and for both $j = 1$ and 2 ,

$$(19) \quad L(F_j; F) \leq L(CI; F_j) \leq C(\cdot) L(F_j; F)$$

where $C(\cdot)$ is a constant depending only on \cdot . Consequently $L(CI; F_j) \leq \left(\frac{z_{1-\frac{\alpha}{2}}}{n}\right) L(F_j; F)$.

Remark 3. It is shown in Cai and Low (2004) that the ordered modulus is concave. It follows that, if $b \geq 1$, then for all $\alpha > 0$,

$$(20) \quad \begin{aligned} b!_+(\alpha; F; G) &= \max(b!(\alpha; F; G); b!(\alpha; G; F)) \\ &\leq \max(b!(\alpha; F; G); b!(\alpha; G; F)) \\ &\leq b!_+(\alpha; F; G). \end{aligned}$$

It then follows from the bounds given in (7) and (17) and inequality (20) that the constant $C(\cdot)$ in (19) can be taken as

$$C(\cdot) = \frac{9 + 4z_{1-\frac{\alpha}{2}}}{(1-\frac{\alpha}{2})z}.$$

2.3. Discussion. In nonparametric function estimation the goal of adaptive estimation is often framed in terms of achieving optimality results simultaneously over a collection of parameter spaces \mathbb{F}_j 's. The benchmark for success is given by how well one could do if the parameter space is completely specified. We termed any such confidence interval strongly adaptive.

So far, attention has focused on constructing adaptive confidence procedures which attain the lower bound on expected length given in Theorem 1. This bound gives the best one can do in this adaptive confidence interval problem. The lower bound however may differ quite dramatically from the

maximum expected length if the parameter space F_j is prespecified. In particular suppose, as is common, that the between class modulus of continuity is Hölderian. That is, the modulus satisfies

$$\omega_+(F_i; F_j) = C_{i,j} n^{q_{i,j}} (1 + o(1)); \quad 1 \leq i, j \leq 2;$$

for some constants $C_{i,j} > 0$ and $0 < q_{i,j} \leq 1$. Such is the case in the examples given in Section 3.2 and also in many other commonly treated problems. When the modulus $\omega_+(F; G)$ is Hölderian write $q(F; G)$ for the exponent of the modulus. That is,

$$\omega_+(F; G) = n^{q(F; G)}.$$

Also set $q(G) = q(G; G)$.

Without loss of generality, assume $q(F_1) \geq q(F_2)$. Throughout the remainder of the paper C is used to denote a generic constant which may vary from place to place and set $F = F_1 \cup F_2$. Note that $q(F_1; F) = \min\{q(F_1); q(F_1; F_2)\}$ and $q(F) = \min\{q(F_1); q(F_2); q(F_1; F_2)\}$. In this setup strongly adaptive confidence intervals exist if and only if $q(F_1; F) = q(F_1)$ or equivalently $q(F_1) \geq q(F_1; F_2)$.

There are four cases of interest.

Case 1. $q(F_2) \geq q(F_1) \geq q(F_1; F_2)$. In this case $q(F_1; F) = q(F_1)$ and strongly adaptive confidence intervals exist. These intervals have maximum expected length which can attain the same optimal rate of convergence as the minimum confidence interval over known F_1 . Specific shape restricted examples are given in Section 3.2 which illustrate this case and more general theory.

Case 2. $q(F_1; F_2) = q(F_2) < q(F_1)$. In this case $q(F_1; F) < q(F_1)$ and thus strongly adaptive confidence intervals do not exist. Adaptive confidence intervals of level $1 - \alpha$ over F_1 and F_2 have maximum expected length over F_1 which satisfies

$$(21) \quad L(CI; F_1) \sim \frac{1}{2} \omega_+^2\left(\frac{Z}{n}; F_1; F\right) n^{q(F)=2}.$$

In contrast, if it is known that $f \in F_1$, $1 - \alpha$ level confidence intervals can be constructed which satisfy

$$L(CI; F_1) \sim C n^{q(F_1)=2} \sim C n^{q(F)=2}.$$

Hence from this point of view the cost of adaptation is substantial. The rate of convergence of the maximum expected length of CI over F_1 is the same as that for the maximum expected length over F .

Example 1. Consider estimating the linear functional $Tf = f(0)$ over Lipschitz classes based on the Gaussian observations given in (1). For $0 < \frac{1}{2} < \alpha < \beta < \frac{1}{2}$, the Lipschitz function class over the interval $[a; b]$ is defined as

$$(22) \quad F(\cdot; M; [a; b]) = \{f : [\frac{1}{2}; \frac{1}{2}] \rightarrow \mathbb{R}; |f(x) - f(y)| \leq M|x - y| \text{ for } x, y \in [a; b]\}.$$

It is also convenient to write $F(\cdot; M)$ for $F(\cdot; M; [\frac{1}{2}; \frac{1}{2}])$.

Let $0 < \alpha < \beta < 1$, set $F_i = F(\cdot; M)$ for $i = 1, 2$. In this case standard calculations as, for example, outlined in Cai and Low (2002) show that $\mathbb{E}(\|F_1\|) = C n^{2\alpha-1} (1 + o(1))$ and $\mathbb{E}(\|F_1; F_2\|) = C n^{2\alpha-2} (1 + o(1))$. Hence

$$q(F_1; F) = q(F_1; F_2) = \frac{2\alpha}{2\alpha+1} < q(F_1) = \frac{2\alpha}{2\alpha+1}.$$

Case 3. $q(F_2) < q(F_1; F_2) < q(F_1)$. In this case $q(F_1; F) < q(F_1)$ and strongly adaptive confidence intervals do not exist. Any 1 -level adaptive confidence interval CI over F_1 and F_2 , must have maximum expected length of CI over F_1 satisfying

$$(23) \quad \mathbb{E}_F(L(CI; F_1)) \geq \frac{1}{2} \mathbb{E}_F \left(\frac{Z}{n} ; F_1; F \right) \geq \frac{n^{q(F_1; F_2)-2}}{n^{q(F_1)-2}}.$$

The cost of adaptation in this case is that the rate of convergence of the maximum expected length of CI over F_1 is slower than that if it is known that $f \in F_1$ but faster than for the maximum expected length over F_2 . An example for this case can be given as follows.

Example 2. Suppose that the white noise with drift process (1) is observed and that the linear functional $Tf = f(0)$. Let the Lipschitz class $F(\cdot; M; [a; b])$ be defined as above and let D be the set of all decreasing functions on $[\frac{1}{2}; \frac{1}{2}]$. Set

$$F_D(\cdot; M_1; \cdot; M_2) = F(\cdot; M_1; [\frac{1}{2}; 0]) \setminus F(\cdot; M_2; [0; \frac{1}{2}]) \setminus D.$$

Let $F_1 = F_D(\cdot; M_1; \cdot; M_2)$ and $F_2 = F_D(\cdot; N_1; \cdot; N_2)$ with $1 > \alpha > \beta > 0$. Then as in Cai and Low (2002) it is easy to check that

$$(24) \quad \begin{aligned} \mathbb{E}(\|F_1\|) &= C n^{2\alpha-1} (1 + o(1)); \\ \mathbb{E}(\|F_2\|) &= C n^{2\beta-1} (1 + o(1)); \\ \mathbb{E}(\|F_1; F_2\|) &= C n^{2\alpha-2} (1 + o(1)); \\ \mathbb{E}(\|F_2; F_1\|) &= C n^{2\beta-2} (1 + o(1)); \end{aligned}$$

Note that in this case $! ("; F_1; F_2) \notin ! ("; F_2; F_1)(1 + o(1))$. Since $\alpha_1 > \alpha_2$, it then follows from (24) that

$$q(F_1; F) = q(F_1; F_2) = \frac{2\alpha_2}{2\alpha_2 + 1}.$$

Hence $0 < q(F_2) < q(F_1; F) < q(F_1) < 1$.

Case 4. $q(F_1; F_2) < q(F_2) < q(F_1)$. In this case, strongly adaptive confidence intervals do not exist and the cost of adaptation is extraordinary. If f is known to be in F_i , one can attain the rate of convergence $n^{q(F_i)=2}$ for them aximum expected length of the optimal level confidence interval over F_i . Without the information level adaptive confidence intervals over F_1 and F_2 must have maximum expected length over F_i at least of order $n^{-q(F_1; F_2)=2}$. An example is given below.

Example 3. Once again consider the white noise model with $Tf = f(0)$. Let

$$F(\alpha_1; M_1; \alpha_2; M_2) = F(\alpha_1; M_1; [\frac{1}{2}; 0]) \setminus F(\alpha_2; M_2; [0; \frac{1}{2}])$$

and consider $0 < \alpha_2 < \alpha_1 < 1$ and $0 < \alpha_1 < \alpha_2 < 1$. Set $F_1 = F(\alpha_1; M_1; \alpha_2; M_2)$ and $F_2 = F(\alpha_1; N_1; \alpha_2; N_2)$. Standard calculations show that $! ("; F_1) = C n^{2\alpha_1=(2\alpha_1+1)}(1 + o(1))$ and $! ("; F_2) = C n^{2\alpha_2=(2\alpha_2+1)}(1 + o(1))$. The between class modulus is given as

$$(25) \quad ! ("; F_1; F_2) = C n^{2\alpha_2=(2\alpha_2+1)}(1 + o(1))$$

where $\alpha = \max(\alpha_1; \alpha_2)$.

When $\alpha_1 > \alpha_2$, the quantity in (25) equals α_1 and hence

$$q(F_1; F_2) = \frac{2\alpha_1}{2\alpha_1 + 1}.$$

Therefore in this case $q(F_1; F_2) < \min(q(F_1); q(F_2))$:

3. Adaptation over nested parameter spaces. Section 2 gave the adaptation theory for two convex parameter spaces. This theory can be extended to more general collections of parameter spaces. In this section the focus is on adaptation over a collection of a finite or countably infinite number of nested convex parameter spaces, $F_1 \supset F_2 \supset \dots \supset F_k$, where in the case of $k = 1$, F_1 denotes $\bigcup_{i=1}^k F_i$. The objective is, for a given linear functional Tf , to construct variable length confidence intervals which have coverage probability of at least $1 - \alpha$ over F_k and which simultaneously minimize the expected length over each of the parameter spaces F_j . A target for these

expected lengths has been provided by the lower bound given in Theorem 1, namely

$$(26) \quad L(F_j; F_k) \geq \frac{1}{2} \left(1 + \frac{Z_{\frac{\alpha}{2}}}{\sqrt{n}}; F_j; F_k \right)$$

where $1 + \frac{Z_{\frac{\alpha}{2}}}{\sqrt{n}}; F_j; F_k$ is the between class modulus as given in (5).

The major result of this section is to show that adaptive confidence intervals exist and to construct such adaptive intervals. As in Section 2.2 the construction of these adaptive confidence procedures relies on the ordered modulus $1 + \frac{Z_{\frac{\alpha}{2}}}{\sqrt{n}}; F_i; F_j$ as given in (4). For $1 \leq i, j \leq k$ set $l_{i,j} = 1 + \frac{Z_{\frac{\alpha}{2}}}{\sqrt{n}}; F_i; F_j$ and let $\hat{T}_{i,j}$ be linear estimators with variances and biases bounded as in (13)–(15).

The confidence procedure is built in two steps. In the first step for each $1 \leq j \leq k$ an interval is constructed which controls the coverage probability over F_k and which also has expected length over F_j within a constant factor of the lower bound given by (26). In the second step these intervals are combined to create a single interval which maintains coverage while simultaneously attaining an expected length over every F_j within a fixed constant factor of the lower bound given in (26).

For the first step define the confidence intervals CI_j as follows. For $1 \leq j \leq k$ set $l_j = 1 + \frac{Z_{\frac{\alpha}{2}}}{\sqrt{n}}; F_j; F_k$ and define CI_j by

$$(27) \quad CI_j = \frac{\hat{T}_{j,k} + \hat{T}_{k,j}}{2} - f(\hat{T}_{j,k} - \hat{T}_{k,j})_+ + 2 l_j g; \\ \frac{\hat{T}_{j,k} + \hat{T}_{k,j}}{2} + f(\hat{T}_{j,k} - \hat{T}_{k,j})_+ + 2 l_j g;$$

Lemma 2 shows that these intervals have guaranteed coverage over F_k and near optimal expected length over F_j .

Remark 4. This interval is designed for $0 < \alpha \leq 0.2$. If $0.2 < \alpha \leq 0.5$ all subsequent results hold with minor modifications, as noted in later remarks, when the interval is replaced by

$$(28) \quad CI_j = \frac{\hat{T}_{j,k} + \hat{T}_{k,j}}{2} - f(\hat{T}_{j,k} - \hat{T}_{k,j})_+ + 3 l_j g; \\ \frac{\hat{T}_{j,k} + \hat{T}_{k,j}}{2} + f(\hat{T}_{j,k} - \hat{T}_{k,j})_+ + 3 l_j g;$$

Lemma 2. For $0 < \alpha < 0.2$, the confidence interval CI_j defined in (27) has coverage probability of at least $1 - \frac{\alpha}{2}$ for all $f \in F_k$ and satisfies

$$(29) \quad \mathbb{E} L(CI_j; F_j) \leq 2 \int_{-\infty}^{\infty} \frac{1}{2} z^2 e^{-\frac{z^2}{2}} + \frac{4}{2} \int_{-\infty}^{\infty} \frac{1}{8} z^2 e^{-\frac{z^2}{2}} + 4 \int_{-\infty}^{\infty} z e^{-\frac{z^2}{2}} dz$$

Remark 5. For $0.2 < \alpha < 0.5$ the interval given in (28) satisfies the same coverage but has expected length bounded by $10 \int_{-\infty}^{\infty} \frac{1}{n} z e^{-\frac{z^2}{2}} dz$.

In the following proof, and throughout the rest of the paper, write Z for a standard Normal random variable.

Proof of Lemma 2. Lemma 2 gives a bound on both coverage probability and expected length. First consider coverage probability. It is easy to see that the interval CI_j contains the interval CI_j defined as

$$(30) \quad CI_j = [\hat{T}_{k;j} - 2j; \hat{T}_{j;k} + 2j]$$

where the interval CI_j is taken to be the empty set whenever the left endpoint of the above interval is larger than the right endpoint. First note that for $f \in F_k$, $\mathbb{E} \hat{T}_{k;j} = T f - \frac{1}{2} j_{k;j}$ and that $\mathbb{E} \hat{T}_{j;k} = T f + \frac{1}{2} j_{j;k}$. Let

$$Z_{k;j} = \frac{\hat{T}_{k;j} - T f - (1/2) j_{k;j}}{j_{k;j}^{1/2}},$$

$$Z_{j;k} = \frac{\hat{T}_{j;k} - T f + (1/2) j_{j;k}}{j_{j;k}^{1/2}}.$$

Then for any $f \in F_k$ it follows from (14) and (15) that $Z_{k;j}$ has a Normal distribution with mean less than or equal to zero and variance bounded by 1, and $Z_{j;k}$ has a Normal distribution with mean greater than or equal to zero and variance bounded by 1. Note that $j = \max(j_{k;j}, j_{j;k})$. Hence for $f \in F_k$,

$$P(T f \notin CI_j) = P(T f \notin CI_j)$$

$$P(Z_{k;j} > \frac{j}{j_{k;j}^{1/2}}) \leq \frac{1}{2} e^{-\frac{j}{2}} + P(Z_{j;k} < -\frac{j}{j_{j;k}^{1/2}}) \leq \frac{1}{2} e^{-\frac{j}{2}}$$

$$2P(Z > \frac{3}{2}) \leq \frac{3}{2} e^{-\frac{9}{8}} \leq \frac{3}{2} e^{-\frac{3}{2}}$$

Note that for a fixed $\alpha > 1$, it is easy to verify that $g(z) = P(Z > z) = P(Z > \alpha z)$ is a strictly decreasing function of z for $z > 0$ and for $\alpha = 0.2$,

$$2P(Z > \frac{3}{2} z) \leq \frac{2}{7} :$$

Hence, $P(Tf \notin CI_j) \leq \frac{2}{7}$ and so the claim of the required coverage probability has been established.

Now turn to the bound on expected length given in (29) for which the following technical lemma is needed.

Lemma 3. Let $X \sim N(\mu; \sigma^2)$ with $\mu \geq 0$ and $0 < \sigma < \mu$. Then

$$(31) \quad E X \mathbb{1}(X > 0) = \mu - \frac{\sigma^2}{2\mu} + \frac{\sigma^2}{2} \exp\left(-\frac{\mu^2}{2\sigma^2}\right) :$$

Proof. It is easy to check by taking partial derivatives that $E X \mathbb{1}(X > 0)$ is an increasing function of both μ and σ . Hence

$$E_{\mu, \sigma} X \mathbb{1}(X > 0)$$

$$\begin{aligned} &= \frac{1}{\sigma} \int_0^\infty x \exp\left(-\frac{(x-\mu)^2}{2\sigma^2}\right) dx \\ &= \frac{1}{\sigma} \int_0^\infty x \exp\left(-\frac{(x-\mu)^2}{2\sigma^2}\right) dx + \frac{\sigma}{2} \int_0^\infty y \exp\left(-\frac{y^2}{2}\right) dy \\ &= \mu - \frac{\sigma^2}{2\mu} + \frac{\sigma^2}{2} \exp\left(-\frac{\mu^2}{2\sigma^2}\right) : \end{aligned}$$

Now note that for $f \in F_j$,

$$E(\hat{T}_{j;k} - \hat{T}_{k;j}) = \frac{1}{j} \text{ and } \text{Var}(\hat{T}_{j;k} - \hat{T}_{k;j}) = \frac{4}{j^2} ;$$

and so from Lemma 3 it follows that

$$(32) \quad E(\hat{T}_{j;k} - \hat{T}_{k;j})_+ = \frac{1}{2j} + \frac{2}{j^2} \exp\left(-\frac{1}{8j^2}\right)$$

and hence (29) is satisfied.

Lemma 2 shows that the interval CI_j has guaranteed coverage over F_k and near optimal expected length over F_j . Before turning to the construction of an adaptive confidence interval we state a simple preliminary lemma. The proof is straightforward and not given here.

Lemma 4. Let $0 < \mu_1 < \mu_2 < \dots < \mu_k$ be a sequence of monotonically increasing positive numbers. Then there exists a unique subsequence $j_1 < j_2 < \dots < j_m$ with $j_m = k$, such that for all $1 \leq i \leq m$,

$$(33) \quad \mu_{j_i} \geq 2\mu_{j_{i-1}} \text{ and } \mu_{j_i} < 2\mu_{j_{i-1}} \text{ for all } j_{i-1} < j < j_i$$

where we set $j_0 = 0$ and $\mu_0 = 0$.

The construction of the adaptive confidence interval proceeds as follows. Once again for $1 \leq j \leq k$, set $J_j = \lfloor \frac{Z_{\alpha/2}}{n} \rfloor; F_j; F_k$. Let $j_1 < j_2 < \dots < j_m < k$ be the subsequence satisfying (33). Let \hat{j} be the index of the shortest interval among all the CI_{j_i} for $1 \leq i \leq m$. More precisely,

$$\hat{j} = \arg \min_{j_i; 1 \leq i \leq m} L(CI_{j_i});$$

Then the adaptive confidence interval for Tf is defined by

$$(34) \quad CI = CI_{\hat{j}};$$

The following theorem shows that CI is a $1 - \alpha$ level adaptive confidence interval over the collection $\{F_j; j = 1, \dots, k\}$.

Theorem 2. The confidence interval CI defined in (34) has coverage probability of at least $1 - \alpha$ for all $f \in F_k$, that is, $CI \subset I_{f, F_k}$ and satisfies

$$(35) \quad L(F_j; F_k) - L(CI; F_j) \leq \frac{16Z_{\alpha/2}}{(1 - \alpha)Z} L(F_j; F_k)$$

simultaneously for all $1 \leq j \leq k$. Moreover,

$$(36) \quad L(CI; F_{j_i}) \leq 8 \lfloor \frac{Z_{\alpha/2}}{n} \rfloor; F_{j_i}; F_k$$

for all $1 \leq i \leq m$, and for all $1 \leq j \leq k$

$$(37) \quad L(CI; F_j) \leq 16 \lfloor \frac{Z_{\alpha/2}}{n} \rfloor; F_j; F_k :$$

The proof of Theorem 2 rests on the following important technical lemma. Recall that Lemma 2 gives a lower bound on coverage over F_k and an upper bound on expected length over F_j . Lemma 5 shows, in a precise way, that if CI_j has a large expected length it must have high coverage probability.

Lemma 5. If $f \in F_k$ and

$$P(Tf \notin CI_j) > 2P(Z \leq \frac{1}{4}(\alpha + 3)Z_{\alpha/2});$$

then

$$E(\hat{T}_{j,k} - \hat{T}_{k,j}) \leq \frac{1}{2} Z_{\alpha/2};$$

Proof. First note that

$$\begin{aligned} P(Tf \notin CI_j) &= P(Tf \leq \frac{\hat{T}_{j,k} + \hat{T}_{k,j}}{2} - \hat{T}_{j,k} \text{ or } Tf \geq \frac{\hat{T}_{j,k} + \hat{T}_{k,j}}{2} + \hat{T}_{k,j} + 2Z_{\alpha/2}) \\ &= P(Tf \leq \frac{\hat{T}_{j,k} + \hat{T}_{k,j}}{2} - \hat{T}_{j,k}) + P(Tf \geq \frac{\hat{T}_{j,k} + \hat{T}_{k,j}}{2} + \hat{T}_{k,j} + 2Z_{\alpha/2}) : \end{aligned}$$

Now note that

$$\frac{1}{2}j - \frac{1}{2}E(\hat{T}_{jk} - \hat{T}_{k;j}) = E\frac{\hat{T}_{jk} + \hat{T}_{k;j}}{2} - Tf - \frac{1}{2}j + \frac{1}{2}E(\hat{T}_{jk} - \hat{T}_{k;j}):$$

Let $X = \frac{\hat{T}_{jk} + \hat{T}_{k;j}}{2} - Tf - \hat{T}_{jk} - \hat{T}_{k;j} + 2j$. Suppose that

$$E(\hat{T}_{jk} - \hat{T}_{k;j}) > j:$$

Then

$$E(X) = \frac{1}{2}(j+3)j \quad \text{and} \quad \text{Var}(X) = \frac{4}{z_{=2}^2}j^2:$$

Hence

$$\begin{aligned} P(Tf - \frac{\hat{T}_{jk} + \hat{T}_{k;j}}{2} - \hat{T}_{jk} - \hat{T}_{k;j} + 2j) = P(X \leq 0) \\ P(Z \leq \frac{1}{4}(j+3)z_{=2}) : \end{aligned}$$

Similarly,

$$P(Tf - \frac{\hat{T}_{jk} + \hat{T}_{k;j}}{2} + (\hat{T}_{jk} - \hat{T}_{k;j} + 2j)) = P(Z \leq \frac{1}{4}(j+3)z_{=2}) :$$

Hence,

$$P(Tf \notin CI_j) = 2P(Z \leq \frac{1}{4}(j+3)z_{=2}):$$

Proof of Theorem 2. Note that it suffices to prove (36) since (37) follows immediately from (36) and (35) is a direct consequence of (20), (7) and (37). For (36) assume without loss of generality that $j \geq j_{-1}$ for all $1 \leq j \leq k$; otherwise we can work along the subsequence. First note that since CI is the shortest of all the CI_j confidence intervals Lemma 2 yields that the expected length of CI satisfies

$$\begin{aligned} L(CI; F_j) &= L(CI_j; F_j) \\ (38) \quad &\leq \frac{1}{2}z_{=2} + P\frac{4}{2z_{=2}} \exp\left[-\frac{1}{8}z_{=2}^2 + 4j\right] \\ &\leq 8j: \end{aligned}$$

Now turn to the proof of coverage. Note that

$$\begin{aligned} P(Tf \notin CI) &= \sum_{j=1}^{X^k} P(Tf \notin CI_j \mid \hat{j} = j) \\ (39) \quad &\leq \sum_{j=1}^{X^k} \inf_{j=1} P(Tf \notin CI_j); P(\hat{j} = j)g: \end{aligned}$$

For $l = 0$, denote $d(l) = 2P(\mathbb{Z} \leq \frac{1}{4}(l+6)z_{=2})$. Note that $d(0) = 2P(\mathbb{Z} \leq \frac{3}{2}z_{=2}) = \frac{2}{7}$. For $l = 1$ let

$$(40) \quad A_1 = \{j : d(l) < P(Tf \neq C I_j) \mid d(l-1)g\}$$

and let $j(l) = \min\{j : j \in A_1\}$. Note that it follows from Lemma 2 that $\sup_{l \geq 1} A_1 = \{j \mid j \geq 1\}$. Then by Lemma 5

$$(41) \quad E(\hat{T}_{j(l),k} - \hat{T}_{k;j(l)}) = (l+3)j(l) :$$

Note that $\text{Var}(\hat{T}_{j(l),k} - \hat{T}_{k;j(l)}) = \frac{4}{z_{=2}^2} j(l)^2$, so

$$P(L(C I_{j(l)}) > 4j(l)) = P(\hat{T}_{j(l),k} - \hat{T}_{k;j(l)} > 2(l+1)j(l)) \\ = P(\mathbb{Z} \leq (\frac{5}{2} - \frac{1}{2}l)z_{=2}) :$$

Since $j \geq 2j-1$, it follows that, for any integer $m > 0$,

$$P(\hat{j} \geq j(l) + m) = P(L(C I_{j(l)}) > 4j(l) + m) \\ = P(L(C I_{j(l)}) > 4^{m/2} j(l)) \\ = P(\mathbb{Z} \leq (2^{m/2} - \frac{5}{2} - \frac{1}{2}l)z_{=2}) \\ = 1_m :$$

Let $j = \min\{j(l) : l = 1, \dots, 8\}$. For $m = 3$ and $l = 1, \dots, 8$, $1_m = P(\mathbb{Z} \leq \frac{1}{2}(11-l)z_{=2})$. If $j = j(1)$, then

$$\sum_{j=j}^{\infty} \min_{j=j} P(Tf \neq C I_j); P(\hat{j} = j)g = d(0) + d(0) + d(0) + \dots \\ = \frac{6}{7} + P(\mathbb{Z} \leq 5z_{=2}) :$$

Similarly, if $j = j(l)$ for some $2 \leq l \leq 8$, then

$$\sum_{j=j}^{\infty} \min_{j=j} P(Tf \neq C I_j); P(\hat{j} = j)g = d(l-1) + d(0) + d(0) + \dots \\ = \frac{6}{7} + P(\mathbb{Z} \leq 5z_{=2}) :$$

Hence

$$(42) \quad \sum_{j=j}^{\infty} \min_{j=j} P(Tf \neq C I_j); P(\hat{j} = j)g = \frac{6}{7} + P(\mathbb{Z} \leq 5z_{=2}) :$$

The following simple lemma can be used to bound $P(\mathbb{Z} \leq 5z_{=2})$.

Lemma 6. Let $Z \sim N(0, 1)$ and let $a > 0$ and $b > 0$ be two constants. Then

$$P(Z \leq a + b) \leq \exp(-ab + \frac{1}{2}b^2)P(Z > a):$$

Applying Lemma 6 with $a = z_{=2}$ and $b = 4z_{=2}$, it follows that

$$P(Z \leq 5z_{=2}) = P(Z \leq z_{=2} + 4z_{=2}) \\ \leq \exp(-12z_{=2}^2) \leq \frac{1}{2} - \frac{1}{14}:$$

Therefore

$$P(Tf \notin CI) \leq \sum_{j=1}^X m \inf_{j \in A_1} P(Tf \notin CI_j); P(\hat{J} = j)g \\ \leq \frac{13}{14} + \sum_{j \in A_1} m \inf_{j \in A_1} P(Tf \notin CI_j); P(\hat{J} = j)g:$$

For $l \geq 9$, let m_1 be the smallest integer satisfying $2^{m_1} \geq \frac{1}{4}(3l+7)$. Then $m_1 \leq \log_2(3l+7) + 1$. Recall that for $j \in A_1$, $P(Tf \notin CI_j) \leq 2P(Z \leq \frac{1}{4}(l+3)z_{=2})$. Now note that

$$P(\hat{J} = j(l+m_1)) \leq \frac{1}{2^{m_1}} P(Z \leq \frac{1}{4}(l+3)z_{=2}):$$

So, for $l \geq 9$,

$$\sum_{j \in A_1} m \inf_{j \in A_1} P(Tf \notin CI_j); P(\hat{J} = j)g \\ \leq m_1 \cdot 2P(Z \leq \frac{1}{4}(l+3)z_{=2}) + \frac{1}{2^{m_1}} \\ (2m_1 + 1)P(Z \leq \frac{1}{4}(l+3)z_{=2}):$$

So

$$\sum_{j \in A_1} m \inf_{j \in A_1} P(Tf \notin CI_j); P(\hat{J} = j)g \\ \leq \sum_{j \in A_1} (2 \log_2(3l+7) + 1)P(Z \leq \frac{1}{4}(l+3)z_{=2}):$$

Lemma 6 yields

$$P(Z \leq \frac{1}{4}(l+3)z_{=2}) \leq P(Z \leq z_{=2} + \frac{1}{4}(l-1)z_{=2}) \\ \leq \exp(-\frac{1}{4}(l-1) + \frac{1}{32}(l-1)^2 z_{=2}^2) \leq \frac{1}{2}:$$

Hence,

$$\begin{aligned} & \sum_{j=9}^{\infty} \inf_{T \in \mathcal{C}(I_j)} P(\hat{J} = j) g \\ & \leq \sum_{j=9}^{\infty} \frac{1}{2} (2 \log_2(31 + 7) - 1) \exp \left(-\frac{1}{4} (1 - 1) + \frac{1}{32} (1 - 1)^2 \right) z_{=2}^2 \\ & \leq \sum_{j=8}^{\infty} \frac{1}{2} (2 \log_2(31 + 10) - 1) \exp \left(-\frac{z_{=2}^2}{32} \right) \exp \left(-\frac{z_{=2}^2}{4} \right) 1 : \end{aligned}$$

It is easy to see that for $l \geq 8$, $(2 \log_2(31 + 10) - 1) \exp \left(-\frac{(z_{=2}^2/32)^2}{1} \right)$ is strictly decreasing and

$$(2 \log_2(31 + 10) - 1) \exp \left(-\frac{z_{=2}^2}{32} \right) \geq \frac{1}{2} :$$

So,

$$\sum_{j=9}^{\infty} \inf_{T \in \mathcal{C}(I_j)} P(\hat{J} = j) g \leq \frac{1}{4} \sum_{j=8}^{\infty} \exp \left(-\frac{z_{=2}^2}{4} \right) 1 = \frac{1}{14} :$$

Hence,

$$P(T \notin \mathcal{C}(I)) \leq \frac{13}{14} + \frac{1}{14} = 1 :$$

3.1. Adaptation over nearly nested parameter spaces. In some common cases of interest such as Hölder spaces, Sobolev spaces and Besov spaces, the parameter spaces are not exactly nested, but have nested structure in terms of the moduli of continuity. Theorem 2 can be generalized to such nearly nested parameter spaces.

Denote by $\mathcal{C}(\text{Hull}(F))$ the convex hull of a parameter set F . Let F_i , $i = 1, \dots, k$, be convex parameter spaces and for any integer $1 \leq m \leq k$ let $G_m = \bigcap_{i=1}^m F_i$. Suppose the following condition, which is trivially satisfied if F_i are nested, holds.

Condition C. For $1 \leq j \leq k$ and some constants $C_2, C_1 > 0$,

$$\|\cdot\|_{\mathcal{C}(\text{Hull}(G_j)); \mathcal{C}(\text{Hull}(G_k))} \leq C_1 \|\cdot\|_{G_j; G_k} \leq C_2 \|\cdot\|_{F_j; F_k}$$

and

$$\|\cdot\|_{\mathcal{C}(\text{Hull}(G_k)); \mathcal{C}(\text{Hull}(G_j))} \leq C_1 \|\cdot\|_{G_k; G_j} \leq C_2 \|\cdot\|_{F_k; F_j}$$

for all $0 < \epsilon < \epsilon_0$.

Similarly to the nested case for $1 \leq i, j \leq k$, set $\hat{\theta}_{i,j}^0 = ! \left(\frac{Z_{i,j} = 2}{n}; C \text{ Hull}(G_i); C \text{ Hull}(G_j) \right)$, and once again Cai and Low (2004) give a construction of linear estimators $\hat{T}_{i,j}^0$ which have variance bounded by

$$\text{Var}(\hat{T}_{i,j}^0) \leq \frac{1}{Z_{i,j}^2} !_{i,j}^{0,2}$$

and bias which satisfies

$$\inf_{f \in \mathcal{F}_j} (E(\hat{T}_{i,j}^0) - T(f)) \leq \frac{1}{2} !_{i,j}^0$$

and

$$\sup_{f \in \mathcal{F}_i} (E(\hat{T}_{i,j}^0) - T(f)) \leq \frac{1}{2} !_{i,j}^0.$$

Set $\hat{\theta}_j^0 = ! \left(\frac{Z_j = 2}{n}; C \text{ Hull}(G_j); C \text{ Hull}(G_k) \right)$ and define the confidence intervals CI_j as earlier. When $0 < \alpha \leq 0.2$, let

$$(43) \quad CI_j = \begin{cases} \frac{\hat{T}_{j,k}^0 + \hat{T}_{k,j}^0}{2} - f(\hat{T}_{j,k}^0 - \hat{T}_{k,j}^0)_+ + 2 \hat{\theta}_j^0 g; \\ \frac{\hat{T}_{j,k}^0 + \hat{T}_{k,j}^0}{2} + f(\hat{T}_{j,k}^0 - \hat{T}_{k,j}^0)_+ + 2 \hat{\theta}_j^0 g \end{cases}$$

and when $0.2 < \alpha \leq 0.5$ let

$$(44) \quad CI_j = \begin{cases} \frac{\hat{T}_{j,k}^0 + \hat{T}_{k,j}^0}{2} - f(\hat{T}_{j,k}^0 - \hat{T}_{k,j}^0)_+ + 3 \hat{\theta}_j^0 g; \\ \frac{\hat{T}_{j,k}^0 + \hat{T}_{k,j}^0}{2} + f(\hat{T}_{j,k}^0 - \hat{T}_{k,j}^0)_+ + 3 \hat{\theta}_j^0 g \end{cases}$$

Following the argument given in the nested case let $\hat{\theta}_{j_i}^0$ be a subsequence of $\hat{\theta}_j^0$ satisfying (33) and let $\hat{j} = \arg \min_{j_i \in 1 \leq i \leq m} L(CI_{j_i})$ be the index of the shortest interval along the subsequence and define the adaptive confidence interval for $T(f)$ by

$$(45) \quad CI = CI_{\hat{j}}.$$

As stated precisely in the following result this confidence interval is adaptive over the parameter spaces $\mathcal{F}_j : j = 1, \dots, k$.

Proposition 3. Suppose Condition C holds. Then the confidence interval CI_j defined in (45) has coverage probability of at least $1 - \alpha$ for all $f \in \mathcal{F} = \bigcup_{j=1}^k \mathcal{F}_j$ and satisfies the lower bound on expected length,

$$(46) \quad L(\mathcal{F}_j; F) \leq L(CI; F_j) \leq C(\alpha) L(\mathcal{F}_j; F);$$

simultaneously for all $1 \leq j \leq k$, where the constant $C(\alpha)$ only depends on α and is independent of k . In other words, $L(CI; F_j) \leq ! \left(\frac{Z_j = 2}{n}; \mathcal{F}_j; F \right)$ for all $1 \leq j \leq k$.

We omit the proof of Proposition 3 since it essentially follows a similar path to that of Theorem 2.

3.2. Examples. Theorem 2 and Proposition 3 have established general adaptation results for collections of nested or nearly nested parameter spaces. In this section a couple of examples are given which illustrate this general theory.

Suppose that we observe the white noise with drift process (1) and that the linear functional is point evaluation. For convenience take $Tf = f(0)$. Let D be the set of all decreasing functions on $[\frac{1}{2}, \frac{1}{2}]$ and let $F_D(\cdot; M) = F(\cdot; M) \setminus D$ be the collection of monotonically decreasing Lipschitz functions where $F(\cdot; M)$ is the Lipschitz class defined in (22).

For integer $j \geq 1$ let $M_j = 2^{j(2+1)} \frac{1}{n}$ and let $G = \bigcap_{j=1}^{\infty} F_D(\cdot; M_j)$. Standard calculations as in, for example, Donoho and Liu (1987), yield

$$(47) \quad \begin{aligned} \mathbb{P}(\cdot; F_D(\cdot; M); G) &= \mathbb{P}(\cdot; G; F_D(\cdot; M)) \\ &= (2+1)^{1=(2+1)} M^{1=(2+1)} n^{2=(2+1)} \end{aligned}$$

for $M = (2+1)^{1=2}$. Let $\mathbb{P}_j = \mathbb{P}(\cdot; F_D(\cdot; M_j); G)$. Then it is easy to see that $\mathbb{P}_{j+1} = 2 \mathbb{P}_j$ and hence the adaptive confidence interval given in (34) has coverage probability over G of at least 1 and satisfies

$$(48) \quad L(CI; F_D(\cdot; M_j)) \leq 6(2+1)^{1=(2+1)} M_j^{1=(2+1)} Z_{\alpha=2}^{2=(2+1)} n^{2=(2+1)};$$

Furthermore, for any $M > 0$,

$$(49) \quad L(CI; F_D(\cdot; M)) \leq 12(2+1)^{1=(2+1)} M^{1=(2+1)} Z_{\alpha=2}^{2=(2+1)} n^{2=(2+1)}$$

for all sufficiently large n .

Another common problem in function estimation is to adapt over smoothness classes. For fixed $M \geq 0$, the classes $F_D(\cdot; M_1) \supset F_D(\cdot; M_2)$ whenever $0 < \alpha_2 < \alpha_1 \leq 1$. Let $G^0 = \bigcap_{\alpha=0}^1 F_D(\cdot; M)$. Then once again standard calculations yield

$$(50) \quad \begin{aligned} \mathbb{P}(\cdot; F_D(\cdot; M); G^0) &= \mathbb{P}(\cdot; G^0; F_D(\cdot; M)) \\ &= (2+1)^{1=(2+1)} M^{1=(2+1)} n^{2=(2+1)}; \end{aligned}$$

Now let $1 = \alpha_1 > \alpha_2 > \dots$ be the sequence such that

$$1 + \frac{Z_{\alpha=2}}{n} F_D(\cdot; M_{j+1}); G^0 = 2 \left(1 + \frac{Z_{\alpha=2}}{n} F_D(\cdot; M_j); G^0 \right);$$

Then the adaptive confidence interval given in (34) has coverage probability over G^0 of at least 1 and satisfies

$$(51) \quad L(CI; F_D(\cdot; M_j)) \leq 6(2_{j+1})^{1=(2_{j+1})} M^{1=(2_{j+1})} Z_{\alpha=2}^{2_{j+1}=(2_{j+1})} n^{2_{j+1}=(2_{j+1})};$$

Furthermore, for any $0 < \epsilon < 1$,

$$(52) \quad L(CI; F_D(\cdot; M)) \leq 12(2 + 1)^{1=(2+1)} M^{1=(2+1)} Z_{=2}^{2=(2+1)} n^{-(2+1)}$$

for all sufficiently large n .

4. Adaptation over a general collection of convex parameter spaces. Section 3 focused on collections of nested parameter spaces. It has been shown that the between class modulus of continuity completely characterizes the optimal expected length of adaptive confidence intervals. One particularly interesting feature of the nested case is that the optimal expected length of the confidence intervals does not depend on the number of parameter spaces in the collection.

The nested case, although interesting, is somewhat special. In this section general finite collections of convex parameter spaces are considered. In this general setting the theory is more complicated and in general the number of parameter spaces, say k , may also play a role in the optimal expected length of adaptive confidence intervals. For a fixed and finite number of parameter spaces the optimal expected length of adaptive intervals is still within a constant factor of the between class modulus of continuity. However the constant factor in this case can depend on the number of parameter spaces. We construct adaptive confidence intervals which show that this constant factor does not grow faster than $\sqrt{\log k}$ and we give an example which shows that this factor is sometimes necessary.

Let $\{F_j : j = 1, \dots, k\}$ be a collection of convex spaces with nonempty intersections, that is, $F_i \cap F_j \neq \emptyset$ for all i, j . The objective is to construct an adaptive confidence interval for a linear functional Tf which has guaranteed coverage probability of 1 over $G = \bigcap_{j=1}^k F_j$ and rate optimal expected length over each of the parameter spaces F_j .

The adaptive confidence interval given in this section differs substantially from that given in the nested case. However, the general strategy for constructing adaptive confidence intervals in this setup is similar to that of the nested case. In particular, a key step is to first construct an interval which has optimal expected length over one of the parameter spaces while attaining coverage probability over the union of the parameter spaces.

4.1. Constrained optimal expected length confidence intervals. As mentioned above, it is convenient to construct a confidence interval which has shortest possible expected length over a given F_j while maintaining coverage probability over $G = \bigcap_{j=1}^k F_j$.

First note that for any confidence interval $CI \subset \mathbb{R}$, Theorem 1 yields a target for the expected length

$$(53) \quad L(CI; F_j) \leq \frac{1}{2} \left(1 + \frac{Z}{n} \right) \mathbb{E} \left[\frac{Z}{n} ; F_j ; G \right]$$

As in Section 2.2, for $1 \leq i, j \leq k$ set $I_{ij} = I(\frac{z_{\alpha/2}}{n}; F_i; F_j)$ and let \hat{T}_{ij} be a linear estimator which has variance bounded by $\frac{1}{z_{\alpha/2}^2} I_{ij}^2$ and bias which satisfies

$$(54) \quad \inf_{f \in F_j} (E(\hat{T}_{ij}) - T f) \leq \frac{1}{2} I_{ij}$$

and

$$(55) \quad \sup_{f \in F_i} (E(\hat{T}_{ij}) - T f) \leq \frac{1}{2} I_{ij}.$$

As a first step in the construction of adaptive confidence intervals, define CI_{ij} by

$$(56) \quad CI_{ij} = [\inf_i \hat{T}_{ij} - \frac{3}{2} I_{ij} g; \max_i \hat{T}_{ji} + \frac{3}{2} I_{ji} g] :$$

The following result shows that this confidence interval attains the lower bound on the maximum expected length over F_j given in (53) and satisfies the constraint that it has the minimum coverage of $1 - \alpha$ for all $f \in G$.

Proposition 4. Let $F_j, j = 1, \dots, k$, be convex parameter spaces with $F_i \setminus F_j \neq \emptyset$ for all i, j and let $G = \bigcup_{j=1}^k F_j$. Let the interval CI_{ij} be defined as in (56). Then $CI_{ij} \subset I_{ij}$ and CI_{ij} has expected length over F_j satisfying

$$(57) \quad L(F_j; G) - L(CI_{ij}; F_j) \leq \frac{8^p \overline{\log(k+1)} + 4z_{\alpha/2}}{(1-\alpha)z} L(F_j; G) :$$

Remark 6. It follows from (59) that the expected length of the confidence interval CI_{ij} is rate optimal as $n \rightarrow \infty$ as long as k remains fixed.

Proof of Proposition 4. First consider the coverage probability of the interval CI_{ij} . Suppose $f \in F_m$ for some $1 \leq m \leq k$. Note that the interval CI_{ij} contains

$$CI_{m,ij} = [\hat{T}_{m,ij} - \frac{3}{2} I_{m,ij}; \hat{T}_{j,m} + \frac{3}{2} I_{j,m}] :$$

The derivation below shows that the interval $CI_{m,ij}$ has correct coverage probability. First note that for $f \in F_m, E(\hat{T}_{m,ij}) - T f \leq \frac{1}{2} I_{m,ij}$ and that $E(\hat{T}_{j,m}) - T f \leq \frac{1}{2} I_{j,m}$. Let

$$X_{m,ij} = \frac{\hat{T}_{m,ij} - T f - (1-\alpha) I_{m,ij}}{I_{m,ij} z_{\alpha/2}} ;$$

$$X_{j,m} = \frac{\hat{T}_{j,m} - T f + (1-\alpha) I_{j,m}}{I_{j,m} z_{\alpha/2}} ;$$

Then for any $f \in F_m$ it follows from (54) and (55) that $X_{m,j}$ has a Normal distribution with mean less than or equal to zero and variance bounded by 1 and $X_{j,m}$ has a Normal distribution with mean greater than or equal to zero and variance bounded by 1. Hence, for $f \in F_m$,

$$\begin{aligned} P(|T_f| \leq C I_{j,\cdot}) &= P(|T_f| \leq C I_{m,j}) \\ &= P(X_{m,j} \leq z_{=2} \text{ and } X_{j,m} \leq z_{=2}) \\ &= 1 - P(X_{m,j} > z_{=2}) - P(X_{j,m} > z_{=2}) \\ &= 1 - 2\Phi(-z_{=2}) \end{aligned}$$

So for any $f \in G$, $P(|T_f| \leq C I_{j,\cdot}) \geq 1 - 2\Phi(-z_{=2})$ and thus coverage has been established.

The bounds on the expected length of these intervals can now be obtained by using the following technical lemma from Dudley [(1999), pages 56 and 57].

Lemma 7. Let X_1, X_2, \dots, X_k be normally distributed random variables with mean 0 and variance σ^2 . Then

$$(58) \quad E \max_{1 \leq i \leq k} |X_i| \leq \sigma \left(2 + \frac{4 + \log 4}{\log(3/2)} \right)^{1/2} \sqrt{\log(k+1)}.$$

Let

$$j = 1 + \frac{z_{=2}}{n} F_j; G = \max_{1 \leq i \leq k} |X_{ij}|; |X_{ji}|$$

It is easy to see that the length of the interval $C I_{j,\cdot}$ is bounded by

$$L(C I_{j,\cdot}) = \max_i (\hat{T}_{j,i} - T_f)_+ + \max_i (T_f - \hat{T}_{i,j})_+ + 3 \cdot j$$

Now note that if $f \in F_j$, then for any $i \in j$,

$$a_{j,i} = E(\hat{T}_{j,i} - T_f) \leq \frac{1}{2} |X_{ji}|$$

and

$$b_{i,j} = E(T_f - \hat{T}_{i,j}) \leq \frac{1}{2} |X_{ij}|$$

Also note that for any real numbers x and y , $(x+y)_+ \leq (x)_+ + (y)_+$. So for $f \in F_j$ the expected length of $C I_{j,\cdot}$ satisfies

$$E L(C I_{j,\cdot}) = E \max_i (\hat{T}_{j,i} - T_f)_+ + E \max_i (T_f - \hat{T}_{i,j})_+ + 3 \cdot j$$

$$= E \max_i (a_{j,i})_+ + E \max_i (b_{i,j})_+ + 3 \cdot j$$

$$+ E \max_i (b_{i,j})_+ + (\hat{T}_{i,j} - T f(b_j))_+ g + 3 j$$

$$E \max_i (\hat{T}_{j,i} - \hat{T} f(a_{j,i}))_+ + E \max_i (\hat{T}_{i,j} - \hat{T}_j(b_j))_+ + 4 j:$$

It then follows from Lemma 7 that

$$E_f(L(CI_{j,})) \leq \frac{2}{z_{=2}} j^2 + \frac{4 + \log 4}{\log(3=2)} \frac{1=2p}{\log(k+1)} + 4 j$$

$$8 \frac{p}{z_{=2}} \frac{\log(k+1)}{\log(k+1)} + 4 j$$

and it follows by taking the supremum over F_j that

$$(59) \quad L(CI_{j,}; F_j) \leq 8 \frac{p}{z_{=2}} \frac{\log(k+1)}{\log(k+1)} + 4 j + \frac{z_{=2}}{p} \frac{1}{n} F_j; G :$$

The proposition now follows by combining (20), (7) and (59).

4.2. Adaptive confidence intervals. The intervals $CI_{j,}$ constructed in the last section have near optimal expected length over F_j but do not control the expected length over other F_i . In this section adaptive confidence intervals over $F_j: 1 \leq j \leq k$ are formed by intersecting such intervals. For a fixed k , the resulting interval has rate optimal expected length over every parameter space F_j for all $1 \leq j \leq k$. A Bonferroni approach is applied to the intervals of Section 4.1 to yield an adaptive confidence interval.

More specifically, define the confidence interval CI by

$$(60) \quad CI = \bigcap_{j=1}^k CI_{j,=k}$$

where $CI_{j,}$ are given in (56). The following theorem shows that this confidence interval has guaranteed coverage probability and also has near optimal expected length over F_j for each $1 \leq j \leq k$.

Theorem 3. Let $F_j, j = 1, \dots, k$, be convex parameter spaces with $F_i \setminus F_j \neq \emptyset$ for all i, j and let $G = \bigcup_{j=1}^k F_j$. Let the interval CI be given as in (60). Then $CI \subset I_{\alpha,G}$ and CI satisfies

$$(61) \quad L(F_j; G) \leq L(CI; F_j) \leq \frac{12z_{=2k}}{(1=2)^z} L(F_j; G)$$

for all $1 \leq j \leq k$.

Proof. The results follow easily from Proposition 4. For any $f \in G$, Proposition 4 shows that

$$P(f \in \bigcap_{j=1}^k C I_{j, \leq k}) \geq 1 - \frac{1}{k}.$$

Hence, for any $f \in G$,

$$P(f \in \bigcap_{j=1}^k C I_j) = 1 - P(f \notin \bigcap_{j=1}^k C I_j) \leq 1 - \prod_{j=1}^k P(f \notin C I_j) \leq 1 - \prod_{j=1}^k \frac{1}{k} = 1 - \frac{1}{k^k}.$$

For the expected length note that

$$L(C I_j; F_j) \leq L(C I_{j, \leq k}; F_j) \leq 8 \frac{\log(k+1)}{z_{=2k}} + 4! + \frac{z_{=2k}}{P_{=n}}; F_j; G$$

for any $1 \leq j \leq k$. For $0 < \epsilon < 0.5$, calculations show that

$$\frac{8 \log(k+1)}{z_{=2k}} \leq 1$$

and hence

$$(62) \quad L(C I_j; F_j) \leq 12! + \frac{z_{=2k}}{P_{=n}}; F_j; G.$$

The theorem now follows by combining (7), (20) and (62).

Remark 7. It follows from Lemma 6 that $z_{=2k} \leq \frac{2}{z_{=2}^2} \log k + 1 \leq z_2$.

Hence it follows from (62) and (20) that

$$\begin{aligned} L(C I_j; F_j) &\leq 12! + \frac{2}{z_{=2}^2} \log k + 1 + \frac{z_{=2}}{P_{=n}}; F_j; G \\ &\leq 12! + \frac{2}{z_{=2}^2} \log k + 1 + \frac{z_{=2}}{P_{=n}}; F_j; G. \end{aligned}$$

The ratio of the upper bound just given to the lower bound in (53) is thus clearly bounded by a constant multiple of $\frac{2}{P_{=n} \log k}$.

Section 5.2 gives an example of a nearly black object which shows that this $\frac{2}{P_{=n} \log k}$ factor cannot in general be improved.

5. Minimax confidence interval for nonconvex parameter spaces. As mentioned in the Introduction, Donoho (1994) constructed for any convex parameter space F fixed length intervals centered at a new estimators which have length within a small constant factor of the minimax expected length

$L(F)$. Although the focus of the present paper is on adaptation the adaptation theory developed in the previous sections can also be used to yield a minimax theory for parameter spaces that are finite unions of convex parameter spaces. In this section confidence intervals with a specified coverage probability are given which also have near optimal maximum expected length. It is also shown, in contrast to the theory for convex parameter spaces, that optimal confidence intervals centered on a new estimator can have expected length much longer than the expected length of optimal confidence intervals centered at nonlinear estimators.

Let $F_i, i = 1, \dots, k$, be convex parameter spaces with $F_i \not\subset F_j$ for all i, j and let $G = \bigcup_{i=1}^k F_i$. Note that the parameter space G is in general nonconvex. The minimax expected length of confidence intervals $CI \in \mathcal{I}_G$ can be bounded above and below as follows.

Set $0 < \alpha < \frac{1}{2}$ and let CI be a $1 - \alpha$ level confidence interval for all $f \in G = \bigcup_{i=1}^k F_i$. It follows from Theorem 1 that the maximum expected length of $CI \in \mathcal{I}_G$ is bounded below by

$$(63) \quad L(CI; G) \geq \frac{1}{2} \left(1 - \frac{Z_{1-\alpha/2}}{n} \right); G :$$

Upper bounds on the minimax expected length can be obtained by considering the confidence interval CI as defined in (60). As shown in Theorem 3 this interval has coverage probability of at least $1 - \alpha$ over G . In addition, it follows from (61) that the maximum of the expected length of CI over G satisfies

$$(64) \quad \begin{aligned} L(CI; G) &= \max_{1 \leq j \leq k} L(CI; F_j) \\ &\leq \max_{1 \leq j \leq k} \left(\frac{1}{2} + \frac{Z_{1-\alpha/2}}{n} \right); F_j; G \\ &= \frac{1}{2} \left(1 + \frac{Z_{1-\alpha/2}}{n} \right); G : \end{aligned}$$

Hence, (63) and (64) together yield the following result on the minimax expected length of $1 - \alpha$ level confidence intervals over G .

Theorem 4. Let $G = \bigcup_{j=1}^k F_j$, where for $j = 1, \dots, k$ F_j are convex spaces with nonempty intersections and suppose $0 < \alpha < \frac{1}{2}$. Then

$$(65) \quad \frac{1}{2} \left(1 - \frac{Z_{1-\alpha/2}}{n} \right); G \leq L(G) \leq \frac{1}{2} \left(1 + \frac{Z_{1-\alpha/2}}{n} \right); G :$$

Hence, the confidence interval CI attains the optimal rate of convergence for the maximum expected length over the parameter spaces G when the number of convex subspaces is fixed and finite.

The example of confidence intervals in Section 5.2 for a linear functional of nearly black objects shows that the factor of $z_{=2k} \sqrt{\log k}$ in the upper bound of (65) cannot be dropped in general when the number k of convex subspaces grows with n .

5.1. Confidence intervals centered at a \hat{T} estimator. We now consider the performance of confidence intervals centered at a \hat{T} estimator over nonconvex parameter spaces. As mentioned earlier, when the parameter space F is assumed to be fixed and convex, Donoho (1994) and Theorem 1 given in Section 2 together show that the length of the shortest fixed length confidence interval centered on an \hat{T} estimator is within a fixed constant factor of the maximum expected length of the optimal confidence interval. Hence there is relatively little to gain by looking beyond the class of fixed length confidence intervals centered on a \hat{T} estimator.

The following theorem considers the case when the parameter space is nonconvex. Once again let $\text{C Hull}(F)$ denote the convex hull of a parameter space F .

Theorem 5. Consider the white noise model (1) or the sequence model (2). Let \hat{T} be an \hat{T} estimator of Tf and $\tau \geq 0$ a nonnegative random variable. If $CI = [\hat{T} - \tau, \hat{T} + \tau]$ is a (variable length) confidence interval centered at \hat{T} and $CI \subset I_F$, then

$$(66) \quad L(CI; F) \leq C(\tau) \frac{2z_{=2}}{\sqrt{n}}; \text{C Hull}(F)$$

where $C(\tau) > 0$ is a constant depending on τ only. In particular, if the interval CI is of fixed length, then

$$(67) \quad L(CI) \leq \frac{1}{2} \frac{2z_{=2}}{\sqrt{n}}; \text{C Hull}(F) :$$

Proof. It is shown in Cai and Low (2004) that the \hat{T} estimator \hat{T} satisfies

$$\sup_{f \in F} \mathbb{E} \|\hat{T} - Tf\| = \sup_{f \in \text{C Hull}(F)} \mathbb{E} \|\hat{T} - Tf\|$$

It then follows from Theorem 2 of Low (1995) that \hat{T} must satisfy either

$$(68) \quad \sup_{f \in F} \mathbb{E} \|\hat{T} - Tf\| \leq \frac{1}{4} \frac{2z_{=2}}{\sqrt{n}}; \text{C Hull}(F)$$

or

$$(69) \quad \tau \leq \frac{1}{4z_{=2}} \frac{2z_{=2}}{\sqrt{n}}; \text{C Hull}(F) ;$$

where $\hat{\sigma}_f$ denotes the standard deviation of the estimator \hat{T}_f . We now consider the two cases separately. If (68) holds, then for any $\epsilon > 0$, there exists $f \in F$ such that

$$(70) \quad B_f - E\hat{T}_f - T_f \leq \frac{1}{4} \left(\frac{2z_{\alpha/2}}{\sqrt{n}} \right); C \text{ Hull}(F) \quad \epsilon:$$

Since $CI = [\hat{T}_f - \hat{\sigma}_f + \epsilon]$ has minimum coverage probability of at least 1 over F ,

$$\begin{aligned} 1 &= P_f(\hat{T}_f - T_f \leq \epsilon) \\ &= P_f(\hat{T}_f - T_f \leq -B_f) + P_f(\hat{T}_f - T_f \leq -B_f \text{ and } \epsilon > B_f) \\ &= P_f(\hat{T}_f - T_f \leq -B_f) + P(\epsilon > B_f): \end{aligned}$$

Since \hat{T}_f is an unbiased estimator and thus has a normal distribution, it is easy to check that $P_f(\hat{T}_f - T_f \leq -B_f) = \frac{1}{2}$ and hence

$$(71) \quad P(\epsilon > B_f) = \frac{1}{2}:$$

Letting $\epsilon \rightarrow 0$ in (70), it then follows that

$$\begin{aligned} E_f L(CI) &= 2E_f(\epsilon) - 2B_f P(\epsilon > B_f) \\ &= \frac{1}{4} - \frac{1}{2} \left(\frac{2z_{\alpha/2}}{\sqrt{n}} \right); C \text{ Hull}(F): \end{aligned}$$

If (69) holds, we have, for $f \in F$,

$$\begin{aligned} 1 &= P_f(\hat{T}_f - T_f \leq \epsilon) \\ &= P\left(\frac{E\hat{T}_f - T_f}{\hat{\sigma}_f} \leq Z\right) = P\left(\frac{E\hat{T}_f - T_f}{\hat{\sigma}_f} \leq \frac{\epsilon}{\hat{\sigma}_f}\right) \\ &= P\left(\frac{\epsilon}{\hat{\sigma}_f} \leq z_{0.25} + P\left(\frac{\epsilon}{\hat{\sigma}_f} \leq z_{0.25} + \frac{\epsilon}{\hat{\sigma}_f}\right)\right) \\ &= P\left(\frac{\epsilon}{\hat{\sigma}_f} \leq z_{0.25}\right) + P(\epsilon > z_{0.25} \hat{\sigma}_f) \end{aligned}$$

where Z denotes a standard normal random variable. Hence

$$P(\epsilon > z_{0.25} \hat{\sigma}_f) = \frac{1}{2}:$$

Consequently,

$$\begin{aligned} E_f L(CI) &= 2E_f(\epsilon) - 2z_{0.25} \hat{\sigma}_f P(\epsilon > z_{0.25} \hat{\sigma}_f) \\ &= (1 - 2) \left(\frac{z_{0.25}}{4z_{\alpha/2}} \right) \left(\frac{2z_{\alpha/2}}{\sqrt{n}} \right); C \text{ Hull}(F) \end{aligned}$$

$$\frac{1}{10z_{=2}} \log \frac{2z_{=2}}{n}; C \# \text{Hull}(F) :$$

Equation (66) now follows by taking $C(\cdot) = \inf \frac{1}{4} - \frac{1}{2}; \frac{1}{10z_{=2}} g$: Equation (67) for the fixed length case is easier to prove and we omit the proof here.

Remark 8. Theorem 5 shows that the minimax expected length of confidence intervals centered at a new estimator is determined by the modulus of continuity over the convex hull of F , not over F itself. In the case that $\log(F) \neq \log(\# \text{Hull}(F))$, any confidence intervals centered at a new estimator will perform poorly. Such is the case in the near black object example given in the next section.

5.2. Nearly black object. In this section an example is given which shows that the factor $\log k$ in the upper bound of the minimax expected length given in Theorem 4 cannot in general be dropped. It is also shown that confidence intervals centered at a new estimator are far from optimal.

Consider the Gaussian sequence model (2) with the index set $M = \{1, 2, \dots, n\}$, namely

$$(72) \quad Y(i) = f(i) + n^{-1/2} z_i; \quad i = 1, \dots, n;$$

where $z_i \stackrel{\text{iid}}{\sim} N(0, 1)$. The size of the vector, n , is assumed large. We assume that the vector f is sparse: only a small fraction of components are nonzero, and the indices or locations of the nonzero components are not known in advance.

Denote the ℓ_0 quasi-norm by $\|f\|_0 = \text{Card}\{i: f(i) \neq 0\}$. Fix m_n . The collection of vectors with at most m_n nonzero entries is

$$G = \ell_0(m_n) = \{f \in \mathbb{R}^n: \|f\|_0 \leq m_n\}.$$

Assume that m_n is known and $m_n \rightarrow \infty$ where $\frac{m_n}{n} < \frac{1}{2}$:

Such an example is considered in Cai and Low (2004) in the context of minimax estimation. The model, which arises naturally in wavelet analysis, has also been studied in Donoho, Johnstone, Hoch and Stern (1992) and Abramovich, Benjamini, Donoho and Johnstone (2000) for estimating the whole object.

Let the linear functional Tf be given by

$$Tf = \sum_{i=1}^n f(i);$$

and following Cai and Low (2004) let $\mathcal{I}(m_n; n)$ be the class of all subsets of $\{1, \dots, n\}$ of m_n elements and for $I \in \mathcal{I}(m_n; n)$ let

$$F_I = \{f \in \mathbb{R}^n: f(j) = 0 \text{ } \forall j \notin I\}.$$

Note that F_I is an m_n -dimensional subspace spanned by the coordinates in I . These are obviously convex and $G = \bigcup F_I$ where the union is taken over I in the set $I(m_n; n)$. From now on we shall assume that I is in the set $I(m_n; n)$.

Simple calculations show that for all $I, J \in I(m_n; n)$

$$P(F_I; F_J) = \frac{\text{Card}(I \cap J)}{\text{Card}(I)}$$

and consequently

$$P(F_I; G) = P(G; F_I) = P(G) = \frac{P}{2m_n}.$$

Remark 9. It is easy to see that $\text{CHull}(G) = R^n$ and hence $P(\text{CHull}(G)) = \frac{P}{n}$. It follows from Theorem 5 and (66) that any confidence interval with coverage of at least 1 centered at an affine estimator must have maximum expected length bounded from below by a fixed constant not depending on n .

Let k be the number of the m_n -dimensional parameter spaces F_I . Then k is equal to $\binom{n}{m_n}$ and it is easy to see that

$$k = \binom{n}{m_n} \approx \frac{n^{m_n}}{m_n^{m_n}}.$$

The following result gives a lower bound on the expected length of any confidence interval with a minimum coverage probability of 1 over G .

Proposition 5. Suppose that we observe the Gaussian sequence model (72), that $n \geq 4$ and $m_n < n$ with $\frac{m_n}{n} < \frac{1}{2}$. Let $Tf = \frac{1}{n} \sum_{i=1}^n f(i)$ and $0 < \frac{1}{2} < \frac{m_n}{n} < \frac{1}{2}$. Suppose that $CI(Y)$ is a confidence interval for Tf based on (72) and $CI(Y) \in I_G$. Then for all sufficiently large n ,

$$(73) \quad E_0 L(CI(Y)) \geq \frac{1}{2} \left(\frac{m_n}{n} - \frac{1}{2} \log \frac{n}{m_n^2} \right) - \frac{1}{4} \frac{\log k}{n} + P(G)$$

where E_0 denotes expectation under the Gaussian model (72) with $f(i) = 0$ for $i = 1, 2, \dots, n$.

Remark 10. It follows immediately from (73) that the maximum expected length of $CI(Y)$ over G satisfies

$$(74) \quad L(CI(Y); G) \geq C \frac{\log k}{n} + P(G) :$$

Comparing the lower bound (74) for the maximum expected length with the minimum upper bound given in (65) shows that the factor $\sqrt{\log k}$ in the upper bound for the minimum expected length cannot be dropped in general. A similar result also holds for adaptation.

Proof of Proposition 5. In the following proof the calculation of the L_1 distance between a mixture of normals and a given normal distribution follows a similar calculation used in Cai and Low (2004). We include the details of the calculation here for completeness. In the proof we will omit the subscript n and simply write m for m_n . Let f be the joint density of the Gaussian observations given in (72). More specifically f is a multivariate normal density with mean $(f(1); f(2); \dots; f(n))$ and covariance matrix $\frac{1}{n}A_n$ where A_n is the $n \times n$ identity matrix. Fix a constant $\epsilon > 0$. For $I \in \mathcal{I}(m; n)$ let f_I be defined by $f_I(j) = \frac{1}{m} \mathbb{1}(j \in I)$ and let f_0 be the sequence defined by $f_0(j) = 0$ for $j = 1, 2, \dots, n$. Finally let

$$f_I = \frac{1}{m} \sum_{I \in \mathcal{I}(m; n)} f_I.$$

Note that a similar mixture prior was used in Baraud (2002) to give lower bounds in a nonparametric testing problem. Note that for all f_I , $\int f_I = \frac{1}{m}$ and that $\int f_0 = 0$. Note also that if

$$\int f_I \leq \frac{1}{m} \int \sum_{I \in \mathcal{I}(m; n)} f_I \leq 1$$

for all $I \in \mathcal{I}(m; n)$ then it follows that

$$\int f_I \leq \frac{1}{m} \int \sum_{I \in \mathcal{I}(m; n)} f_I \leq 1.$$

Note that

$$\frac{\int f_I^2}{\int f_0^2} = \frac{1}{m^2} \sum_{I \in \mathcal{I}(m; n)} \sum_{I' \in \mathcal{I}(m; n)} \int \frac{f_I f_{I'}}{f_0};$$

and simple calculations show that

$$\int \frac{f_I f_{I'}}{f_0} = \exp(j^2);$$

where j is the number of points in the set $I \cap I'$. It follows that

$$\frac{\int f_I^2}{\int f_0^2} = E \exp(J^2);$$

Hence

$$P_{f_0} \left(0 \in CI \text{ and } m \leq \frac{1}{n} \log \frac{n}{m^2} \right) \leq \frac{1}{2} \epsilon_n.$$

Since CI is an interval the length of this interval must be at least $m \frac{1}{n}$ when both 0 and $m \frac{1}{n}$ are in the interval. Hence for n sufficiently large,

$$E_{f_0} L(CI(Y)) \leq \frac{1}{2} \epsilon_n + \frac{1}{2} \log \frac{n}{m^2}.$$

Now take $\epsilon_n = \frac{1}{2}$. Then for all sufficiently large n ,

$$E_{f_0} L(CI(Y)) \leq \frac{1}{2} + \frac{1}{2} \log \frac{n}{m^2} + \frac{1}{4} \log k + \frac{1}{2} \log \frac{n}{m^2}; G,$$

where k is the number of convex parameter spaces in G .

Remark 11. It follows immediately from Proposition 5 that

$$L(F_I) \leq \frac{1}{2} + \frac{1}{4} \log k + \frac{1}{2} \log \frac{n}{m^2}; F_I; G.$$

Hence the factor of $\frac{1}{2} \log k$ for adaptation in the upper bound of Theorem 3 and the same factor for minimax confidence procedures in Theorem 4 cannot in general be removed.

Acknowledgment. We thank one of the referees for very thorough and useful comments which have helped to improve the presentation of the paper. In particular, some of the notation used in the paper is based on suggestions made by this referee.

REFERENCES

- Abramovich, F., Benjamini, Y., Donoho, D. and Johnstone, I. (2000). Adapting to unknown sparsity by controlling the false discovery rate. Technical Report 2000-19, Dept. Statistics, Stanford Univ.
- Baraud, Y. (2002). Non-asymptotic minimax rates of testing in signal detection. *Bernoulli* 8 577{606. [M R 1935648](#)
- Beran, R. and Dumbgen, L. (1998). Modulation of estimators and confidence sets. *Ann. Statist.* 26 1826{1856. [M R 1673280](#)
- Brown, L. D. and Low, M. G. (1996). Asymptotic equivalence of nonparametric regression and white noise. *Ann. Statist.* 24 2384{2398. [M R 1425958](#)
- Cai, T. and Low, M. (2002). On modulus of continuity and adaptability in nonparametric functional estimation. Technical report, Dept. Statistics, Univ. Pennsylvania.

- Cai, T. and Low, M. (2003). Adaptive estimation and confidence intervals for convex functions and monotone functions. Technical report, Dept. Statistics, Univ. Pennsylvania.
- Cai, T. and Low, M. (2004). Minimax estimation of linear functionals over nonconvex parameter spaces. *Ann. Statist.* 32 552{576. [M R 2060169](#)
- Donoho, D. L. (1994). Statistical estimation and optimal recovery. *Ann. Statist.* 22 238{270. [M R 1272082](#)
- Donoho, D. L., Johnstone, I. M., Hoch, J. C. and Stern, A. S. (1992). Maximum entropy and the nearly black object (with discussion). *J. Roy. Statist. Soc. Ser. B.* 54 41{81. [M R 1157714](#)
- Donoho, D. L. and Liu, R. C. (1987). Geometrizing rates of convergence. I. Technical Report 137, Dept. Statistics, Univ. California, Berkeley.
- Donoho, D. L. and Liu, R. C. (1991). Geometrizing rates of convergence. III. *Ann. Statist.* 19 668{701. [M R 1105839](#)
- Dudley, R. (1999). *Uniform Central Limit Theorems*. Cambridge Univ. Press. [M R 1720712](#)
- Dumbgen, L. (1998). New goodness-of-fit tests and their application to nonparametric confidence sets. *Ann. Statist.* 26 288{314. [M R 1611768](#)
- Feller, W. (1968). *An Introduction to Probability Theory and Its Applications* 1, 3rd ed. Wiley, New York. [M R 228020](#)
- Genovese, C. R. and Wasserman, L. (2002). Confidence sets for nonparametric wavelet regression. Technical report, Dept. Statistics, Carnegie Mellon Univ.
- Hall, P. (1992). Effect of bias estimation on coverage accuracy of bootstrap confidence intervals for a probability density. *Ann. Statist.* 20 675{694. [M R 1165587](#)
- Hardle, W. and Marron, J. S. (1991). Bootstrap simultaneous error bars for nonparametric regression. *Ann. Statist.* 19 778{796. [M R 1105844](#)
- Hengartner, N. W. and Stark, P. B. (1995). Finite-sample confidence envelopes for shape-restricted densities. *Ann. Statist.* 23 525{550. [M R 1332580](#)
- Ibragimov, I. A. and Hasminskii, R. Z. (1984). Nonparametric estimation of the value of a linear functional in Gaussian white noise. *Theory Probab. Appl.* 29 18{32. [M R 739497](#)
- Li, K.-C. (1989). Honest confidence regions for nonparametric regression. *Ann. Statist.* 17 1001{1008. [M R 1015135](#)
- Low, M. G. (1995). Bias-variance tradeoffs in functional estimation problems. *Ann. Statist.* 23 824{835. [M R 1345202](#)
- Low, M. G. (1997). On nonparametric confidence intervals. *Ann. Statist.* 25 2547{2554. [M R 1604412](#)
- Picard, D. and Tribouley, K. (2000). Adaptive confidence interval for pointwise curve estimation. *Ann. Statist.* 28 298{335. [M R 1762913](#)
- Pratt, J. W. (1961). Length of confidence intervals. *J. Amer. Statist. Assoc.* 56 549{567. [M R 125678](#)
- Stark, P. B. (1992). A minimax confidence intervals for a bounded normal mean. *Statist. Probab. Lett.* 13 39{44. [M R 1147637](#)
- Zeytinoglu, M. and Mintz, M. (1984). Optimal fixed size confidence procedures for a restricted parameter space. *Ann. Statist.* 12 945{957. [M R 751284](#)

Department of Statistics
The Wharton School
University of Pennsylvania
Philadelphia, Pennsylvania 19104
USA
e-mail: tcai@wharton.upenn.edu