

EQUIVALENCE THEORY FOR DENSITY ESTIMATION, POISSON PROCESSES AND GAUSSIAN WHITE NOISE WITH DRIFT

By Lawrence D. Brown¹, Andrew V. Carter, Mark G. Low²
 and Cun-Hui Zhang³

University of Pennsylvania, University of California, Santa Barbara,
 University of Pennsylvania and Rutgers University

This paper establishes the global asymptotic equivalence between a Poisson process with variable intensity and white noise with drift under sharp smoothness conditions on the unknown function. This equivalence is also extended to density estimation models by Poissonization. The asymptotic equivalences are established by constructing explicit equivalence mappings. The impact of such asymptotic equivalence results is that an investigation in one of these nonparametric models automatically yields asymptotically analogous results in the other models.

1. Introduction. The purpose of this paper is to give an explicit construction of global asymptotic equivalence in the sense of [Le Cam \(1964\)](#) between a Poisson process with variable intensity and white noise with drift. The construction is extended to density estimation models. It yields asymptotic solutions to both density estimation and Poisson process problems based on asymptotic solutions to white noise with drift problems and vice versa.

Density estimation model. A random vector $V_n^?$ of length n is observed such that $V_n^? = (V_1^?; \dots; V_n^?)$ is a sequence of i.i.d. variables with a common density $f \in \mathcal{F}$.

Received April 2002; revised February 2004.

¹Supported by NSF Grant DMS-99-71751.

²Supported by NSF Grant DMS-03-06576.

³Supported by NSF Grants DMS-01-02529 and DMS-02-03086.

[AMS 2000 subject classifications.](#) Primary 62B15; secondary 62G07, 62G20.

Key words and phrases. Asymptotic equivalence, decision theory, local limit theorem, quantile transform, white noise model.

This is an electronic reprint of the original article published by the [Institute of Mathematical Statistics](#) in [The Annals of Statistics](#), 2004, Vol. 32, No. 5, 2074{2097. This reprint differs from the original in pagination and typographic detail.

Poisson process. A random vector of random length fN ; X_N is observed such that N is a Poisson variable with $EN = n$ and that given $N = m$, $X_N = X_m$ ($X_1; \dots; X_m$) is a sequence of i.i.d. variables with a common density $f \in F$. The resulting observations are then distributed as a Poisson process with intensity function nf .

White noise. A Gaussian process $Z_n^? = Z_n^?(t); 0 \leq t \leq 1$ is observed such that

$$(1.1) \quad Z_n^?(t) = \int_0^t \frac{B^?(x)}{f(x)} dx + \frac{B^?(t)}{2^n}; \quad 0 \leq t \leq 1;$$

with a standard Brownian motion $B^?(t)$ and an unknown probability density function $f \in F$ in $[0; 1]$.

Asymptotic equivalence. For any two experiments $\mathcal{E}_1$ and $\mathcal{E}_2$ with a common parameter space Θ , $(\mathcal{E}_1; \mathcal{E}_2; \epsilon)$ denotes Le Cam's distance [cf., e.g., [Le Cam \(1986\)](#) or [Le Cam and Yang \(1990\)](#)] defined as

$$(\mathcal{E}_1; \mathcal{E}_2; \epsilon) = \sup_{\mathcal{L}} \max_{j=1,2} \sup_{(j)} \inf_{(k)} \sup_{\mathcal{P}^{(k)}} \mathbb{E}^{(j)} L(\mathcal{L}; (j)) - \mathbb{E}^{(k)} L(\mathcal{L}; (k));$$

where (a) the first supremum is taken over all decision problems with loss function $kL, k_1 \leq 1$, (b) given the decision problem and $j = 1, 2, k = 3 - j$ ($k = 2$ for $j = 1$ and $k = 1$ for $j = 2$) the "maximum" value of the maximum difference in risks over $\mathcal{L}$ is computed over all (randomized) statistical procedures (j) for $\mathcal{L}$, and (c) the expectations $\mathbb{E}^{(j)}$ are evaluated in experiments $\mathcal{E}_\nu$ with parameter $\nu = j; k$. The statistical interpretation of the Le Cam distance is as follows: If $(\mathcal{E}_1; \mathcal{E}_2; \epsilon) < \epsilon$, then for any decision problem with $kL, k_1 \leq 1$ and any statistical procedure (j) with the experiment $\mathcal{E}_j, j = 1, 2$, there exists a (randomized) procedure (k) with $\mathcal{E}_k, k = 3 - j$, such that the risk of (k) evaluated in $\mathcal{E}_k$ nearly matches (within ϵ) that of (j) evaluated in $\mathcal{E}_j$.

Two sequences of experiments $f_{1,n}; n \geq 1$ and $f_{2,n}; n \geq 1$, with a common parameter space F , are asymptotically equivalent if

$$(\mathcal{E}_{1,n}; \mathcal{E}_{2,n}; F) \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

The interpretation is that the risks of corresponding procedures converge.

A key result of [Le Cam \(1964\)](#) is that this equivalence of experiments can be characterized using random transformations between the probability spaces. A random transformation, $T(X; U)$ which maps observations X into the space of observations Y (with possible dependence on an independent, uninformative random component U) also maps distributions in $\mathcal{P}_1$ to approximations of the distributions in $\mathcal{P}_2$ via $P^{(1)}T = P^{(2)}$. For the mapping between the Poisson and Gaussian processes we shall restrict ourselves

to transformations T with deterministic inverses, $T^{-1}(T(X;U)) = X$. The experiments are asymptotically equivalent if the total-variation distance between $P^{(2)}$ and the distribution of T under $P^{(1)}$ converges to 0 uniformly in $\mathcal{F}$. As explained in [Brown and Low \(1996\)](#) and [Brown, Cai, Low and Zhang \(2002\)](#), knowing an appropriate T allows explicit construction of estimation procedures in $\mathcal{F}_1$ by applying statistical procedures from $\mathcal{F}_2$ to $T(X;U)$.

In general, asymptotic equivalence also implies a transformation from the $P^{(2)}$ to the $P^{(1)}$ and the corresponding total-variation distance bound. However, in the case of the equivalence between the Poisson process and white noise with drift, by requiring that the transformation be invertible, we have saved ourselves a step. The transformation in the other direction is T^{-1} , and

$$\|P^{(1)}T - P^{(2)}\|_k = \|P^{(1)}TT^{-1} - P^{(2)}T^{-1}\|_k = \|P^{(1)} - P^{(2)}T^{-1}\|_k.$$

Therefore, it is sufficient if $\sup_k \|P^{(1)}T - P^{(2)}\|_k \rightarrow 0$.

The equivalence mappings T_n constructed in this paper from the sample space of the Poisson process to the sample space of the white noise are invertible randomized mappings such that

$$(1.2) \quad \sup_{f \in \mathcal{F}} H_f(T_n(N; X_N); Z_n^2) \rightarrow 0$$

under certain conditions on the family $\mathcal{F}$. Here $H_f(Z_1; Z_2)$ denotes the Hellinger distance of stochastic processes or random vectors Z_1 and Z_2 living in the same sample space, when the true unknown density is f . Since T_n are invertible randomized mappings, $T_n(N; X_N)$ are sufficient statistics for the Poisson processes and their inverses T_n^{-1} are necessarily many-to-one deterministic mappings. Similar considerations apply for the mapping of the density estimation problem to the white noise with drift problem, although in that case there are two mappings, one from the density estimation to the white noise with drift model and another from the white noise with drift model back to the density estimation model. These mappings are given in [Section 2](#).

There have recently been several papers on the global asymptotic equivalence of nonparametric experiments. [Brown and Low \(1996\)](#) established global asymptotic equivalence of the white noise problem with unknown drift f to a nonparametric regression problem with deterministic design and unknown regression f when f belongs to a Lipschitz class with smoothness index $> \frac{1}{2}$. It has also been demonstrated that such nonparametric problems are typically asymptotically nonequivalent when the unknown f belongs to larger classes, for example, with smoothness index $\leq \frac{1}{2}$. [Brown and Low \(1996\)](#) showed the asymptotic nonequivalence between the

white noise problem and nonparametric regression with deterministic design for $\frac{1}{2}$, [Efroymov and Samarov \(1996\)](#) showed that the asymptotic equivalence may fail when $\gamma < \frac{1}{4}$. [Brown and Zhang \(1998\)](#) showed the asymptotic nonequivalence for $\gamma > \frac{1}{2}$ between any pair of the following four experiments: white noise, density problem, nonparametric regression with random design, and nonparametric regression with deterministic design. In [Brown, Cai, Low and Zhang \(2002\)](#) the asymptotic equivalence for nonparametric regression with random design was shown under Besov constraints which include Lipschitz classes with any smoothness index $\gamma > \frac{1}{2}$. [Gramma and Nussbaum \(1998\)](#) solved the fixed-design nonparametric regression problem for nonnormal errors. [Milstein and Nussbaum \(1998\)](#) showed that some diffusion problems can be approximated by discrete versions that are nonparametric autoregression models, and [Golubev and Nussbaum \(1998\)](#) established a discrete Gaussian approximation to the problem of estimating the spectral density of a stationary process.

Most closely related to this paper is the work in [Nussbaum \(1996\)](#) where global asymptotic equivalence of the white noise problem to the nonparametric density problem with unknown density $g = f^2 = 4$ is shown. In this paper the global asymptotic equivalence was established under the following smoothness assumption: f belongs to the Lipschitz classes with smoothness index $\gamma > \frac{1}{2}$.

The parameter spaces. The class of functions F will be assumed throughout to be densities with respect to Lebesgue measure on $[0; 1]$ that are uniformly bounded away from 0. The smoothness conditions on F can be described in terms of Haar basis functions of the densities. Let

$$(1.3) \quad c_{k,\gamma} = c_{k,\gamma}(f) = \int_{I_{k,\gamma}} f(x) dx; \quad \gamma = 0, \dots, 2^k - 1; k = 0, 1, \dots;$$

be the Haar coefficients of f , where

$$(1.4) \quad c_{k,\gamma} = 2^{k/2} (\mathbb{1}_{I_{k+1,2\gamma}} - \mathbb{1}_{I_{k+1,2\gamma+1}})$$

are the Haar basis functions with $I_{k,\gamma} = [\gamma 2^{-k}; (\gamma+1) 2^{-k}]$. The convergence of the Hellinger distance in (1.2) is established via an inequality in [Theorem 3](#) in terms of the tails of the Besov norms $\|f\|_{B_{2,2,2}^{\gamma}}$ and $\|f\|_{B_{2,4,4}^{\gamma}}$ of the Haar coefficients $c_{k,\gamma} = c_{k,\gamma}(f)$ in (1.3).

The Besov norm $\|f\|_{B_{p,q}^{\gamma}}$ for the Haar coefficients, with smoothness index γ and shape parameters p and q , are defined by

$$(1.5) \quad \|f\|_{B_{p,q}^{\gamma}}^p = \sum_{k=0}^{\infty} \left(\sum_{\gamma=0}^{2^k-1} |c_{k,\gamma}|^p \right)^{q/p} = \sum_{k=0}^{\infty} \left(\sum_{\gamma=0}^{2^k-1} |c_{k,\gamma}|^p \right)^{q/p} 2^{k(1-\frac{1}{p})} 2^{k(1-\frac{1}{q})} = \sum_{k=0}^{\infty} \left(\sum_{\gamma=0}^{2^k-1} |c_{k,\gamma}|^p \right)^{q/p} 2^{k(1-\frac{1}{p})} 2^{k(1-\frac{1}{q})} :$$

Let f_k be the piecewise average of f at resolution level k , that is, the piecewise constant function defined by

$$(1.6) \quad f_k = f_k(t) = \sum_{j=0}^{2^k-1} \mathbb{1}_{[t_j, t_{j+1})} \int_{t_j}^{t_{j+1}} f(t) dt, \quad t \in [0, 1].$$

Since $\|f_k - f_{k+1}\|_p^p = \sum_{j=0}^{2^k-1} \int_{t_j}^{t_{j+1}} |f(t) - \int_{t_j}^{t_{j+1}} f(t) dt| dt = \sum_{j=0}^{2^k-1} \int_{t_j}^{t_{j+1}} |f(t) - f_k(t)| dt$, (1.5) can be written as

$$\|f_k - f_{k+1}\|_{p,q} = \left(\sum_{j=0}^{2^k-1} \int_{t_j}^{t_{j+1}} |f(t) - f_k(t)|^q dt \right)^{1/q};$$

and its tail at resolution level $k_0 = 0$ is $\|f - f_{k_0}\|_{p,q}, k_0 = 0$, with

$$(1.7) \quad \|f - f_{k_0}\|_{p,q}^q = \sum_{k=k_0}^{\infty} \left(\sum_{j=0}^{2^k-1} \int_{t_j}^{t_{j+1}} |f(t) - f_k(t)|^q dt \right)^{1/p};$$

Let $B(p, q)$ be the Besov space

$$B(p, q) = \{f : \|f - f_{k_0}\|_{p,q} < \infty\}.$$

The following two theorems on the equivalence of white noise with drift, density estimation and Poisson estimation models are corollaries of our main result, Theorem 3, which bounds the squared Hellinger distance between particular invertible randomized mappings of the Poisson process and white noise with drift models. The randomized mappings are given in Section 2. Proofs of these theorems are given in the Appendix.

Theorem 1. Let $Z_n, f_n; X_N, g$ and V_n be the Gaussian process, Poisson process and density estimation experiments, respectively. Suppose that H is compact in both $B(1/2; 2; 2)$ and $B(1/2; 4; 4)$ and that $H = \{f : \int_0^1 f(x) dx > 0\}$ for some $\epsilon_0 > 0$. Then

$$(1.8) \quad \lim_{n \rightarrow \infty} (Z_n; f_n; X_N, g; H) = 0$$

and

$$(1.9) \quad \lim_{n \rightarrow \infty} (Z_n; V_n; H) = 0.$$

Our construction also shows that asymptotic equivalence holds for a class F if F is bounded in the Lipschitz norm with smoothness index α and compact in the Sobolev norm with smoothness index β such that $\alpha + \beta > \frac{1}{2}$.

For $0 < \alpha < 1$ the Lipschitz norm $\|f\|_{\text{Lip}}^{(\alpha)}$ and Sobolev norm $\|f\|_{\text{Sob}}^{(\alpha)}$ are defined by

$$\|f\|_{\text{Lip}}^{(\alpha)} = \sup_{0 \leq x < y \leq 1} \frac{|f(x) - f(y)|}{|x - y|^\alpha}; \quad \|f\|_{\text{Sob}}^{(\alpha)} = \left(\sum_{n=1}^{\infty} n^{2\alpha} |\hat{f}_n(f)|^2 \right)^{1/2};$$

where $\hat{f}_n(f) = \int_0^1 f(x) e^{in2\pi x} dx$ are the Fourier coefficients of f .

Theorem 2. Let $Z_n^?$, $fN; X_N g$ and $V_n^?$ be the Gaussian process, Poisson process and density estimation experiments, respectively, and let F be bounded in the Lipschitz norm with smoothness index α and compact in the Sobolev norm with smoothness index β . Suppose $F = f f : \inf_{0 < x < 1} f(x) > 0$ for some $\epsilon_0 > 0$. Then if $\alpha + \beta > 1$, $\frac{3}{4}$ or $\beta > \frac{1}{2}$,

$$\lim_{n \rightarrow \infty} (Z_n^?; fN; X_N g; F) = 0$$

and

$$\lim_{n \rightarrow \infty} (Z_n^?; V_n^?; F) = 0.$$

2. The equivalence mappings. This section describes in detail the mappings which provide the asymptotic equivalence claimed in this paper. The fact that these mappings yield asymptotic equivalence is a consequence of our major result, Theorem 3. The construction is broken into several stages.

From observations of the white noise (1.1), define random vectors

$$(2.1) \quad \bar{Z}_k^? = f\bar{Z}_{k,\cdot}^?; 0 \leq \cdot < 2^k g; \quad \bar{Z}_{k,\cdot}^? = 2^k Z^? \frac{\cdot + 1}{2^k} - Z^? \frac{\cdot}{2^k};$$

$$(2.2) \quad W_k^? = fW_{k,2,\cdot}^?; 0 \leq \cdot < 2^k g; \quad W_{k,2,\cdot}^? = W_{k,2^?+1}^? (\bar{Z}_{k,2^?}^? - \bar{Z}_{k,2^?+1}^?) = 2^?;$$

Let k_0, k_m be suitable integers with $\lim_{m \rightarrow \infty} k_{0,m} = 1$. Following Brown, Cai, Low and Zhang (2002), we construct equivalence mappings by finding the counterparts of $\bar{Z}_{k_0}^?$ and $W_{k_0}^?$, $k > k_0$, with the Poisson process $(N; X_N)$, to strongly approximate the Gaussian variables.

It can be easily verified from (1.1) that $f\bar{Z}_{k_0,\cdot}^?; 0 \leq \cdot < 2^{k_0} g; W_{k,2,\cdot}^?; 0 \leq \cdot < 2^{k-1} g; k > k_0$ are uncorrelated normal random variables with

$$(2.3) \quad \begin{aligned} E \bar{Z}_{k,\cdot}^? &= h_{k,\cdot} = 2^k \int_{I_{k,\cdot}} h; \quad h = \int_0^1 f; \\ \text{Var}(\bar{Z}_{k,\cdot}^?) &= \int_{I_{k,\cdot}} 2^k = (4n); \end{aligned}$$

for $\cdot = 0; \dots; 2^k - 1$, and for $\cdot = 0; \dots; 2^{k-1} - 1$,

$$(2.4) \quad \begin{aligned} E W_{k,2,\cdot}^? &= \frac{1}{2} (h_{k,2^?} - h_{k,2^?+1}) = \int_{I_{k-1,\cdot}} f; \\ \text{Var}(W_{k,2,\cdot}^?) &= \int_{I_{k-1,\cdot}} 2^{k-1}; \end{aligned}$$

Let $\mathcal{U} = \{U_{k,i}; k = k_0, \dots, 0\}$ be a sequence of i.i.d. uniform variables in $[1/2, 1/2]$ independent of $(N; X_N)$. For $k = 0, 1, \dots$ and $i = 0, \dots, 2^k - 1$ define

$$(2.5) \quad N_{k,i} \sim \text{Poisson}(f_{k,i}; 0 \leq i < 2^k); \quad N_{k,i} \sim \sum_{i=1}^2 X_{i-1} I_{k,i}; g;$$

We shall approximate $\bar{Z}_k$ in (2.1) in distribution by

$$(2.6) \quad \bar{Z}_k \sim \sum_{k=0}^{\infty} \text{sgn}(N_{k,i} + U_{k,i}) \frac{1}{\sqrt{N_{k,i} + U_{k,i}}}$$

at the initial resolution level $k = k_0$. Since $N_{k,i}$ are Poisson variables with

$$(2.7) \quad E N_{k,i} = \frac{n}{2^k} f_{k,i} = \frac{f_{k,i}}{4 \cdot 2^k}; \quad f_{k,i} = \sum_{i=1}^2 f; I_{k,i};$$

by Taylor expansion and central limit theory

$$\bar{Z}_{k,i} \sim \sum_{k=0}^{\infty} \frac{1}{\sqrt{N_{k,i}}} + \frac{N_{k,i} - k_{k,i}}{2 \cdot 2^{k/2}} \sim N\left(\frac{1}{\sqrt{f_{k,i}}}, \frac{2}{k}\right)$$

as $k \rightarrow \infty$, compared with (2.3). Note that $\frac{1}{\sqrt{f_{k,i}}} \sim h_{k,i}$ under suitable smoothness conditions on f , in view of (2.3) and (2.7). The Poisson variables $N_{k,i}$ can be fully recovered from $\bar{Z}_{k,i}$, while the randomization turns $N_{k,i}$ into continuous variables.

Approximation of $W_{k,i}$ for $k > k_0$ is more delicate, since the central limit theorem is not sufficiently accurate at high resolution levels. Let F_m be the cumulative distribution function of the independent sum of a binomial variable $X_{m,i=2}$ with parameter $(m; \frac{1}{2})$ and a uniform variable U in $[1/2, 1/2]$,

$$(2.8) \quad F_m(x) = P(X_{m,i=2} + U \leq x); g;$$

with F_0 being the uniform distribution in $[1/2, 1/2]$. Let Φ be the $N(0,1)$ cumulative distribution. We shall approximate W_k by using a quantile transformation of randomized versions of the Poisson random variables. More specifically, let

$$(2.9) \quad W_{k,i} \sim \text{Poisson}(f_{k,i}; 0 \leq i < 2^k); \quad W_{k,i} \sim \sum_{i=1}^2 (F_{N_{k-1,i}}(N_{k-1,i} + U_{k,i}))$$

with $W_{k-1,i} \sim \text{Poisson}(f_{k-1,i}; i = 0, \dots, 2^{k-1} - 1)$, and the Φ in (2.3). Given $N_{k-1,i} = m$,

$$(2.10) \quad N_{k,i} \sim \text{Bin}(m; p_{k,i}); \quad p_{k,i} = \frac{R_{I_{k-1,i}} f}{\sum_{i=1}^2 f} = \frac{f_{k,i}}{f_{k,i} + f_{k,i+1}};$$

so that $W_{k,i}$ is distributed exactly according to $N(0, \frac{2}{k-1})$ for $p_{k,i} = \frac{1}{2}$, compared with (2.4). Thus, the distributions of $W_{k,i}$ and $W_{k,i}^?$ are close

at high resolution levels as long as f is sufficiently smooth, even for small $N_{k-1} = m$.

The equivalence mappings T_n , with randomization through $\mathcal{U}$, are defined by

$$T_n : f_N ; X_N ; \mathcal{U} \rightarrow W_{[k_0, 1)} ; Z_n = fZ_n(t) : 0 \leq t \leq 1g;$$

where for $k_0 \leq k-1$, $W_{[k_0, k)} = f\bar{Z}_{k_0} ; W_j ; k_0 < j < kg$, and $\bar{Z}_k$ and W_k are as in (2.6) and (2.9). The inverse of T_n is a deterministic many-to-one mapping defined by

$$T_n^{-1} : Z_n \rightarrow W_{[k_0, 1)} ; (N ; X_N ?);$$

where for $k_0 \leq k-1$, $W_{[k_0, k)} = f\bar{Z}_{k_0}^? ; W_j^? ; k_0 < j < kg$.

Remark 1. One need only carry out the above construction to $k = k_1 : 2^{k_1} > n$ since we shall assume that $f \in B(\frac{1}{2}; 2; 2)$ and then the observations $W_{[k_0, k)}^? = f\bar{Z}_{k_0}^? ; W_j^? ; k_0 < j < kg$ and $W_{[k_0, k)} = f\bar{Z}_{k_0} ; W_j ; k_0 < j < kg$ are asymptotically sufficient for the Gaussian process and Poisson process experiments. See Brown and Low (1996) for a detailed argument in the context of nonparametric regression.

Mappings for the density estimation model. The constructive asymptotic equivalence between density estimation experiments and Gaussian experiments is established by first randomizing the density estimation experiment to an approximation of the Poisson process and then applying the randomized mapping as given above. Set $k = \sup_{f \in H} kf - f_k k_{1=2; 2; 2}^2$ and note that since H is compact in $B(1=2; 2; 2)$, $k \neq 0$. Now let k_0 be the smallest integer such that $4^{k_0} = n$ and divide the unit interval into subintervals of equal length with length equal to 2^{-k_0} . Let f_n be the corresponding histogram estimate based on $V_n^?$. Now note that since functions $f \in H$ are bounded below by $\mu_0 > 0$ it follows that

$$(2.11) \quad \int_0^1 \frac{p}{(f_n - p)^2} - \int_0^1 \frac{p}{(f_n - p - \bar{f})^2} = \int_0^1 \frac{p}{(f_n - p - \bar{f})^2} \frac{(p - \bar{f}_n + p - \bar{f})^2}{\mu_0} = \int_0^1 \frac{(f_n - f)^2}{\mu_0};$$

Now

$$(2.12) \quad E \int_0^1 (f_n - f)^2 = E \int_0^1 (f_n - f_{k_0})^2 + \int_0^1 (f - f_{k_0})^2$$

and simple calculations show that the histogram estimate f_n satisfies $E f_n(x) = f_{k_0}(x)$ and $\text{Var} f_n(x) = f_{k_0}(x) \frac{2^{k_0}}{n}$. Hence,

$$(2.13) \quad n^{1=2} E \int_0^1 (f_n - f_{k_0})^2 = n^{1=2} \frac{2^{k_0}}{n} \rightarrow 2^{1=2 - k_0} \neq 0;$$

$$(2.14) \quad n^{1=2} \begin{matrix} Z & 1 \\ 0 & \end{matrix} (f - f_{k_0})^2 - \frac{1}{1=2} k f - f_{k_0} k_{1=2;2}^{1=2} k_0 ! 0:$$
$$(2.15) \quad n^{1/2} \sup_{f \in \mathcal{H}} E_0^{\mathbb{Z}^1} \left(\frac{1}{f_n} \overline{f} \right)^2 \rightarrow 0;$$

Now generate N , a Poisson random variable with expectation n and independent of V_n^2 . If $N > n$ generate $N - n$ conditionally independent observations $V_{n+1}^2, \dots, V_N^2$ with common density f_n . Finally let $(N; X_N) = (N; V_1^2, V_2^2, \dots, V_N^2)$ and write $R_{\frac{1}{n}}$ for this randomization from V_n^2 to $(N; X_N)$.

A map from the Poisson number of independent observations back to the fixed number of observations is obtained similarly. This time let $\hat{f}_n$ be the histogram estimator based on $(N; X_N)$. If $N < n$ generate $n - N$ additional conditionally independent observations with common density f_n . It is also easy to check that

$$(2.16) \quad n^{1/2} \sup_{f \in \mathcal{H}} E \left(\frac{1}{n} \sum_{i=1}^n \left(\hat{f}_n - \bar{f} \right)^2 \right) \rightarrow 0:$$

$$R_n^2 : (N; X_N) \rightarrow V_n :$$

Carter (2002) provides a transformation from a 2^{k_0} -dimensional multino-mial to a multivariate normal as in Theorem 4 such that the total-variation distance between the distributions is $O(k_0 2^{k_0 n^{-1/2}})$. The transformation is similar to ours in that it adds uniform noise and then uses the square root as a variance-stabilizing transformation. However, the covariance structure

of the multinomial complicates the issue and necessitates using a multi-resolution structure similar to the one applied here to the conditional experiments. The [Carter \(2002\)](#) result can be used in place of Theorem 4 to get a slightly weaker bound on the error in the approximation in Theorem 3 (because of the extra k_0 factor) when the total number of observations is fixed. This is enough to establish Theorem 2 if the inequalities bounding γ_1 and γ_2 are changed to strictly greater than. It is also enough to establish Theorem 1 if H is a Besov space with $s > \frac{1}{2}$. [Carter \(2000\)](#) also showed that a somewhat more complicated transformation leads to a deficiency bound on the normal approximation to the multinomials without the added k_0 factor.

3. Main theorem. The theorems in Section 1 on the equivalence of white noise with drift experiments and Poisson process experiments are consequences of the following theorem which uniformly bounds the Hellinger distance between the randomized mappings described in Section 2.

Theorem 3. Suppose in $f_{0 < x < 1}(x) = f_0 > 0$. Let $W_{[k_0, k]}^? \sim \overline{fZ}_{k_0}^?; W_j^?; k_0 < j < kg$ with the variables in (2.1) and (2.2), and $W_{[k_0, k]} \sim \overline{fZ}_{k_0}; W_j; k_0 < j < kg$ with the variables in (2.6) and (2.9). Then there exist universal constants C, D_1 and D_2 such that for all $k_1 > k_0$,

$$H^2(W_{[k_0, k_1]}^?; W_{[k_0, k_1]}) \leq \frac{C}{f_0} \frac{4^{k_0}}{n} + \frac{D_1}{f_0^2} \sum_{k=k_0}^{k_1} 2^{k-1} \sum_{k_j=0}^{2^{k-1}} \gamma_{k_j}^2 + \frac{D_2}{f_0^3} \frac{n}{4^{k_0}} \sum_{k=k_0}^{k_1} 2^{3k-1} \sum_{k_j=0}^{2^{k-1}} \gamma_{k_j}^4,$$

$$\frac{C}{f_0} \frac{4^{k_0}}{n} + \frac{D_1}{f_0^2} k f_{k_0} k_{1=2;2;2}^2 + \frac{D_2}{f_0^3} \frac{n}{4^{k_0}} k f_{k_0} k_{1=2;4;4}^4;$$

where γ_{k_j} are the Haar coefficients of f as in (1.3), f_k is as in (1.6) and $k_{1=2;p;p}$ are the Besov norms in (1.5).

Remark 3. Here the universal constant C is the same as the one in Theorem 4, while $D_1 = \frac{3D}{8} + 2$ and $D_2 = \frac{D}{9} + \frac{8}{3}$ for the D in Theorem 5.

The proof of Theorem 3 is based on the inequalities established in Sections 4 and 5 for the normal approximation of Poisson and Binomial variables. Some additional technical lemmas are given in the Appendix.

Let $X_{m,p}$ be a Binomial variable, X be a Poisson variable with mean λ , and $\mathcal{U}$ be a uniform variable in $[\frac{1}{2}, \frac{1}{2}]$ independent of $X_{m,p}$ and X . Define

$$(3.1) \quad g_{m,p}(x) = \frac{d}{dx} P f^{-1}(F_m(X_{m,p} + \mathcal{U})) - x g$$

with the F_m in (2.8) and the $N(0;1)$ distribution function Φ , and define

$$(3.2) \quad g(x) = \frac{d}{dx} P \{ \text{sgn}(X + U) = x \} = \frac{d}{dx} P \{ X + U \leq x \} = g(x).$$

Write f_b for the density of $N(b;1)$ variables.

Proof of Theorem 3. Let $g_{[k_0, k]}^2(w_{[k_0, k]})$ and $g_{[k_0, k]}(w_{[k_0, k]})$ be the joint densities of $W_{[k_0, k]}^?$ and $W_{[k_0, k]}$, $g_k^2(w_k)$ be the joint density of $W_k^?$, and $g_k(w_k | w_{[k_0, k]})$ be the conditional joint density of W_k given $W_{[k_0, k]}$. Since $W_k^?$ is independent of $W_{[k_0, k]}^?$,

$$\frac{P \{ g_{[k_0, k]}^2(w_{[k_0, k]}) \}}{g_{[k_0, k]}^2(w_{[k_0, k]})} = \frac{P \{ g_{[k_0, k+1]}^2(w_{[k_0, k+1]}) \}}{g_{[k_0, k+1]}^2(w_{[k_0, k+1]})} = \frac{P \{ g_{[k_0, k]}^2(w_{[k_0, k]}) \}}{g_{[k_0, k]}^2(w_{[k_0, k]})} (1 - \frac{P \{ g_k^2(w_k) \}}{g_k^2(w_k)});$$

so that the Hellinger distance can be written as

$$(3.3) \quad \begin{aligned} H_f^2(W_{[k_0, k_1]}^?; W_{[k_0, k_1]}) &= 2^{-1} \int \frac{P \{ g_{[k_0, k_1]}^2(w_{[k_0, k_1]}) \}}{g_{[k_0, k_1]}^2(w_{[k_0, k_1]})} \\ &= 2^{-1} \int \frac{P \{ g_{[k_0, k_0+1]}^2(w_{[k_0, k_0+1]}) \}}{g_{[k_0, k_0+1]}^2(w_{[k_0, k_0+1]})} \\ &\quad + \sum_{k_0 < k < k_1} \int \frac{P \{ g_{[k_0, k]}^2(w_{[k_0, k]}) \}}{g_{[k_0, k]}^2(w_{[k_0, k]})} (1 - \frac{P \{ g_k^2(w_k) \}}{g_k^2(w_k)}) \\ &= H_f^2(\bar{Z}_{k_0}^?; \bar{Z}_{k_0}) + \sum_{k_0 < k < k_1} \int \frac{P \{ g_{[k_0, k]}^2(w_{[k_0, k]}) \}}{g_{[k_0, k]}^2(w_{[k_0, k]})} H^2(g_k^?; g_k): \end{aligned}$$

At the initial resolution level k_0 , $N_{k_0, i}$ are independent Poisson variables by (2.5), so that $\bar{Z}_{k_0, i}$ are independent. This and the independence of $\bar{Z}_{k_0, i}^?$ from (2.1) imply

$$H_f^2(\bar{Z}_{k_0}^?; \bar{Z}_{k_0}) = \sum_{i=0}^{2^{k_0}-1} H_f^2(\bar{Z}_{k_0, i}^?; \bar{Z}_{k_0, i}):$$

By (2.6) and (3.2) $\bar{Z}_{k_0, i} =_{k_0}$ have densities $g_{k_0, i}$, while $\bar{Z}_{k_0, i}^? =_{k_0}$ are $N(h_{k_0, i}; 1)$ variables by (2.3). Thus, Theorem 4 can be used to obtain

$$H_f^2(\bar{Z}_{k_0, i}^?; \bar{Z}_{k_0, i}) = H_f^2(g_{k_0, i}; h_{k_0, i}) = \frac{C}{k_0} + \frac{1}{2} \int \frac{P \{ g_{k_0, i} \}}{g_{k_0, i}} \frac{h_{k_0, i}^2}{k_0}.$$

Since $f_{k, i} = f_{k, i}(4^{-k})$ by (2.7) and $2^k = 2^{k-2}n$ by (2.3), the above calculation yields

$$H_f^2(\bar{Z}_{k_0}^?; \bar{Z}_{k_0}) \leq C \sum_{i=0}^{2^{k_0}-1} \frac{2^{k_0}}{nf_{k_0, i}} + \sum_{i=0}^{2^{k_0}-1} \frac{2^{k_0}}{2^{k_0}} \left(\frac{P \{ f_{k_0, i} \}}{f_{k_0, i}} - h_{k_0, i} \right)^2$$

(3.4)

$$C \frac{2^{2k_0}}{n_0^2} + \sum_{i=0}^{2^{k_0}-1} \frac{n_0^{2k_0}}{2^{n_0^3}} \sum_{I_{k_0,i}} (f - f_{k_0,i})^2$$

by Lemma 1 (i) and the bound $f \leq n_0$.

For $k > k_0$ and $0 \leq i < 2^{k-1}-1$, define

$$(3.5) \quad p_{k,i} = \frac{p_{m_{k,i/2}, (2p_{k,i/2}-1)}}{p_{k-1,i} (2p_{k,i/2}-1)};$$

where $p_{k,i/2}$ are as in (2.10), $p_{k,i} = f_{k,i}$, $n=2^k$ are as in (2.7), and the functions $m_{k,i/2} = m_{k,i/2}(w_{[k_0,k]})$ are defined by $N_{k-1,i} = m_{k,i/2}(W_{[k_0,k]})$. At a fixed resolution level $k > k_0$, and for $i = 0, \dots, 2^{k-1}-1$, $N_{k,i/2}$ are independent binomial variables conditionally on $W_{[k_0,k]}$, so that by (2.9) and (3.1) $W_{k,i/2} = N_{k-1,i}$ are independent variables with densities $g_{m_{k,i/2}, p_{k,i/2}}$ under the conditional density g_k . In addition, $W_{k,i/2}$ are independent normal variables with variance $\frac{2}{k-1}$ under g_k . Thus,

$$(3.6) \quad H^2(g_k, g_k) = \sum_{i=0}^{2^{k-1}-1} H^2(g_{m_{k,i/2}, p_{k,i/2}}, g_{m_{k,i/2}, p_{k,i/2}});$$

by (2.4), where $\frac{2}{k-1} = E W_{k,i/2} = \frac{p_{k-1,i}}{4n} h_{k-1,i}$. It follows from Theorem 5 and (3.5) that for fixed $w_{[k_0,k]}$,

$$(3.7) \quad H^2(g_{m_{k,i/2}, p_{k,i/2}}, g_{m_{k,i/2}, p_{k,i/2}}) \leq D(p_{k,i/2}, \frac{1}{2})^2 + m_{k,i/2}(p_{k,i/2}, \frac{1}{2})^4 + \frac{(p_{k,i/2}, \frac{2}{k-1})^2}{2};$$

Furthermore, it follows from Lemma 3 that

$$\begin{aligned} & \sum_{i=0}^{2^{k-1}-1} \frac{p_{m_{k,i/2}, p_{k,i/2}}^2}{g_{[k_0,k]}^2 g_{[k_0,k]}} (p_{m_{k,i/2}, p_{k,i/2}} - p_{k-1,i})^2 \\ & \leq \sum_{i=0}^{2^{k-1}-1} \frac{p_{m_{k,i/2}, p_{k,i/2}}^2}{g_{[k_0,k]} (p_{m_{k,i/2}, p_{k,i/2}} - p_{k-1,i})^4} \\ & = E \left(\frac{p_{m_{k,i/2}, p_{k,i/2}}^2}{N_{k-1,i} (p_{k-1,i})^4} \right) \leq 2; \end{aligned}$$

so that by (3.5),

$$\sum_{i=0}^{2^{k-1}-1} \frac{p_{m_{k,i/2}, p_{k,i/2}}^2}{g_{[k_0,k]}^2 g_{[k_0,k]}} (p_{k,i/2}, \frac{2}{k-1})^2 \leq 4(2p_{k,i/2}-1)^2 + 2(p_{k,i/2}, \frac{2}{k-1})^2;$$

Similarly, $\sum_{i=0}^{2^{k-1}-1} \frac{p_{m_{k,i/2}, p_{k,i/2}}^2}{g_{[k_0,k]}^2 g_{[k_0,k]}} m_{k,i/2}^2 \leq E N_{k-1,i}^2 = k-1+1=2$. Thus, by (3.7),

$$(3.8) \quad \sum_{i=0}^{2^{k-1}-1} \frac{p_{m_{k,i/2}, p_{k,i/2}}^2}{g_{[k_0,k]}^2 g_{[k_0,k]}} H^2(g_{m_{k,i/2}, p_{k,i/2}}, g_{m_{k,i/2}, p_{k,i/2}})$$

$$4D(p_{k,i/2}, \frac{1}{2})^2 + D(p_{k-1,i}, p_{k,i/2}, \frac{1}{2})^4 + (p_{k,i/2}, \frac{2}{k-1})^2;$$

with $D_1 = 3D = 8 + 2$. Now, by (2.10) and (1.3),

$$(3.9) \quad p_{k;2}, \quad \frac{1}{2} = \frac{\sum_{I_{k;2}} \frac{f}{2} \sum_{I_{k;2}+1} \frac{f}{f}}{\sum_{I_{k-1}} \frac{f}{f}} = \frac{p_{2^{k-1}}}{2f_{k-1}};$$

so that by (3.5), (2.7), the definition of $g_{k;2}$ in (3.6) and Lemma 1 (ii),

$$(3.10) \quad \begin{aligned} g_{k;2}, \quad g_{k;2} &= \frac{\sum \frac{p_{2^{k-1}}}{n f_{k-1}} \frac{2^{k-1}}{f_{k-1}}}{2^{k-1}} \frac{p_{2^{k-1}}}{4n} h_{k-1}, \\ &= \frac{p_{2^{k-1}}}{4n} \frac{2^{k-1}}{f_{k-1}} h_{k-1}, \\ &= \frac{p_{2^{k-1}}}{4n 2^{(k-1)=2-1}} \frac{2^{k-1}}{f_{k-1}^{3=2}} (f - f_{k-1})^2; \end{aligned}$$

Inserting (3.9) and (3.10) into (3.8) and summing over i via (3.6), we find

$$(3.11) \quad \begin{aligned} & \sum \frac{p_{2^{k-1}}}{g_{[k_0;k]} g_{[k_0;k]} H^2(g_k; g_k)} \\ & \stackrel{2^k X^{1-1}}{=} \sum \frac{p_{2^{k-1}}}{g_{[k_0;k]} g_{[k_0;k]} H^2(g_{m_{k;2}}, p_{k;2}, i_{k;2})} \\ & \stackrel{2^k X^{1-1}}{=} \sum \left(4D_1 \frac{2^{k-2}}{8f_{k-1}^2} + D_{k-1} \frac{4^{k-4}}{64f_{k-1}^4} \right. \\ & \quad \left. + \frac{n2^k}{2f_{k-1}^3} \sum_{I_{k-1}} (f - f_{k-1})^2 \right) \\ & \stackrel{2^k X^{1-1}}{=} \sum \left(\frac{D_1}{n_0^2} 2^{k-1} \frac{2}{f_{k-1}} + \frac{D}{16} + 1 \frac{n2^{k-1}}{n_0^3} \sum_{I_{k-1}} (f - f_{k-1})^2 \right); \end{aligned}$$

due to $i_{k;2} = n f_{k;2} = 2^k$ in (2.7) and $\sum_{I_{k;2}} \frac{2^k}{f_{k;2}} (f - f_{k;2})^2$.

Finally, inserting (3.4) and (3.11) into (3.3) and then using Lemma 2 yields

$$\begin{aligned} & H_f^2(W_{[k_0;k_1+1]}, W_{[k_0;k_1+1]}) \\ & \leq C \frac{2^{2k_0}}{n n_0} + \frac{D_1}{n_0^2} \sum_{k=k_0}^{k_X-2} 2^k \sum_{i=0}^{2^k-1} \frac{2^k}{f_{k;2}} \\ & \quad + \frac{D}{16} + \frac{3}{2} \sum_{k=k_0}^{k_X-2} \sum_{i=0}^{2^k-1} \frac{n2^k}{n_0^3} \sum_{I_{k;2}} (f - f_{k;2})^2 \end{aligned}$$

Let f and f denote the densities of $X^e + \mathcal{U}$ and X^e , respectively. Since $t(\cdot)$ is invertible, $H^2(X^e; \mathbb{Z}) = H^2(X^e + \mathcal{U}; X^e) = 2 \int_{\mathbb{R}^p} \frac{f^2}{f}$, so that it suffices to show

$$(4.5) \quad A^Z P_{ff} = 1 - \frac{C}{C_{\infty}}; \quad \lim_{f \rightarrow 1} C = \frac{7}{192}.$$

Since $\mathcal{U}$ is uniform, $f(x) = e^{-j}$ on $[j-1, j]$, so that

$$(4.6) \quad A = \sum_{j=0}^{\infty} f(j) \sum_{j=1}^{\infty} f(x) f(j) g^{1=2} dx:$$

Since $t^0(x) = j^{1=2}$, by (4.4) $f(x) = j^{1=2} \cdot (t(x) - 2^{p-})$. This gives

$$\frac{f(x)}{f(j)} = \frac{\exp \left[-\frac{(2^p \bar{x} - 2^p)^2}{2} \right]}{2^p \exp \left[-\frac{(2^p j - 2^p)^2}{2} \right]} = \exp \left[-\frac{(2^p \bar{x} - 2^p)^2 - (2^p j - 2^p)^2}{2} \right]; \quad j - \frac{1}{2} \leq x < j + \frac{1}{2};$$

for $j \geq 1$, in view of the Stirling formula $j! = \frac{p}{2} j^{j+1/2} \exp(-j - \frac{1}{2j})$, where

$$(4.7) \quad j(x) = \frac{p}{x} - \frac{p-1}{x^2} - \frac{\log x}{4} + \frac{1}{2} + \frac{j}{2} \log \frac{j}{x} + \frac{\log j}{4} - \frac{j}{2} + \frac{j}{2}$$

with $1 = (12j + 1) < j < 1 = (12j)$, for $j = 1; 2; \dots$. Now, by the mean-value theorem,

$$\int_{j-1}^{j+1} \frac{f(x)}{f(j)} dx = \int_{j-1}^{j+1} \exp \left[-\frac{1}{2} \left(\frac{f'(j)}{f(j)} (x-j) + \frac{f''(j)}{2f(j)} (x-j)^2 \right) \right] dx$$

for some $j_j \leq \frac{1}{2}$, with

$$(4.8) \quad \begin{matrix} s- \\ 0 \\ j \end{matrix}(\mathbf{x}) = \frac{1}{\mathbf{x}} - \frac{1}{4\mathbf{x}}; \quad \begin{matrix} p- \\ 0 \\ j \end{matrix}(\mathbf{x}) = \frac{1}{2\mathbf{x}^{3=2}} + \frac{1}{4\mathbf{x}^2}:$$

Since $\exp[-\frac{1}{2}(x_j - j)^2]$ is symmetric about j , it follows that

$$(4.9) \quad \int_{j-1=2}^{j+1=2} \frac{f(x)}{f(j)} dx = \int_{j-1=2}^{j+1=2} \exp \left[j(j) + \frac{f''(x_j)(x-j)^2}{2} + \sum_{k=3}^{\infty} \frac{f^{(k)}(j)(x-j)^{2k}}{(2k)!} \right] dx :$$

Now, we shall take uniform Taylor expansions of $\mathbf{g}_j$ and their derivatives in

$$J_{fj:j} = 1j^{2=5}g:$$

By (4.7), $f_j(j) = f(j) + f_{j=2}$ with

$$f(x) = \left(\frac{x}{j} - 1\right)^2 + \frac{1}{2} \frac{x}{j} + \frac{x}{2} \log x:$$

Since $f(1) = f_0(1) = f_0(1) = 0$, $f_0(1) = 1/4$ and $f_0(1) = \frac{7}{8}$,

$$\frac{j}{4} = \frac{(j-1)^3}{3! \cdot 3} - \frac{7}{8} \frac{(j-1)^4}{4! \cdot 4} (1 + o(1)) = o(1):$$

Since $1 = (12j + 1) < j < 1 = (12j)$, $f_{j=2} = (1 + o(1)) = (24) = o(1)$. Thus,

$$f_j(j) = \frac{(j-1)^3}{24 \cdot 2} - \frac{7}{8} \frac{(j-1)^4}{24 \cdot 3} (1 + o(1)) + \frac{1 + o(1)}{24} = o(1)$$

uniformly in J as $j \rightarrow \infty$. Similarly, by (4.8) and $f_j(j) = \frac{1}{2}$,

$$f_j(j)g^2 = (1 + o(1)) \frac{(j-1)^2}{4 \cdot 2} + \frac{o(1)}{2} = o(1);$$

$$f_j(x_j) = \frac{1 + o(1)}{2} = o(1):$$

These expansions and (4.9) imply that uniformly in J ,

$$\begin{aligned} & \sum_{j=1}^{\infty} \frac{f_j(x)}{f_j(j)} dx \\ &= \sum_{j=1}^{\infty} \left(1 + f_j(j) + f_j(x_j) + (f_j(j))^2 g \frac{(x-j)^2}{2} \right. \\ & \quad \left. + o(1) \frac{x^2}{k+1} \right) \\ &= 1 + \frac{(j-1)^3}{24 \cdot 2} - \frac{7}{8} \frac{(j-1)^4}{24 \cdot 3} + \frac{1}{24} + \frac{1}{24} \frac{1}{2} + \frac{(j-1)^2}{4 \cdot 2} \\ & \quad + o(1) \frac{x^2}{k+1}; \end{aligned}$$

as $\sum_{j=1}^{\infty} (x-j)^2 dx = \frac{1}{12}$. Since $f_j(j)$ is the Poisson probability mass function of X^j ,

$$\begin{aligned} & \sum_{j=2}^{\infty} f_j(j) \sum_{j=1}^{\infty} \frac{f_j(x)}{f_j(j)} dx \\ (4.10) \quad &= 1 + \frac{1}{24} - \frac{7}{8} \frac{3}{24} + \frac{1}{24} - \frac{1}{96} + \frac{o(1)}{192} = 1 - \frac{7 + o(1)}{192} \end{aligned}$$

as $\sum_{j \neq J}^P f(j) = 1 + o(1) = 1$. Note that $E(X^3) = 3$ and $E(X^4) = 3^2 + 1$. Hence, (4.5) follows from (4.6), (4.10) and the fact that

$$\sum_{j \neq J}^X f(j) \sum_{j=1}^Z \frac{f(x)}{f(j)} \stackrel{1=2}{=} \int \frac{f(x)}{f(j)} dx = \int \frac{f(x)}{f(j)} dx = o\left(\frac{1}{n}\right);$$

5. Approximation of binomial variables. The strong approximation of a normal by a binomial depends on the cumulative distribution function F_m in (2.8). The addition of the independent uniform $\mathcal{U}$ in (2.8) to the binomial $X_{m,p}$ makes the c.d.f. continuous and thus F_m is a one-to-one function on $(-\frac{1}{2}; m + \frac{1}{2})$ that maps symmetric binomials to standard normals.

Let ϕ_b be the $N(b; 1)$ density and $g_{m,p}$ be the probability density of

$$(5.1) \quad F_m(X_{m,p} + \mathcal{U}); \quad X_{m,p} \sim \text{Bin}(m, p);$$

as in (3.1), where $\mathcal{U}$ is an independent uniform on $[-\frac{1}{2}; \frac{1}{2}]$.

Theorem 5. There is a constant $C_1 > 0$ such that, for all $m \geq 0$,

$$(5.2) \quad H^2(g_{m,p}; \phi_b) = \int \left(\frac{g_{m,p}(z)}{\phi_b(z)} - 1 \right)^2 dz \leq C_1 \left(\frac{b^2}{m} + \frac{b^8}{m^2} \right);$$

where $b = \left(\sum_{m=2}^P \log(p/(1-p)) \right)$. Consequently,

$$(5.3) \quad H^2(g_{m,p}; \phi) \leq \left(\frac{1}{2} \right)^2 + m \left(\frac{1}{2} \right)^4 + \frac{\left(\sum_{m=2}^P (2p-1) \right)^2}{2};$$

Proof. The case when $m = 0$ is trivial because $X = 0$ with probability 1 and therefore $g_{0,p}$ is exactly an $N(0; 1)$. Thus, the following assumes that $m \geq 1$.

It follows from (3.1) that

$$(5.4) \quad g_{m,p}(z) = p^j (1-p)^{m-j} 2^m \phi_0(z);$$

where $j = j(z)$ is the integer between 0 and m such that

$$(5.5) \quad F_m(j - \frac{1}{2}) \leq z < F_m(j + \frac{1}{2});$$

Let $\epsilon = \log(p/q)$ so that

$$\log \frac{g_{m,p}(z)}{\phi_0(z)} = j \frac{m}{2} + \frac{m \log(4pq)}{2};$$

and the second term can be approximated by

$$(5.6) \quad \frac{2}{4} - \frac{4}{24} \log(4pq) = \log \frac{2 + e + e}{4} - \frac{2}{4} + \frac{4}{32};$$

Let $h_1(\cdot) = (2 + e^{-\frac{1}{j}} + e^{-\frac{1}{j^2}}) = 4$. The second inequality in (5.6) follows from $\log(h_1(\cdot)) \leq \log(1 + \frac{1}{j^2} + \frac{1}{j^4}) \leq \frac{1}{j^2} + \frac{1}{j^4} = 32$. The first inequality in (5.6) follows from $h_1(\cdot) \leq 1 + \frac{1}{j^2} + \frac{1}{j^4} = 24$ for $j \geq 4$, and from $\log(h_1(\cdot)) \leq \frac{1}{j^2} + \frac{1}{j^4} = 4$ for $j > 4$. Now, let

$$(5.7) \quad z^0 = z^0(z) = \frac{j(z)}{P_{m=2}^{\infty}} \quad \text{and} \quad b = \frac{P_m}{2}:$$

Then for some $e^{-1} = 24$ $h_2(\cdot) \leq 32$ the log ratio is

$$\log \frac{g_{m,p}(z)}{g_{m,p}^0(z)} = z^0 b + \frac{b^2}{2} + h_2(\cdot) m^{-4}:$$

The log ratio of normals with different means is $\log(g_{m,p}^0/b) = -z^0 b + \frac{b^2}{2} = 2$.

Therefore the ratio with respect to the normal with mean b is

$$(5.8) \quad \log \frac{g_{m,p}}{g_{m,p}^0} = h_2(\cdot) m^{-4} - b(z - z^0); \quad |h_2(\cdot)| \leq \frac{1}{24}:$$

Since $y \log(x=y) \leq x - y \leq x \log(x=y)$, for all positive x and y ,

$$\frac{1}{2} P_{m,p} \log \frac{g_{m,p}^0}{g_{m,p}} \leq P_{m,p} \log \frac{g_{m,p}^0}{g_{m,p}} \leq \frac{1}{2} P_{m,p} \log \frac{g_{m,p}^0}{g_{m,p}};$$

so that by (5.8),

$$(5.9) \quad H^2(g_{m,p}; g_{m,p}^0) \leq \frac{1}{4} \int_0^Z \log \frac{g_{m,p}^0}{g_{m,p}} (g_{m,p}^0 + g_{m,p}) dz + \frac{m^{-4}}{24} + \frac{b^2}{2} \int_0^Z (z - z^0)^2 (g_{m,p}^0 + g_{m,p}) dz:$$

It follows from [Carter and Pollard \(2004\)](#) that the difference between z and $z^0 = z^0(z)$ is bounded by

$$(5.10) \quad |z - z^0| \leq \begin{cases} C_2 (m^{-1/2} + m^{-1/3}) & \text{for all } z \\ C_2 (m^{-1/2} + m^{-1/3}) & \text{if } |z| \leq \frac{1}{2m}; \end{cases}$$

for some constant C_2 . Thus,

$$(5.11) \quad \int_0^Z (z - z^0)^2 g_{m,p} dz \leq 2C_2^2 \left(\frac{1}{m} + \frac{1}{m^2} \int_0^Z g_{m,p} dz + \int_{|z| > 2m} \frac{z^6}{m^2} g_{m,p} dz \right):$$

Since $\int_0^R g_{m,p} dz = (j - m^{-2}) P_m = P_m - jg$,

$$\int_0^Z z^6 g_{m,p} dz = E \left[\frac{X_{m,p}^6}{P_m} \right] = O \left(1 + m^{-3} \frac{1}{2} \right) = O(1 + b^6)$$

uniformly in (m, p) . It follows from (5.4) that

$$\int_{|z| > 2m} z^6 g_{m,p} dz \leq 2^m \int_{|z| > 2m} z^6 g_{m,p} dz = O(2^m m^{-6} e^{-m}) = O(m^{-1}):$$

The above two inequalities and (5.11) imply

$$\int_{\mathbb{Z}} (z - z^0)^2 \mathfrak{g}_m \mathfrak{p} dz \leq 2C_2^2 O(1+m + b^6 m^2);$$

Similarly, $\int_{\mathbb{R}} (z - z^0)^2 \mathfrak{h}_b dz \leq 2C_2^2 O(1+m + b^6 m^2)$. Inserting these two inequalities into (5.9) yields (5.2) in view of (5.7).

Now let us prove (5.3). The Hellinger distance is bounded by 2, so that $b^8 m^2$ in (5.2) can be replaced by $b^4 m$ and it suffices to consider $\frac{1}{2}j - \frac{1}{4}$ for the proof of (5.3). By inspecting the infinite series expansion of $\log(\frac{p}{q}) = \log(1+x) - \log(1-x)$ for $x = 2p - 1$, we find that for $\frac{1}{2}j - \frac{1}{4}$, $j \log(\frac{p}{q}) \leq \frac{8}{3} 2^{p-1} j$ and $j \log(\frac{p}{q}) \geq 4(p - \frac{1}{2})j - \frac{8}{9} 2^{p-1} j^3$. These inequalities, respectively, imply

$$\frac{b^2}{m} + \frac{b^4}{m^2} \leq \frac{16}{9} (2p - 1)^2 + \frac{256}{81} m (2p - 1)^4$$

and $\frac{1}{m} (2p - 1)^2 \leq \frac{16}{81} m 2^{p-1} j - \frac{4}{81} m 2^{p-1} j^3$, in view of the definition of b , which then imply (5.3) via (5.2) and the fact that $H^2(\mathfrak{p}_b, \mathfrak{p}) = (b^2)^2 = 4$.

APPENDIX

A.1. The Tusnady inequality. The coupling of symmetric binomials and normal maps the integers j onto intervals $[j; j+1]$ such that the normal $(m=2; m=4)$ probability in the interval is equal to the binomial probability at $\frac{m}{j} 2^{-j}$. Taking the standardized values

$$z_j = \frac{2(j - \frac{m}{2})}{\sqrt{\frac{m}{2}}}; \quad u_j = \frac{2(j - 1 - \frac{m}{2})}{\sqrt{\frac{m}{2}}};$$

Carter and Pollard (2004) showed that for $m=2 < j < m$ and certain universal finite constants C

$$C \frac{u_j + 1}{m} \leq z_j - u_j \leq 1 + 2 \frac{u_j^2}{m} - \frac{u_j}{m} \leq \frac{\log(1 + u_j^2/m)}{2cu_j} \leq C + \frac{u_j + \log m}{m}$$

where $c = \sqrt{2 \log 2}$ and ϕ is an increasing function with $\phi(0) = 1/2$ and $\phi(1) = \log 2 - 1/2$.

This immediately implies that

$$(A.1) \quad \mathfrak{p}_j - u_j j \leq \frac{C_0}{m} (j j^3 + \log m) \leq 8 \frac{u_j^2}{m} + \frac{1}{2}$$

for a certain universal constant $C_0 < 1$. We shall prove (5.10) here based on (A.1). Because of the symmetry in both distributions, it is only necessary to consider $z > 0$.

It follows from (5.5) and (5.7) that

$$z_j - z < z_{j+1} - z \quad () \quad u_j - z^0 = z^0(z) < u_{j+1} :$$

Let $z_j - z < z_{j+1} - z$. Since $u_{j+1} - u_j = 2^{-\frac{p}{m}}$, for $u_{j+1}^2 - u_j^2 = 2^{-\frac{p}{m}}$ (A.1) implies

$$(A.2) \quad z^0_j - z_j - u_j \leq z_{j+1} - u_{j+1} + \frac{2}{m} C_0^0 \frac{1}{m} + \frac{z_j^3 \wedge z_j^{03}}{m} :$$

Since u_j and z_j are both increasing in j , it follows that $(z \wedge z^0) = \frac{p}{m}$ are uniformly bounded away from zero for $u_{j+1} = 2^{-\frac{p}{m}}$, so that

$$(A.3) \quad z^0_j - z_j - u_j \leq z_{j+1} - u_{j+1} + \frac{2}{m} C_0^0 \frac{z_j^3 \wedge z_j^{03}}{m}$$

for $(m+1) = \frac{p}{m} = u_{m+1} - u_{j+1} = 2^{-\frac{p}{m}}$ and $z = \frac{p}{2m}$. Since $z - z^0 = z - u_{m+1} = \frac{p}{2m}$ for $z > \frac{p}{2m}$, (A.2) and (A.3) imply

$$z^0_j - z_j = \begin{cases} C_2 (m^{-1/2} + m^{-1} z_j^3); & \text{for all } z_j \\ C_2 (m^{-1/2} + m^{-1} z_j^{03}); & \text{if } z_j \leq \frac{p}{2m}; \end{cases}$$

for a certain universal $C_2 < 1$, that is, (5.10).

A.2. Technical lemmas. The following three lemmas simplify the rest of the proof of Theorem 3.

Lemma 1. (i) Let $f_{k,i}$, and $h_{k,i}$, be as in (2.7) and (2.3): Then

$$(A.4) \quad 0 \leq \frac{p}{f_{k,i}} - h_{k,i} \leq 2^{k-1} f_{k,i}^{3/2} (f - f_{k,i})^2 :$$

(ii) Let k_i , be the Haar coefficients of f as in (1.3): Then

$$(A.5) \quad h_{k,i} \leq \frac{p^{k_i}}{2 f_{k,i}} \leq 2^{k-1} f_{k,i}^{3/2} (f - f_{k,i})^2 :$$

Proof. Let $T = (f - f_{k,i}) = f_{k,i}$. 1. By algebra,

$$\frac{p}{1+T} - 1 = \frac{T}{1 + \frac{T}{1+T}} = \frac{T}{2} \frac{T^2}{2(1 + \frac{T}{1+T})^2} :$$

It follows from (2.3) and (2.7) that

$$\begin{aligned} h_{k,i} &= 2^k \frac{p}{f_{k,i}} - \frac{p}{1+T} \\ &= 2^k \frac{p}{f_{k,i}} - 1 + \frac{f - f_{k,i}}{2f_{k,i}} - \frac{(f - f_{k,i})^2}{2f_{k,i}^2(1 + \frac{f - f_{k,i}}{1+T})^2} ; \end{aligned}$$

which implies (A.4) as $2^{k^R}_{I_{k_i}} = 1$ and by (2.7) $^R_{I_{k_i}}(f - f_{k_i}, \cdot) = 0$. For (ii) we have

$$\begin{aligned} Z_{h_{k_i}} &= \frac{p}{f_{k_i}} \cdot \frac{p}{1+T} \\ &= \frac{p}{f_{k_i}} \cdot \frac{f}{1 + \frac{f}{2f_{k_i}}} \cdot \frac{(f - \frac{f_{k_i}}{p})^2}{2f_{k_i}^2 (1 + \frac{f}{2f_{k_i}})^2} ; \end{aligned}$$

which implies (A.5) as $\sum_{k=1}^R j_{k,i} = 0$ and $\sum_{k=1}^R j_{k,i}^2 \leq 2^k$ by (1.4).

Lemma 2. Let $\kappa_{k,\cdot}$ be the Haar coefficients in (1.3) and $f_{k,\cdot}$ be as in (2.7): Then

$$x^{\frac{1}{k}} \cdot x^{\frac{2k-1}{k}} = x^{\frac{k+2k-1}{k}} = x^{\frac{3k-1}{k}} \quad (\text{f}) \quad f_{k;'}^2 = \frac{2^{ck_0} x^{\frac{1}{k}} \cdot x^{\frac{2k-1}{k}}}{(1 - 2^c)^2} \cdot x^{\frac{1}{k}} \cdot x^{\frac{2k-1}{k}} = \frac{2^{ck_0} x^{\frac{3k-1}{k}}}{(1 - 2^c)^2}, \quad 8 < c > 0:$$

Proof. De ne

$$I_{i;j;k} = \begin{cases} 1; & \text{if } I_{i;j} = I_{k;}, \\ 0; & \text{otherwise.} \end{cases}$$

Since $\sum_{j=1}^p |a_{ij;j,k}|^2 = 2^{i-k}$ for $i \leq k$, using Cauchy-Schwarz twice yields

$$\begin{aligned} Z &= \sum_{I_k;'} (f_k)^2 = \sum_{i=k}^1 \sum_{j=0}^{2^i-1} X_{i,j;k}'^2 \\ &= \sum_{i=k}^1 \sum_{j=0}^{2^{ic}-1} \sum_{k'=0}^{2^{ic}-1} X_{i,j;k}'^2 X_{i,k';j}^2 \\ &= \sum_{i=k}^1 \sum_{j=0}^{2^{ic}-1} \sum_{k'=0}^{2^{ic}-1} X_{i,j;k}'^2 X_{i,k';j}^2 \\ &= \frac{\sum_{i=k}^1 \sum_{j=0}^{2^{ic}-1} X_{i,j;k}'^2}{1=2^c} \sum_{j=0}^{2^{ic}-1} X_{i,j;k}'^2 : \end{aligned}$$

Since $\sum_{i,j;k}^k p_{i,j;k} = 1$ for $i \leq k$, the above inequality implies

$$\sum_{k=k_0}^1 2^k \frac{X^{1+2^k}}{1-2^{2^k}} \sum_{i=k}^1 2^{i(1+c)} \sum_{j=0}^{2^k-1} X^{2^j} \sum_{i;j;k;\backslash}^4$$

Lemma 3. Let X be a Poisson random variable with mean λ . Then

Proof. Since $E(X^e)^4 = (3 + 1)$,

A.3. Proof of Theorem 1. First note that

and

Note also that since for any randomization T and random X and Y , $H(TX;TY) \leq H(X;Y)$, it follows that

and

For the class H and the randomizations R_n^1 and R_n^2 it follows from (2.15), (2.16) and the proof of Proposition 3 on page 508 of Le Cam (1986) that

and

Hence (1.9) and (1.8) will follow once

$$(A.6) \quad \sup_{f \in H} H(T_n(N; X_N); Z_n^?) \rightarrow 0$$

is established.

By Theorem 3, for (A.6) to hold it is sufficient to show that

$$\sup_{f \in H} \frac{4^{k_0}}{n} + kf_{k_0} k_{1=2;2;2}^2 + \frac{n}{4^{k_0}} kf_{k_0} k_{1=2;4;4}^4 \rightarrow 0:$$

If the class of functions H is a compact set in the Besov spaces, then the partial sums converge uniformly to 0,

$$\sup_{f \in H} kf_{k_0} k_{1=2;p;p} \rightarrow 0$$

for $p = 2$ or 4 as $k \rightarrow \infty$. This implies that there is a sequence $k \rightarrow \infty$ such that $\sup_{f \in H} kf_{k_0} k_{1=2;4;4}^4 \rightarrow 0$. To be specific, let

$$k = \sup_{f \in H} kf_{k_0} k_{1=2;4;4}^2:$$

It is necessary to choose the sequence of integers $k_0(n)$ that will be the critical dimension that divides the two techniques. Let k_0 be the smallest integer such that $\frac{4^{k_0}}{n} \rightarrow 0$. Therefore, $k_0(n) \rightarrow \infty$, and as $n \rightarrow \infty$,

$$\begin{aligned} \sup_{f \in H} \frac{4^{k_0}}{n} + kf_{k_0} k_{1=2;2;2}^2 + \frac{n}{4^{k_0}} kf_{k_0} k_{1=2;4;4}^4 \\ \sup_{f \in H} 4^{-k_0} + kf_{k_0} k_{1=2;2;2}^2 + \frac{1}{4^{k_0}} kf_{k_0} k_{1=2;4;4}^4 \rightarrow 0: \end{aligned}$$

A.4. Proof of Theorem 2. Theorem 2 follows from Theorem 1 and the fact that the Lipschitz and Sobolev spaces described are compact in the Besov spaces.

The Lipschitz class is equivalent to $B_{\infty,1;1}$ and therefore is compact in $B_{1=2;p;p}$ if $p > \frac{1}{2}$. The Sobolev class is equivalent to $B_{2;2;2}$ and

$$kf_{k_0} k_{2;2;2}^2 \leq C \sum_n \int_n (f)^2 n^2;$$

where C depends only on γ . Thus if F is compact in $\text{Sobolev}(\gamma)$ for $\gamma > \frac{1}{2}$ then it is compact in $B_{1=2;2;2}$.

Further restrictions are required to show that the $\text{Sobolev}(\gamma)$ class is compact in $B_{1=2;4;4}$. If $kf_k \in C_{(L)}^{(L)}$, then $kf_k - f_{k+1} k_1 \in C_{(L)}^{(L)} 2^{-k}$, so that

$$\begin{aligned} kf_{k_0} k_{1=2;4;4}^4 &\leq C_{(L)}^2 \sum_{k=k_0}^{\infty} 2^{k2(L)} \int_k - f_{k+1} \int^2 dx \\ &= C_{(L)}^2 kf_{k_0} k_{(L)}^2_{(1);2;2}: \end{aligned}$$

Therefore, for F bounded in Lipschitz(1), a compact Sobolev(1) set is also compact in $B_{1=2;4;4}$ if $\alpha = 1$.

Finally, if F is compact in Sobolev(1), $\alpha = 1$, then it immediately follows from the Sobolev embedding theorem that the function is bounded in Lipschitz(1=4) [e.g., Folland (1984), pages 270 and 273], and it follows that F is compact in $B_{1=2;4;4}$.

Acknowledgments. We thank the referees and an Associate Editor for several suggestions which led to improvements in the final manuscript.

REFERENCES

- Brown, L.D., Cai, T., Low, M.G. and Zhang, C.H. (2002). A asymptotic equivalence theory for nonparametric regression with random design. *Ann. Statist.* 30 688{707. [M R 1922538](#)
- Brown, L.D. and Low, M.G. (1996). A asymptotic equivalence of nonparametric regression and white noise. *Ann. Statist.* 24 2384{2398. [M R 1425958](#)
- Brown, L.D. and Zhang, C.H. (1998). A asymptotic nonequivalence of nonparametric experiments when the smoothness index is 1=2. *Ann. Statist.* 26 279{287. [M R 1611772](#)
- Carter, A.V. (2000). A asymptotic equivalence of nonparametric experiments. Ph.D. dissertation, Yale Univ.
- Carter, A.V. (2002). Efficiency distance between multinomial and multivariate normal experiments. *Ann. Statist.* 30 708{730. [M R 1922539](#)
- Carter, A.V. and Pollard, D. (2004). Tusnady's inequality revisited. *Ann. Statist.* 32. To appear.
- Efromovich, S. and Samarov, A. (1996). A asymptotic equivalence of nonparametric regression and white noise has its limits. *Statist. Probab. Lett.* 28 143{145. [M R 1394666](#)
- Folland, G.B. (1984). *Real Analysis*. Wiley, New York. [M R 767633](#)
- Golubev, G. and Nussbaum, M. (1998). A asymptotic equivalence of spectral density and regression estimation. Technical report, Weierstrass Institute, Berlin.
- Gramma, I. and Nussbaum, M. (1998). A asymptotic equivalence for nonparametric generalized linear models. *Probab. Theory Related Fields* 111 167{214. [M R 1633574](#)
- Le Cam, L. (1964). Sufficiency and approximate sufficiency. *Ann. Math. Statist.* 35 1419{1455. [M R 207093](#)
- Le Cam, L. (1986). *Asymptotic Methods in Statistical Decision Theory*. Springer, New York. [M R 856411](#)
- Le Cam, L. and Yang, G. (1990). *Asymptotics in Statistics*. Springer, New York. [M R 1066869](#)
- Milstein, G. and Nussbaum, M. (1998). Diffusion approximation for nonparametric autoregression. *Probab. Theory Related Fields* 112 535{543. [M R 1664703](#)
- Nussbaum, M. (1996). A asymptotic equivalence of density estimation and Gaussian white noise. *Ann. Statist.* 24 2399{2430. [M R 1425959](#)

L.D.Brown
M.G.Low
Department of Statistics
The W harton School
University of Pennsylvania
Philadelphia, Pennsylvania 19104-6340
U S A
e-mail: lbrown@wharton.upenn.edu
e-mail: lowm@wharton.upenn.edu

A.V.Carter
Department of Statistics
and Applied Probability
University of California, Santa Barbara
Santa Barbara, California 93106-3110
U S A
e-mail: carter@pstat.ucsb.edu

C.-H. Zhang
Department of Statistics
504 Hill Center, Busch Campus
Rutgers University
Piscataway, New Jersey 08854-8019
U S A