

Long range action in networks of chaotic elements

Michael Blank*, Leonid Bunimovich†

January 14, 2022

Abstract. We show that under certain simple assumptions on the topology (structure) of networks of strongly interacting chaotic elements a phenomenon of long range action takes place, namely that the asymptotic (as time goes to infinity) dynamics of an arbitrary large network is completely determined by its boundary conditions. This phenomenon takes place under very mild and robust assumptions on local dynamics with short range interactions. However, we show that it is unstable with respect to arbitrarily weak local random perturbations.

1 Introduction

Despite a number of impressive results in the field of coupled map lattices (CML) up to now only the situation when the interaction between individual systems is very weak is reasonably well understood from the mathematical point of view (see a review of recent results, e.g. in [4, 5]). In that case it is known that under mild technical assumptions the asymptotic behavior of a CML is very close to the behavior of a direct product of individual uncoupled systems. Note that counterexamples demonstrating very different types of behavior under arbitrary weak interactions are known as well (see [3] for details). On the other hand, numerical studies (see, e.g. [7]) suggest that when the interactions becomes stronger a kind of phase transition from one to several invariant SBR measures takes place. Despite a number of efforts during last ten years to give a mathematical description of this phenomenon no rigorous results for infinite systems have been obtained. Probably one of the reasons is that all mathematical results in the field are of perturbative nature and corresponding methods do not work well for large perturbations. Another problem here is the necessity to work with infinite collections of individual maps. Therefore, it is harder to study corresponding functional spaces.

In this paper we shall follow a more “physical” approach to the analysis of CMLs’ dynamics in a strong interactions case. Indeed, a general receipt in statistical physics to an analysis of an infinitely large system is to consider its restriction to a finite volume (box) with certain boundary conditions and to study what happens when the size of the box grows to infinity. The case when the dynamics inside of the (sufficiently large) box is not influenced by boundary conditions corresponds to the situation known as the case (range) of weak interactions (or as the absence of phase transitions). We shall be mainly interested in the opposite situation when the asymptotic (as time goes to infinity) dynamics inside of the box (with an arbitrary large diameter) is completely determined by the boundary conditions and which is reasonably to call the long range action (LRA). As we shall see the LRA takes place under sufficiently strong interactions in networks whose topologies satisfy the property which we call *unidirectional* and which generalizes well known one-dimensional CMLs with unidirectional interactions [8].

The paper is organized as follows. In the next section we give some technical definitions and formulate our main result about the existence of the LRA in a reasonably large class of CMLs (Theorem 2.1). Sections 3 and 4 contain the proof of this result and the proof of the equivalence of the notion of the unidirectional interaction to the absence of cycles in the directed connectivity graph of interactions in

*Russian Academy of Sciences, Inst. for Information Transmission Problems, and Observatoire de la Cote d’Azur, e-mail: blank@obs-nice.fr

†School of Mathematics, Georgia Institute of Technology; e-mail: bunimovh@math.gatech.edu

a network. Section 5 is dedicated to the discussion of the topology of networks corresponding to CMLs under study and to the influence of different types of boundary conditions on the existence of LRA. In Section 6 we study random perturbations of local dynamics and show that the LRA phenomenon is stochastically unstable. Finally in Section 7 we briefly address some open questions and compare our results to some situations known in the theory of synchronization.

2 The setup

Let $T : X \rightarrow X$ be a map from a compact metric space (X, ρ) into itself satisfying the Lipschitz condition with the contracting constant Λ_T , i.e.

$$\Lambda_T := \sup_{x \neq y \in X} \frac{\rho(Tx, Ty)}{\rho(x, y)} < \infty, \quad (2.1)$$

(e.g. an Anosov map on a finite-dimensional torus with the Euclidean metric). Then the pair (T, X) defines a dynamical system. Consider an infinite collection of dynamical systems (T_i, X) with $i \in \mathbb{Z}^d$ and a certain interaction between them. We always assume that

$$\Lambda_{T_i} \leq \Lambda_T < \infty$$

for all $i \in \mathbb{Z}^d$ and denote the direct product of the systems (T_i, X) by $(\vec{T}, \vec{X})$. The maps T_i we shall call *local* maps. Elements $\vec{x}$ of the infinite dimensional space $\vec{X}$ will be called *vectors* and their coordinates will be denoted as $x_i := (\vec{x})_i \in X$ with $i \in \mathbb{Z}^d$.

Definition 2.1 By the *interaction* on the lattice $\vec{X} := X^{\mathbb{Z}^d}$ we shall mean the map $\vec{\mathcal{I}} : \vec{X} \rightarrow \vec{X}$ preserving the ‘diagonal’, i.e. $\vec{\mathcal{I}}(C\vec{e}) = C\vec{e}$ for any constant C and the vector $\vec{e} \in \vec{X}$ with all entries equal to 1. We restrict ourselves to the case of continuous interactions satisfying the assumption that there exists a constant $\Lambda_{\vec{\mathcal{I}}} < 1$ such that

$$\rho((\vec{\mathcal{I}}\vec{x})_\ell, (\vec{\mathcal{I}}\vec{y})_\ell) \leq \Lambda_{\vec{\mathcal{I}}} \sum_{i \in \mathbb{Z}^d} \rho(x_i, y_i) \quad (2.2)$$

for any $\ell \in \mathbb{Z}^d$ and any vectors $\vec{x}, \vec{y} \in \vec{X}$.

In particular, this means that for any pair of vectors which differ at only one coordinate the map $\vec{\mathcal{I}}$ is a pure contraction. It is tempting to assume that the interaction $\vec{\mathcal{I}}$ is a pure contraction in the product metric, which would simplify a lot the further analysis. At the first sight it seems that the difference between the assumption (2.2) and the pure contraction is not essential. However a closer look shows that the latter contradicts to the definition of the interaction which says that the map $\vec{\mathcal{I}}$ preserves the ‘diagonal’, i.e. the distance between the vectors $\vec{\mathcal{I}}(C_1\vec{e})$ and $\vec{\mathcal{I}}(C_2\vec{e})$ is the same as the distance between the vectors $C_1\vec{e}$ and $C_2\vec{e}$ for any constants C_1, C_2 .

The *strength* of the interactions we shall measure by the difference $1 - \Lambda_{\vec{\mathcal{I}}}$. Indeed, without the interactions one has $\Lambda_{\vec{\mathcal{I}}} = 1$. A “typical” example: $\vec{\mathcal{I}}$ is a linear map represented by an infinite matrix with normalized rows. Local interactions correspond to the case when only terms near the diagonal of the matrix differ from zero. Again it is useful to have in mind a “standard” example of the local interaction (often called a “diffusive coupling”) on a one-dimensional lattice when after the interactions the new value at the site i is calculated according to the formula:

$$x_i \rightarrow (1 - 2c)x_i + c(x_{i-1} + x_{i+1}). \quad (2.3)$$

Here the interaction is parametrized by the value $0 \leq c \leq 1/2$ and $\Lambda_{\vec{\mathcal{I}}} = \max(1 - 2c, c)$.

Another important and well known example describes unidirectional interactions on a one-dimensional lattice defined by the formula:

$$x_i \rightarrow (1 - c)x_i + cx_{i-1}. \quad (2.4)$$

In this case $0 \leq c \leq 1$ and $\Lambda_{\vec{T}} = \max(1 - c, c)$.

Both these examples belong to the class of *short range* interactions when subsystems interact only if the Euclidian distance between the corresponding sites do not exceed a certain threshold. To emphasize that our results do not come as a consequence of long range interactions (when subsystems being arbitrary far apart may interact with each other) we shall restrict the consideration only to short range interactions.

Definition 2.2 The superposition $\vec{T} := \vec{I} \circ \vec{T}$ of the interaction and the direct product of local maps defines the dynamical system $(\vec{T}, \vec{X})$ which is called a *coupled map lattice* (CML).

Definition 2.3 A *box* $\mathcal{B}$ is a bounded subset of the integer lattice $\mathbb{Z}^d$.

To restrict the dynamics to the box $\mathcal{B}$ we define it as follows:

$$\left(\vec{T}|_{\mathcal{B}} \vec{x}\right)_i := \begin{cases} \left(\left(\vec{I} \circ \vec{T}\right) \vec{x}\right)_i & \text{if } i \in \mathcal{B} \\ x_i & \text{otherwise.} \end{cases} \quad (2.5)$$

Definition 2.4 A collection of values of $\vec{x}$ outside of the box $\mathcal{B}$ is called *boundary conditions* and is denoted by $\vec{x}|_{\mathcal{B}^c}$, where by $\mathcal{B}^c$ we denote the complement to the set $\mathcal{B}$.

If we choose certain boundary conditions the map $\vec{T}|_{\mathcal{B}}$ defines a finite dimensional dynamics inside a finite box $\mathcal{B}$. The collection of functions $x_i(t) := \left(\vec{T}|_{\mathcal{B}}^t \vec{x}(0)\right)_i$ with $i \in \mathcal{B}$ and $t \in \mathbb{Z}_+$ will be called a *solution* of our problem corresponding to the initial data $\vec{x}(0)$. Clearly $\vec{x}(0)$ can be decomposed into the genuine initial data $\vec{x}|_{\mathcal{B}}(0)$ and the boundary conditions $\vec{x}|_{\mathcal{B}^c}$.

Definition 2.5 By the *long range action* (LRA) we shall mean that for a given box $\mathcal{B}$ and given boundary conditions $\vec{x}|_{\mathcal{B}^c}$ there exists a vector $\vec{x}|_{\mathcal{B}}$ (called a *limit solution*) such that for any initial data $\vec{x}|_{\mathcal{B}}(0)$ the corresponding solution $\vec{x}(t)$ converges pointwise with time t to $\vec{x}|_{\mathcal{B}}$.

The main issue here is that the boundary conditions define the long term dynamics inside the box $\mathcal{B}$ in a unique way contrary to the usual situation when the dynamics inside the box is asymptotically independent on the boundary conditions. On the other hand, we do not assume that different boundary conditions lead to different solutions. The point is that if the local maps are not bijective a pair of very different boundary conditions may lead to the same limit solution. On the other hand, under some simple additional assumptions (of non-degeneracy type) close but different boundary conditions lead to different limit solutions.¹

Definition 2.6 We shall say that the interaction $\vec{I}$ in the box $\mathcal{B}$ is *unidirectional* if there is a positive function $\mathcal{N} : \mathcal{B} \rightarrow \mathbb{Z}^1$ such that

- $\forall i \neq j \in \mathcal{B}$ we have $\mathcal{N}(i) \neq \mathcal{N}(j)$;
- $\forall \vec{x} \in \vec{X}$ and $\forall i \in \mathcal{B}$ the value $(\vec{I}\vec{x})_i$ does not depend on the values x_j for all j with $\mathcal{N}(j) > \mathcal{N}(i)$.

In other words the function $\mathcal{N}$ is an enumeration according to which sites in $\mathcal{B}$ with larger numbers do not ‘interact’ with sites with smaller numbers.

It is convenient to describe a topology of interactions in terms of connectivity graphs.

Definition 2.7 A *connectivity graph* $G := G(\mathcal{B}, \vec{I})$ of the interaction $\vec{I}$ in the box $\mathcal{B}$ is a directed graph G , whose vertices correspond to the sites of $\mathcal{B}$. Besides the edge (i, j) connecting the vertex $i \in G$ to the vertex $j \in G$ belongs to G if and only if the value $(\vec{I}\vec{x})_j$ depends on the value x_i . The latter means that the value $(\vec{I}\vec{x})_j$ assumes at least two different values when we are changing x_i assuming that all other coordinates of $\vec{x}$ are fixed.

¹See a discussion in the end of Section 3.




Figure 1: Examples of unidirectional connectivity graphs

To take into account the boundary conditions we include to the graph G all edges leading from $\mathcal{B}^c$ to the points belonging to the box $\mathcal{B}$ and extend the function $\mathcal{N}$ to these sites setting its value to 0 on them.

For each site $\ell \in \mathcal{B}$ we denote by $\Gamma(\ell)$ the collection of sites in the box $\mathcal{B}$ from which there are directed paths in the graph G leading to ℓ , and denote by $L(\ell)$ the length of the shortest path in G leading to ℓ from the points outside the box $\mathcal{B}$. If there are no such paths we set $L(\ell) = \infty$.

Now we are ready to formulate the main result of the paper.

Theorem 2.1 *Let the interaction in a finite box $\mathcal{B}$ be unidirectional and let for any $i \in \mathcal{B}$ there be a path in $G(\mathcal{B}, \tilde{\mathcal{I}})$ from a site outside of the box $\mathcal{B}$ to i . Then for any boundary conditions the LRA takes place whenever $\Lambda_{\tilde{\mathcal{I}}} \Lambda_T < 1$.*

Note the local maps may be both regular and chaotic here. The simplest example of the unidirectional interaction is a linear one-dimensional (i.e. $d = 1$) system of unidirectionally interacting maps with the interaction described by the relation (2.4). Of course, if this situation would exhaust the list of examples there would be no reason to discuss them at all. It turned out that there is a much bigger class of systems with effectively unidirectional interactions. This class of interactions have a very simple and intuitive interpretation in terms of a connectivity graph.

Theorem 2.2 *An interaction is unidirectional if and only if the corresponding directed connectivity graph has no cycles.*

Two examples of graphs satisfying this property are depicted in Fig. 1. The left panel in Fig. 1 corresponds to the interaction similar to the one which is well known under the name Toom's North-East voting model in statistical physics (see e.g. [6]), while the right one demonstrates a bit more complex topology of interactions. Observe that in both examples the graphs may be continued periodically to an arbitrary large rectangular box.

3 Proof of Theorem 2.1

First we prove the following technical result.

Lemma 3.1 *Let $\{v_i(t)\}_{i=1}^N$ be a collection of functions $v_i(t) : \mathbb{Z}_+^1 \rightarrow X$ with the parameter $t \in \mathbb{Z}_+^1$ and let the function $u(t) : \mathbb{Z}_+^1 \rightarrow X$ be defined by the following (non-autonomous) relation:*

$$u(t+1) := \Phi(\tau_0 u(t), \tau_1 v_1(t), \tau_2 v_2(t), \dots), \quad (3.1)$$

where $u(0) := u_0 \in X$, the maps $\tau_i : X \rightarrow X$ satisfy the inequality $\Lambda_{\tau_i} \leq \Lambda_T$ for all i , and the map $\Phi : X^{N+1} \rightarrow X$ satisfy the inequality

$$\rho(\Phi(z_0, z_1, \dots), \Phi(z'_0, z'_1, \dots)) \leq \Lambda_\Phi \sum_{i=0}^N \rho(z_i, z'_i) \quad (3.2)$$

for any two sequences $\{z_i\}$, $\{z'_i\}$. Then whenever $\Lambda_\Phi \Lambda_T < 1$ there exists a function $\hat{u}(t) : \mathbb{Z}_+^1 \rightarrow X$ (depending on v_i) such that for any $u_0 \in X$ the solution $u(t)$ of the equation (3.1) with $u(0) = u_0$ converges with time at exponential rate to $\hat{u}(t)$, i.e. $\rho(u(t), \hat{u}(t)) \leq C\gamma^t$ for all $t \geq 0$ and some $\gamma < 1$, $C < \infty$. Moreover, if there exists a vector $\vec{v} \in X^N$ such that $\max_{1 \leq i \leq N} \rho(v_i(t), v_i) \leq \hat{C}\hat{\gamma}^t$ for all $t \geq 0$ and some $\hat{\gamma} < 1$, $\hat{C} < \infty$, then the limit $\hat{u} := \lim_{t \rightarrow \infty} \hat{u}(t)$ exists as well.

Proof. Choose two different initial values $u_0, u'_0 \in X$ and denote the corresponding solutions of the equation (3.1) by $u(t)$ and $u'(t)$ respectively. Then

$$\begin{aligned} & \rho(u(t+1), u'(t+1)) \\ &= \rho(\Phi(\tau_0 u(t), \tau_1 v_1(t), \tau_2 v_2(t), \dots), \Phi(\tau_0 u'(t), \tau_1 v_1(t), \tau_2 v_2(t), \dots)) \\ &\leq \Lambda_\Phi \rho(\tau_0 u(t), \tau_0 u'(t)) \leq \Lambda_\Phi \Lambda_T \rho(u(t), u'(t)) \leq \dots \leq (\Lambda_\Phi \Lambda_T)^t \rho(u_0, u'_0) \xrightarrow{t \rightarrow \infty} 0, \end{aligned}$$

which proves the first assertion with $C := \max_{x,y \in X} \rho(x, y)$ and $\gamma := \Lambda_\Phi \Lambda_T < 1$.

To prove the second assertion consider

$$\begin{aligned} \rho(u(t+1), u(t)) &= \rho(\Phi(\tau_0 u(t), \tau_1 v_1(t), \tau_2 v_2(t), \dots), \Phi(\tau_0 u(t-1), \tau_1 v_1(t-1), \tau_2 v_2(t-1), \dots)) \\ &\leq \Lambda_\Phi \Lambda_T \left(\rho(u(t), u(t-1)) + \sum_{i=1}^N \rho(v_i(t), v_i(t-1)) \right). \end{aligned}$$

Using the notation $\gamma := \Lambda_\Phi \Lambda_T < 1$ and continuing this estimate backward in time we get

$$\begin{aligned} \rho(u(t+1), u(t)) &\leq \gamma^2 \rho(u(t-1), u(t-2)) + \gamma^2 \sum_{i=1}^N \rho(v_i(t-1), v_i(t-2)) \\ &\quad + \gamma \sum_{i=1}^N \rho(v_i(t), v_i(t-1)) \\ &\leq \dots \leq \gamma^t \rho(u(1), u(0)) + \sum_{k=1}^t \gamma^k \sum_{i=1}^N \rho(v_i(t-k+1), v_i(t-k)) \\ &\leq \gamma^t \rho(u(1), u(0)) + 2N\hat{C} \sum_{k=1}^t \gamma^k \hat{\gamma}^{t-k+1} \\ &\leq \gamma^t \rho(u(1), u(0)) + 2N\hat{C}t (\max(\gamma, \hat{\gamma}))^{t+1} \xrightarrow{t \rightarrow \infty} 0 \end{aligned}$$

at exponential rate (since $\max(\gamma, \hat{\gamma}) < 1$), which yields the result. $\square$

Proof of Theorem 2.1. Choose any element $i \in \mathcal{B}$. According to the definition of the map $\vec{\mathcal{T}}|_{\mathcal{B}}$ and to the property of the unidirectional interaction the value $(\vec{\mathcal{T}}|_{\mathcal{B}}\vec{x})_i$ depends only on x_i and the values at sites with smaller indices according to the ordering given by the function $\mathcal{N}$.

Our strategy is to apply Lemma 3.1 consecutively (according to the enumeration $\mathcal{N}$) to all elements of the box $\mathcal{B}$ starting from the ‘smallest’ one (say $i \in \mathcal{B}$) having the number 1. In that case the corresponding value $(\vec{\mathcal{T}}|_{\mathcal{B}}\vec{x})_i$ depends only on x_i and the values at sites outside of the box $\mathcal{B}$ given by boundary conditions. Thus $x_i(t)$ is defined by the relation of type (3.1) with $\Lambda_\Phi := \Lambda_{\vec{\mathcal{T}}}$. Hence by Lemma 3.1 $x_i(t)$ converges to a certain limit $\hat{x}_i$ as time $t \rightarrow \infty$.

Then we apply Lemma 3.1 to the site corresponding to the second smallest value of $\mathcal{N}$, etc. Observe that the exponential rate of convergence needed to apply the second assertion of Lemma 3.1 follows from the first one.

It is important that in any given step of the procedure the values at sites with smaller indices (according to the enumeration $\mathcal{N}$) whose entries enter into the formula (3.1) and belong to the corresponding collection Γ do not depend on the value of the site under consideration. $\square$

Corollary 3.2 *Under the assumptions of Theorem 2.1 the limit solution $\vec{x}$ depends continuously on the boundary conditions.*

Let us study how a limit solution depends on changes in boundary conditions. To simplify calculations we consider here only the case of linear one-dimensional interactions given by (2.4) and identical one-dimensional maps $T_i \equiv T$ acting on a unit circle. We assume also that the boundary conditions are homogeneous, i.e. $x_i = v \ \forall i \in \mathcal{B}^c$, and in some neighborhood U of the point $v \in X$ the map T is linear and expanding, i.e. $Tx = ax + b$ with $|a| > 1$ for all $x \in U$. Following the same argument as in the proof of Theorem 2.1 we start with the site corresponding to the smallest positive value of the function $\mathcal{N}$. Denote the value of the corresponding limit solution (existing by Theorem 2.1) at this site by u . Then u satisfies the relation

$$u = (1 - c)Tu + cTv .$$

Substituting the local linear representation of T we get

$$u = (1 - c)(au + b) + c(av + b) ,$$

or

$$u = \frac{cav + b}{ca - (a - 1)} .$$

Choose another homogeneous boundary condition $v' \in U$. Then the corresponding limit solution can be written as

$$u' = \frac{cav' + b}{ca - (a - 1)} .$$

Therefore

$$|u - u'| = \frac{ca}{ca - (a - 1)} \cdot |v - v'|$$

and the constant $\frac{ca}{ca - (a - 1)} > 1$ since $|a| > 1$ and $(1 - c)a \leq \Lambda_{\vec{T}}\Lambda_T < 1$. Note that we need to assume here that $u, u' \in U$, so the neighborhood U should be large enough.

Continuing this procedure for sites with larger $\mathcal{N}$ -indices we see that the distance between limit solutions corresponding to the homogeneous boundary conditions v and v' grows exponentially with distance from the boundary of the box (at least until one of limit solutions leaves U).

4 Proof of Theorem 2.2

First assume that the interaction in $\mathcal{B}$ is unidirectional but the graph $G(\mathcal{B})$ has a cycle contrary to our claim. Then when we are moving along this directed cycle the enumeration function $\mathcal{N}(\cdot)$ can only grow which contradicts to the existence of the cycle.

Assume now that the directed graph $G(\mathcal{B})$ has no cycles. We construct the enumeration function $\mathcal{N}(\cdot)$ as follows. We shall say that a vertex $i \in \mathcal{B}$ is a *starting* one if there is no $j \in \mathcal{B}$ such that the edge $(j, i) \in G$, and *free* if there is no $j \in \mathcal{B}$ such that $(i, j) \in G$.

Clearly due to the absence of cycles in G we deduce that the set of starting vertices is non empty.

We start the enumeration procedure by choosing some starting vertex say $i \in \mathcal{B}$ and setting $\mathcal{N}(i) := 1$.

Assume now that we already have enumerated M vertices from $\mathcal{B}$. Then we normalize the function $\mathcal{N}$ in such a way that for each $k = 1, 2, \dots, M$ its k -th largest value becomes equal to k (i.e. it is an exact numbering).

Choose an arbitrary path $i_1, i_2, \dots, i_n$ in the graph G satisfying the properties that the function $\mathcal{N}$ is not defined on all vertices of this path, the vertex i_1 is starting or $\exists j \in \mathcal{B} : (j, i_1) \in G$ and $\mathcal{N}(j)$ is well defined, and either (a) i_n is free or (b) $\exists j' \in \mathcal{B} : (i_n, j') \in G$ and $\mathcal{N}(j')$ is well defined.

If the condition (a) is satisfied we set

$$\mathcal{N}(i_k) := M + k, \quad k = 1, 2, \dots, n,$$

otherwise if the condition (b) is satisfied we set

$$\mathcal{N}(i_k) := \mathcal{N}(j') - 1 + \frac{k}{n+1}, \quad k = 1, 2, \dots, n.$$

After this we normalize the function $\mathcal{N}$, choose a new path, etc., until no non-enumerated elements left in $\mathcal{B}$.

Our aim was to define the function $\mathcal{N}$ on this path in such a way that the unidirectional interaction property should hold with respect to previously numbered vertices. It is straightforward to check that the enumeration obtained on each step of our construction does satisfy this property. $\square$

5 Discussion of boundary conditions of different types

The result of Theorem 2.1 may suggest that at least in the case of homogeneous boundary conditions the mechanism leading to the LRA phenomenon is a simple propagation of the boundary conditions through the boundary into a box $\mathcal{B}$. In general this is not the case and the limit solution $\vec{\hat{x}}$ can be rather inhomogeneous in space. However this propagation phenomenon still may take place as we shall demonstrate in the following result. Assume that all local maps T_i are identical and denote this common local map by T .

Lemma 5.1 *Let the map T have a fixed point $\xi \in X$ (i.e. $T\xi = \xi$) and suppose that the boundary conditions are fixed at the value ξ , i.e. $x_i(0) := \xi \forall i \in \mathcal{B}^c$. Then under the conditions of Theorem 2.1 for any initial data $\vec{x}|_{\mathcal{B}}(0)$ the solution $\vec{x}(t)$ will converge with time to the constant solution with $(\vec{\hat{x}})_i \equiv \xi$ for all $i \in \mathcal{B}$.*

Proof. First, according to Theorem 2.1 there exists the only one limit solution $\vec{\hat{x}}$. On the other hand, by definition the interaction $\vec{\mathcal{I}}$ preserves the “diagonal” and hence any sequence with identical coordinates is preserved by $\vec{\mathcal{I}}$. Therefore the vector with identical coordinates $(\vec{\hat{x}})_i \equiv \xi$ for all $i \in \mathcal{B}$ is invariant for both the map and the interaction and is, indeed, the only solution to our problem. $\square$

It turns out that the solutions depend rather sensitively on the way how one defines the corresponding boundary conditions.

Let us start with a very mild change in the definition of the interaction between the sites inside of the box $\mathcal{B}$ and the outer ones. Previously we used the convention that in order to calculate the value $(\vec{\mathcal{T}}_{|\mathcal{B}}\vec{x})_i(t+1)$ one first applies the map $\vec{T}$ at all sites and then applies the interaction $\vec{\mathcal{I}}$ to the result of the previous operation. Contrary to this, assume that we apply the map only to the sites from the box $\mathcal{B}$ (but not to the outer sites) and denote the map describing the corresponding system by $\vec{\mathcal{T}}_{|\mathcal{B}}^{(0)}$. From the point of view of Theorem 2.1 this change does not matter much and its claim remains valid for the modified map $\vec{\mathcal{T}}_{|\mathcal{B}}^{(0)}$. Indeed, it is straightforward to check that the modified map $\vec{\mathcal{T}}_{|\mathcal{B}}^{(0)}$ with certain boundary conditions $\vec{x}'_{|\mathcal{B}^c}$ is equivalent to the original map $\vec{\mathcal{T}}_{|\mathcal{B}}$ with the boundary conditions $x_i \in T_i^{-1}x'$ for all $i \in \mathcal{B}^c$. Of course, we assume here that the preimages do exist, i.e. $T_i^{-1}x' \neq \emptyset$. On

the other hand, the result about the propagation of homogeneous boundary conditions (Lemma 5.1) is no longer valid. The problem is that the homogeneous vector $\vec{x}$ immediately becomes inhomogeneous under the action of $\vec{T}_{|\mathcal{B}}^{(0)}$.

The boundary conditions considered so far can be called *frozen* because they do not change in time. We shall consider now two other types of boundary conditions. The first of them, called *free*, corresponds to the situation when after each application of the map the values at the sites outside of the box $\mathcal{B}$ do change under T , i.e.

$$\left(\vec{T}_{|\mathcal{B}}^{(\text{free})} \vec{x}\right)_i := \begin{cases} \left((\vec{\mathcal{I}} \circ \vec{T}) \vec{x}\right)_i & \text{if } i \in \mathcal{B} \\ T_i x_i & \text{otherwise.} \end{cases} \quad (5.1)$$

Theorem 5.1 *Let the interaction in a finite box $\mathcal{B}$ be unidirectional, the boundary conditions are free, for any $i \in \mathcal{B}$ there be a path in $G(\mathcal{B}, \vec{\mathcal{I}})$ from a site outside of the box $\mathcal{B}$ to i , and let $\Lambda_{\vec{\mathcal{I}}} \Lambda_T < 1$. Then for any (initial) boundary conditions $\vec{x}_{\mathcal{B}^c}(0)$ there exists a vector-function $\vec{\hat{x}}(t)$ such that*

$$\rho\left(\left(\vec{T}_{|\mathcal{B}}^{(\text{free})}\right)^t \vec{x}(0), \vec{\hat{x}}(t)\right) \xrightarrow{t \rightarrow \infty} 0.$$

for any initial data $\vec{x}_{\mathcal{B}}(0)$.

Proof. Observe first that Lemma 3.1 has been proven for time dependent entries. Now Theorem 5.1 follows essentially from the same argument as the one used in the proof of Theorem 2.1. This proves the existence of the vector-function $\vec{\hat{x}}(t)$ which is the limit solution (depending on time now) to the problem with free boundary conditions. In fact the only difference to the frozen case is that the limit solution is no longer a constant. $\square$

A completely different situation takes place if one considers periodic boundary conditions. Of course, in this case a box $\mathcal{B}$ must have a rectangular shape² (otherwise the periodic continuation is not well defined). Then the dynamics inside the box is completely determined by the initial data inside the box $\mathcal{B}$. However, in distinction to the previous cases a limit solution might not be unique. Assume for simplicity that $T_i \equiv T$ for all i and the map T has two different periodic points ξ and η with non-intersecting trajectories. Consider two different initial data: $\vec{x}_i(0) := \xi$, $\vec{x}'_i(0) := \eta$ for $i \in \mathcal{B}$. Then the functions $\vec{x}(t) := \vec{T}^t \xi$ and $\vec{x}'(t) := \vec{T}^t \eta$ describes two different time-periodic solutions to the problem having no common limit points. In fact, we even do not need any specific property of the interaction here and use only the invariance of the diagonal.

6 Stochastic perturbations of local dynamics

Conditions under which the existence of the LRA has been proven in Theorem 2.1 look very robust and certainly remain valid under small deterministic perturbations of local dynamics.³ Our aim in this section is to show that the situation changes drastically if instead of deterministic perturbations one considers random ones. For a general introduction to random perturbations of chaotic dynamical systems we refer the reader to [2]. However, since the interaction map $\vec{\mathcal{I}}$ strictly speaking is not only non-contracting but non-hyperbolic as well, one cannot use general results here and everything has to be done “by hands”.

From now on we assume that all the assumptions of Theorem 2.1 are satisfied and hence the LRA phenomenon takes place for the unperturbed system. To simplify notation and to emphasize that the stochastic instability of the LRA is not caused by some artificial assumptions, throughout this section we assume that the local maps coincide (i.e. $T_i \equiv T$) and the local phase space X is a unit d -dimensional flat torus with the Euclidean metric on it. Note however that all results that we shall discuss remain valid when the maps T_i are different but satisfy to assumptions we shall make for T , and one can replace the unit torus by a smooth compact d -dimensional manifold.

²or a more general polygon which one can fill in by reflections the entire plane.

³Namely, the vector $\vec{\hat{x}}$ depends continuously on the local maps T_i .

Definition 6.1 A *stochastic perturbation* of a dynamical system (T, X) is a Markov chain with the same phase space X defined by the transition probabilities $Q(x, A)$ to jump from a point $x \in X$ to a Borel set $A \subset X$. By a *stochastically perturbed* system we mean a superposition of the original system and a perturbation, i.e. a new Markov chain defined by the transition probabilities $Q(Tx, A)$.

Definition 6.2 The *amplitude* of the stochastic perturbation is defined as the smallest number ε such that $Q(Tx, A) = 0$ whenever $\min_{y \in A} \rho(x, y) > \varepsilon$ for all $x \in X$. Therefore we shall use the notation Q_ε to emphasize that the amplitude of the perturbation is ε . We shall assume also that the transition probabilities are translationally invariant, i.e. for any $x, z \in X$ and a Borel set $A \subseteq X$

$$Q(x, A) \equiv Q(x + z, A + z) . \quad (6.1)$$

Note that the phase space X is a torus here and hence the addition is well defined.

To describe both deterministic and stochastic counterparts of the dynamics in the same terms it is useful to consider their action on the space of probabilistic measures $\mathcal{M}(X)$ on X . To distinguish between the pointwise action and the action on measures we shall use the upper index star for the latter (e.g. T^* stands for the action on measures for the map T). These objects are defined as follows:

$$T^*\mu(A) := \mu(T^{-1}A), \quad Q_\varepsilon^*\mu(A) := \int Q_\varepsilon(x, A)d\mu(x)$$

for any probabilistic measure $\mu \in \mathcal{M}(X)$ and any Borel set $A \subseteq X$. According to the previous definition an action of the stochastically perturbed system on measures is a product of these operators: $Q_\varepsilon^*T^*$, while the action of the stochastically perturbed CML on measures can be written as

$$\vec{T}_\varepsilon^* := \vec{T}^* \vec{Q}_\varepsilon^* \vec{T}^* : \mathcal{M}(\vec{X}) \rightarrow \mathcal{M}(\vec{X}) ,$$

where $\vec{Q}_\varepsilon^*$ stands for the direct product of the operators Q_ε^* corresponding to local stochastic perturbations.

Definition 6.3 A solution $\vec{\mu}_\varepsilon \in \mathcal{M}(\vec{X})$ to the equation $\vec{T}_\varepsilon^* \vec{\mu} = \vec{\mu}$ is called an *invariant measure* for the stochastically perturbed system.

In distinction to our previous results which were concerned mainly with the stability type analysis we will need a number of assumptions to prove the instability against stochastic perturbations. Let us introduce the following notation. For $\mu \in \mathcal{M}(X)$ denote by $\text{supp}(\mu)$ a *support* of the measure μ which defined as the intersection of all closed subsets of X of full μ -measure. We define also the collection of *projection operators* π_k acting on the space $\mathcal{M}(\vec{X})$ where the index $k \in \mathbb{Z}^d$ corresponds to the k -th component of the direct product space $\vec{X}$, i.e. $\pi_k \vec{\mu}$ stands for the projection of the measure $\vec{\mu} \in \mathcal{M}(\vec{X})$ to the k -th component of the direct product of $\mathcal{M}(X)$.

We shall use the notation $B_r(x)$ for the ball of radius r centered at a point $x \in \mathcal{B}(X)$ and δ_x for the δ -measure at point x . To control a spread out of an initial measure $\vec{\mu} \in \mathcal{M}(\vec{X})$ under the action of the dynamics we introduce the following notion and a list of assumptions.

Definition 6.4 By *upper/lower spreads* $S_\pm(\mu)$ of a measure $\mu \in \mathcal{M}(X)$ we shall mean the radius of the smallest ball which contains support of the measure μ in the case of $S_+(\mu)$, and the radius of the largest ball contained in the support of the measure μ in the case of $S_-(\mu)$.

The first of those functionals characterizes the maximal distance between points in the support of a measure, while the second one controls its concentration properties.

We assume that there are constants $\Lambda_T \geq \lambda_T > 1$ and $\lambda_Q, a, b, \sigma > 0$ such that

$$\lambda_T \rho(x, y) \leq \rho(Tx, Ty) \leq \Lambda_T \rho(x, y) \quad \forall x, y : \rho(x, y) \leq \sigma , \quad (6.2)$$

$$\text{supp}(Q_\varepsilon^* \delta_x) \supseteq B_{\lambda_Q \varepsilon}(x) \quad \forall x \in X , \quad (6.3)$$

$$S_+(\pi_\ell \vec{\mathcal{I}}^* \vec{\mu}) \geq a S_+(\pi_\ell \vec{\mu}) + b \sum_{\ell'} S_+(\pi_{\ell'} \vec{\mu}) \quad (6.4)$$

for any $\ell \in \mathcal{B}$. The summation $\sum_{\ell'}$ here and in the sequel is taken over all sites $\ell' \in \Gamma(\ell)$ such that $(\ell', \ell) \in G$. We assume also that the maximal degree of vertices of the graph G is bounded by $K < \infty$.

Theorem 6.1 *Let the local maps, stochastic perturbations and the interactions satisfy the assumptions (6.2), (6.3), (6.4) and we consider a finite box $\mathcal{B}$ with certain frozen boundary conditions. Assume additionally that*

$$a\lambda_T < 1, \quad (a+b)\lambda_T \geq 1. \quad (6.5)$$

Then the stochastically perturbed system in the box $\mathcal{B}$ admits an absolutely continuous invariant measure $\vec{\mu}_\varepsilon \in \mathcal{M}(\vec{X})$, and there exists a constant $\gamma > 0$ (which do neither depend on $\varepsilon > 0$ nor on $\mathcal{B}$) such that

$$S_-(\pi_\ell \vec{\mu}_\varepsilon) \geq \gamma L(\ell) \varepsilon$$

for each site $\ell \in \mathcal{B}$ and small enough $\varepsilon > 0$.

In other words the lower spread of the invariant measure grows at least linearly with the ‘distance’ from the boundary $L(\ell)$. Note the difference to a ‘diffusive’ growth which one might expect from random perturbations.

Let now demonstrate that the assumptions (6.2 – 6.5) are natural. First note that the assumption (6.2) together with the assumption $\lambda_T > 1$ imply only that the map T is locally expanding. Thus, e.g. the dyadic map $Tx := 2x \bmod 1$ defined on a unit circle X considered as the interval $[0, 1)$ with the metric ρ on it defined as the length of the shortest arc between points will satisfy them if we set $\lambda_T = \Lambda_T = 2$ and $\sigma = 1/3$. On the other hand, the assumption $a\lambda_T < 1$ is necessary to be consistent with the LRA phenomenon in an unperturbed system (Theorem 2.1).

To understand the reason behind the assumption (6.4) consider the simplest linear one-dimensional interaction defined by the relation (2.4), for which we have $a = 1 - c$ and $b = c$. Then (6.4) and the inequality $(a+b)\lambda_T \geq 1$ clearly hold true.

The assumption (6.3) gives the information about the local stochastic perturbations of a point mass and clearly is satisfied in the case of independent local stochastic perturbations uniformly distributed in an ε -neighborhood of a point with $\lambda_Q = 1$.

Proof of Theorem 6.1. Since the construction we shall apply is somewhat involved we start with the main scheme of the proof. Similar to the construction of the limit solution in the proof of Theorem 2.1 we shall construct the invariant measure $\vec{\mu}_\varepsilon$ step by step starting from the first site according to the enumeration $\mathcal{N}$ and adding each time only one site according to this enumeration.⁴ Using that the dynamics on already considered sites do not depend on the dynamics of the new one (but not vice versa) we may (and will) consider the interaction with those sites as an additional stochastic perturbation. On this stage our main goal is to show that the action on measures of the stochastically perturbed system is a contraction in a suitable metric in the space of measures. To this end we shall show that the expansion rate of the operator T^* is proportional to Λ_T , while for stochastic perturbation Q^* it is close to 1. The key point is to show that the expansion rate of projections of the operator $\vec{\mathcal{I}}^*$ are proportional to $\Lambda_{\vec{\mathcal{I}}}$, which together with the assumption $\Lambda_T \Lambda_{\vec{\mathcal{I}}} < 1$ guarantees the contraction of the operator $\vec{\mathcal{T}}_\varepsilon^*$ and hence the convergence in time to the unique invariant measure. On the next step of the construction we shall obtain an upper bound for the upper spread of the projection of the invariant measure $\vec{\mu}_\varepsilon$ to the ℓ -th subsystem as a function of the ‘distance’ to the boundary $L(\ell)$ and then using it we shall get the lower bound for the lower spread of this projection.

We introduce the so called Hutchinson metrics in the space of probabilistic measures $\mathcal{M}(X)$ defined by the relation

$$\rho^*(\mu, \nu) := \sup_{\Lambda_\varphi \leq 1} \left(\int \varphi d\mu - \int \varphi d\nu \right),$$

⁴In fact, the limiting solution constructed in the proof of Theorem 2.1 describes the support of the invariant measure $\vec{\mu}_0$ of the unperturbed system.

where the supremum is taken over all continuous functions $\varphi : X \rightarrow \mathbb{R}^1$ satisfying the assumption that their Lipschitz constants Λ_φ (defined by the relation (2.1)) do not exceed 1.

Lemma 6.1 *Under the assumptions of Theorem 6.1 for any pair of measures $\mu, \nu \in \mathcal{M}(X)$ we have*

$$\rho^*(T^*\mu, T^*\nu) \leq \Lambda_T^\alpha \rho^*(\mu, \nu) , \quad (6.6)$$

$$\rho^*(Q_\varepsilon^*\mu, Q_\varepsilon^*\nu) \leq \rho^*(\mu, \nu) , \quad (6.7)$$

$$\rho^*(\pi_\ell \vec{\mathcal{I}}^* \vec{\mu}, \pi_\ell \vec{\mathcal{I}}^* \vec{\nu}) \leq \Lambda_{\vec{\mathcal{I}}} \rho^*(\mu, \nu) , \quad (6.8)$$

where $\vec{\mu}, \vec{\nu} \in \mathcal{M}(\vec{X})$ are arbitrary measures satisfying the assumption that $\pi_\ell \vec{\mu} = \mu$ and $\pi_\ell \vec{\nu} = \nu$.

Proof. By definition

$$\begin{aligned} \rho^*(T^*\mu, T^*\nu) &= \sup_{\Lambda_\varphi \leq 1} \left(\int \varphi dT^*\mu - \int \varphi dT^*\nu \right) \\ &= \sup_{\Lambda_\varphi \leq 1} \left(\int \varphi \circ T d\mu - \int \varphi \circ T d\nu \right) . \end{aligned}$$

Since $\rho(Tx, Ty) \leq \Lambda_T \rho(x, y)$ for any pair of points $x, y \in X$ we have

$$|\varphi \circ T(x) - \varphi \circ T(y)| \leq \Lambda_\varphi \rho(Tx, Ty) \leq \Lambda_\varphi \Lambda_T \rho(x, y) .$$

Thus setting $\psi := \Lambda_T^{-1} \varphi \circ T$ we get $\Lambda_\psi \leq \Lambda_\varphi$ and hence

$$\rho^*(T^*\mu, T^*\nu) \leq \Lambda_T \sup_{\Lambda_\psi \leq 1} \left(\int \psi d\mu - \int \psi d\nu \right) = \Lambda_T \rho^*(\mu, \nu) ,$$

which proves inequality (6.6).

Let us deal now with stochastic perturbations.

$$\begin{aligned} \rho^*(Q_\varepsilon^*\mu, Q_\varepsilon^*\nu) &= \sup_{\Lambda_\varphi \leq 1} \left(\int \varphi dQ_\varepsilon^*\mu - \int \varphi dQ_\varepsilon^*\nu \right) \\ &= \sup_{\Lambda_\varphi \leq 1} \left(\int Q_\varepsilon \varphi d\mu - \int Q_\varepsilon \varphi d\nu \right) . \end{aligned}$$

Using the translation invariance of $Q_\varepsilon(\cdot, \cdot)$ we get

$$\begin{aligned} |Q_\varepsilon \varphi(x) - Q_\varepsilon \varphi(y)| &= \left| \int \varphi(z) Q_\varepsilon(x, dz) - \int \varphi(z) Q_\varepsilon(y, dz) \right| \\ &= \left| \int \varphi(z) Q_\varepsilon(x, dz) - \int \varphi(z - x + y) Q_\varepsilon(y, dz - x + y) \right| \\ &= \left| \int \varphi(z) Q_\varepsilon(x, dz) - \int \varphi(z - x + y) Q_\varepsilon(x, dz) \right| \\ &\leq \int |\varphi(z) - \varphi(z - x + y)| \cdot Q_\varepsilon(x, dz) \leq \Lambda_\varphi \rho(x, y) . \end{aligned}$$

Thus setting $\psi := Q_\varepsilon \varphi$ we obtain

$$\rho^*(Q_\varepsilon^*\mu, Q_\varepsilon^*\nu) \leq \sup_{\Lambda_\psi \leq 1} \left(\int \psi d\mu - \int \psi d\nu \right) = \rho^*(\mu, \nu) ,$$

which proves the inequality (6.7).

To prove the remaining inequality (6.8) we introduce the following notation. For a given $x \in X$ let $\vec{x}_{(x_\ell=x)} \in \vec{X}$ mean a vector whose ℓ -th coordinate coincides with x , and for $\mu_\ell := \pi_\ell \vec{\mu} \in \mathcal{M}(X)$

denote by $\mu_\ell^\perp$ the projection of the measure $\vec{\mu} \in \mathcal{M}(\vec{X})$ to all subsystems except the ℓ -th one (i.e. we integrate over the ℓ -th coordinate). Then

$$\int \varphi(x) d\pi_\ell \vec{\mathcal{I}}^* \vec{\mu}(x) = \int \left(\int \varphi((\vec{\mathcal{I}} \vec{x}_{(x_\ell=x)})) d\mu_\ell^\perp(\vec{x}_{(x_\ell=x)}) \right) d\mu_\ell(x).$$

Consider a function

$$\psi(x) := \Lambda_{\vec{\mathcal{I}}}^{-1} \int \varphi((\vec{\mathcal{I}} \vec{x}_{(x_\ell=x)})) d\mu_\ell^\perp(\vec{x}_{(x_\ell=x)}).$$

Then

$$\Lambda_{\vec{\mathcal{I}}} \cdot |\psi(x) - \psi(y)| \leq \int |\varphi((\vec{\mathcal{I}} \vec{x}_{(x_\ell=x)})) - \varphi((\vec{\mathcal{I}} \vec{x}_{(x_\ell=y)}))| d\mu_\ell^\perp(\vec{x}_{(x_\ell=x)}),$$

since $\mu_\ell^\perp(\vec{x}_{(x_\ell=x)}) = \mu_\ell^\perp(\vec{x}_{(x_\ell=y)})$ because the measure $\mu_\ell^\perp$ does not depend on x_ℓ .

On the other hand, by the inequality 2.2 we have

$$\rho(\vec{\mathcal{I}} \vec{x}_{(x_\ell=x)}, \vec{\mathcal{I}} \vec{x}_{(x_\ell=y)}) \leq \Lambda_{\vec{\mathcal{I}}} \rho(x, y),$$

which implies that

$$|\psi(x) - \psi(y)| \leq \Lambda_{\vec{\mathcal{I}}}^{-1} \Lambda_{\vec{\mathcal{I}}} \rho(x, y) \int d\mu_\ell^\perp(\vec{x}_{(x_\ell=x)}) = \rho(x, y).$$

Therefore using the same argument as in the previous estimates we obtain the inequality (6.8). $\square$

For a given $\ell \in \mathcal{B}$ the set $\Gamma(\ell)$ consists of a finite number of elements connected to ℓ (directly or indirectly⁵). Consider a subsystem consisted only of the sites belonging to $\Gamma(\ell)$. The restricted dynamics of this subsystem does not depend on the dynamics of the other parts of the system.

Collecting inequalities (6.6), (6.7), (6.8) we see that the operator corresponding to the stochastically perturbed system is strictly contracting in the ρ^* metric, which (due to the compactness of this space) immediately implies that for any initial measure $\vec{\mu} \in \mathcal{M}(\vec{X})$ we have:

$$\rho^*(\pi_\ell(\vec{\mathcal{T}}_\varepsilon^*)^t \vec{\mu}, \pi_\ell \vec{\mu}_\varepsilon) \xrightarrow{t \rightarrow \infty} 0.$$

Going through all sites of the box $\mathcal{B}$ we construct the invariant measure $\vec{\mu}_\varepsilon$ for the entire stochastically perturbed system restricted to the box $\mathcal{B}$ with the given frozen boundary conditions.

Now we are ready to obtain estimates from above for the upper spread $S_+(\pi_\ell \vec{\mu}_\varepsilon)$. It is straightforward to check that under (6.2) for any measure $\mu \in \mathcal{M}(X)$ whose upper spread satisfies the inequality $S_+(\mu) \leq \sigma/\Lambda_T$ we have

$$S_+(T^* \mu) \leq \Lambda_T S_+(\mu). \quad (6.9)$$

The relation (2.2) implies that if $S_+(\pi_\ell \mu), S_+(\pi_{\ell'} \mu) \leq \sigma/\Lambda_T$ then

$$S_+(\pi_\ell \vec{\mathcal{I}}^* \vec{\mu}) \leq \Lambda_{\vec{\mathcal{I}}} S_+(\pi_\ell \vec{\mu}) + \Lambda_{\vec{\mathcal{I}}} \sum_{\ell'} S_+(\pi_{\ell'} \vec{\mu}). \quad (6.10)$$

Since the amplitude of the stochastic perturbations is bounded from above by $\varepsilon > 0$, one has

$$S_+(Q_\varepsilon^* \mu) \leq S_+(\mu) + \varepsilon. \quad (6.11)$$

The dynamics at the site ℓ' does not depend on the dynamics at the site ℓ (but not vice versa) therefore we may choose an initial measure $\vec{\mu} \in \mathcal{M}(\vec{X})$ satisfying the following conditions:

$$S_+(\pi_\ell \vec{\mu}) \leq \sigma/(K\Lambda_T), \quad \pi_k \vec{\mu} \equiv \pi_k \vec{\mu}_\varepsilon \quad \forall k : \mathcal{N}(k) < \mathcal{N}(\ell).$$

⁵through some other elements of $\mathcal{B}$

Then by the inequalities (6.9, 6.10, 6.11) we have:

$$S_+(\pi_\ell \vec{\mathcal{T}}_\varepsilon^* \vec{\mu}) \leq \Lambda_{\vec{\mathcal{I}}}(\Lambda_T S_+(\pi_\ell \vec{\mu}) + \varepsilon) + \Lambda_{\vec{\mathcal{I}}} \sum_{\ell'} (\Lambda_T S_+(\pi_{\ell'} \vec{\mu}) + \varepsilon). \quad (6.12)$$

Denote $q := \Lambda_{\vec{\mathcal{I}}} \Lambda_T$, $p := \vec{\mathcal{I}}(K+1)$. Applying the inequality (6.12) t times and using that $S_+(\pi_{\ell'} \vec{\mathcal{T}}_\varepsilon^* \vec{\mu}) \equiv S_+(\pi_{\ell'} \vec{\mu})$ we get

$$S_+(\pi_\ell (\vec{\mathcal{T}}_\varepsilon^*)^t \vec{\mu}) \leq q^t S_+(\pi_\ell \vec{\mu}) + (q \sum_{\ell'} S_+(\pi_{\ell'} \vec{\mu}) + p\varepsilon) \cdot \frac{1 - q^t}{1 - q}.$$

Now passing to the limit as $t \rightarrow \infty$ and using that $q = \Lambda_{\vec{\mathcal{I}}} \Lambda_T < 1$ (by the condition of Theorem 2.1) we obtain the desired estimate from above:

$$S_+(\pi_\ell \vec{\mu}_\varepsilon) \leq \frac{q}{1 - q} \sum_{\ell'} S_+(\pi_{\ell'} \vec{\mu}_\varepsilon) + \frac{p\varepsilon}{1 - q}.$$

Thus the upper spread of the ℓ -th projection of the invariant measure cannot exceed a constant in power $L(l)$ multiplied by ε . Observe that for each $t \geq 1$ the inequality $D(\pi_\ell (\vec{\mathcal{T}}_\ell^*)^t \vec{\mu}) \leq \sigma$ holds automatically since $q < 1$.

Now whence we have established the control over the upper spread of $\vec{\mu}_\varepsilon$ we will obtain an estimate from below for the lower spread of this measure.

Under (6.2) for any measure $\mu \in \mathcal{M}(X)$ with $S_-(\mu) \leq \sigma/\Lambda_T$ we have:

$$S_-(T^* \mu) \geq \lambda_T S_-(\mu). \quad (6.13)$$

The relation (6.3) implies

$$S_-(Q_\varepsilon^* \mu) \geq S_-(\mu) + \lambda_Q \varepsilon. \quad (6.14)$$

Making use of the inequalities (6.13), (6.14) and (6.4) for any measure $\mu \in \mathcal{M}(X)$ with $S_-(\mu) \leq \sigma/\Lambda_T$ and a measure $\vec{\mu} \in \mathcal{M}(\vec{X})$ with $\pi_\ell \vec{\mu} = \mu$ we get:

$$S_-(\pi_\ell \vec{\mathcal{T}}_\varepsilon^* \vec{\mu}) \geq a(\lambda_T S_-(\pi_\ell \vec{\mu}) + \lambda_Q \varepsilon) + b(\lambda_T \sum_{\ell'} S_-(\pi_{\ell'} \vec{\mu}) + \lambda_Q \varepsilon). \quad (6.15)$$

Denote $\alpha := a\lambda_T < 1$, $\beta := b\lambda_T \geq 1 - \alpha$ (by the assumption (6.5)) and $\kappa := (a + b)\lambda_Q$. Now, applying (6.15) to $(\vec{\mathcal{T}}_\varepsilon^*)^t$ we have

$$S_-(\pi_\ell (\vec{\mathcal{T}}_\varepsilon^*)^t \vec{\mu}) \geq \alpha^t S_-(\pi_\ell \vec{\mu}) + (\beta \sum_{\ell'} S_-(\pi_{\ell'} \vec{\mu}) + \kappa \varepsilon) \cdot \frac{1 - \alpha^t}{1 - \alpha}.$$

Passing to the limit and using that $\beta/(1 - \alpha) \geq 1$ we obtain the final estimate

$$S_-(\pi_\ell \vec{\mu}_\varepsilon) \geq \sum_{\ell'} S_-(\pi_{\ell'} \vec{\mu}_\varepsilon) + \frac{\kappa \varepsilon}{1 - \alpha}.$$

Theorem 6.1 follows by setting $\gamma := \kappa \varepsilon / (1 - \alpha)$. □

7 Conclusion

We have shown that the absence of loops in directed connectivity graphs of networks of dynamical elements results in appearance of a long range action provided that interactions between local dynamical systems (elements of a network) are sufficiently strong. If local systems have a regular (contractive) dynamics than the LRA appears already for arbitrarily weak interactions (conditions of Theorem 2.1 clearly hold in this case).

One expects that this restriction to the network topology is by no means a necessary one. In principle, we see two ways to overcome this problem. The first of them is to use some rather general ergodic-theoretical results (see e.g. [9]) to recode interaction potentials with one-sided dependence to the ones with the two-sided dependence. It is known that if those potentials are homological to each other than they generate the same Gibbs measures (corresponding to our invariant measures). The problem here is that it is hard to describe the set of constructed potentials corresponding to general non-unidirectional interactions.

Another possibility is to consider non-unidirectional interactions as small perturbations to the unidirectional ones. This approach seems to be more promising but one inevitably needs to make a number of additional assumptions about the unperturbed system to ensure that the unidirectional interactions are stronger than the perturbations.

There is yet another possibility related to the assumption that the interaction in the network is close to a mean-field model, i.e. a large part of the network is interacting with each its element. The main disadvantage of this assumption is that it is not completely clear how to extend this property for infinite networks.

It is worth mentioning that instead of the lattice $\mathbb{Z}^d$ one can consider a countable ordered collection of vertices. The short range interactions in this context would mean that subsystems corresponding to vertices interact only if the distance between them according to the above mentioned order does not exceed a certain threshold. Clearly all our results remain valid in this setting.

The problems similar to the ones discussed in this paper were studied in the context of the synchronization theory. More specifically, we are talking here about a complete synchronization of dynamical systems [8]. The complete synchronization means that asymptotically, as time goes to infinity, the differences in the behavior of all elements in the network disappear. In particular, in [10] it has been shown that in the case of the network described by the ordinary differential equations whose directed connectivity graph is a tree the synchronization phenomenon takes place under strong enough interactions, while under the assumption that the connectivity graph is “strongly connected” (i.e. the interaction is close to a mean-field model) this has been proven in [1] (see also references in these papers and the recent monograph [8]). In these publications, in distinction to our approach, a synchronization phenomenon is studied in closed finite networks consisting of identical elements and the synchronized system may demonstrate chaotic behavior.

Acknowledgements

The research by M.B. was partially supported by Russian Foundation for Fundamental Research and French Ministry of Education grants. L.B. was partially supported by the NSF grant DMS - 0140165 and by the Humboldt foundation.

References

- [1] Belykh V.N., Belykh I.V., Hasler M., Connection graph stability method for synchronized coupled chaotic systems, *Physica D* 195:1/2(2004), 159-187.
- [2] Blank M., Discreteness and continuity in problems of chaotic dynamics, Amer. Math. Soc., Providence, Rhode Island (1997).
- [3] Blank M., Bunimovich L., Multicomponent dynamical systems: SRB measures and phase transitions, *Nonlinearity*, 16:1(2003), 387-401.
- [4] Bunimovich L., Coupled Map Lattices: at the age of maturity, *Lect. Notes in Physics* (to appear).
- [5] Keller G., Liverani C., Coupled map lattices without cluster expansion, *Discr. Cont. Dynam. Sys.* 11 (2004), 325-335.
- [6] Liggett T. M., *Interacting Particle Systems*, Springer, 2005.
- [7] Miller J., Huse D.A., Macroscopic equilibrium from microscopic irreversibility in a chaotic coupled-map lattice, *Phys. Rev. E*, 48(1993), 2528-2535.
- [8] Pikovsky A., Rosenblum M., Kurths J., *Synchronization: A universal concept in nonlinear sciences*. Cambridge Univ. Press, 2001.
- [9] Sinai Ya.G, Gibbs measures in ergodic theory, *Russian Math. Surveys*, 27 (1972), 21-69.
- [10] Wu Chai Wah, Synchronization in networks of nonlinear dynamical systems coupled via a directed graph, *Nonlinearity* 18:3 (2005) 1057-1064.