

SMOOTHING EFFECT OF QUENCHED DISORDER ON POLYMER DEPINNING TRANSITIONS

GIAMBATTISTA GIACOMIN AND FABIO LUCIO TONINELLI

ABSTRACT. We consider general disordered models of pinning of directed polymers on a defect line. This class contains in particular the $(1+1)$ -dimensional interface wetting model, the disordered Poland–Scheraga model of DNA denaturation and other $(1+d)$ -dimensional polymers in interaction with flat interfaces. We consider also the case of copolymers with adsorption at a selective interface. Under quite general conditions, these models are known to have a (de)localization transition at some critical line in the phase diagram. In this work we prove in particular that, as soon as disorder is present, the transition is at least of second order, in the sense that the free energy is differentiable at the critical line, so that the order parameter vanishes continuously at the transition. On the other hand, it is known that the corresponding non-disordered models can have a first order (de)localization transition, with a discontinuous first derivative. Our result shows therefore that the presence of the disorder has really a smoothing effect on the transition. The relation with the predictions based on the Harris criterion is discussed.

2000 *Mathematics Subject Classification:* 60K35, 82B41, 82B44

Keywords: Quenched disorder, Smoothing of Delocalization Transition, Directed Polymers, Pinning and Wetting Models, Poland–Scheraga Models, Copolymers, Harris Criterion.

1. INTRODUCTION AND MODELS

Quenched disorder is expected to smooth phase transitions in many situations. This is for instance the case of ferromagnetic spin systems in dimension $d \leq 2$ (if the spins are discrete) or in dimension $d \leq 4$ (if they have rotation symmetry): when a random magnetic field is present the Imry–Ma argument [20], made rigorous by Aizenman and Wehr [1], implies that these models do not exhibit, at any temperature, the first order phase transition with associated spontaneous magnetization which characterizes the corresponding pure models. For the analogous phenomenon for SOS effective interface models see [6].

In the present work, under some conditions on the disorder distribution, we prove that a similar effect takes place in models of directed polymers in random media exhibiting a localization/delocalization transition on a defect line. Such a transition may be of first or higher order in the corresponding pure, i.e. non-disordered, cases and we show that, as soon as disorder is present, the transition is at least of second order. It is important to emphasize that the mechanism inducing the smoothing of the transition in our case is very different from the Imry–Ma one, and it is rather based on an estimate of the probability that the polymer visits rare but very favorable regions where the disorder produces a large positive fluctuation of the partition function.

Date: March 29, 2022.

We will consider mainly two classes of models: random pinning (or wetting) models and random copolymers at selective interfaces. In pinning models [2, 10, 13, 24, 27] the typical situation one has in mind is that of a directed path in $(1+d)$ dimensions, which receives a reward (or a penalty) at each intersection with a 1-dimensional region (a *defect line*, in the physical language) according to whether the charge present on the line at the intersection point is positive or negative. On the other hand the copolymer model, whose study was initiated in [14] in the theoretical physical literature (see [23] and references therein for updated physics developments) and in [5, 25] in the mathematical one, aims at modeling a $(1+1)$ -dimensional, directed heteropolymer containing both hydrophobic and hydrophilic components, in presence of an interface (the line $S = 0$ in the notations of Section 1.2) separating two solvents, situated in the upper and lower half-planes ($S > 0$ and $S < 0$). One solvent favors the hydrophilic components and the other favors the hydrophobic ones, i.e., if the charge of the n -th monomer is positive (negative), this monomer tends to be in the upper (lower) half-plane.

The main question one would like to answer is whether or not the interaction induces a localization of the polymer along the defect line or interface.

1.1. Random pinning and wetting models. Let $S := \{S_n\}_{n=0,1,\dots}$ be a homogeneous process on an arbitrary set that contains a point 0, with $S_0 = 0$ and law \mathbf{P} . For $\beta \geq 0, h \in \mathbb{R}$ and $\omega = \{\omega_n\}_{n=1,2,\dots} \in \mathbb{R}^{\mathbb{N}}$, one introduces the probability measure

$$\frac{d\mathbf{P}_{N,\omega}}{d\mathbf{P}}(S) = \frac{1}{Z_{N,\omega}} \exp\left(\sum_{n=1}^N (\beta\omega_n - h) \mathbf{1}_{S_n=0}\right) \mathbf{1}_{S_N=0}. \quad (1.1)$$

The choice of setting $S_N = 0$ is just for technical convenience, see Remark 1.1 below. Let us set $\tau_0 = 0$ and, for $i \in \mathbb{N} := \{1, 2, \dots\}$, $\tau_i = \inf\{n > \tau_{i-1} : S_n = 0\}$ if $\tau_{i-1} < +\infty$ and $\tau_i = +\infty$ if $\tau_{i-1} = +\infty$. We assume that $\tau := \{\tau_i\}_i$ is the sequence of partial sums of an IID sequence of random variables taking values in $\mathbb{N} \cup \{\infty\}$ with discrete density $K(\cdot)$:

$$K(n) = \mathbf{P}(\tau_1 = n). \quad (1.2)$$

This is of course the case if S is a Markov chain and for definiteness one should keep this case in mind. In order to avoid trivialities, we assume that $K(\infty) < 1$ and for the model to be defined we assume also that there exists $s \in \mathbb{N}$ such that $\mathbf{P}(S_N = 0) > 0$ for every $N \in s\mathbb{N}$. Therefore, starting with (1.1), we always implicitly assume that $N \in s\mathbb{N}$, when N is the length of the polymer.

Moreover the cases we will consider are such that there exist $\alpha \geq 1$ such that

$$\liminf_{n \rightarrow \infty} \frac{\log K(sn)}{\log n} \geq -\alpha, \quad (1.3)$$

and $K(n) = 0$ if $n \notin s\mathbb{N}$. We advise the reader who feels uneasy with the weak requirement (1.3) to focus on the case of $K(n)$ behaving like $n^{-\alpha}$, possibly times a *slowly varying function* (see Appendix A). Note that, for $\alpha < 2$, the return times to zero of S are not integrable. We stress that we have introduced s to account for the possible periodicity of S : of course the most natural example is that of the simple random walk on \mathbb{Z} , $\mathbf{P}(S_i - S_{i-1} = 1) = \mathbf{P}(S_i - S_{i-1} = -1) = 1/2$ and $\{S_i - S_{i-1}\}_i$ IID, for which (1.3) holds with $s = 2$ and $\alpha = 3/2$ [11, Ch. III]. Another example is that of the simple random walk on \mathbb{Z}^d , $d \geq 2$: in this case, $\alpha = d/2$.

We can rewrite $Z_{N,\omega}$, and in fact the model itself, in terms of the sequence τ : the partition function in (1.1) is

$$Z_{N,\omega} := \mathbf{E} [\exp(\mathcal{H}_{N,\omega}(S)); \tau_{\mathcal{N}_N} = N], \quad (1.4)$$

with $\mathcal{N}_N := \sup\{i : \tau_i \leq N\}$ and

$$\mathcal{H}_{N,\omega}(S) = \sum_{i=1}^{\mathcal{N}_N} (\beta\omega_{\tau_i} - h). \quad (1.5)$$

The disordered pinning model, which we consider here, is obtained assuming that the sequence ω is chosen as a typical realization of an IID sequence of random variables with law \mathbb{P} , still denoted by $\omega = \{\omega_n\}_n$. We assume finiteness of exponential moments:

$$\mathbb{E}[\exp(t\omega_1)] < \infty, \quad (1.6)$$

for $t \in \mathbb{R}$ and, without loss of generality, $\mathbb{E}\omega_1 = 0$ and $\mathbb{E}\omega_1^2 = 1$. Further assumptions on \mathbb{P} will be formulated in Section 2, where we state our main results.

Under the above assumptions on the disorder the *quenched free energy* of the model exists, namely the limit

$$\mathbb{F}(\beta, h) := \lim_{N \rightarrow \infty} \frac{1}{N} \log Z_{N,\omega}, \quad (1.7)$$

exists $\mathbb{P}(d\omega)$ -almost surely and in the $\mathbb{L}^1(\mathbb{P})$ sense. The existence of this limit can be proven via standard super-additivity arguments based on Kingman's sub-additive ergodic theorem [22] (we refer for example to [2, 5, 16] for details). This approach yields also automatically the so called *self-averaging* property of the free energy, that is the fact that $\mathbb{F}(\beta, h)$ is non random. A simple but fundamental observation is that

$$\mathbb{F}(\beta, h) \geq 0. \quad (1.8)$$

The proof of such a result is elementary:

$$\frac{1}{N} \mathbb{E} \log Z_{N,\omega} \geq \frac{1}{N} \mathbb{E} \log \mathbf{E} \left[\exp \left(\sum_{i=1}^{\mathcal{N}_N} (\beta\omega_{\tau_i} - h) \right); \tau_1 = N \right] \quad (1.9)$$

$$= \frac{1}{N} (\beta \mathbb{E}[\omega_N] - h) + \frac{1}{N} \log \mathbf{P}(\tau_1 = N) \xrightarrow{N \rightarrow \infty} 0, \quad (1.10)$$

where we have used assumption (1.3). The proof of (1.8) suggests the following partition of the parameter space (or *phase diagram*):

- The localized region: $\mathcal{L} = \{(\beta, h) : \mathbb{F}(\beta, h) > 0\}$;
- The delocalized region: $\mathcal{D} = \{(\beta, h) : \mathbb{F}(\beta, h) = 0\}$.

We set $h_c(\beta) := \sup\{h : \mathbb{F}(\beta, h) > 0\}$, and we will call $h_c(\beta)$ the *critical point*. Since $\mathbb{F}(\beta, \cdot)$ is not increasing and continuous, $\mathcal{D} = \{(\beta, h) : h \geq h_c(\beta)\}$. It is rather easy to see that $|h_c(\beta)| < \infty$, see e.g. [2] and [16].

Remark 1.1. It would be of course as natural to consider the model with partition function $Z_{N,\omega}^f := \mathbf{E} [\exp(\mathcal{H}_{N,\omega}(S))]$. It is therefore worth to stress that by standard

arguments, see e.g. [16], one sees that

$$\begin{aligned} Z_{N,\omega} &\leq Z_{N,\omega}^{\mathbf{f}} \leq \max_{\substack{n \in s\mathbb{N}: \\ n \leq N}} \frac{\sum_{j=n}^{\infty} K(j)}{K(n)} \exp(|\beta\omega_N - h|) Z_{N,\omega} \\ &\leq \frac{1}{\min_{\substack{n \in s\mathbb{N}: \\ n \leq N}} K(n)} \exp(|\beta\omega_N - h|) Z_{N,\omega}, \end{aligned} \tag{1.11}$$

uniformly in $N \in s\mathbb{N}$ and ω . In particular, by (1.3), the free energy is unaffected by the presence of the constraint $\tau_{N_N} = N$: we stick to the constrained case because it simplifies some technical steps of the proofs.

Remark 1.2. For ease of exposition we have used the generic denomination of *pinning model*, but our framework, as it is possibly clearer when the partition function is cast in the form (1.4), includes a variety of models, like for example the Poland–Scheraga model of DNA denaturation, for which theoretical arguments on models with excluded volume interactions suggest a value of α larger than 2 [21]: note that whether the transition in the disordered case is of first or higher order is a crucial and controversial issue in the field, see for example [9, 8, 15]. We stress also that, since we can choose $K(\infty) > 0$, it is rather easy to see that also the disordered (1 + 1)–dimensional interface wetting models [10, 13] enter the general class we are considering.

It is known that the nonrandom case $\beta = 0$ is exactly solvable and that, according to the law of τ_1 , the transition at $h_c(0) = \log(1 - K(\infty))$ can be either of first or higher order. For completeness and to match our set–up, we give a quick self–contained analysis of this case in Appendix A.

1.2. Random copolymers at a selective interface. In the case of the copolymer model, the natural setting is to assume, in addition, that the state space of the process $\{S_n\}_n$ is \mathbb{Z} , that the law \mathbf{P} is invariant under the transformation $S \rightarrow -S$ and that $(S_i - S_{i-1}) \in \{-1, 0, +1\}$ for every $i \geq 1$. For instance, if $\{S_i - S_{i-1}\}_i$ is a sequence of IID variables then it is a classical result [12, Ch. XII.7] that $K(n) = c(1 + o(1))n^{-3/2}$, for large values of $n \in s\mathbb{N}$ (so $\alpha = 3/2$): c is a constant that can be expressed in terms of $\mathbf{P}(S_1 - S_0 = 0)$ and of course $s = 1$ unless $\mathbf{P}(S_1 - S_0 = 0) = 0$. The copolymer model is defined introducing

$$\mathcal{H}_{N,\omega}(S) = \frac{1}{2} \sum_{n=1}^N (\beta\omega_n + h) \text{sign}(S_n), \tag{1.12}$$

with the convention that $\text{sign}(0) = +1$, and the corresponding partition function

$$Z_{N,\omega} := \mathbf{E}[\exp(\mathcal{H}_{N,\omega}(S)); S_N = 0]. \tag{1.13}$$

We may assume without loss of generality that both β and h are nonnegative. The factor 1/2 in (1.12) is introduced just for convenience (see Section 3.2). As in Section 1.1, the corresponding random model is obtained by choosing ω as a realization of an IID sequence of centered random variables of unit variance and finite exponential moments. In analogy with (1.7) and (1.8), the limit

$$f(\beta, h) := \lim_{N \rightarrow \infty} \frac{1}{N} \log Z_{N,\omega}, \tag{1.14}$$

exists $\mathbb{P}(d\omega)$ -almost surely and in the $\mathbb{L}^1(\mathbb{P})$ sense and one can prove as in (1.9) that $F(\beta, h) := f(\beta, h) - h/2 \geq 0$. Therefore, also in this case we can partition the phase diagram into a localized and a delocalized phase, as

- The localized region: $\mathcal{L} = \{(\beta, h) : F(\beta, h) > 0\}$;
- The delocalized region: $\mathcal{D} = \{(\beta, h) : F(\beta, h) = 0\}$.

Also in this case, we set $h_c(\beta) := \sup\{h : F(\beta, h) > 0\}$ and we observe that $\mathcal{D} = \{(\beta, h) : h \geq h_c(\beta)\}$.

Many results have been proven for copolymers at selective interfaces, but they are almost always about the case of simple random walks. However they can be extended in a rather straightforward way to the general case we consider here (we omit the details also because they are not directly pertinent to the content of this paper). Above all we have that $0 < h_c(\beta) < \infty$ for every $\beta > 0$. Even more, the explicit bounds carry over to the general S we consider here: the upper bound [5] is even independent of the choice of the law of S , while the lower bound [4] depends on α .

We take this opportunity to stress also that various results about path behavior in the two regions are available, both for the pinning problem and for the copolymer, and, once again, mostly in the simple random walk case $\alpha = 3/2$. For instance, it is known that in \mathcal{L} long excursions of the walk from the line $S = 0$ are exponentially suppressed and the fraction of sites where the polymer crosses the line remains nonzero in the thermodynamic limit (e.g., cf [16, 25] and references therein, and [18]). On the other hand, in the interior of \mathcal{D} the number of intersections is, in a suitable sense, $O(\log N)$ [17] and, for the copolymer, the number of steps where $\text{sign}(S_n) = -1$ is also $O(\log N)$ [17].

2. SMOOTHING OF THE DEPINNING TRANSITION

While the free energy can be proven to be infinitely differentiable with respect to all of its parameters in the region \mathcal{L} [18], no results are available about its regularity at the critical point, apart from the obvious fact that F is continuous since it is convex. The result of the present paper partly fills this gap, showing that the transition is at least of second order, as soon as disorder is present.

In order to state our main theorem, we need some further assumptions on the disorder variables ω . We will consider two distinct cases:

C1: *Bounded random variables.* The random variable ω_1 is bounded,

$$|\omega_1| \leq M < \infty. \quad (2.1)$$

C2: *Unbounded continuous random variables.* The law of ω_1 has a density $P(\cdot)$ with respect to the Lebesgue measure on \mathbb{R} , and there exists $0 < R < \infty$ such that

$$\int_{\mathbb{R}} P(y+x) \log \left(\frac{P(y+x)}{P(y)} \right) dy \leq Rx^2 \quad (2.2)$$

in a neighborhood of $x = 0$. This is true in great generality whenever $P(\cdot)$ is positive, for example when the disorder is Gaussian and, more generally, whenever $P(\cdot) = \exp(-V(\cdot))$, with $V(\cdot)$ a polynomial bounded below.

Then, one has:

Theorem 2.1. *Under condition C1 or C2, both for the copolymer and for the pinning model, for every $0 < \beta < \infty$ there exists $0 < c(\beta) < \infty$, possibly depending on \mathbb{P} , such that for every $1 \leq \alpha < \infty$*

$$F(\beta, h) \leq \alpha c(\beta)(h_c(\beta) - h)^2 \quad (2.3)$$

if $h < h_c(\beta)$.

Although the above result, coupled of course with (1.8), seems to be in the same spirit as the rounding effect proven by Aizenman and Wehr [1] for the two-dimensional Random Field Ising Model, the physical mechanisms of smoothing are deeply different in the two cases. While [1] is based on a rigorous version of the Imry-Ma argument [20] (i.e., a comparison between the effect of boundary conditions and of disorder fluctuations in the bulk due to the random magnetic field) in our case the boundary conditions play no role at all and everything is based on an energy-entropy argument inspired by [4].

Remark 2.2. It is important to observe that, as explained in Appendix A, in the pinning case the deterministic model, $\beta = 0$, has a *first order phase transition* whenever $\sum_{n \in \mathbb{N}} nK(n) < +\infty$ [2], in particular when $\alpha > 2$. Theorem 2.1 therefore shows that the disorder has really a smoothing effect on the transition. But more than that is true: for $\alpha \in (3/2, 2)$, $F(0, h)$ at $h_c(0)$ is strictly less regular than $F(\beta, h)$ at $h_c(\beta)$. In fact $F(0, h) > (h_c(0) - h)^{2-\delta}$ for $\delta \in (0, (2\alpha - 3)/(\alpha - 1))$ and small values of $h_c(0) - h > 0$ (for sharper results, see Appendix A). Notice that this is in agreement with the so-called *Harris criterion* [19], which predicts that arbitrarily weak disorder modifies the nature of a second-order phase transition as soon as the critical exponent of the specific heat in the pure case is positive. In the present situation, this condition corresponds just to $\alpha > 3/2$. The Harris criterion also predicts that the critical behavior does not change if $\alpha < 3/2$, which is compatible with Theorem 2.1. Rigorous work connected to the Harris criterion, in the Ising model context, may be found in [7].

As one realizes easily from the proof of Theorem 2.1 given in Section 3, the constant $c(\beta)$ in (2.3) can be very large (of order $O(\beta^{-2})$) for β small. This is rather intuitive: for $\beta \rightarrow 0$ one approaches the deterministic situation, where the transition can be of first order.

Remark 2.3. In the theoretical physics literature, the (de)localization transition is claimed to be in some cases of order higher than two [28], or even of infinite order [27, 23]. The method we present here, which is rather insensitive to the details of the model, does not allow to prove more than second order in general. It is likely that finer results require model-specific techniques.

Remark 2.4. *A generalized model: copolymers with adsorption.* It is also possible to consider copolymer models with an additional pinning interaction [26], as

$$\mathcal{H}_{N, \omega, \tilde{\omega}}(S) = \sum_{n=1}^N (\beta_1 \omega_n - h_1) \mathbf{1}_{S_n=0} + \frac{1}{2} \sum_{n=1}^N (\beta_2 \tilde{\omega}_n - h_2) \text{sign}(S_n). \quad (2.4)$$

Here, both ω and $\tilde{\omega}$ are sequences of IID centered random variables with unit variance and finite exponential moments. In addition, one assumes ω to be independent from $\tilde{\omega}$. This model corresponds to the situation where the interface between the two solvents is not neutral. While for simplicity we will not present details for this model, we sketch here what happens in this case. In analogy with the previous models, one can partition the phase diagram (i.e., the space of the parameters $\beta_1, h_1, \beta_2, h_2$) into a localized and a delocalized

region, separated by a *critical surface*. In this case, Theorem 2.1 is easily generalized to give that the free energy $F(\beta_1, h_1, \beta_2, h_2)$ has continuous first derivatives with respect to h_1, h_2 when these parameters approach the critical surface from the localized region.

3. PROOF OF THE SMOOTHING EFFECT

3.1. The pinning case. In this section we prove Theorem 2.1 for the pinning case, and in the next one we explain how the proof can be immediately extended to the copolymer model.

The key idea, in analogy with [17], is to introduce a new free energy where the fraction of sites where the polymer comes back to zero is fixed. In other words, recalling that

$$\mathcal{N}_N = |\{1 \leq n \leq N : S_n = 0\}| \quad (3.1)$$

one introduces, for $m \in [0, 1]$,

$$\begin{aligned} \phi(\beta, m) &= \lim_{\varepsilon \searrow 0} \lim_{N \rightarrow \infty} \frac{1}{N} \mathbb{E} \log \hat{Z}_{N, \omega}(\beta; m, \varepsilon) \\ &:= \lim_{\varepsilon \searrow 0} \lim_{N \rightarrow \infty} \frac{1}{N} \mathbb{E} \log \mathbf{E} \left(e^{\beta \sum_{n=1}^N \omega_n \mathbf{1}_{S_n=0}} \mathbf{1}_{(\mathcal{N}_N/N) \in [m-\varepsilon, m+\varepsilon]} \mathbf{1}_{S_N=0} \right). \end{aligned} \quad (3.2)$$

Note that the limit is well defined, since $\mathbb{E} \log \hat{Z}_{N, \omega}(\beta; m, \varepsilon)$ is super-additive in N , thanks to the IID assumption on the increments of the sequence of return times τ , and non-increasing for $\varepsilon \searrow 0$. Therefore the limit $\phi_\varepsilon(\beta, m)$ of $(1/N) \mathbb{E} \log \hat{Z}_{N, \omega}(\beta; m, \varepsilon)$, as $N \rightarrow \infty$, exists, as well as the second limit $\phi_\varepsilon(\beta, m) \searrow \phi(\beta, m)$ as $\varepsilon \searrow 0$. Notice moreover that (3.2) holds also without taking the expectation, as for $(1/N) \log Z_{N, \omega}$: in this case of course the limit $N \rightarrow \infty$ has to be taken in the $\mathbb{P}(d\omega)$ -a.s. sense. Moreover, it is immediate to realize that $\phi(\beta, 0) = 0$ and that

$$\phi(\beta, m) \leq F(\beta, 0) \leq \beta, \quad (3.3)$$

so that $\phi(\beta, m)$ is always bounded above. Finally, always thanks the IID property of the differences of successive return times to zero, it is easy to show that $\phi(\beta, \cdot)$ is concave:

$$\phi(\beta, m) \geq x\phi(\beta, m_1) + (1-x)\phi(\beta, m_2) \quad (3.4)$$

if $0 \leq x \leq 1$ and $m = xm_1 + (1-x)m_2$. By exploiting the $\mathbb{P}(d\omega)$ -a.s. convergence of $(1/N) \log \hat{Z}_{N, \omega}(\beta; m, \varepsilon)$ to the nonrandom limit $\phi_\varepsilon(\beta, m)$ and the subsequent convergence for $\varepsilon \searrow 0$, one deduces that $F(\beta, h)$ and $\phi(\beta, m)$ are related by a Legendre transform:

$$F(\beta, h) = \sup_{m \in [0, 1]} (\phi(\beta, m) - hm). \quad (3.5)$$

In turn, this allows to identify $h_c(\beta)$ in terms of $\phi(\beta, \cdot)$ as

$$h_c(\beta) = \inf \left\{ h : \sup_{m \in [0, 1]} (\phi(\beta, m) - hm) = 0 \right\}. \quad (3.6)$$

The key technical step in the proof of Theorem 2.1 is the following:

Theorem 3.1. *Under condition C1 or C2, for every $0 < \beta < \infty$ there exists $0 < \mathbf{C}(\beta) < \infty$ such that, for every $1 \leq \alpha < \infty$,*

$$\phi(\beta, m) - h_c(\beta)m \leq -\frac{\mathbf{C}(\beta)}{\alpha} m^2 \quad (3.7)$$

if $0 \leq m \leq 1$.

Proof of Theorem 2.1. It is an immediate consequence of Theorem 3.1 and (3.5). In fact, by (3.7) we have

$$\phi(\beta, m) - hm \leq -\frac{\mathbf{C}(\beta)}{\alpha}m^2 + (h_c(\beta) - h)m, \quad (3.8)$$

for every $m \in [0, 1]$. Taking the supremum over m on both sides of (3.8), by (3.5) we obtain

$$\begin{aligned} \mathbb{F}(\beta, h) &\leq \sup_{m \in [0, 1]} \left(-\frac{\mathbf{C}(\beta)}{\alpha}m^2 + (h_c(\beta) - h)m \right) \\ &\leq \sup_{m \geq 0} \left(-\frac{\mathbf{C}(\beta)}{\alpha}m^2 + (h_c(\beta) - h)m \right) = \frac{\alpha}{4\mathbf{C}(\beta)}(h_c(\beta) - h)^2. \end{aligned} \quad (3.9)$$

Theorem 2.1, pinning \square

We go now to the proof of Theorem 3.1. We will consider first the case in which ω_1 satisfies condition **C2**, because it technically lighter. The two cases differ only in the first part of the proof (that is, up to Remark 3.2 below), where the probability of a rare event is estimated from below by changing the law of the disorder and by evaluating the corresponding relative entropy price. In the case of **C2** it is sufficient to *shift* (i.e. to *translate*) the distribution of the disorder variables, and the relative entropy estimate implied by (2.2) fits well the rest of the proof. Under assumption **C1**, instead, we have to *tilt* the law of ω and the arising expressions need to be re-worked, see Lemma 3.4 below, before stepping to the second part of the proof.

Proof of Theorem 3.1 under assumption C2. Due to the concavity of $\phi(\beta, \cdot)$, it is enough to prove (3.7) for $0 < m \leq c_1$, with $c_1 = c_1(\beta) > 0$, not depending on α . We define

$$\gamma(\beta, m) := -\phi(\beta, m) + h_c(\beta)m + c_2 \frac{\beta^2 m^2}{\alpha} > 0, \quad (3.10)$$

where $c_2 > 0$ is a constant depending only on \mathbb{P} , which will be chosen later. We stress that the term containing c_2 has been added simply because a priori one knows only that $-\phi(\beta, m) + h_c(\beta)m \geq 0$, cf. (3.6), and it turns out to be technically practical to work with $\gamma(\beta, m) > 0$. For $\ell \in \mathbb{N}$ we define also

$$A_{\ell, m, \varepsilon} = \left\{ \omega \in \mathbb{R}^\ell : \log \hat{Z}_{\ell, \omega}(\beta; m, \varepsilon) - h_c(\beta)m\ell \geq \gamma(\beta, m)\ell \right\}. \quad (3.11)$$

Moreover for $0 < m \leq c_1$, we let $\tilde{\mathbb{P}}$ be the law obtained from \mathbb{P} shifting the distribution of $\omega_1, \dots, \omega_\ell$ so that

$$\tilde{\mathbb{E}}[\omega_j] = 8 \frac{\gamma(\beta, m)}{\beta m}. \quad (3.12)$$

Note that

$$\lim_{m \rightarrow 0} \gamma(\beta, m)/m = 0, \quad (3.13)$$

since otherwise, by convexity of $\gamma(\beta, \cdot)$, one has $\gamma(\beta, m) \geq \varepsilon m$ for some $\varepsilon > 0$ and every m , that is $\mathbb{F}(\beta, h_c(\beta) - \varepsilon) = 0$, which is in contrast with the definition of $h_c(\beta)$.

We now show that the event

$$E := \{ \omega : (\omega_1, \dots, \omega_\ell) \in A_{\ell, m, \varepsilon} \}, \quad (3.14)$$

becomes $\tilde{\mathbb{P}}$ -typical for ℓ large.

We first observe that, thanks to the constraint $\mathcal{N}_\ell/\ell \geq m - \varepsilon$, and assuming that $\varepsilon \leq m/2$, one has

$$\tilde{\mathbb{E}} \left(\frac{1}{\ell} \log \hat{Z}_{\ell,\omega}(\beta; m, \varepsilon) - h_c(\beta)m \right) \geq 4\gamma(\beta, m) + \mathbb{E} \left(\frac{1}{\ell} \log \hat{Z}_{\ell,\omega}(\beta; m, \varepsilon) - h_c(\beta)m \right). \quad (3.15)$$

By (3.2) and (3.10), there exist $\varepsilon_0(m) > 0$ and $\ell_0(\varepsilon, m) \in s\mathbb{N}$ such that

$$\tilde{\mathbb{E}} \left(\frac{1}{\ell} \log \hat{Z}_{\ell,\omega}(\beta; m, \varepsilon) - h_c(\beta)m \right) \geq 2\gamma(\beta, m), \quad (3.16)$$

for $\varepsilon \leq \varepsilon_0, \ell \geq \ell_0$. This in turn implies that

$$\tilde{\mathbb{P}}(E) \geq \tilde{\mathbb{P}} \left(\log \hat{Z}_{\ell,\omega}(\beta; m, \varepsilon) - \tilde{\mathbb{E}} \log \hat{Z}_{\ell,\omega}(\beta; m, \varepsilon) \geq -\gamma(\beta, m)\ell \right), \quad (3.17)$$

which is greater than, say, $1/2$ for ℓ sufficiently large, since $\gamma(\beta, m) > 0$ and (recall (3.2) and discussion following that formula)

$$\frac{1}{\ell} \log \hat{Z}_{\ell,\omega}(\beta; m, \varepsilon) - \frac{1}{\ell} \tilde{\mathbb{E}} \log \hat{Z}_{\ell,\omega}(\beta; m, \varepsilon) \xrightarrow{\ell \rightarrow \infty} 0, \quad (3.18)$$

in $\tilde{\mathbb{P}}$ -probability.

The price of shifting \mathbb{P} to $\tilde{\mathbb{P}}$ is directly estimated by using the assumption **C2**, cf. (2.2), and recalling (3.13): assuming that $m \leq c_1(\beta)$, with $c_1(\beta)$ sufficiently small, one obtains the estimate

$$\mathbb{H}(\tilde{\mathbb{P}}|\mathbb{P}) := \mathbb{E} \left(\frac{d\tilde{\mathbb{P}}}{d\mathbb{P}} \log \frac{d\tilde{\mathbb{P}}}{d\mathbb{P}} \right) \leq 64R \left(\frac{\gamma(\beta, m)}{\beta m} \right)^2 \ell, \quad (3.19)$$

and by applying the relative entropy inequality

$$\log \left(\frac{\mathbb{P}(E)}{\tilde{\mathbb{P}}(E)} \right) \geq -\frac{1}{\tilde{\mathbb{P}}(E)} \left(\mathbb{H}(\tilde{\mathbb{P}}|\mathbb{P}) + e^{-1} \right), \quad (3.20)$$

we obtain

$$\tilde{p} := \mathbb{P}(E) \geq \frac{1}{2} \exp \left(-128R (\gamma(\beta, m)/(\beta m))^2 \ell \right), \quad (3.21)$$

for large ℓ .

Remark 3.2. Inequality (3.20) holds whenever the measures $\mathbb{P}, \tilde{\mathbb{P}}$ are absolutely continuous with respect to each other and for every event E of nonzero measure. It is a simple consequence of Jensen inequality: since $r \log r \geq -e^{-1}$ for every $r > 0$, one has

$$\begin{aligned} \log \left(\frac{\mathbb{P}(E)}{\tilde{\mathbb{P}}(E)} \right) &= \log \tilde{\mathbb{E}} \left(\frac{d\mathbb{P}}{d\tilde{\mathbb{P}}} \Big| E \right) \geq \tilde{\mathbb{E}} \left(\log \left(\frac{d\mathbb{P}}{d\tilde{\mathbb{P}}} \right) \Big| E \right) \\ &= -\frac{1}{\tilde{\mathbb{P}}(E)} \mathbb{E} \left(\frac{d\tilde{\mathbb{P}}}{d\mathbb{P}} \log \frac{d\tilde{\mathbb{P}}}{d\mathbb{P}} \mathbf{1}_E \right) \geq -\frac{1}{\tilde{\mathbb{P}}(E)} \left(\mathbb{E} \left(\frac{d\tilde{\mathbb{P}}}{d\mathbb{P}} \log \frac{d\tilde{\mathbb{P}}}{d\mathbb{P}} \right) + e^{-1} \right), \end{aligned} \quad (3.22)$$

which is just (3.20).

We now apply an energy–entropy argument similar to that of [4] which, in the present case, roughly consists in selecting only those polymer trajectories which visit the rare stretches where the disorder configuration is such to produce a sufficiently large positive fluctuation of the partition function. Of course the precise definition of these rare stretches is directly related to the event E . This selection strategy gives a lower bound on the free

energy, which implies (3.7). More precisely, we consider a system of length $k\ell$, with $k \in \mathbb{N}$, $\ell \in s\mathbb{N}$ and we divide it into blocks $B_j = \{j\ell + 1, j\ell + 2, \dots, (j+1)\ell\}$ of length ℓ , with $j = 0, \dots, k-1$. For a given realization of ω , we denote by $\mathcal{I}(\omega)$ the ordered set of nonnegative integers

$$\mathcal{I}(\omega) = \{j_1, \dots, j_{|\mathcal{I}(\omega)|}\} := \{0 \leq j \leq k-1 : (\omega_{j\ell+1}, \dots, \omega_{(j+1)\ell}) \in A_{\ell, m, \varepsilon}\}, \quad (3.23)$$

where it is understood that $\varepsilon \leq \varepsilon_0(m)$, $\ell \geq \ell_0(m, \varepsilon)$ and $m \leq c_1$ as above, so that (3.21) holds. We bound the partition function $Z_{N, \omega}$ below by inserting in the average over the paths the constraint that $S_i \neq 0$ whenever $i \in B_j$ with $j \notin \mathcal{I}(\omega)$, that $S_i = 0$ whenever $i = j\ell$ or $i = (j+1)\ell$ with $j \in \mathcal{I}(\omega)$, and that, for every $j \in \mathcal{I}(\omega)$,

$$\frac{|\{i \in B_j : S_i = 0\}|}{\ell} \in [m - \varepsilon, m + \varepsilon]. \quad (3.24)$$

In this way, recalling the definition of $A_{\ell, m, \varepsilon}$ and assuming that $\varepsilon \leq \gamma(\beta, m)/(2|h_c(\beta)|)$,




FIGURE 1. A typical trajectory contributing to the lower bound (3.25). In this example $k = 22$, $\ell = 8$, $\mathcal{I}(\omega) = \{4, 10, 17\}$. Note that $S_n \neq 0$ for n in a block B_j with $j \notin \{4, 10, 17\}$, except at the boundary with a block B_j with $j \in \{4, 10, 17\}$, since by construction $S_n = 0$ at the boundaries of these blocks. Inside B_j , $j \in \mathcal{I}(\omega)$, the path moves with relative freedom, but it is bound to touch the line $S = 0$ approximately $m\ell$ times (see the text for the choice of m).

one obtains

$$\frac{1}{k\ell} \log Z_{k\ell, \omega}(\beta, h_c(\beta)) \geq \frac{1}{k} |\mathcal{I}(\omega)| \frac{\gamma(\beta, m)}{2} + \frac{1}{k\ell} \log \prod_{\substack{r=0, \dots, |\mathcal{I}(\omega)| \\ L_r \neq 0}} K(L_r), \quad (3.25)$$

where we recall that $K(n)$ is the \mathbf{P} -probability that first return to 0 of an excursion of the free process occurs at step n , as in (1.2), while L_r 's, the (possibly vanishing) lengths of the excursions of the process between two blocks $B_i, B_{i'}$ with $i, i' \in \mathcal{I}(\omega)$, are defined as

$$L_r = \ell |j_{r+1} - j_r - 1|, \quad (3.26)$$

with the convention that $j_0 = -1$, $j_{|\mathcal{I}(\omega)|+1} = k$, see Figure 1. Taking the expectation with respect to the disorder and using (1.3), one obtains then

$$\frac{1}{k\ell} \mathbb{E} \log Z_{k\ell, \omega}(\beta, h_c(\beta)) \geq \frac{\gamma(\beta, m)}{2} \tilde{p} - \frac{2\alpha}{k\ell} \mathbb{E} \sum_{r=0}^{|\mathcal{I}(\omega)|} \mathbf{1}_{L_r \neq 0} \log L_r, \quad (3.27)$$

for ℓ sufficiently large. At this point, as in [4], one uses Jensen's inequality and the concavity of the logarithm to get

$$\begin{aligned} \frac{1}{k\ell} \mathbb{E} \log Z_{k\ell, \omega}(\beta, h_c(\beta)) &\geq \frac{\gamma(\beta, m)}{2} \tilde{p} - \frac{2\alpha}{k\ell} \mathbb{E} \left[(|\mathcal{I}(\omega)| + 1) \log \left(\frac{\sum_{r=0}^{|\mathcal{I}(\omega)|} L_r}{|\mathcal{I}(\omega)| + 1} \right) \right] \\ &\geq \frac{\gamma(\beta, m)}{2} \tilde{p} - 2\alpha \mathbb{E} \left[\frac{(|\mathcal{I}(\omega)| + 1)}{k\ell} \log \left(\frac{k\ell}{|\mathcal{I}(\omega)| + 1} \right) \right]. \end{aligned} \quad (3.28)$$

Since the disorder variables in the distinct blocks $\{B_j\}_j$ are independent, the law of large numbers implies

$$\frac{|\mathcal{I}(\omega)|}{k} \xrightarrow{k \rightarrow \infty} \tilde{p} \quad (3.29)$$

$\mathbb{P}(d\omega)$ -a.s., so that, recalling (3.21),

$$\begin{aligned} 0 = \mathbb{F}(\beta, h_c(\beta)) &= \lim_{k \rightarrow \infty} \frac{1}{k\ell} \mathbb{E} \log Z_{k\ell, \omega}(\beta, h_c(\beta)) \\ &\geq \tilde{p} \frac{\gamma(\beta, m)}{2} - \frac{2\alpha \tilde{p}}{\ell} \log \frac{\ell}{\tilde{p}} \geq \tilde{p} \left(\frac{\gamma(\beta, m)}{2} - 256\alpha R \left(\frac{\gamma(\beta, m)}{\beta m} \right)^2 - 2\alpha \frac{\log(2\ell)}{\ell} \right). \end{aligned} \quad (3.30)$$

Since ℓ is arbitrary, one obtains

$$\gamma(\beta, m) \geq \frac{\beta^2 m^2}{512\alpha R} \quad (3.31)$$

which is the desired inequality and (3.7), provided that c_2 in (3.10) satisfies $c_2 < (512R)^{-1}$.
Theorem 3.1, C2 \square

Remark 3.3. It is easy to check that, in the Gaussian case $\omega_1 \sim \mathcal{N}(0, 1)$, Theorem 2.1 holds with $c(\beta) = c_3 \beta^{-2}$, c_3 a suitable positive constant. Indeed, in this case the estimate (2.2) holds for every $x \in \mathbb{R}$ (and $R = 1/2$) and therefore one can take $c_1 = 1$ in the proof of Lemma 3.1, so that (3.31) implies (3.7) with $\mathbf{C}(\beta) = c_4 \beta^2$. Of course, the same is true, up to constants, for every \mathbb{P} such that inequality (2.2) holds uniformly for $x \in \mathbb{R}$.

Proof of Theorem 3.1 under assumption C1. The proof proceeds as in case C2, up to the definition of the law $\tilde{\mathbb{P}}$, but in this case the law obtained by shifting \mathbb{P} in general has an infinite entropy with respect to \mathbb{P} . Therefore, in this case we define $\tilde{\mathbb{P}}$ rather by tilting the law of the first ℓ variables:

$$\frac{d\tilde{\mathbb{P}}}{d\mathbb{P}}(\omega) = \frac{1}{z^\ell} \exp \left(u \sum_{n=1}^{\ell} \omega_n \right), \quad (3.32)$$

where $u \geq 0$ will be chosen later and $z = z(u) = \mathbb{E} e^{u\omega_1}$. Let, for $\ell \in s\mathbb{N}$,

$$\psi_\ell(u) = \frac{1}{\ell} \tilde{\mathbb{E}} \log \hat{Z}_{\ell, \omega}(\beta; m, \varepsilon) \quad (3.33)$$

and observe that

$$\psi_\ell(0) = \frac{1}{\ell} \mathbb{E} \log \hat{Z}_{\ell, \omega}(\beta; m, \varepsilon).$$

Then, one has

Lemma 3.4. *There exist $u_0(\beta) > 0$ and $c_0(\beta) > 0$, possibly depending on \mathbb{P} , such that for every $0 < \beta < \infty$, $1 \leq \alpha < \infty$ the following holds: for every $m \in (0, 1]$, if $\varepsilon \leq m/2$ and $0 \leq u \leq u_0$ we have*

$$\psi_\ell(u) - \psi_\ell(0) \geq c_0 \beta m u. \quad (3.34)$$

Lemma 3.4 will be proven below. To proceed with the proof of Theorem 3.1, we choose $u = 4\gamma(\beta, m)/(\beta m c_0)$ and notice that u is certainly smaller than u_0 if $m \leq c_1(\beta)$ with c_1 sufficiently small (see (3.13)). Then, choosing $0 < m \leq c_1$ and $\varepsilon \leq m/2$, (3.34) implies that (3.15) is valid also in the present case. On the other hand, it is immediate to verify that (3.19) still holds, with R replaced by some $R'(\beta, M)$ and $\gamma(\beta, m)$ by $\gamma(\beta, m)/c_0$. The rest of the proof proceeds exactly as in the case **C2**.

Theorem \square 3.1, **C1**

Proof of Lemma 3.4. Note that

$$\partial_u \psi_\ell(u) = \frac{1}{\ell} \sum_{i=1}^{\ell} \tilde{\mathbb{E}} \left[\omega_i \log \hat{Z}_{\ell, \omega}(\beta; m, \varepsilon) \right] - \ell \xi \psi_\ell(u), \quad (3.35)$$

where $\xi = \xi(u) = \tilde{\mathbb{E}} \omega_1$. The first term in the right-hand side of (3.35) can be rewritten (with some abuse of notation) as

$$\begin{aligned} & \frac{\xi}{\ell} \sum_{i=1}^{\ell} \tilde{\mathbb{E}} \left[\log \hat{Z}_{\ell, \omega}(\beta; m, \varepsilon) \Big|_{\omega_i=0} \right] + \frac{\beta}{\ell} \sum_{i=1}^{\ell} \tilde{\mathbb{E}} \left[\omega_i \int_0^{\omega_i} dy \hat{\mathbf{E}}_{\ell, \omega}(\mathbf{1}_{S_i=0}) \Big|_{\omega_i=y} \right] \\ &= \ell \xi \psi_\ell(u) + \frac{\beta}{\ell} \sum_{i=1}^{\ell} \tilde{\mathbb{E}} \left[(\omega_i - \xi) \int_0^{\omega_i} dy \hat{\mathbf{E}}_{\ell, \omega}(\mathbf{1}_{S_i=0}) \Big|_{\omega_i=y} \right], \end{aligned} \quad (3.36)$$

where

$$\hat{\mathbf{E}}_{\ell, \omega}(\cdot) = \frac{\mathbf{E} \left[\cdot \exp \left(\sum_{i=1}^{\mathcal{N}_\ell} \beta \omega_{\tau_i} \right) \mathbf{1}_{S_\ell=0} \mathbf{1}_{(\mathcal{N}_\ell/\ell) \in [m-\varepsilon, m+\varepsilon]} \right]}{\hat{Z}_{\ell, \omega}(\beta; m, \varepsilon)} \quad (3.37)$$

and obviously the first term in (3.36) cancels the second one in the right-hand side of (3.35). Next, observe that the following identity holds:

$$\hat{\mathbf{E}}_{\ell, \omega}(\mathbf{1}_{S_i=0}) \Big|_{\omega_i=y} = \frac{e^{\beta(y-y')} \hat{\mathbf{E}}_{\ell, \omega}(\mathbf{1}_{S_i=0}) \Big|_{\omega_i=y'}}{1 + (e^{\beta(y-y')} - 1) \hat{\mathbf{E}}_{\ell, \omega}(\mathbf{1}_{S_i=0}) \Big|_{\omega_i=y'}}. \quad (3.38)$$

Now recall (2.1), so that it is sufficient to consider y and y' such that $|y - y'| \leq 2M$, and from (3.38) we obtain

$$e^{-2\beta M} \hat{\mathbf{E}}_{\ell, \omega}(\mathbf{1}_{S_i=0}) \Big|_{\omega_i=y'} \leq \hat{\mathbf{E}}_{\ell, \omega}(\mathbf{1}_{S_i=0}) \Big|_{\omega_i=y} \leq e^{+2\beta M} \hat{\mathbf{E}}_{\ell, \omega}(\mathbf{1}_{S_i=0}) \Big|_{\omega_i=y'}. \quad (3.39)$$

We can use this inequality to bound below the last term in (3.36). We have in fact

$$\tilde{\mathbb{E}} \left[\omega_i \int_0^{\omega_i} dy \hat{\mathbf{E}}_{\ell, \omega}(\mathbf{1}_{S_i=0}) \Big|_{\omega_i=y} \right] \geq e^{-2\beta M} \eta \tilde{\mathbb{E}} \hat{\mathbf{E}}_{\ell, \omega}(\mathbf{1}_{S_i=0}) \Big|_{\omega_i=0} \geq e^{-4\beta M} \eta \tilde{\mathbb{E}} \hat{\mathbf{E}}_{\ell, \omega}(\mathbf{1}_{S_i=0}),$$

where $\eta = \eta(u) = \tilde{\mathbb{E}} \omega_1^2$, while

$$\tilde{\mathbb{E}} \int_0^{\omega_i} dy \hat{\mathbf{E}}_{\ell, \omega}(\mathbf{1}_{S_i=0}) \Big|_{\omega_i=y} \leq M e^{2\beta M} \tilde{\mathbb{E}} \hat{\mathbf{E}}_{\ell, \omega}(\mathbf{1}_{S_i=0}). \quad (3.40)$$

Therefore, recalling the constraint $(\mathcal{N}_\ell/\ell) \in [m - \varepsilon, m + \varepsilon]$ in the definition of $\hat{\mathbf{P}}_{\ell,\omega}(\cdot)$, one has the following lower bound:

$$\partial_u \psi_\ell(u) \geq \beta(m - \varepsilon)\eta e^{-4M\beta} - \beta(m + \varepsilon)\xi M e^{2\beta M}. \quad (3.41)$$

Now choose $\varepsilon \leq m/2$ and notice that $\xi = u + O(u^2)$ for $u \ll 1$, while $\eta = 1 + O(u)$. Therefore, there exists $u_0(\beta, M) > 0$ such that, for $0 \leq u \leq u_0$ and for every m , the following holds:

$$\partial_u \psi_\ell(u) \geq \beta m \frac{e^{-4M\beta}}{4} - 2\beta m u M e^{2\beta M} \geq \beta m \frac{e^{-4M\beta}}{8} =: c_0 \beta m. \quad (3.42)$$

An integration in u concludes the proof of (3.34).

Lemma 3.4 \square

3.2. The copolymer case. In order to prove Theorem 2.1 for the copolymer model (1.12), it is convenient to start from the observation that one can rewrite the limit free energy as

$$\mathbb{F}(\beta, h) = \lim_{N \rightarrow \infty} \frac{1}{N} \mathbb{E} \log \mathbf{E} \left[\exp \left(- \sum_{n=1}^N (\beta \omega_n + h) \Delta_n \right) \mathbf{1}_{S_N=0} \right], \quad (3.43)$$

where $\Delta_n = \mathbf{1}_{\text{sign}(S_n)=-1}$. Comparing this expression for the copolymer free energy with (1.1), it is clear that in the present case the role of $\mathbf{1}_{S_n=0}$ is played by Δ_n . The proof then proceeds exactly like in the pinning case, with the only differences that in the definition (3.2) of $\phi(\beta, m)$ the constraint $(\mathcal{N}_N/N) \in [m - \varepsilon, m + \varepsilon]$ has to be replaced by

$$\frac{|\{1 \leq n \leq N : \Delta_n = 1\}|}{N} \in [m - \varepsilon, m + \varepsilon] \quad (3.44)$$

and that, in the energy–entropy argument, the path is required to satisfy $S_i > 0$ (and not just $S_i \neq 0$) whenever $i \in B_j$ with $j \notin \mathcal{I}(\omega)$. This implies that $K(L)$ in (3.25) has to be replaced by $K(L)/2$, which has the effect of adding a negative term of order $O(\tilde{p}/\ell)$ in the lower bound (3.30), which is negligible for ℓ sufficiently large.

Theorem 3.1, copolymer \square

APPENDIX A. THE NON-DISORDERED PINNING MODEL

For $\beta = 0$, the free energy (1.7) may be identified explicitly with the following procedure. First we consider the equation

$$\sum_{n \in \mathbb{N}} K(n) \exp(-bn) = \exp(h), \quad (\text{A.1})$$

and we look for a solution $b > 0$, which exists only if $\sum_{n \in \mathbb{N}} K(n) > \exp(h)$, in which case it is unique. Then if we set $\tilde{K}(n) := \exp(-h - bn)K(n)$, $\tilde{K}(\cdot)$ is a discrete probability

density and one can write

$$\begin{aligned} \mathbf{E}[\exp(-h\mathcal{N}_N); \tau_{\mathcal{N}_N} = N] &= \sum_{j=1}^N \sum_{\substack{(\ell_1, \dots, \ell_j) \in \mathbb{N}^j \\ \sum_{i=1}^j \ell_i = N}} \prod_{i=1}^j \exp(-h)K(\ell_i) \\ &= \exp(bN) \sum_{j=1}^N \sum_{\substack{(\ell_1, \dots, \ell_j) \in \mathbb{N}^j \\ \sum_{i=1}^j \ell_i = N}} \prod_{i=1}^j \tilde{K}(\ell_i) =: \exp(bN) G_N, \end{aligned} \tag{A.2}$$

and one easily sees that G_N is the probability that the random walk which starts at 0 and takes positive integer IID jumps with law $\tilde{K}(\cdot)$ hits the site N . It is a classical fundamental result of renewal theory that $\lim_N G_N = 1/\sum_{n \in \mathbb{N}} n\tilde{K}(n)$ [11, Ch. XIII]. This of course implies that $b = F(0, h)$. On the other hand, if (A.1) admits no positive solution, by proceeding as in (A.1) and by setting simply $\tilde{K}(n) := \exp(-h)K(n)$, so that $\tilde{K}(\cdot)$ is a sub-probability density, one easily sees that $F(0, h) = 0$. So equation (A.1) contains all the information about the free energy.

Let us then observe that (A.1) has a positive solution if and only if $h < \log(1 - K(\infty))$ and therefore $h_c(0) = \log(1 - K(\infty)) \leq 0$ ($h_c(\beta)$ being defined before Remark 1.1). The behavior at criticality can be extracted from (A.1) in a rather straightforward way too, but of course we need to make precise the requirement on $K(\cdot)$ beyond the lower bound (1.3):

- *The case $\Sigma := \sum_{n \in \mathbb{N}} nK(n) < \infty$.* This is a necessary and sufficient condition for the transition to be of first order. More precisely, for $\Sigma \in (0, \infty]$

$$F(0, h) = \frac{\exp(h_c(0))}{\Sigma} (h_c(0) - h) + o((h_c(0) - h)), \quad \text{for } h \nearrow h_c(0). \tag{A.3}$$

Formula (A.3) follows since by Dominated Convergence if $\Sigma < \infty$ then

$$\sum_{n \in \mathbb{N}} K(n) \exp(-bn) = \exp(h_c(0)) - b\Sigma + o(b) \tag{A.4}$$

for $b \searrow 0$, while by a direct estimate $\lim_{b \searrow 0} b^{-1} \sum_{n \in \mathbb{N}} K(n)[1 - \exp(-bn)] = \infty$ if $\Sigma = \infty$. Note that the condition $\Sigma < \infty$ holds, in particular, in the case where

$$K(n) = L(n)/n^\alpha \tag{A.5}$$

for large $n \in \mathbb{N}$, with $\alpha > 2$ and $L(\cdot)$ a function varying slowly at infinity, i.e., a positive function such that $\lim_{r \rightarrow \infty} L(xr)/L(r) = 1$ for every $x > 0$ (see [3] for more details on slowly varying functions).

- *The case $\Sigma = \infty$.* We set $\bar{K}(n) = \sum_{j=n+1}^{\infty} K(j)$ (by this we mean the sum over $j \in \mathbb{N}$ such that $j > n$: $K(\infty)$ is not included in the sum) and assume that

$$\sum_{j=1}^N \bar{K}(j) = \hat{L}(N)N^{2-\alpha},$$

for some function $\hat{L}(\cdot)$ which is slowly varying at infinity. This is true, in particular, in the case (A.5). By the easy (Abelian) part of the classical Tauberian Theorem [12, Ch.XIII.5, Theorem 2] we have that $\sum_{n=0}^{\infty} \exp(-bn)\bar{K}(n) = c'b^{\alpha-2}\hat{L}(1/b)(1 +$

$o(1))$ as $b \searrow 0$, with $c' = c'(\alpha) > 0$. Therefore

$$\begin{aligned} \sum_{n=1}^{\infty} e^{-bn} K(n) &= \bar{K}(0) + (e^{-b} - 1) \sum_{n=0}^{\infty} e^{-bn} \bar{K}(n) \\ &= \bar{K}(0) - c'b^{\alpha-1} \hat{L}(1/b)(1 + o(1)), \end{aligned} \quad (\text{A.6})$$

and

$$F(0, h) = (h_c(0) - h)^{1/(\alpha-1)} \tilde{L}(1/(h_c(0) - h)), \quad \text{for } h \nearrow h_c(0), \quad (\text{A.7})$$

with $\tilde{L}(\cdot)$ a slowly varying function (see [3, (1.5.1) and Theorem 1.5.12]). It is therefore clear that the transition is of second order for $\alpha \in (3/2, 2]$ (we emphasize, for the case $\alpha = 2$, that we are assuming that $\Sigma = +\infty$) and it is of higher order for $\alpha < 3/2$. The value $\alpha = 3/2$ is borderline and the order of the transition depends then on the slowly varying function $\hat{L}(\cdot)$. In the case of one dimensional symmetric random walks with IID increments taking values in $\{-1, 0, +1\}$, $\alpha = 3/2$ and $L(n) = c(1 + o(1))$ for n large, c a positive constant, and therefore the transition is really of second order and not higher.

ACKNOWLEDGMENTS

We would like to thank Bernard Derrida, Thomas Garel, Cécile Monthus and David Mukamel for interesting discussions. This research has been conducted in the framework of the GIP-ANR project JC05_42461 (*POLINTBIO*).

REFERENCES

- [1] M. Aizenman and J. Wehr, *Rounding effects of quenched randomness on first-order phase transitions*, Comm. Math. Phys. **130** (1990), 489–528.
- [2] K. S. Alexander and V. Sidoravicius, *Pinning of polymers and interfaces by random potentials*, preprint (2005). Available on: arXiv.org e-Print archive: math.PR/0501028
- [3] N. H. Bingham, C. M. Goldie and J. L. Teugels, *Regular Variation*, Cambridge University Press, Cambridge, 1987.
- [4] T. Bodineau and G. Giacomin, *On the localization transition of random copolymers near selective interfaces*, J. Statist. Phys. **117** (2004), 801–818.
- [5] E. Bolthausen and F. den Hollander, *Localization transition for a polymer near an interface*, Ann. Probab. **25** (1997), 1334–1366.
- [6] A. Bovier and C. Külske, *There are no nice interfaces in $(2+1)$ -dimensional SOS models in random media*, J. Statist. Phys. **83** (1996), 751–759.
- [7] J. T. Chayes, L. Chayes, D. S. Fisher and T. Spencer, *Correlation Length Bounds for Disordered Ising Ferromagnets*, Commun. Math. Phys. **120**, 501–523 (1989).
- [8] B. Coluzzi, *Numerical study on a disordered model for DNA denaturation transition*, arXiv:cond-mat/0504080v1 (2005).
- [9] D. Cule and T. Hwa, *Denaturation of heterogeneous DNA*, Phys. Rev. Lett. **79** (1997), 2375–2378.
- [10] B. Derrida, V. Hakim and J. Vannimenus, *Effect of disorder on two-dimensional wetting*, J. Statist. Phys. **66** (1992), 1189–1213.
- [11] W. Feller, *An introduction to probability theory and its applications*, Vol. I, Third edition, John Wiley & Sons, Inc., New York–London–Sydney, 1968.
- [12] W. Feller, *An introduction to probability theory and its applications*, Vol. II, Second edition, John Wiley & Sons, Inc., New York–London–Sydney, 1971.
- [13] G. Forgacs, J. M. Luck, Th. M. Nieuwenhuizen and H. Orland, *Wetting of a Disordered Substrate: Exact Critical behavior in Two Dimensions*, Phys. Rev. Lett. **57** (1986), 2184–2187.
- [14] T. Garel, D. A. Huse, S. Leibler and H. Orland, *Localization transition of random chains at interfaces*, Europhys. Lett. **8** (1989), 9–13.

- [15] T. Garel and C. Monthus, *Numerical study of the disordered Poland–Scheraga model of DNA denaturation*, J. Stat. Mech., Theory and Experiments (2005), P06004.
- [16] G. Giacomin, *Localization phenomena in random polymer models*, preprint (2004). Available online: <http://www.proba.jussieu.fr/pageperso/giacomin/pub/publicat.html>
- [17] G. Giacomin and F. L. Toninelli, *Estimates on path delocalization for copolymers at selective interfaces*, Probab. Theor. Rel. Fields **133** (2005), 464–482.
- [18] G. Giacomin and F. L. Toninelli, *The localized phase of disordered copolymers with adsorption*, preprint (2005). Available on: arXiv.org e-Print archive: math.PR/0510047
- [19] A. B. Harris, *Effect of random defects on the critical behaviour of Ising models*, J. Phys. C **7** (1974), 1671–1692.
- [20] Y. Imry and S.–K. Ma, *Random–Field Instability of the Ordered State of Continuous Symmetry*, Phys. Rev. Lett. **35** (1975), 1399–1401.
- [21] Y. Kafri, D. Mukamel and L. Peliti, *Why is the DNA denaturation transition first order?*, Phys. Rev. Lett. **85** (2000), 4988–4991.
- [22] J. F. C. Kingman, *Subadditive ergodic theory*, Ann. Probab. **1** (1973), 882–909.
- [23] C. Monthus, *On the localization of random heteropolymers at the interface between two selective solvents*, Eur. Phys. J. B **13** (2000), 111–130.
- [24] N. Petrelis, *Polymer pinning at an interface*, preprint (2005). Available on: arXiv.org e-Print archive: math.PR/0504464
- [25] Ya. G. Sinai, *A random walk with a random potential*, Theory Probab. Appl. **38** (1993), 382–385.
- [26] C. E. Soteros and S. G. Whittington, *The statistical mechanics of random copolymers*, J. Phys. A: Math. Gen. **37** (2004), R279–R325.
- [27] L.–H. Tang and H. Chaté, *Rare–Event Induced Binding Transition of Heteropolymers*, Phys. Rev. Lett. **86** (2001), 830–833.
- [28] A. Trovato and A. Maritan, *A variational approach to the localization transition of heteropolymers at interfaces*, Europhys. Lett. **46** (1999), 301–306.

LABORATOIRE DE PROBABILITÉS DE P 6 & 7 (CNRS U.M.R. 7599) AND UNIVERSITÉ PARIS 7 – DENIS DIDEROT, U.F.R. MATHÉMATIQUES, CASE 7012, 2 PLACE JUSSIEU, 75251 PARIS CEDEX 05, FRANCE

Home page: <http://www.proba.jussieu.fr/pageperso/giacomin/GBpage.html>

E-mail address: giacomini@math.jussieu.fr

LABORATOIRE DE PHYSIQUE, ENS LYON (CNRS U.M.R. 5672), 46 ALLÉE D’ITALIE, 69364 LYON CEDEX 07, FRANCE

Home page: <http://perso.ens-lyon.fr/fabio-lucio.toninelli>

E-mail address: fltonine@ens-lyon.fr