

# Likelihood inference for incompletely observed stochastic processes: ignorability conditions

Daniel Commenges and Anne Gégout-Petit

*Daniel Commenges, Epidemiology and Biostatistics Research Center, INSERM, Université*

*Victor Segalen Bordeaux 2, 146 rue Léo Saignat, Bordeaux, 33076, France*

*Anne Gégout-Petit, IMB. INRIA project CQFD, Université Victor Segalen Bordeaux 2, 3,*

*Place de la Victoire, 33076 BORDEAUX Cedex*

*e-mail:* [daniel.commenges@isped.u-bordeaux2.fr](mailto:daniel.commenges@isped.u-bordeaux2.fr); [anne.petit@u-bordeaux2.fr](mailto:anne.petit@u-bordeaux2.fr)

<https://sites.google.com/site/danielcommenges/>

**Abstract:** We develop a study of ignorability and conditions thereof for likelihood inference in the framework of stochastic processes. We define a coarsening model for processes which includes discrete-time observations as well as censored continuous-time observations and applies to continuous state-space processes as well as counting processes. For preparing the work we recall formulas for manipulating marginal and conditional likelihood ratios (which can apply to stochastic processes). Ignorability is defined in terms of local equality of two likelihood ratios. We give static conditions of ignorability and then dynamical conditions which are more interpretable. We illustrate the use of the dynamical conditions of ignorability in problems of censoring, missing data and joint modelling.

**AMS 2000 subject classifications:** Primary 62M99; secondary 62F99.

**Keywords and phrases:** coarsening, censoring, counting processes, ignorability, likelihood, longitudinal data, missing data, Radon-Nikodym derivatives, stochastic processes.

## Contents

1	Introduction . . . . .	2
2	Full, marginal and conditional likelihood . . . . .	4
3	A coarsening model for processes . . . . .	6

3.1	Time coarsening: the $(R_t)$ process . . . . .	6
3.2	Extension to vertical coarsening . . . . .	9
4	Ignorability . . . . .	10
4.1	The sigma-field representation for incomplete data . . . . .	10
4.2	Model, notations and observed likelihood . . . . .	12
4.3	Definition of ignorability . . . . .	13
5	Static conditions of ignorability for the TCMP . . . . .	14
6	Dynamical conditions of ignorability . . . . .	17
7	Applications . . . . .	20
7.1	Right-censoring of counting processes with time-dependent co- variable . . . . .	20
7.2	Longitudinal markers (continuous state-space processes) . . . . .	22
7.2.1	Missing data at fixed observation times . . . . .	22
7.2.2	Random observation times . . . . .	22
7.2.3	Vertically coarsened observations . . . . .	23
7.3	Joint modeling . . . . .	23
8	Conclusion . . . . .	24
	References . . . . .	25
A	CAR(TCMP)-loc . . . . .	27
B	Proof of Theorem 6.3 . . . . .	29
C	Extension to left-continuous $R$ . . . . .	30

## 1. Introduction

Incomplete data are very common in statistics: when the mechanism leading to incomplete data (m.l.i.d.) is fixed a relatively simple likelihood can be written in general. Often the m.l.i.d. can not be considered as fixed and the question arises whether it can still be ignored. Rubin (27) introduced the concept of ignorability for the simplest case in which the observation is a sequence of random variables and some of them are missing; he gave conditions under which inference based on the assumption of fixed m.l.i.d. was valid, even in the case when in fact it

was not fixed. He established a typology of cases of missing data and showed in particular that the m.l.i.d. was ignorable for likelihood inference in the case of missing at random (MAR) observations.

In the framework of survival analysis the most frequent cases of incomplete data is right-censoring (9; 21) and interval censoring (25). Conditions under which the conventional likelihood for right-censored survival data was valid have been studied (19; 23). Andersen et al. (2) developed the concept of independent censoring in the counting process framework. Heitjan and Rubin (14) also proposed some less conventional incomplete data cases which they called “coarsening”. This topic was also studied by Jacobsen and Keiding (18), Gill et al. (12) and Nielsen (24). The problematic of non-ignorable m.l.i.d. has prompted the development of joint models, in which the m.l.i.d. was included, for instance in a model proposed by Diggle and Kenward (10); see (28) for a recent example.

The aim of this paper is to study ignorability in the context of stochastic processes: these processes may be counting processes but also continuous state-space processes, such as diffusion processes. For giving a rigorous treatment of that topic, we will need to rely on basic probability tools. First we will speak in terms of likelihood ratio which is defined as a Radon-Nikodym derivative: this enables to manipulate likelihood ratios for the observation of stochastic processes (for a review see (5)). Local equalities of  $\sigma$ -fields and of random variables will play an important role in the very definition of ignorability and in the proofs and we recall a probability result described in (20):

**Lemma 1.1.** *Let the  $\sigma$ -fields  $\mathcal{F}, \mathcal{G} \subset \mathcal{A}$  and functions  $\xi, \eta \in L^1$  be such that  $A \cap \mathcal{F} = A \cap \mathcal{G}$  and  $\xi = \eta$  a.s. on some set  $A \in \mathcal{F} \cap \mathcal{G}$ . Then  $E[\xi|\mathcal{F}] = E[\eta|\mathcal{G}]$  a.s. on  $A$ .*

Also, results on the likelihood of point processes due to Jacod (16) will play a key role in several proofs.

We begin in section 2 by recalling the general definition of the likelihood ratio and of marginal and conditional likelihoods which are valid for stochastic processes; a set of useful formulae is given. In section 3 a coarsening model for

stochastic processes is given: it is represented by a stochastic process  $R = (R_t)$  which indicates at each time whether the process of interest,  $X$ , is observed or not; if the process of interest is multivariate,  $R$  can be multivariate. Then in section 4 we present a formulation of the incomplete observation problem based on  $\sigma$ -fields. If  $R$  is random the likelihood should include it, which complicates the inference problem. We give a definition of ignorability: essentially the mechanism leading to incomplete data is ignorable if the full likelihood is equal to the likelihood of what is observed of  $X$  in a model where the response process  $R$  is fixed and equal to its observed value. In section 5 we give static conditions of ignorability: the main condition called CAR(TCMP) (Definition 3) is that the likelihood of  $R$  given  $X$  depends only on observed quantities. In section 6 we give a dynamical condition called CAR(DYN) (Definition 5) which is more interpretable and usable than the general ones in some contexts: it says that the law of the response process only depends on what has been observed up to time  $t$ ; this is made rigorous by expressing this in terms of the equality of the compensator of the  $R$  process in a filtration including  $X$  and in the observed filtration. Finally section 7 illustrates the use of the theory in survival models, longitudinal data and joint models. Moreover we briefly illustrates different points along the paper by an example drawn from AIDS epidemiology. Section 8 is a short conclusion.

## 2. Full, marginal and conditional likelihood

Consider a measurable space  $(\Omega, \mathcal{F})$  and a family of measures  $\{P_\theta\}_{\theta \in \Theta}$  absolutely continuous relatively to a dominant measure  $P_{\theta_0}$ . For  $\mathcal{X}$  a sub- $\sigma$ -field of  $\mathcal{F}$  the likelihood ratio on  $\mathcal{X}$  is defined by:

$$\mathcal{L}_\mathcal{X}^{\theta/\theta_0} = \frac{dP_\theta}{dP_{\theta_0}|_\mathcal{X}} \quad \text{a.s.}$$

where  $\frac{dP_\theta}{dP_{\theta_0}|_\mathcal{X}}$  is the Radon-Nikodym derivative of  $P_\theta$  relatively to  $P_{\theta_0}$ . Recall that  $\frac{dP_\theta}{dP_{\theta_0}|_\mathcal{X}}$  is the  $\mathcal{X}$ -measurable random variable such that  $P_\theta(F) = \int_F \frac{dP_\theta}{dP_{\theta_0}|_\mathcal{X}} dP_{\theta_0}, F \in \mathcal{X}$ .

$\mathcal{X}$ . If  $\mathcal{X} \subset \mathcal{Y} \subset \mathcal{F}$  we have the fundamental formula (30):

$$\mathcal{L}_{\mathcal{X}}^{\theta/\theta_0} = \mathbb{E}_{\theta_0}[\mathcal{L}_{\mathcal{Y}}^{\theta/\theta_0} | \mathcal{X}] \quad \text{a.s.}$$

Note that because conditional expectations and likelihood ratios are defined a.s., all the equalities involving them are to be understood as a.s., even if this is not specified for sake of notational simplicity.

When  $\mathcal{F}$  is generated by two random elements  $X$  and  $Y$  and denote by  $\mathcal{X}$  and  $\mathcal{Y}$  the  $\sigma$ -fields they generate respectively; thus  $\mathcal{F} = \mathcal{X} \vee \mathcal{Y}$ ; we will note  $\mathcal{L}_{\mathcal{F}}^{\theta/\theta_0} = \mathcal{L}_{\mathcal{X} \vee \mathcal{Y}}^{\theta/\theta_0} = \mathcal{L}_{\mathcal{X}, \mathcal{Y}}^{\theta/\theta_0}$ . The likelihoods  $\mathcal{L}_{\mathcal{X}}^{\theta/\theta_0}$  and  $\mathcal{L}_{\mathcal{Y}}^{\theta/\theta_0}$  are called marginal likelihoods, and are linked to the full likelihood by the conditional expectations:  $\mathcal{L}_{\mathcal{X}}^{\theta/\theta_0} = \mathbb{E}[\mathcal{L}_{\mathcal{X}, \mathcal{Y}}^{\theta/\theta_0} | \mathcal{X}]$  and  $\mathcal{L}_{\mathcal{Y}}^{\theta/\theta_0} = \mathbb{E}[\mathcal{L}_{\mathcal{X}, \mathcal{Y}}^{\theta/\theta_0} | \mathcal{Y}]$ , as derives from the fundamental formula.

Conditional likelihoods can also be defined (see (15)); for brevity we do not recall the definition. The conditional likelihood ratio of  $Y$  given  $\mathcal{X}$  will be denoted  $\mathcal{L}_{\mathcal{Y}|\mathcal{X}}^{\theta/\theta_0}$ . The following properties will be used in this paper:

- i)  $\mathcal{L}_{\mathcal{Y}|\mathcal{X}}^{\theta/\theta_0}$  is  $\mathcal{X} \vee \mathcal{Y}$  – measurable
- ii)  $\forall A \in \mathcal{Y}, \quad \mathbb{E}_{\theta} [I_A | \mathcal{X}] = \mathbb{E}_{\theta_0} [I_A \mathcal{L}_{\mathcal{Y}|\mathcal{X}}^{\theta/\theta_0} | \mathcal{X}], \text{ a.s.}$
- iii)  $\mathcal{L}_{\mathcal{Y}, \mathcal{X}}^{\theta/\theta_0} = \mathcal{L}_{\mathcal{Y}|\mathcal{X}}^{\theta/\theta_0} \mathcal{L}_{\mathcal{X}}^{\theta/\theta_0} \quad \text{a.s.}$
- iv)  $\mathcal{Y} \subset \mathcal{X} \implies \mathcal{L}_{\mathcal{Y}, \mathcal{X}}^{\theta/\theta_0} = \mathcal{L}_{\mathcal{X}}^{\theta/\theta_0} \iff \mathcal{L}_{\mathcal{Y}|\mathcal{X}}^{\theta/\theta_0} = 1 \iff \mathcal{L}_{\mathcal{X}|\mathcal{Y}}^{\theta/\theta_0} = \frac{\mathcal{L}_{\mathcal{X}}^{\theta/\theta_0}}{\mathcal{L}_{\mathcal{Y}}^{\theta/\theta_0}}$
- v)  $\mathcal{L}_{\mathcal{X}_1, \dots, \mathcal{X}_m}^{\theta/\theta_0} = \mathcal{L}_{\mathcal{X}_1}^{\theta/\theta_0} \prod_{k=2}^m \mathcal{L}_{\mathcal{X}_k | \mathcal{X}_1, \dots, \mathcal{X}_{k-1}}^{\theta/\theta_0}$
- vi)  $\mathcal{X} \vee \mathcal{Y} = \mathcal{X} \vee \mathcal{Z} \implies \mathcal{L}_{\mathcal{Y}|\mathcal{X}}^{\theta/\theta_0} = \mathcal{L}_{\mathcal{Z}|\mathcal{X}}^{\theta/\theta_0}$

Note that ii) is the generalization of the main property of the likelihood ratio to conditional expectations; thus  $\mathcal{L}_{\mathcal{Y}|\mathcal{X}}^{\theta/\theta_0}$  deserves its name of conditional likelihood ratio; it also implies  $\mathbb{E}_{\theta_0} [\mathcal{L}_{\mathcal{Y}|\mathcal{X}}^{\theta/\theta_0} | \mathcal{X}] = 1$ . Note also that under the assumption of a family of equivalent measures, all the likelihoods are strictly positive a.s.

### 3. A coarsening model for processes

#### 3.1. Time coarsening: the $(R_t)$ process

We define a Time-Coarsening Model for Processes (TCMP) which is a general scheme where a process can be observed on a set of times, the set being possibly random. In many real studies, in particular in epidemiology, we would have to consider a sample of  $n$  independent “subjects”, to each of whom a process  $X^i = (X_t^i) = (X_t^i)_{t \geq 0}$  would be associated. Since the likelihood would be the product of the individual likelihoods, it is sufficient to consider only one process. We first consider a process  $X = (X_t)$  where  $X_t$  takes values in  $\mathfrak{R}$ , then we will extend the model to a multivariate process. The main objective is to describe observation schemes for processes in continuous time  $t$ , but  $t$  may also be discrete so that the results can be applied to finite collections of random variables. We shall consider a response indicator process  $R = (R_t)$  taking value 1 at  $t$  if  $X_t$  is observed and 0 otherwise; this a generalization of the response indicator variable introduced by Rubin (27). This unifies different concepts of censoring and observation of longitudinal data. Particular cases are:

- (i) right-censored survival data: case where  $X$  is a 0 – 1 counting process and  $R_t = 1_{t \leq C}$ , where  $C$  is a censoring variable;
- (ii) left-censored survival data: case where  $X$  is a 0 – 1 counting process and  $R_t = 1$  if  $t \geq C$ , 0 otherwise;
- (iii) interval-censored survival data : case where  $X$  is a 0 – 1 counting process and  $R_t = 1$  if  $t \in \{V_1, V_2, \dots, V_m\}$ , 0 otherwise.
- (iv) Repeated measurements: case where  $X$  has a continuous state space and  $R_t = 1$  if  $t \in \{V_1, V_2, \dots, V_m\}$ , 0 otherwise.

Note that  $C$  in cases (i) and (ii), and  $V_j, j = 1, \dots, m$  in (iii) and (iv) are random variables. Cases (i) and (ii) illustrate a situation where  $R$  is either right- or left-continuous at each jump time and correspond to observation in continuous time on some windows (see Figure 1); case (iii) and (iv) correspond to observations in discrete time. In the latter cases  $R_t = 1$  only on a finite or denumerable set of the half line  $[0, +\infty[$ . The TCMP allows to represent a large




FIG 1. *right-censoring*




FIG 2. *Continuous monitoring followed by discrete-time visits*

number of non-standard observation schemes. For instance the subjects can be observed on windows separated by periods where no observation are taken. In most applications the process  $X$  will be either observed continuously on some periods or observed only at discrete time points, but the TCMP can represent

a mixing of the two types of observation for the same process: for instance a subject could be observed in continuous time when he is at hospital and at discrete times when he has left hospital (see Figure 2). Also the process  $X$  is not necessarily a 0-1 counting process but may be for instance a more general counting process, allowing recurrent events, or a process with continuous state-space like a diffusion process.

**Example 1** (Vital status, AIDS status). *The vital status, represented by a counting process, is in general observed in continuous time up to a censoring date (the end of the study). This is the conventional right-censoring case, case (i). The status with respect of a disease is often observed in discrete time. The AIDS status for an HIV infected person can be represented by a counting process  $X$ . Assume that the AIDS status is observed at visits at the hospital at times  $V_1, V_2, \dots, V_m$ . This is a priori case (iii), that is,  $R_t = 1$  if  $t \in \{V_1, V_2, \dots, V_m\}$ , 0 otherwise.*

That the TCMP includes censoring or coarsening models for random variables is obvious from the fact that to each random variable  $Y$  we can associate a counting process  $X_t = 1_{\{Y \geq t\}}$ . We can define a coarsening model for a random variable  $Y$  by partitions  $\mathcal{P}$  defined by intervals  $A_j$  and indicator  $\nu_j$  which take values 1 or 0: if  $\nu_j = 1$ ,  $Y$  is exactly observed on  $\{Y \in A_j\}$ , if  $\nu_j = 0$  it is only observed that  $Y$  falls in  $A_j$ . This is equivalent to the TCMP for  $X_t$  defined by  $R_t = \sum \nu_j 1_{A_j}(t)$ . In order to construct a random mechanism we can take a random partition, for instance determined by a set of random variables.

The model can be extended to the case of multivariate processes, as may be required by the observation of several processes on the same “subject”. So, we may consider that  $X_t$  takes values in  $\mathbb{R}^d, d \geq 1$  and we may also consider a multidimensional response process  $(R_t)$ .

**Example 2** (Joint vital and AIDS statuses). *Consider a joint model of AIDS and vital status. The process  $X = (X_1, X_2)$  is a bivariate counting process,  $X_1$  representing AIDS and  $X_2$  representing vital statuses respectively. The observation scheme can be represented by the bivariate response indicator process*



$R = (R_1, R_2)$ , with  $R_{1t} = 1_{t \leq C}$  and  $R_{2t} = 1$  if  $t \in \{V_1, V_2, \dots, V_m\}$ , 0 otherwise.

**Remark.** The  $(R_t)$  process is a generalization of the response indicator variables of Rubin (27); van der Laan and Robins (29) used similar response indicator processes for the case of counting processes in the right-censored and interval-censored cases. It is different from the filtering process proposed in Andersen et al. (2) (section III.4): for instance neither left-censoring nor discrete observation times can be treated with Andersen et al. filtering process. See also Arjas, Haara and Norros (4) and Arjas and Haara (3) for a detailed treatment of the filtering problem in the framework of marked point processes. The  $(R_t)$  process can also be considered as a special case of the auxiliary random variable  $G$  used by Heitjan and Rubin (14) or Jacobsen and Keiding (18) if we interpret this random variable as a random element in a Skorohod space.

### 3.2. Extension to vertical coarsening

In addition to time-coarsening (which could be seen as “horizontal coarsening”), “vertical coarsening” occurs if for  $t$  such that  $R_t = 1$ ,  $X_t$  is not completely observed, but a only coarsened version  $Y_t = g(X_t, \varepsilon_t)$  where  $\varepsilon_t$  is observed; here it is assumed for simplicity that the processes  $(\varepsilon_t)$  and  $(X_t)$  are independent as well as the processes  $(\varepsilon_t)$  and  $(R_t)$ . Two cases, which are important in applications, are described: “fixed vertical coarsening” and noisy observation.

Fixed vertical coarsening is as follows: when  $R_t = 1$ ,  $X_t$  is not completely observed, but according to a fixed m.l.i.d.. This would apply to left-censored observation of a biological marker due to a detection limit (such as HIV-RNA), as exemplified in section 7.2.3. In that case  $(\varepsilon_t) = 0$  and  $Y_t = (1_{X_t > \eta}, \tilde{X}_t)$ , where  $\tilde{X}_t = \max(X_t, \eta)$ .

It often occurs that only noisy observations are available. When  $X_t$  is quantitative (i.d. can take any real value, case (iv) of section 3.1), a conventional model is that the observation that we make at  $t \in \{V_1, V_2, \dots, V_m\}$  is noisy: typically we observe  $Y(t) = X_t + \varepsilon_t$ , with  $\varepsilon_t$  representing a measurement error independent of all the other variables of the problem. This could also be applied

in the case where  $X_t$  is binary but there are classification errors.

**Example 3** (CD4 counts).  $X$  represents the concentration of  $T$  CD4 lymphocytes in an HIV infected person, and we have noisy observations of it  $Y_j = X_{V_j} + \varepsilon_j$ ,  $j = 1, \dots, m$ . These noisy discrete-time observations are called the CD4 counts; see section 7.2.2.

In the following the theory will be developed for the TCMP without vertical censoring; see Remark 3 of section 6 for the extension of the results to that case.

## 4. Ignorability

### 4.1. The sigma-field representation for incomplete data

A model for the random element  $X$  is a family of measures  $\{P_\theta\}_{\theta \in \Theta}$  on a  $\sigma$ -field  $\mathcal{X}$  generated by  $X$  (for us,  $X$  will be a stochastic process:  $X_t$  takes values in  $\mathbb{R}^d$  while the path of  $X$  is an element of a Skorohod space). We will assume that the measures in the family are equivalent and take  $P_{\theta_0}$  as the reference measure. If we observe  $\mathcal{X}$  we will use the likelihood  $\mathcal{L}_{\mathcal{X}}^{\theta/\theta_0}$  for inference about  $\theta$ . We will represent the observed events by a  $\sigma$ -field  $\mathcal{O}$ . A general definition of incomplete data is:  $\mathcal{X} \not\subset \mathcal{O}$ . A simple case of incomplete data is when only a sub- $\sigma$ -field of  $\mathcal{X}$  has been observed:  $\mathcal{O} \subset \mathcal{X}$ . A particular case occurs when the m.l.i.d. can be represented by a TCMP with  $R$  a deterministic function. We shall denote by  $\{R = r\}$  the event  $\{R_t = r_t, t \geq 0\}$  where  $r_t$  is a particular path (an element of the Skorohod space for instance). If  $R$  is deterministic there is a value (a path)  $r$  such that  $\{R = r\} = \Omega$ .

**Example 4** (Fixed right-censoring).  $R_t = 1$  for  $t < c$  and  $R_t = 0$  for  $t \geq c$ , where  $c$  is fixed.  $R$  takes the fixed value  $r$  such that  $r_t = 1_{t \leq c}$ . In such a case we have  $\mathcal{O} = \mathcal{X}^r = \sigma(X_t, t \geq 0 : r_t = 1) = \sigma(r_t X_t, t \geq 0)$ . In that case we have  $\mathcal{L}_{\mathcal{O}} = \mathcal{L}_{\mathcal{X}^r}$  which is in general relatively easy to compute.

**Example 5** (Fixed right-censoring: the survival case). If  $X$  is a counting process, Jacod's formula (16) can be used to obtain the likelihood; in the case where

$X$  is a 0 – 1 counting process with only one jump time  $T$ . The observed  $\sigma$ -field  $\mathcal{O}$  can be seen as generated by the variables  $\delta = 1_{\{T \leq c\}}$  and  $\tilde{T} = \min(T, c)$ . The likelihood takes the form

$$\frac{\alpha_\theta^\delta(\tilde{T}) \exp[-\int_0^{\tilde{T}} \alpha_\theta(u) du]}{\alpha_0^\delta(\tilde{T}) \exp[-\int_0^{\tilde{T}} \alpha_0(u) du]},$$

where  $\alpha_\theta(\cdot)$  (resp.  $\alpha_0(\cdot)$ ) are the risk functions under  $P_\theta$  (resp.  $P_0$ ).

Feigin (11) has given the likelihood for a diffusion process observed in continuous time. If a process is observed at fixed times  $v_1, \dots, v_m$  the likelihood can be computed as the likelihood for observation of the vector of random variables  $X_{v_1}, \dots, X_{v_m}$ ; if  $X$  is a 0 – 1 counting process this case has been denoted “interval censoring” and the likelihood is simple to compute (see (25), (1)); if  $X$  is a Gaussian process, the vector  $X_{v_1}, \dots, X_{v_m}$  has a normal distribution which makes the likelihood easy to compute.

If  $R$  is not fixed, the above definition of  $\mathcal{O}$  is meaningless; we must include  $R$  in the description of the problem. We shall consider a larger  $\sigma$ -field  $\mathcal{F} = \mathcal{X} \vee \mathcal{R}$ , where  $\mathcal{R}$  is the  $\sigma$ -field generated by  $R$  for right- or left-continuous processes that is:  $\mathcal{R} = \sigma(R_t, t \geq 0)$ ; if  $R$  takes value 1 at only a finite (or denumerable) number of times (corresponding to the discrete observation case) we can take  $\mathcal{R}$  as generated by the counting process counting the number of observation times. We consider that  $R$  is observed (see Remark 2) so that a representation of  $\mathcal{O}$  is  $\mathcal{O} = \sigma(R_t X_t, R_t, t \geq 0)$ . In section 6 which develops a dynamical approach to the problem, we shall define adequate filtrations; for instance if  $R$  is càdlàg (right-continuous with left-hand limits) the observed filtration  $(\mathcal{O}_t)$  will be the family of  $\sigma$ -fields  $\mathcal{O}_t = \sigma(R_u X_u, R_u, 0 \leq u \leq t)$ . We have of course  $\mathcal{O} \subset \mathcal{F}$ ; generally we have  $\mathcal{X} \not\subset \mathcal{O}$  (incomplete data) and  $\mathcal{O} \neq \mathcal{X}^r$  (the observation is not a predetermined subset of values of  $X$ ).

**Remark 1.** We might think that we could define an interesting  $\sigma$ -field by  $\sigma(R_t X_t, t \geq 0)$  which could take the role of the notation  $x_{obs}$  used in most of the literature in missing data (for instance (22)); however if  $X_t \neq 0$  for all  $t$ , the latter  $\sigma$ -field is equal to  $\mathcal{O}$ . When  $R$  is random it is not possible to disentangle

the observed part of  $X$  from  $R$ ; only the realized value of  $R$  effects a partition between  $x_{obs}$  and  $x_{mis}$ . This is in fact the meaning of  $\mathcal{X}^r$  which is the observed part of  $\mathcal{X}$  when  $R = r$ .

**Remark 2.** It is natural to say that  $R$  is observed: for each  $t$  we know whether we observe  $X_t$  or not. There is however an important case where this natural assumption does not hold: in survival analysis,  $X$  is a 0 – 1 counting process, so after a jump has been observed there is no need for observation anymore; so we may ignore whether we would have observed the process if it had been necessary. The simplest way to get out of this problem is to put  $R_t = 1$  if  $X_t$  is known. More generally assume that there is an absorbing state  $a$  such that if  $X_t = a$ ,  $X_{t+u} = a$  for  $u > 0$ ; define the  $\mathcal{O}_t$ -stopping time  $T = \inf\{t : X_t = a \text{ and } R_t = 1\}$ . By convention put  $R_{T+u} = 1, u > 0$ . This part of the law of  $R$  is anyway unidentifiable. In the remaining of the paper we will consider that  $R$  is observed.

**Remark 3.** If  $X_t$  is multivariate and its components may be differently coarsened, then  $R_t$  is multivariate and  $R_t X_t$  must be interpreted as the scalar product of the two vectors.

#### 4.2. Model, notations and observed likelihood

From now on, a model for the random element  $(X, R)$  is a family of measures  $\{P_{(\theta, \psi)}\}_{(\theta, \psi) \in \Theta \times \Psi}$  on a measurable space  $(\Omega, \mathcal{F})$ .  $X$  (resp.  $R$ ) takes values in a measurable space  $(\Xi, \xi)$  (resp.  $(\Gamma, \rho)$ ). For us  $X$  and  $R$  will be d-dimensional càdlàg stochastic processes, so  $(\Xi, \xi)$  and  $(\Gamma, \rho)$  are Skorohod spaces endowed with their Borel  $\sigma$ -fields. The parameter spaces  $\Theta$  and  $\Psi$  need not be finite dimensional. We will assume that the measures in the family are equivalent and take  $P_{(\theta_0, \psi_0)}$  as the reference measure.  $P_\theta$  is the restriction of  $P_{(\theta, \psi)}$  to  $\mathcal{X}$ : that is, the marginal probability of  $X$  does not depend on  $\psi$ . The additional parameter  $\psi$  will be considered as a nuisance parameter. We assume implicitly a "Non-Informativeness" assumption in the coarsening mechanism, which is :

$$P_{(\theta_1, \psi)}(A|\mathcal{X}) = P_{(\theta_2, \psi)}(A|\mathcal{X}), \text{ a.s., } A \in \mathcal{R} \text{ for all } \theta_1, \theta_2, \psi \quad (4.1)$$

In words, the conditional probability of  $R$  given  $X$  does not depend on  $\theta$ . This has an important consequence in terms of likelihood ratio. The latter assumption can also be written as:  $E_{(\theta_1, \psi)}(1_A | \mathcal{X}) = E_{(\theta_2, \psi)}(1_A | \mathcal{X})$ ,  $A \in \mathcal{R}$ , which remembering property ii) of the conditional likelihood is equivalent to  $E_{(\theta_0, \psi_0)}(1_A \mathcal{L}_{\mathcal{R}|\mathcal{X}}^{(\theta_1, \psi)/(\theta_0, \psi_0)} | \mathcal{X}) = E_{(\theta_0, \psi_0)}(1_A \mathcal{L}_{\mathcal{R}|\mathcal{X}}^{(\theta_2, \psi)/(\theta_0, \psi_0)} | \mathcal{X})$ . This in turn implies that  $\mathcal{L}_{\mathcal{R}|\mathcal{X}}^{(\theta_1, \psi)/(\theta_0, \psi_0)} = \mathcal{L}_{\mathcal{R}|\mathcal{X}}^{(\theta_2, \psi)/(\theta_0, \psi_0)}$  and we will denote this common value  $\mathcal{L}_{\mathcal{R}|\mathcal{X}}^{\psi/\psi_0}$ . Moreover it can be proved (and is intuitive) that  $\mathcal{L}_{\mathcal{R}|\mathcal{X}}^{\psi/\psi_0} = 1$ , a.s..

From now on, we fix our notation for  $\mathcal{O}$  and  $\mathcal{X}^r$  by  $\mathcal{O} = \sigma(R_t X_t, R_t, t \geq 0)$  and for  $r = (r_t)$  a deterministic path of  $R$ ,  $\mathcal{X}^r = \sigma(X_t, t \geq 0 : r_t = 1) = \sigma(r_t X_t, t \geq 0)$ . Note that (4.1) remains true even if  $A \in \mathcal{O}$ . The inferential likelihood is  $\mathcal{L}_{\mathcal{O}}^{(\theta, \psi)/(\theta_0, \psi_0)}$  (which is  $\mathcal{O}$ -measurable and thus can be computed from the observations) and the fundamental property yields:

$$\mathcal{L}_{\mathcal{O}}^{(\theta, \psi)/(\theta_0, \psi_0)} = E_{(\theta_0, \psi_0)}[\mathcal{L}_{\mathcal{X}, \mathcal{R}}^{(\theta, \psi)/(\theta_0, \psi_0)} | \mathcal{O}] = E_{(\theta_0, \psi_0)}[\mathcal{L}_{\mathcal{R}|\mathcal{X}}^{\psi/\psi_0} \mathcal{L}_{\mathcal{X}}^{\theta/\theta_0} | \mathcal{O}]$$

**Example 6** (Random right-censoring in the survival case). *In the survival case we can distinguish the computation of the likelihood on the observed events  $\{\delta = 1\}$  and  $\{\delta = 0\}$ . On  $\{\delta = 1\}$   $X$  is completely observed so that we can put  $\mathcal{L}_{\mathcal{X}}^{\theta/\theta_0}$  out of the conditional expectation getting  $\mathcal{L}_{\mathcal{O}}^{(\theta, \psi)/(\theta_0, \psi_0)} = E_{(\theta_0, \psi_0)}[\mathcal{L}_{\mathcal{R}|\mathcal{X}}^{\psi/\psi_0} | \mathcal{O}] \mathcal{L}_{\mathcal{X}}^{\theta/\theta_0}$ . Since  $\mathcal{L}_{\mathcal{R}|\mathcal{X}}^{\psi/\psi_0} = \frac{P_{\psi}(C > T)}{P_{\psi_0}(C > T)}$  it is known on  $\{\delta = 1\}$  and can be put out of the conditional expectation. Moreover it does not depend on the parameter of interest  $\theta$ , so we could get rid of it for inference about  $\theta$ . On  $\{\delta = 0\}$  however, the conditional expectation is  $E_{(\theta_0, \psi_0)}[\mathcal{L}_{\mathcal{R}|\mathcal{X}}^{\psi/\psi_0} \frac{f_{\theta}(T)}{f_{\theta_0}(T)} | C, T > C]$ , where  $f_{\theta}(\cdot)$  is the density function of  $T$ ; since  $X$  is not completely observed none of the terms can get out of it without additional assumptions.*

#### 4.3. Definition of ignorability

If the m.l.i.d. (represented by  $R$ ) is random it may still be tempting to ignore it, treating it as fixed, and use for inference  $\mathcal{L}_{\mathcal{X}^r}$  which is relatively easy to compute and does not depend on  $\psi$ .

**Definition 1.** *In the TCMP, the likelihood ratio ignoring the m.l.i.d. is the like-*

likelihood ratio  $\mathcal{L}_{\mathcal{X}^r}^{\theta/\theta_0}$  obtained under the assumption that  $R$  is fixed at its observed value  $r$ .

When the fixed m.l.i.d. assumption does not hold, the question arises to know in which cases (if ever)  $\mathcal{L}_{\mathcal{X}^r}^{\theta/\theta_0}$  leads to the same inference about  $\theta$  as  $\mathcal{L}_{\mathcal{O}}^{\theta, \psi_0/\theta_0, \psi_0}$ , if the true value of  $\psi$  was known to be  $\psi_0$  or  $\mathcal{L}_{\mathcal{O}}^{\theta, \psi/\theta_0, \psi_0}$  if  $\psi$  had to be estimated.

For defining ignorability we face the problem that both Radon-Nikodym derivatives and conditional expectations are defined almost surely; for some results we must restrict the theoretical framework to measures giving a non null probability to a denumerable set of trajectories of  $R$ .

**Assumption A1** There exists a denumerable set  $(r_1, r_2, \dots)$  such that  $P_\theta(R = r_i) > 0$  and  $\sum_i P_\theta(R = r_i) = 1$ , for all  $\theta$ .

This is a theoretical limitation but this has no impact on application since in practice the times are always rounded. Some of the results below will need this restriction, other will not.

**Definition 2** (Ignorability). *The m.l.i.d. will be called **ignorable on  $r$**  if  $\mathcal{L}_{\mathcal{O}}^{(\theta, \psi)/(\theta_0, \psi_0)} = U \mathcal{L}_{\mathcal{X}^r}^{\theta/\theta_0}$  a.s. on  $\{R = r\}$  for all  $\theta$ , whatever  $\psi$  and  $\psi_0$ , where  $U$  is random variable not depending on  $\theta$ . It will be called **ignorable** if assumption A1 holds and the m.l.i.d. is ignorable on  $r$  for all values of  $r$ .*

**Example 7** (Random right-censoring in the survival case). *In the survival case on  $\{\delta = 0\}$  and if we observe  $C = c$  the likelihood ignoring the m.l.i.d. is  $\mathcal{L}_{\mathcal{X}^r}^{\theta/\theta_0} = S_\theta(\tilde{T})/S_{\theta_0}(\tilde{T})$ . Thus, ignorability holds if  $E_{(\theta_0, \psi_0)}[\mathcal{L}_{\mathcal{R}|\mathcal{X}}^{\psi/\psi_0} \frac{f_\theta(T)}{f_{\theta_0}(T)} | C, T > C] = U \frac{S_\theta(\tilde{T})}{S_{\theta_0}(\tilde{T})}$ , where  $S_\theta(\cdot)$  is the survival function of the distribution of  $T$  under  $P_\theta$ .*

**Remark.** If the m.l.i.d. is ignorable it is then obvious that  $\mathcal{L}_{\mathcal{X}^r}^{\theta/\theta_0}$  and  $\mathcal{L}_{\mathcal{O}}^{(\theta, \psi_0)/(\theta_0, \psi_0)}$  lead to the same inference about  $\theta$ .

## 5. Static conditions of ignorability for the TCMP

We give a first fact which does not seem to have been noted previously in a general context.

**Fact.** Ignorability on  $\{R_t = 1, t \geq 0\}$  always holds.

**Proof.** On the event  $\{R_t = 1, t \geq 0\}$ ,  $X$  and  $R$  are observed so that  $\mathcal{F} = \mathcal{O}$ ; thus from Lemma 1.1 we have  $\mathcal{L}_{\mathcal{O}}^{(\theta, \psi)/(\theta_0, \psi_0)} = \mathbb{E}_{(\theta_0, \psi_0)}(\mathcal{L}_{\mathcal{F}}^{(\theta, \psi)/(\theta_0, \psi_0)} | \mathcal{O}) = \mathbb{E}_{(\theta_0, \psi_0)}(\mathcal{L}_{\mathcal{F}} | \mathcal{F}) = \mathcal{L}_{\mathcal{F}}^{(\theta, \psi)/(\theta_0, \psi_0)} = \mathcal{L}_{\mathcal{R}|\mathcal{X}}^{\psi/\psi_0} \mathcal{L}_{\mathcal{X}}^{\theta/\theta_0}$  (the last equality comes from property iii) of the likelihood); for  $\psi = \psi_0$  we retrieve the likelihood  $\mathcal{L}_{\mathcal{X}}^{\theta/\theta_0}$  for the complete observation of  $X$  and ignorability holds on  $\{R_t = 1, t \geq 0\}$ .  $\square$

We shall now study “static” conditions of ignorability, in contrast with the “dynamic” conditions of the next section. Gill & al (12) have introduced two conditions of ignorability: CAR(REL) (Relative Coarsening At Random) and CAR(ABS) (Absoluter Coarsening At Random); these conditions were further studied by Nielsen (24). We give two conditions of ignorability in the TCMP framework. The first one is an adaption of CAR(REL); the second one is stronger but original and doesn’t imply CAR(ABS).

**Definition 3** (CAR(TCMP)). *We will say that CAR(TCMP) holds for the couple  $(X, R)$  if  $\mathcal{L}_{\mathcal{R}|\mathcal{X}}^{\psi/\psi_0}$  is  $\mathcal{O}$ -measurable for all  $(\theta, \psi)$ .*

**Example 8** ( $R$  and  $X$  independent). *If  $R$  and  $X$  are independent  $\mathcal{L}_{\mathcal{R}|\mathcal{X}}^{\psi/\psi_0} = \mathcal{L}_{\mathcal{R}}^{\psi/\psi_0}$ ; since  $R$  is observed  $\mathcal{L}_{\mathcal{R}}^{\psi/\psi_0}$  is  $\mathcal{O}$ -measurable.*

With our notations the condition CAR(REL) is “there exists a version of  $\mathcal{L}_{\mathcal{O}|\mathcal{X}}^{\psi/\psi_0}(o, x)$  such that the mapping  $x \rightarrow \mathcal{L}_{\mathcal{O}|\mathcal{X}}^{\psi/\psi_0}(o, x)$  is constant for all  $x$  compatible with  $o$ ”, where  $o$  is an elementary event of  $\mathcal{O}$ , i.e.  $o = (r, rx)$ . First note that with our assumptions,  $x$  is compatible with  $o = (r, y)$  if  $rx = y$  and second note that  $\mathcal{L}_{\mathcal{R}|\mathcal{X}}(r; x) = \mathcal{L}_{\mathcal{O}|\mathcal{X}}(o; x)$  if  $o = (r, rx)$  (by property vi) and because  $\mathcal{R} \vee \mathcal{X} = \mathcal{O} \vee \mathcal{X}$ . So in the TCMP setting the condition CAR(REL) becomes “there exists a version of  $\mathcal{L}_{\mathcal{R}|\mathcal{X}}^{\psi/\psi_0}(o, x)$  such that : for all  $r$  and  $(x, x')$  verifying  $rx = rx'$  then  $\mathcal{L}_{\mathcal{R}|\mathcal{X}}^{\psi/\psi_0}(r, x) = \mathcal{L}_{\mathcal{R}|\mathcal{X}}^{\psi/\psi_0}(r, x')$ ”.

**Theorem 5.1.** *CAR(TCMP) is equivalent to CAR(REL) in the TCMP setting.*

**Proof.** If CAR(REL) is true then for all  $(r, x)$ ,  $\mathcal{L}_{\mathcal{R}|\mathcal{X}}^{\psi/\psi_0}(r; x) = \mathcal{L}_{\mathcal{R}|\mathcal{X}}^{\psi/\psi_0}(r; rx)$  (by taking  $x' = rx$  in the above formula) and then  $\mathcal{L}_{\mathcal{R}|\mathcal{X}}^{\psi/\psi_0}$  is a function of  $(R, RX)$  and so  $\mathcal{L}_{\mathcal{R}|\mathcal{X}}^{\psi/\psi_0}$  is  $\mathcal{O}$ -measurable that is CAR(TCMP). Conversely, if  $\mathcal{L}_{\mathcal{R}|\mathcal{X}}^{\psi/\psi_0}$  is

$\mathcal{O}$ -measurable, there is a version of it which is constant on the atoms of  $\mathcal{O}$  i.e. on the set of the form  $O_{r,y} = \{(r,x) \text{ such that } x \text{ verifies } rx = y\}$  and we have  $\text{CAR}(\text{REL})$ .  $\square$

The next theorem shows that  $\text{CAR}(\text{TCMP})$  implies a factorization of the likelihood in two parts : one which depends on  $\psi$  and the second one on  $\theta$ ; this may be called “weak ignorability”.

**Theorem 5.2** (Factorization). *If the couple  $(R, X)$  satisfies  $\text{CAR}(\text{TCMP})$  then we have  $\mathcal{L}_{\mathcal{O}}^{(\theta,\psi)/(\theta_0,\psi_0)} = \mathcal{L}_{\mathcal{R}|\mathcal{X}}^{\psi/\psi_0} \mathbb{E}_{(\theta_0,\psi_0)}(\mathcal{L}_{\mathcal{X}}^{\theta/\theta_0}|\mathcal{O})$  and  $\mathbb{E}_{(\theta_0,\psi_0)}(\mathcal{L}_{\mathcal{X}}^{\theta/\theta_0}|\mathcal{O})$  does not depend on  $\psi_0$ .*

**Proof.** A proof can be obtained using the previous theorem and the fact that  $\text{CAR}(\text{REL})$  has been proved to imply a similar factorization theorem. However a direct proof is quite simple in our formalism. From the decomposition formula we have  $\mathcal{L}_{\mathcal{F}}^{(\theta,\psi)/(\theta_0,\psi_0)} = \mathcal{L}_{\mathcal{R}|\mathcal{X}}^{\psi/\psi_0} \mathcal{L}_{\mathcal{X}}^{\theta/\theta_0}$  and we can pull  $\mathcal{L}_{\mathcal{R}|\mathcal{X}}^{\psi/\psi_0}$  out of the conditional expectation using  $\text{CAR}(\text{TCMP})$  thus obtaining  $\mathcal{L}_{\mathcal{O}}^{(\theta,\psi)/(\theta_0,\psi_0)} = \mathcal{L}_{\mathcal{R}|\mathcal{X}}^{\psi/\psi_0} \mathbb{E}_{(\theta_0,\psi_0)}(\mathcal{L}_{\mathcal{X}}^{\theta/\theta_0}|\mathcal{O})$ . It only remains to prove that the last term does not depend on  $\psi_0$ . We have

$$\begin{aligned} \mathcal{L}_{\mathcal{O}}^{(\theta,\psi)/(\theta_0,\psi_0)} &= \mathcal{L}_{\mathcal{O}}^{(\theta,\psi)/(\theta_0,\psi_1)} \mathcal{L}_{\mathcal{O}}^{(\theta_0,\psi_1)/(\theta_0,\psi_0)} \\ &= \mathcal{L}_{\mathcal{R}|\mathcal{X}}^{\psi/\psi_1} \mathbb{E}_{(\theta_0,\psi_1)}[\mathcal{L}_{\mathcal{X}}^{\theta/\theta_0}|\mathcal{O}] \mathcal{L}_{\mathcal{R}|\mathcal{X}}^{\psi_1/\psi_0} \mathbb{E}_{(\theta_0,\psi_0)}[\mathcal{L}_{\mathcal{X}}^{\theta_0/\theta_0}|\mathcal{O}] \\ &= \mathcal{L}_{\mathcal{R}|\mathcal{X}}^{\psi/\psi_0} \mathbb{E}_{(\theta_0,\psi_1)}[\mathcal{L}_{\mathcal{X}}^{\theta/\theta_0}|\mathcal{O}] \end{aligned}$$

$\square$

As a result, we can use  $\mathbb{E}_{(\theta_0,\psi_0)}(\mathcal{L}_{\mathcal{X}}^{\theta/\theta_0}|\mathcal{O})$  for inference on  $\theta$ , getting rid of the nuisance parameter  $\psi$ .

**Example 9** (Random right-censoring in the survival case). *Continuing Example 7 we have on  $\delta = 0$ , the likelihood if  $\text{CAR}(\text{TCMP})$  holds can be written:  $\mathbb{E}_{(\theta_0,\psi_0)}[\mathcal{L}_{\mathcal{R}|\mathcal{X}}^{\psi/\psi_0} \frac{f_{\theta}(T)}{f_0(T)}|C, T > C] = \mathcal{L}_{\mathcal{R}|\mathcal{X}}^{\psi/\psi_0} \mathbb{E}_{(\theta_0,\psi_0)}[\frac{f_{\theta}(T)}{f_0(T)}|C, T > C] = \mathcal{L}_{\mathcal{R}|\mathcal{X}}^{\psi/\psi_0} \frac{S_{\theta}(\tilde{T})}{S_{\theta_0}(\tilde{T})}$ . Thus the observed likelihood is proportional to the likelihood ignoring the m.l.i.d.; it is strictly equal if we take  $\psi = \psi_0$ , our definition of ignorability.*

The following condition allows obtaining a more precise result locally.



**Definition 4** (CAR(TCMP)-loc). *We will call CAR(TCMP)-loc on  $r$  the condition:  $\mathcal{L}_{\mathcal{R}|\mathcal{X}}^{(\theta,\psi)/(\theta_0,\psi_0)} = \mathcal{L}_{\mathcal{R}|\mathcal{X}^r}^{(\theta,\psi)/(\theta_0,\psi_0)}$  a.s. on  $\{R = r\}$ .*

We have:

**Theorem 5.3.** *CAR(TCMP)-loc on  $r$  implies ignorability on  $r$ .*

Details about this concept and proof of Theorem A are given in Appendix 1.

## 6. Dynamical conditions of ignorability

Assume that we can represent  $(R_t)$  by a marked point process  $(N_t)$ ; if the component  $R_h$  of  $R$  takes value 1 at isolated points  $V_{h,1}, V_{h,2}, \dots$  (discrete-time observations), then its associated counting process  $N_{ht}^d$  is defined by  $N_{ht}^d = \sum_{j=1}^{\infty} 1_{\{V_{h,j} \leq t\}}$ . If the component  $R_{h,t}$  is càdlàg, it can be written as  $R_{h,t} = \sum_{k=0}^{\infty} 1_{[V_{h,2k}, V_{h,2k+1}[}$  and its associate counting process  $N_{ht}^c = \sum_{j=0}^{\infty} 1_{\{V_{h,j} \leq t\}}$ . If the component  $R_h$  of  $R$  is a mixing of the two kinds of observation, (discrete- and continuous-time) then at points of discontinuities we have to add a mark which indicates if the jump affects the discrete part of  $R_h$  represented by  $N_{ht}^d$  or the càdlàg one represented by  $N_{ht}^c$ . In all cases  $R$  can be represented by a marked point process  $(N_t)$ . Denote by  $(\mathcal{N}_t)$  the self-exciting filtration of  $(N_t)$ . We define the filtration  $(\mathcal{O}_t)$  as the family of  $\sigma$ -fields  $\mathcal{O}_t = \sigma(N_u, R_u X_u, 0 \leq u \leq t)$ . Let us call  $\Lambda^{\mathcal{O},N} = (\Lambda_t^{\mathcal{O},N})$  and  $\Lambda^{\mathcal{F}^*,N} = (\Lambda_t^{\mathcal{F}^*,N})$  the compensators of  $N$  in the filtrations  $(\mathcal{O}_t)$  and  $(\mathcal{F}_t^*)$  respectively and for probability  $P_{\theta,\psi}$ , where  $(\mathcal{F}_t^*)$  is the family of  $\sigma$ -fields  $\mathcal{F}_t^* = \mathcal{X} \vee \mathcal{O}_t, t \geq 0$ . The compensators generally depend on  $(\theta, \psi)$  but we omit this for notational simplicity; the compensators for  $P_{\theta_0,\psi_0}$  will be denoted  $\Lambda_0$ . In the following we will assume that there exists a fixed time  $\tau$  such that  $R_{\tau+u} = R_\tau$  and that there is no explosion of the process on  $[0, \tau]$  (i.e.  $\sum_{t \leq \tau} \Delta R_t < \infty$ ) (thus  $\Lambda_{\tau+u}^N = \Lambda_\tau^N, u \geq 0$ , for any filtration).

**Definition 5** (CAR(DYN)). *We will denote CAR(DYN) the condition:*

$\forall (\theta, \psi)$  we have under  $P_{(\theta,\psi)}$  :  $(\Lambda_t^{\mathcal{O},N}) = (\Lambda_t^{\mathcal{F}^*,N})$ , (up to indistinguishability).

**Remark 1.** This is an absolute condition in the sense of Gill et al. (12) and Nielsen (24) since the CAR(DYN) criterion is defined for each probability separately while the definition of CAR(TCMP) bears on Radon-Nikodym derivatives.

**Remark 2.** For this condition to lead really to ignorability the model for the mechanism leading to missing data must be well specified. That is, the condition  $(\Lambda_t^{\mathcal{O},N}) = (\Lambda_t^{\mathcal{F}^*,N})$  must hold under the true probability.

**Example 10** (Discrete visit times). *Suppose that  $N$  represents visit times. This process has the compensators  $(\Lambda_t^{\mathcal{O},N})$  and  $(\Lambda_t^{\mathcal{F}^*,N})$  in the filtrations  $(\mathcal{O}_t)$  and  $(\mathcal{F}_t^*)$  respectively. CAR(DYN) holds if these compensators are equal. In words, the next visit time depends only on what has been observed at previous visit times. For instance if we are studying the concentration of CD4 T lymphocytes, this is the case if the next visit time is fixed as a function of the observed CD4 counts (see section 7.2.2).*

It appears that CAR(DYN) is stronger than CAR(TCMP).

**Theorem 6.1.** *CAR(DYN) implies CAR(TCMP).*

**Proof.** Let us write the likelihood for  $N$  and  $X$ . The filtration  $(\mathcal{F}_t^*)$  is the self-generated filtration for  $N$  when  $\mathcal{F}_0^* = \mathcal{X}$ . Thus we have using Jacod's formula  $\mathcal{L}_{\mathcal{F}^*} = \mathcal{L}_{\mathcal{X}}^{\theta/\theta_0} \phi(\Lambda_u^{\mathcal{F}^*,N}, \Lambda_{0u}^{\mathcal{F}^*,N}, N_u, u \geq 0)$ , where

$$\phi(\Lambda, \Lambda_0, N) = \phi(\Lambda_u, \Lambda_{0u}, N_u, 0 \leq u) = \prod_{t \geq 0} \prod_h \left( \frac{d\Lambda_h}{d\Lambda_{0h} t} \right)^{\Delta N_{ht}} \frac{\prod_{t \geq 0: \Delta N_{.t} \neq 1} (1 - d\Lambda_{.t})}{\prod_{t \geq 0: \Delta N_{.t} \neq 1} (1 - d\Lambda_{0.t})},$$

where  $N_{.} = \sum_h N_h$  and  $\Lambda_{.} = \sum_h \Lambda_h$ . Identifying with the decomposition  $\mathcal{L}_{\mathcal{F}^*} = \mathcal{L}_{\mathcal{X}}^{\theta/\theta_0} \mathcal{L}_{\mathcal{R}|\mathcal{X}}^{\psi/\psi_0}$  we find that  $\mathcal{L}_{\mathcal{R}|\mathcal{X}}^{\psi/\psi_0} = \phi(\Lambda^{\mathcal{F}^*,N}, \Lambda_0^{\mathcal{F}^*,N}, N)$ . With CAR(DYN) this is equal to  $\phi(\Lambda^{\mathcal{O},N}, \Lambda_0^{\mathcal{O},N}, N)$  which is  $\mathcal{O}$ -measurable, and thus CAR(TCMP) holds.  $\square$

**Theorem 6.2.** *CAR(DYN) implies ignorability.*

**Proof.** We have to prove that CAR(DYN) implies CAR(TCMP)-loc for all  $r$ ; then using Theorem A we will have that CAR(DYN) implies ignorability.

Let us define  $\mathcal{F}_t^r = \mathcal{X}^r \vee \mathcal{N}_t$  and  $\mathcal{F}^r = \mathcal{X}^r \vee \mathcal{N}$  then for all  $t \in [0, \tau]$ ,  $\mathcal{F}_t^r \subseteq \mathcal{F}_t^*$ . As seen in the previous proof,  $\mathcal{L}_{\mathcal{R}|\mathcal{X}}^{(\theta, \psi)/(\theta_0, \psi_0)} = \phi(\Lambda^{\mathcal{F}^*, N}, \Lambda_0^{\mathcal{F}^*, N}, N)$  and  $\mathcal{L}_{\mathcal{R}|\mathcal{X}^r}^{(\theta, \psi)/(\theta_0, \psi_0)} = \phi(\Lambda^{\mathcal{F}^r, N}, \Lambda_0^{\mathcal{F}^r, N}, N)$ . So the proof will be complete if  $\forall t \in [0, t]$ ,  $\Lambda_t^{\mathcal{F}^r, N} 1_{\{R=r\}} = \Lambda_t^{\mathcal{F}^*, N} 1_{\{R=r\}}$ . We get

$$\Lambda_t^{\mathcal{F}^r, N} 1_{\{R=r\}} = E[\Lambda_t^{\mathcal{F}^*, N} | \mathcal{F}_t^r] 1_{\{R=r\}} \quad (6.1)$$

$$= E[\Lambda_t^{\mathcal{O}, N} 1_{\{R_{\cdot \wedge t} = r_{\cdot \wedge t}\}} | \mathcal{F}_t^r] 1_{\{R=r\}} \quad (6.2)$$

$$= \Lambda_t^{\mathcal{O}, N} 1_{\{R=r\}} \quad (6.3)$$

(6.1) is due to the innovation theorem, (6.2) is due to CAR(DYN) and the fact that  $\{R_{\cdot \wedge t} = r_{\cdot \wedge t}\} \subset \{R=r\}$  and  $\{R_{\cdot \wedge t} = r_{\cdot \wedge t}\} \in \mathcal{O}_t \cap \mathcal{F}_t^r$ . At last the local inclusion  $\{R_{\cdot \wedge t} = r_{\cdot \wedge t}\} \cap \mathcal{O}_t \subseteq \{R_{\cdot \wedge t} = r_{\cdot \wedge t}\} \cap \mathcal{F}_t^r$  implies the  $\mathcal{F}_t^r$ -measurability of  $\Lambda_t^{\mathcal{O}, N} 1_{\{R_{\cdot \wedge t} = r_{\cdot \wedge t}\}}$  and (6.3). Using again CAR(DYN), we have the desired equality.  $\square$

**Corollary 6.1.** *If  $R$  is  $\mathcal{O}_t$ -predictable, CAR(DYN) holds and thus ignorability holds.*

**Proof.** The associated counting process  $N$  is itself  $\mathcal{F}_t^*$ -predictable. Thus the Doob-Meyer decomposition is  $N_t = N_t + 0$  in both  $(\mathcal{O}_t)$  and  $(\mathcal{F}_t^*)$ . It follows that  $\Lambda^{\mathcal{F}^*, N} = \Lambda^{\mathcal{O}, N} = N$  which is CAR(DYN). By Theorem 6.2 and A ignorability holds.  $\square$

In the framework of counting processes, Andersen et al. (2) have proposed a criterion of independent right-censoring. We adapt their criterion to right-continuous censoring processes and we restrict to the univariate case for simplicity. In that case  $R_t = 1_{\{t < C\}}$  where  $C$  is a censoring variable and  $N_t = 1 - R_t$ . Let  $X$  be a counting process and  $\Lambda^{\mathcal{X}, X}$  its compensator in the self-generated filtration  $(\mathcal{X}_t)$ . Let  $(\mathcal{F}_t)$  the filtration generated by both  $X$  and  $N$ . We have independent right-censoring if the compensator of  $X$  is the same in the filtration including information on the censoring that is:  $\Lambda^{\mathcal{F}, X} = \Lambda^{\mathcal{X}, X}$ . In the context where both the independent censoring and CAR(DYN) apply, the following theorem says that they are equivalent.

**Theorem 6.3** (Equivalence of independent censoring and CAR(DYN)). *Let  $X$  be a counting process which admits a càg (left-continuous) intensity  $\lambda^{\mathcal{X}, X}$  in the self-generated filtration  $(\mathcal{X}_t)$ . Consider a right-continuous right-censoring process of  $X$  satisfying CAR(DYN); then this is an independent censoring. Inversely, independent censoring implies CAR(DYN).*

The proof is given in Appendix 2.

In some situations it is natural to consider response indicator processes which are left-continuous; case II right-censoring is an example. This is developed in Appendix 3.

**Remark 3.** All the theoretical results can be extended to TCMP with vertical coarsening (section 3.2) by replacing  $X_t$  by  $Y_t = g(X_t, \varepsilon_t)$ . In particular the observed  $\sigma$ -field at time  $t$  becomes  $\mathcal{O}_t = \sigma(R_t, R_t Y_t)$ .

## 7. Applications

### 7.1. Right-censoring of counting processes with time-dependent covariable

We consider the modelling of independent counting processes  $W^i = (W_t^i)$  with possibly time-dependent explanatory variables  $Z^i = (Z_t^i)$ . We observe  $X^i = (W^i, Z^i)$ ,  $i = 1, \dots, n$  through a mechanism specified by  $R = (R^{W^1}, \dots, R^{W^n}; R^{Z^1}, \dots, R^{Z^n})$ . The  $(Z_t^i)$  are supposed to be completely observed so that  $R_t^{Z^i} = 1$  for all  $t$ . We consider a family of probability measures  $\{P^{\theta, \gamma, \psi}\}$ , where  $\theta$  is the parameter of interest which parameterizes the dynamics of  $(W_t^i)$  given the value of the explanatory variables, that is, the intensity of  $(W_t^i)$  in the filtration  $(\mathcal{X}_t^i)$  (defined as the family of  $\sigma$ -fields  $\mathcal{X}_t^i = \mathcal{W}_t^i \vee \mathcal{Z}_t^i$ ) depends on  $\theta$  only. Let  $\gamma$  parameterize the marginal law of  $(Z_t^i)$ . More precisely, it is assumed that the compensator of  $Z_t^i$  is the same in the filtration  $(\mathcal{X}_t^i)$  and in the filtration  $(\mathcal{Z}_t^i)$ ; in other word it is assumed that  $Z$  is WCLI of  $W$ ; see Commenges and Gégout-Petit (7). If this is not the case, then we must resort to joint modelling. Consider that assumption A1 holds; for right-censoring this means that the set of times at which observation

may be stopped is denumerable; this is not a limitation in practice: for instance in an epidemiological cohort we may say that observation may be stopped each day at a fixed hour (we generally do not have a precision better than one day). If ignorability holds for  $(X, R)$ , we have on  $\{R = r\}$ ,  $\mathcal{L}_{\mathcal{X}^r}^{\theta, \gamma / \theta_0, \gamma_0} = \mathcal{L}_{\mathcal{O}}^{\theta, \gamma, \psi_0 / \theta_0, \gamma_0, \psi_0}$ . But we have also  $\mathcal{L}_{\mathcal{X}^r}^{\theta, \gamma / \theta_0, \gamma_0} = \mathcal{L}_{\mathcal{W}^r | Z^r}^{\theta / \theta_0} \mathcal{L}_{Z^r}^{\gamma / \gamma_0}$ , so that in terms of inference about  $\theta$  we only need to compute  $\mathcal{L}_{\mathcal{W}^r | Z^r}^{\theta / \theta_0}$ . The conclusion is that although we can use the conditional likelihood of  $W$  given  $Z$ , we may still consider the ignorability condition for  $(X, R)$ , that is, the response indicator process may depend on both observed  $W$  and  $Z$ . This discussion is related to the so-called “reduced model” introduced in (8).

We give a particular example of artificial right censoring of 0 – 1 counting processes which is compatible with ignorability (assuming to simplify that there is no other source of random censoring). Consider a study where  $n$  subjects are potentially followed-up until a time  $t^*$ ; it is assumed that the processes  $(X^i, Z^i)$  are independent and identically distributed. Suppose that at a given time  $t_1$  we make an analysis of the data. Using for instance maximum likelihood estimators in a parametric model we can construct an estimator  $\hat{\theta}_1$  which by definition is  $\mathcal{O}_{t_1}$ -measurable. Let us suppose for simplicity that  $Z_t, t > t_1$  is known at  $t_1$ ; it is then possible to compute an estimator of the probability that subject  $i$  experiences the event before the end of the study  $P_{\hat{\theta}}(W_{t^*}^i = 1 | \mathcal{W}_{t_1}, \mathcal{Z}_{t^*}^i)$ , where  $\mathcal{W}_t$  and  $\mathcal{Z}_t$  denote  $\sigma$ -fields generated by the  $n$  processes up to time  $t$ . For reducing the cost of the study without reducing too much its power, we may decide to follow after  $t_1$  only the subjects for whom  $P_{\hat{\theta}}(W_{t^*}^i = 1 | \mathcal{W}_{t_1}^i, \mathcal{Z}_{t^*}^i) \geq c$ , for some chosen  $c$ . In the TCMP this means putting  $R_t^{W^i} = 0$  for  $t > t_1$  for those subjects with an estimated probability below  $c$ . It is clear that  $R$  is  $\mathcal{O}_t$ -predictable which from Corollary 6.1 implies ignorability on all  $r$  because A1 holds. Note that ignorability is not dependent of the good specification of the model used at  $t_1$ ; of course the validity of the final analysis will depend on the good specification of the model used for it.

## 7.2. Longitudinal markers (continuous state-space processes)

### 7.2.1. Missing data at fixed observation times

This is the classical set-up of “repeated measurements” or “longitudinal data” (in a narrow sense). We consider independent continuous state-space processes  $(X_t^i)$ ,  $i = 1, \dots, n$ ; for simplicity we do not consider covariates. Here  $(X_t^i)$  is planned to be observed at  $v_j^i$ ,  $j = 1, \dots, m$  but there may be missing data. The response indicator processes for  $(X_t^i)$  can be written:  $R_t^{X^i} = \sum_{j=1}^m 1_{\{t=v_j^i\}} S_j^i$ , where  $S_j^i$  are binary variables. The jump at  $v_j^i-$  of the compensator of the counting process associated to  $R^i$  in a filtration  $(\mathcal{G}_t)$  is  $P(S_j^i = 1 | \mathcal{G}_{v_j^i-})$ . CAR(DYN) can thus be expressed as:  $P(S_j^i = 1 | S_1^i, \dots, S_{v_{j-1}^i}^i, \mathcal{X}) = P(S_j^i = 1 | S_1^i, \dots, S_{v_{j-1}^i}^i, S_1^i X_1^i, \dots, S_{v_{j-1}^i}^i X_{v_{j-1}^i}^i)$ , which implies (by taking conditional expectation) that

$$P(S_j^i = 1 | S_1^i, \dots, S_{v_{j-1}^i}^i, \mathcal{X}_{v_j^i-}) = P(S_j^i = 1 | S_1^i, \dots, S_{v_{j-1}^i}^i, S_1^i X_1^i, \dots, S_{v_{j-1}^i}^i X_{v_{j-1}^i}^i);$$

this can intuitively be interpreted in saying that the missing data mechanism may depend (only) on the observed  $W^i$  and on the  $S^i$  up just before time  $v_j^i$ ; this case could be treated with the conventional MAR concept.

### 7.2.2. Random observation times

We may consider determining observation times in order to reduce the costs of a study or to improve the monitoring of patients. Consider the case of a study of the evolution of the concentration of CD4+ T lymphocytes in HIV infected patients, represented by  $(X_t^i)$  for subject  $i$ . Following example 3, the observations are the CD4 counts, that is noisy measurements of  $(X_t^i)$  at discrete visit times:  $Y_{ij} = X_{V_j^i}^i + \varepsilon_{ij}$ . The time of the next visit,  $V_{j+1}^i$  may depend on CD4 count at  $V_j^i$  according to the procedure:  $V_{j+1}^i = V_j^i + g(X_{V_j^i}^i)$ , where  $g(\cdot)$  is a known function.. For instance we could decide to see a patient with a delay of three months if  $CD4 \geq 500$ , two months if  $200 \leq CD4 < 500$  and one month if  $CD4 < 200$ . It is clear that CAR(DYN) would hold in this instance. On the contrary if the visit time was decided based in part on clinical symptoms not

included in the model, or if drop-out could be due to severe clinical events or death (related to CD4 and not included in the model), CAR(DYN) would not hold.

### 7.2.3. Vertically coarsened observations

If we are interested in the evolution of HIV-RNA, we may consider the same issues as above ( $X_t^i$  now representing HIV-RNA), with the additional complexity of a (known) detection limit  $\eta$ : this produces a left-censoring of  $X_t^i$ : if  $R_t^i = 1$  we observe  $Y_t^i = (1_{X_t^i > \eta}, \tilde{X}_t^i)$ , where  $\tilde{X}_t^i = \max(X_t^i, \eta)$  (see (28)). This is a case of what we have called “vertical coarsening” in section 3.2.

### 7.3. Joint modeling

One of the reasons for considering a joint model is precisely to remove the bias due to so called “informative censoring “. Consider as in the example of section 7.2.2 that we are interested in the evolution of the concentration of CD4+ T-lymphocytes represented by  $(T_t^i)$ . Only noisy discrete observation of  $(T_t^i)$ , the CD4 counts, are available. Assume that subjects are lost from follow-up when they develop AIDS. There is a rather strong relationship between concentration of CD4+ T-lymphocytes and the risk of developing AIDS, so that the intensity of the counting process describing drop-out depends on it: CAR(DYN) does not hold in a model which does not include AIDS.

Thus we may consider jointly modelling the concentration of CD4+ T-lymphocytes and AIDS and consider the process  $(X_t^i) = (T_t^i, W_t^i, Z_t^i)$ , where  $(W_t^i)$  is a counting process which represents AIDS and  $(Z_t^i)$  is a multivariate process of explanatory variables (such a joint model has been developed in (13)). We may allow as in section 7.2.2 that the visit times depend on the observed CD4 counts; we may also allow that the probability that the subject drop out after  $V_j^i$  depends on the AIDS status at this visit. The compensator of the associated counting process  $N^i$  will be constant between  $V_j^i$  and  $V_{j+1}^i$  and will make a jump at  $V_{j+1}^i$  equal to one minus the probability of drop-out: if the latter depends on

the observed AIDS status  $W_{V_j}^i$  only, then CAR(DYN) holds; if it depends of unobserved status  $W_t^i$  for  $t \neq V_j$ , then CAR(DYN) does not hold.

## 8. Conclusion

We have proposed a coarsening model for processes (TCMP) and developed a theory of ignorability in this framework. The theory applies to general stochastic processes having discrete or continuous state-space; in particular it applies to both counting processes and diffusion processes. The framework of repeated measurements can be represented as a continuous-time continuous state-space process observed at discrete times. Our results hold even if the observed part of the process of interest  $X$  has a null probability, which is the case in the examples of the previous section. We have given a factorization condition for the likelihood which allows to get rid of the nuisance parameter and may be called weak ignorability; we can define ignorability in a strong sense, that is equality between the correct likelihood and the likelihood ignoring the m.l.i.d. on events  $\{R = r\}$  of non-null probability. This restriction comes from the fact that likelihoods, as Radon-Nikodym derivatives, are not uniquely defined.

We have already applied some of the results presented here to a multi-state model for dementia, institutionalization and death (6). This model can be represented by a three-variate counting process to which we associate the three-variate response process. In this application the observed event of  $X$  is generally of null probability although it is not a singleton; institutionalization can be observed either exactly or in an interval; we showed that the m.l.i.d. could be ignorable. The factorization theorem is also useful for defining a risk for model choice in the context of coarsened observations of multistate processes as is proposed in (8). Finally this scheme of observation is useful for making inference in the framework of a general dynamical model such as proposed by Commenges and Gégout-Petit (7). In these latter papers it is made explicit that there is a true probability  $P^*$  under which the events in the universe are generated. In the present paper we have spoken in terms of model. However it must be clear



that for ignorability to hold really, the conditions must hold under the true probability; in particular the condition in CAR(DYN) must be hold under  $P^*$ .

## References

- [1] ALIOUM A AND COMMENGES, D. (1996). A proportional hazards model for arbitrarily censored and truncated data.. *Biometrics* **52**, 95-107.
- [2] ANDERSEN, P. K., BORGAN, Ø., GILL, R. D. AND KEIDING, N. (1993). *Statistical Models Based on Counting Processes*. New-York: Springer-Verlag. [MR1198884](#)
- [3] ARJAS, E. AND HAARA, P. (1992). Observation scheme and likelihood. *Scandinavian Journal of Statistics* **19**, 111-132. [MR1173594](#)
- [4] ARJAS, E., HAARA, P. AND NORROS, I. (1992). Filtering the histories of a partially observed marked point process. *Stochastic Processes and their Applications* **40**, 225-250. [MR1158025](#)
- [5] BARNDORFF-NIELSEN, O. E. AND SORENSEN, M. (1994) A review of some aspects of asymptotic likelihood theory for stochastic processes. *Int. Statist. Rev.* **62**, 133-165.
- [6] COMMENGES, D. AND GÉGOUT-PETIT, A. (2007). Likelihood for generally coarsened observations from multi-state or counting process models. *Scand. J. Statist.* **34**, 432-450. [MR2346649](#)
- [7] COMMENGES, AND GÉGOUT-PETIT, A. (2009). A general dynamical model with causal interpretation. *J.Roy.Stat.Soc.B* **71**, 719-736.
- [8] COMMENGES, D., JOLY, P., GÉGOUT-PETIT, A. AND LIQUET, B. (2007). Choice between semi-parametric estimators of Markov and non-Markov multi-state models from generally coarsened observations. *Scand. J. Statist.* **34**, 33-52. [MR2325241](#)
- [9] COX, D. R. (1972). Regression models and life tables. *Journal of the Royal Statistical Society, Series B* **34**, 187-220. [MRMR0341758](#)

- [10] DIGGLE, P. J. AND KENWARD, M. G. (1994). Informative Drop-out in Longitudinal Data Analysis. *Applied Statistics* **43**, 49-93.
- [11] FEIGIN, P. D. (1976). Maximum likelihood estimation for continuous-time stochastic processes. *Adv. Appl. Prob.* **8**, 712-736. [MR0426342](#)
- [12] GILL, R. D., VAN DER LAAN, M. J. AND ROBINS, J.M. (1997). Coarsening at random: characterizations, conjectures and counter-examples. *State of the Art in Survival Analysis*, D.-Y. Lin and T.R. Fleming (eds), Springer Lecture Notes in Statistics 123, pp. 255-294.
- [13] GUEDJ, J, THIEBAUT, R AND COMMENGES, D. (2010) Joint modeling of the clinical progression and of the biomarkers dynamics using a mechanistic model. *Biometrics*, DOI: 10.1111/j.1541-0420.2010.01418.x, in press.
- [14] HEITJAN, D. F. AND RUBIN, D. B. (1991). Ignorability and coarse data. *Ann. Statist.* **19**, 2244-2253. [MR1135174](#)
- [15] HOFFMANN-JORGENSEN, J. (1994). *Probability with a view toward statistics*. London: Chapman and Hall.[MR1278485](#)
- [16] JACOD, J. (1975). Multivariate point processes: predictable projection; Radon-Nikodym derivative, representation of martingales. *Z. Wahrsch. Verw. Geb.* **31**, 235-253. [MR0380978](#)
- [17] JACOD, J. AND SHIRYAEV, A. N. (2003) *Limit Theorems for Stochastic Processes*. Springer, New-York. [MR1943877](#)
- [18] JACOBSEN, M. AND KEIDING, N. (1995). Coarsening at random in general sample spaces and random censoring in continuous time. *Annals of Statistics* **23**, 774-786. [MR1345200](#)
- [19] KALBFLEISCH, J. D. AND PRENTICE R. L. (1980). *The Statistical Analysis of Failure Time Data*, Wiley, New-York. [MR0570114](#)
- [20] KALLENBERG, O. (2002). *Foundations of modern probabilities*. Springer Verlag, New-York. [MR1876169](#)
- [21] KAPLAN, E. L. AND MEIER, P. (1958). Non-parametric estimation from incomplete observations, *Journal of the American Statistical Association* **80**, 457-481, 562-563. [MR0093867](#)

- [22] KENWARD M. G. AND MOLENBERGHS G. (1998). Likelihood based frequentist inference when data are missing at random. *Statist. Sci.* **13**, 236-247. [MR1665713](#)
- [23] LAGAKOS, S. (1979). General right-censoring and its impact on the analysis of survival data. *Biometrics* **35**, 139-156.
- [24] NIELSEN, S.F. (2000). Relative coarsening at random. *Statistica Neerlandica* **54**, 79-99.
- [25] PETO, R. (1973). Experimental survival curves for interval-censored data. *Applied Statistics* **22**, 86- 91.
- [26] O’SULLIVAN, F. (1988). Fast computation of fully automated log-density and log-hazard estimators. *SIAM J. Scient. Statist. Comput.* **9**, 363-379. [MR0930052](#)
- [27] RUBIN, D. B. (1976). Inference and missing data. *Biometrika* **63**, 581-592. [MR0455196](#)
- [28] THIÉBAUT, R. , JACQMIN-GADDA H. , BABIKER A. AND COMMENGES D. (2005). Joint Modelling of bivariate longitudinal data with informative drop-out and left-censoring, with application to the evolution of CD4+ cell count and HIV RNA viral load in response to treatment of HIV infection. *Statistics in Medicine* **24**, 65-82. [MR2134496](#)
- [29] VAN DER LAAN, M.J. AND ROBINS, J.M. (2003). Unified methods for censored longitudinal longitudinal data ans causality. Springer. [MR1958123](#)
- [30] WILLIAMS, D. (1991). *Probability with Martingales*. Cambridge University Press. [MR1155402](#)

## Appendix A: CAR(TCMP)-loc

This appendix develops the concept of CAR(TCMP)-loc of Definition 4. First note that this definition has a meaning only for  $r$  such that  $P(\{R = r\}) > 0$ . The TCMP model verifies the condition CAR(ABS) defined by Nielsen (2000)

if  $\forall(\theta, \psi)$ ,  $P_{(\theta, \psi)}$  is CAR(ABS) i.e. the following condition is true :

for  $P_\theta$  a.e.  $x, x'$ , for every  $A \in \mathcal{O}$ ,

$$P_\psi(A \cap D_x \cap D_{x'} | X = x) = P_\psi(A \cap D_x \cap D_{x'} | X = x') \quad (\text{A.1})$$

where  $D_x = \{(r, y) \in (\Xi, \Gamma); ry = rx\} \in \mathcal{O}$ . Note moreover that as pointed by Gill & all (12), if  $P_{(\theta_0, \psi_0)}$  is CAR(ABS) and the model is CAR(REL), then the model is CAR(ABS).

**Theorem A.1.** *If the TCMP is CAR(ABS), then CAR(TCMP)-loc holds on all  $r$ .*

**Proof.** Assume the model is CAR(ABS) and remark that  $\{R = r\} \cap D_x = \{R = r\} \cap D_{rx}$  and that  $\forall(\theta, \psi)$ ,  $P_{(\theta, \psi)}[D_x | X = x] = 1$ , then  $\forall(\theta, \psi)$  for  $P_\theta$  a.e.  $x$  and every  $\mathcal{O}$ -measurable  $Z$ , we get

$$\begin{aligned} \mathbb{E}_{(\theta, \psi)}[1_{\{R=r\}} Z | X = x] &= \mathbb{E}_{(\theta, \psi)}[1_{\{R=r\}} Z 1_{D_x} | X = x] \\ &= \mathbb{E}_{(\theta, \psi)}[1_{\{R=r\}} Z | X = rx] \end{aligned}$$

From this equality, we deduce that  $\forall(\theta, \psi)$  for every  $Z$ ,  $\mathcal{O}$ -measurable,  $\mathbb{E}_{(\theta, \psi)}[1_{\{R=r\}} Z | \mathcal{X}] = \mathbb{E}_{(\theta, \psi)}[1_{\{R=r\}} Z | \mathcal{X}^r]$ . It follows that for every  $A \in \mathcal{R}$ :  $\mathbb{E}_{(\theta_0, \psi_0)}[1_{\{R=r\}} 1_A \mathcal{L}_{\mathcal{R}|\mathcal{X}}^{(\theta, \psi)/(\theta_0, \psi_0)} | \mathcal{X}] = \mathbb{E}_{(\theta, \psi)}[1_{\{R=r\}} 1_A \mathcal{L}_{\mathcal{R}|\mathcal{X}^r}^{(\theta, \psi)/(\theta_0, \psi_0)} | \mathcal{X}]$  which implies CAR(TCMP)-loc.  $\square$

**Proof of Theorem A.** By the iterated decomposition formula (property v), we have  $\mathcal{L}_{\mathcal{X}, \mathcal{R}}^{(\theta, \psi)/(\theta_0, \psi_0)} = \mathcal{L}_{\mathcal{X}, \mathcal{X}^r, \mathcal{R}} = \mathcal{L}_{\mathcal{X}|\mathcal{X}^r, \mathcal{R}}^{(\theta, \psi)/(\theta_0, \psi_0)} \mathcal{L}_{\mathcal{R}|\mathcal{X}^r} \mathcal{L}_{\mathcal{X}^r}$ . We have  $\mathcal{L}_{\mathcal{O}}^{(\theta, \psi)/(\theta_0, \psi_0)} = \mathbb{E}[\mathcal{L}_{\mathcal{X}, \mathcal{R}}^{(\theta, \psi)/(\theta_0, \psi_0)} | \mathcal{O}] = \mathbb{E}[\mathcal{L}_{\mathcal{X}|\mathcal{X}^r, \mathcal{R}}^{(\theta, \psi)/(\theta_0, \psi_0)} \mathcal{L}_{\mathcal{R}|\mathcal{X}^r} \mathcal{L}_{\mathcal{X}^r} | \mathcal{O}]$ . On  $\{R = r\}$ ,  $\mathcal{O} = \mathcal{X}^r \vee \mathcal{R}$  and by Lemma 1.1 we have:

$$\mathcal{L}_{\mathcal{O}}^{(\theta, \psi)/(\theta_0, \psi_0)} = \mathcal{L}_{\mathcal{X}^r}^{\theta/\theta_0} \mathcal{L}_{\mathcal{R}|\mathcal{X}^r}^{(\theta, \psi)/(\theta_0, \psi_0)} \mathbb{E}_{(\theta_0, \psi_0)}[\mathcal{L}_{\mathcal{X}|\mathcal{X}^r, \mathcal{R}}^{(\theta, \psi)/(\theta_0, \psi_0)} | \mathcal{X}^r \vee \mathcal{R}] = \mathcal{L}_{\mathcal{X}^r}^{\theta/\theta_0} \mathcal{L}_{\mathcal{R}|\mathcal{X}^r}^{(\theta, \psi)/(\theta_0, \psi_0)}.$$

If CAR(TCMP)-loc holds we have on  $\{R = r\}$ ,  $\mathcal{L}_{\mathcal{R}|\mathcal{X}^r}^{(\theta, \psi)/(\theta_0, \psi_0)} = \mathcal{L}_{\mathcal{R}|\mathcal{X}}^{(\theta, \psi)/(\theta_0, \psi_0)}$ ; we have  $\mathcal{L}_{\mathcal{R}|\mathcal{X}}^{(\theta, \psi)/(\theta_0, \psi_0)} = \mathcal{L}_{\mathcal{R}|\mathcal{X}}^{\psi/\psi_0}$ . Thus with both conditions we have  $\mathcal{L}_{\mathcal{O}}^{(\theta, \psi)/(\theta_0, \psi_0)} = \mathcal{L}_{\mathcal{X}^r}^{\theta/\theta_0} \mathcal{L}_{\mathcal{R}|\mathcal{X}}^{\psi/\psi_0}$ , a.s. on  $\{R = r\}$  and this implies ignorability.  $\square$

## Appendix B: Proof of Theorem 6.3

The likelihood of the counting process  $(X, N)$  can be written using Jacod's formula as  $\mathcal{L}_{\mathcal{X}, \mathcal{N}} = \phi(\Lambda^{\mathcal{F}, X}, \Lambda_0^{\mathcal{F}, X}, X) \phi(\Lambda^{\mathcal{F}, N}, \Lambda_0^{\mathcal{F}, N}, N)$ . As we have done above we can also write it:  $\mathcal{L}_{\mathcal{X}, \mathcal{N}} = \mathcal{L}_{\mathcal{X}}^{\theta/\theta_0} \phi(\Lambda^{\mathcal{F}^*, N}, \Lambda_0^{\mathcal{F}^*, N}, N)$ . Noting that  $\mathcal{L}_{\mathcal{X}}^{\theta/\theta_0} = \phi(\Lambda^{\mathcal{X}, X}, \Lambda_0^{\mathcal{X}, X}, X)$  and equating the two representations we have:

$$\phi(\Lambda^{\mathcal{F}, X}, \Lambda_0^{\mathcal{F}, X}, X) \phi(\Lambda^{\mathcal{F}, N}, \Lambda_0^{\mathcal{F}, N}, N) = \phi(\Lambda^{\mathcal{X}, X}, \Lambda_0^{\mathcal{X}, X}, X) \phi(\Lambda^{\mathcal{F}^*, N}, \Lambda_0^{\mathcal{F}^*, N}, N).$$

CAR(DYN) says that  $\Lambda^{\mathcal{F}^*, N} = \Lambda^{\mathcal{O}, N}$  and in this right-censoring case we have  $\Lambda^{\mathcal{O}, N} = \Lambda^{\mathcal{F}, N}$  (this is because  $\mathcal{O}_t = \mathcal{F}_t$  on  $\{t \leq C\}$  and  $\Lambda_{C+u}^N = \Lambda_C^N$  whatever the filtration). So if CAR(DYN) holds, the above equation yields  $\phi(\Lambda^{\mathcal{F}, X}, \Lambda_0^{\mathcal{F}, X}, X) = \phi(\Lambda^{\mathcal{X}, X}, \Lambda_0^{\mathcal{X}, X}, X)$ . This must be true almost surely, for all  $(\theta, \psi)$ , and moreover, we still have this equality if we stop the observation at time  $t$  or at a  $(\mathcal{X}_t)$ -stopping time  $T$ . All that we have to prove is that this implies  $\Lambda^{\mathcal{F}, X} = \Lambda^{\mathcal{X}, X}$ , which is independent censoring.

Let us begin with  $X$  a 0–1 counting process and denote its jump time  $T$ . If we stop observation at  $t$ , we have on  $\{T > t\}$ ,  $\frac{\exp \Lambda_t^{\mathcal{F}, X}}{\exp \Lambda_{0t}^{\mathcal{F}, X}} = \frac{\exp \Lambda_t^{\mathcal{X}, X}}{\exp \Lambda_{0t}^{\mathcal{X}, X}}$ ; because of left-continuity, we have also the equality of  $t = T$  and because the intensity is equal to zero after  $T$ , the equality holds for all  $t$  almost surely. Taking log and differentiating we obtain:

$$\lambda_t^{\mathcal{F}, X} - \lambda_{0t}^{\mathcal{F}, X} = \lambda_t^{\mathcal{X}, X} - \lambda_{0t}^{\mathcal{X}, X}. \quad (\text{B.1})$$

The likelihood has a limit when  $t \rightarrow \infty$  and at the limit we have  $\frac{\lambda_T^{\mathcal{F}, X} \exp \Lambda_T^{\mathcal{F}, X}}{\lambda_{0T}^{\mathcal{F}, X} \exp \Lambda_{0T}^{\mathcal{F}, X}} = \frac{\lambda_T^{\mathcal{X}, X} \exp \Lambda_T^{\mathcal{X}, X}}{\lambda_{0T}^{\mathcal{X}, X} \exp \Lambda_{0T}^{\mathcal{X}, X}}$  from which we successively deduce  $\frac{\lambda_T^{\mathcal{F}, X}}{\lambda_{0T}^{\mathcal{F}, X}} = \frac{\lambda_T^{\mathcal{X}, X}}{\lambda_{0T}^{\mathcal{X}, X}}$  and  $\frac{\lambda_t^{\mathcal{F}, X}}{\lambda_{0t}^{\mathcal{F}, X}} = \frac{\lambda_t^{\mathcal{X}, X}}{\lambda_{0t}^{\mathcal{X}, X}}$ , almost surely for all  $t$  on the support of the distribution of  $T$ . Combining this result with (B.1), we obtain  $\lambda_t^{\mathcal{F}, X} = \lambda_t^{\mathcal{X}, X}$  a.s., which for càg processes implies indistinguishability of the intensities and of the cumulative intensities. If the process may have several jumps  $T_1, T_2, \dots$ , we first prove by the same reasoning that we have equality of the intensities on  $\{t \leq T_1\}$ , then using this result and again the same reasoning we have equality on  $]T_1, T_2]$  and so on. All this reasoning is symmetrical so we can also prove that independent censoring implies CAR(DYN).  $\square$

### Appendix C: Extension to left-continuous $R$

In some situations it is natural to consider response indicator processes which are left-continuous; case II right-censoring is an example. In that case it is not possible to directly associate to  $R$  a counting process and hence to apply CAR(DYN). In order to extend the application of CAR(DYN) to such processes, we will consider them as limits of right-continuous processes. Let us still consider the univariate case. Consider for instance the case where  $R_0 = 1$ .  $(R_t)$  may be left-continuous at jumps of odd ranks:  $V_{2j+1}, j \geq 0$ ; the process can be written  $R_t = \sum_{j \geq 0} 1_{[V_{2j}, V_{2j+1}]}$ . Consider the sequence of processes  $R_t^n = (\sum_{j \geq 0} 1_{[V_{2j} \leq t < V_{2j+1}^n]}) \wedge 1$ , defined by:  $R_0^n = 1$ ,  $V_{2j+1}^n = V_{2j+1} + 1/n$ . The limit of  $(R_t^n)$  is  $(R_t)$ .

**Theorem C.1.** *Consider a process  $R = (R_t)$  which is right-continuous at upward jumps and may left-continuous at downward jumps. Consider a sequence of right-continuous processes  $R^n = (R_t^n)$  constructed as above; if each  $(X, R^n)$  satisfies CAR(TCMP)-loc on  $r$  then ignorability holds for  $(X, R)$  on  $r$ .*

**Proof.** We note  $\mathcal{O}^n$ , the observed  $\sigma$ -field associated to  $R^n$ . If  $(X, R^n)$  satisfies CAR(TCMP)-loc on  $r$  then (see the proof of Theorem (A)), we have for all  $n$ : on  $\{R = r\}$ ,  $\mathcal{L}_{\mathcal{O}^n}^{(\theta, \psi)/(\theta_0, \psi_0)} = \mathcal{L}_{\mathcal{R}^n | \mathcal{X}}^{(\theta, \psi)/(\theta_0, \psi_0)} \mathcal{L}_{\mathcal{X}^{r^n}}^{\theta/\theta_0}$ .  $\mathcal{O}^n$  is larger than  $\mathcal{O}$ :  $\mathcal{O} \subset \mathcal{O}^n$  and it is clear that  $\mathcal{O}^n$  is a decreasing sequence of  $\sigma$ -fields:  $\mathcal{O} = \cap_n \mathcal{O}^n = \mathcal{O}^\infty$ . By the Downward Levy Theorem (30) we have:  $\mathcal{L}_{\mathcal{O}^n}^{(\theta, \psi)/(\theta_0, \psi_0)} = E_{(\theta_0, \psi_0)}(\mathcal{L}_{\mathcal{F}}^{(\theta, \psi)/(\theta_0, \psi_0)} | \mathcal{O}^n) \rightarrow E_{(\theta_0, \psi_0)}(\mathcal{L}_{\mathcal{F}}^{(\theta, \psi)/(\theta_0, \psi_0)} | \mathcal{O}) = \mathcal{L}_{\mathcal{O}}^{(\theta, \psi)/(\theta_0, \psi_0)}$  a.s.. Using again the Downward Levy Theorem we get  $\mathcal{L}_{\mathcal{X}^{r^n}}^{\theta/\theta_0} \rightarrow \mathcal{L}_{\mathcal{X}^r}^{\theta/\theta_0}$  a.s. Moreover note that  $\mathcal{R}^n = \mathcal{R}$  because the process  $R^n$  is deterministically defined from  $R$ : this implies that  $\mathcal{L}_{\mathcal{R}^n | \mathcal{X}}^{(\theta, \psi)/(\theta_0, \psi_0)} = \mathcal{L}_{\mathcal{R} | \mathcal{X}}^{\psi/\psi_0}$ . At the limit we have thus:  $\mathcal{L}_{\mathcal{O}}^{(\theta, \psi)/(\theta_0, \psi_0)} = \mathcal{L}_{\mathcal{R} | \mathcal{X}}^{\psi/\psi_0} \mathcal{L}_{\mathcal{X}^r}^{\theta/\theta_0}$  which concludes the proof. A similar result could be obtained for upward jumps.  $\square$

As an example consider the case of Type II right-censoring where we have independent 0 – 1 counting processes  $(X_t^i)$ ,  $i = 1, \dots, n$  and observation is stopped just after observing the  $d^{th}$  event. Thus the response indicator process  $R$

is not independent on the multivariate process  $X$ . In fact we have  $R_t = 1_{\{\bar{X}_t \leq d\}}$ , where  $\bar{X}_t = \sum_i X_t^i$ . This is a case of a left-continuous process which has only one downward jump. Since  $R$  is  $\mathcal{X}$ -measurable  $\mathcal{L}_{\mathcal{R}|\mathcal{X}}^{\psi/\psi_0} = 1$  (by property iv) and is thus obviously  $\mathcal{O}$ -measurable which is CAR(TCMP). Consider now a slightly more sophisticated model which we call randomized Type II censoring in which we may stop observation after each event with a given probability depending of what have been observed. For instance let  $(T_1, T_2, \dots, T_n)$  the times of occurrence of the first, second, ..., events, and let the probability of stopping observation just after  $T_j$  (conditional on having observed  $X$  until  $T_j$ ) be  $\frac{j-1}{j}$ ,  $j = 1, \dots, n$ ; let  $C$  be the jump time of  $R$  ( $C = T_j$ , for some  $j$ ). We consider  $R$  as the limit of the sequence of right-continuous processes  $R^n$  such that  $R_t^n = 1_{\{t \geq C + 1/n\}}$ . We can easily verify that these observation processes satisfy CAR(DYN) (because future values of  $X$  are not used for defining the probability of stopping observation), and thus CAR(TCMP) by Theorem 5; thus  $R$  itself satisfies CAR(TCMP) by Theorem C.1.