

INVARIANT METRICS AND LAPLACIANS ON SIEGEL-JACOBI SPACE

JAE-HYUN YANG

Abstract. In this paper, we compute Riemannian metrics on the Siegel-Jacobi space which are invariant under the natural action of the Jacobi group explicitly and also provide the Laplacians of these invariant metrics. These are expressed in terms of the trace form.

1. Introduction

For a given fixed positive integer n , we let

$$H_n = \{Z \in \mathbb{C}^{(n,n)} \mid Z = {}^t Z; \operatorname{Im} Z > 0\}$$

be the Siegel upper half plane of degree n and let

$$Sp(n; \mathbb{R}) = \{M \in \mathbb{R}^{(2n, 2n)} \mid {}^t M J_n M = J_n\}$$

be the symplectic group of degree n , where

$$J_n = \begin{pmatrix} 0 & E_n \\ E_n & 0 \end{pmatrix}.$$

We see that $Sp(n; \mathbb{R})$ acts on H_n transitively by

$$(1.1) \quad M \cdot Z = (AZ + B)(CZ + D)^{-1};$$

where $M = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in Sp(n; \mathbb{R})$ and $Z \in H_n$:

For two positive integers n and m , we consider the Heisenberg group

$$H_{\mathbb{R}}^{(n,m)} = \left\{ \begin{pmatrix} x & y & z \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \mid x \in \mathbb{R}^{(n,m)}; y \in \mathbb{R}^{(m,n)}; z \in \mathbb{R} \right\}$$

endowed with the following multiplication law

$$\begin{pmatrix} x & y & z \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} x' & y' & z' \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} = \begin{pmatrix} x+x' & y+y' & z+z'+\frac{1}{2}(y, x') \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix};$$

We define the semidirect product of $Sp(n; \mathbb{R})$ and $H_{\mathbb{R}}^{(n,m)}$

$$G^J := Sp(n; \mathbb{R}) \ltimes H_{\mathbb{R}}^{(n,m)}$$

endowed with the following multiplication law

$$\begin{pmatrix} M & (x, y, z) \\ 0 & 1 \end{pmatrix} \begin{pmatrix} M' & (x', y', z') \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} MM' & (x+x', y+y', z+z'+\frac{1}{2}(y, x')) \\ 0 & 1 \end{pmatrix}$$

Key words and phrases. invariant metrics, Siegel-Jacobi space, spectral theory.

2000 Mathematics Subject Classification: Primary 32F45, 32M10.

This work was supported by Inha University Research Grant.

with $M \in M^0 \subset Sp(n; \mathbb{R}); (\begin{smallmatrix} a & b \\ c & d \end{smallmatrix}); (\begin{smallmatrix} 0 & 0 \\ 0 & 0 \end{smallmatrix}) \in H_R^{(n,m)}$ and $(\tilde{\cdot}; \sim) = (\cdot; \cdot) M^0$. We call this group G^J the Jacobi group of degree n and index m . We have the natural action of G^J on $H_n \subset \mathbb{C}^{(m,n)}$ defined by

$$(1.2) \quad M \cdot (\begin{smallmatrix} Z \\ W \end{smallmatrix}); (\begin{smallmatrix} a & b \\ c & d \end{smallmatrix}) = M \cdot \begin{smallmatrix} Z \\ W + Z + \end{smallmatrix} (CZ + D)^{-1};$$

where $M = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in Sp(n; \mathbb{R}); (\begin{smallmatrix} a & b \\ c & d \end{smallmatrix}) \in H_R^{(n,m)}$ and $(Z; W) \in H_n \subset \mathbb{C}^{(m,n)}$. The homogeneous space $H_n \subset \mathbb{C}^{(m,n)}$ is called the Siegel-Jacobi space of degree n and index m . We refer to [2-3], [6-7], [11], [14-21] for more details on materials related to the Siegel-Jacobi space.

For brevity, we write $H_{n,m} := H_n \subset \mathbb{C}^{(m,n)}$. For a coordinate $(Z; W) \in H_{n,m}$ with $Z = (z) \in H_n$ and $W = (w_{kl}) \in \mathbb{C}^{(m,n)}$, we put

$$\begin{aligned} Z &= X + iY; & X &= (x); & Y &= (y) \text{ real}; \\ W &= U + iV; & U &= (u_{kl}); & V &= (v_{kl}) \text{ real}; \\ dZ &= (dz); & d\bar{Z} &= (d\bar{z}); & dY &= (dy); \\ dW &= (dw_{kl}); & d\bar{W} &= (d\bar{w}_{kl}); & dV &= (dv_{kl}); \end{aligned}$$

$$\begin{aligned} \frac{\partial}{\partial Z} &= \frac{1+}{2} \frac{\partial}{\partial x} - i \frac{\partial}{\partial y}; & \frac{\partial}{\partial \bar{Z}} &= \frac{1+}{2} \frac{\partial}{\partial x} + i \frac{\partial}{\partial y}; \\ \frac{\partial}{\partial X} &= \frac{1+}{2} \frac{\partial}{\partial x} - i \frac{\partial}{\partial y}; & \frac{\partial}{\partial Y} &= \frac{1+}{2} \frac{\partial}{\partial x} + i \frac{\partial}{\partial y}; \\ \frac{\partial}{\partial W} &= \begin{pmatrix} \frac{\partial}{\partial w_{11}} & \cdots & \frac{\partial}{\partial w_{m1}} \\ \vdots & \ddots & \vdots \\ \frac{\partial}{\partial w_{1n}} & \cdots & \frac{\partial}{\partial w_{mn}} \end{pmatrix} \begin{pmatrix} 1 \\ \vdots \\ 1 \end{pmatrix} \begin{pmatrix} C \\ \vdots \\ A \end{pmatrix}; & \frac{\partial}{\partial \bar{W}} &= \begin{pmatrix} \frac{\partial}{\partial \bar{w}_{11}} & \cdots & \frac{\partial}{\partial \bar{w}_{m1}} \\ \vdots & \ddots & \vdots \\ \frac{\partial}{\partial \bar{w}_{1n}} & \cdots & \frac{\partial}{\partial \bar{w}_{mn}} \end{pmatrix} \begin{pmatrix} 1 \\ \vdots \\ 1 \end{pmatrix} \begin{pmatrix} C \\ \vdots \\ A \end{pmatrix}; \\ \frac{\partial}{\partial U} &= \begin{pmatrix} \frac{\partial}{\partial u_{11}} & \cdots & \frac{\partial}{\partial u_{m1}} \\ \vdots & \ddots & \vdots \\ \frac{\partial}{\partial u_{1n}} & \cdots & \frac{\partial}{\partial u_{mn}} \end{pmatrix} \begin{pmatrix} 1 \\ \vdots \\ 1 \end{pmatrix} \begin{pmatrix} C \\ \vdots \\ A \end{pmatrix}; & \frac{\partial}{\partial V} &= \begin{pmatrix} \frac{\partial}{\partial v_{11}} & \cdots & \frac{\partial}{\partial v_{m1}} \\ \vdots & \ddots & \vdots \\ \frac{\partial}{\partial v_{1n}} & \cdots & \frac{\partial}{\partial v_{mn}} \end{pmatrix} \begin{pmatrix} 1 \\ \vdots \\ 1 \end{pmatrix} \begin{pmatrix} C \\ \vdots \\ A \end{pmatrix}; \end{aligned}$$

where δ_{ij} denotes the Kronecker delta symbol.

C. L. Siegel [12] introduced the symplectic metric ds_n^2 on H_n invariant under the action (1.1) of $Sp(n; \mathbb{R})$ given by

$$(1.3) \quad ds_n^2 = Y^{-1} dZ Y^{-1} d\bar{Z}$$

and H. Maass [8] proved that the differential operator

$$(1.4) \quad \Delta_n = 4 \left(Y^{-1} \frac{\partial}{\partial Z} Y^{-1} \frac{\partial}{\partial \bar{Z}} \right)$$

is the Laplacian of H_n for the symplectic metric ds_n^2 . Here $\text{tr}(A)$ denotes the trace of a square matrix A .

In this paper, for arbitrary positive integers n and m , we express the G^J -invariant metrics on $H_n \subset \mathbb{C}^{(m,n)}$ and their Laplacians explicitly.

In fact, we prove the following theorems.

Theorem 1.1. For any two positive real numbers A and B , the following metric

$$\begin{aligned} ds_{n,m;A,B}^2 = & A \cdot Y^{-1} dZ \cdot Y^{-1} \overline{dZ} \\ & + B \cdot Y^{-1} {}^t V \cdot Y \cdot Y^{-1} dZ \cdot Y^{-1} \overline{dZ} + Y^{-1} {}^t (dW) \cdot \overline{dW} \\ & - Y \cdot Y^{-1} dZ \cdot Y^{-1} {}^t (\overline{dW}) - Y \cdot Y^{-1} \overline{dZ} \cdot Y^{-1} {}^t (dW) \end{aligned}$$

is a Riemannian metric on $H_{n,m}$ which is invariant under the action (1.2) of the Jacobi group G^J .

Theorem 1.2. For any two positive real numbers A and B , the Laplacian $\Delta_{n,m;A,B}$ of $(H_{n,m}; ds_{n,m;A,B}^2)$ is given by

$$\begin{aligned} \Delta_{n,m;A,B} = & \frac{4}{A} \cdot Y^t \cdot Y \cdot \frac{\partial}{\partial Z} \cdot \frac{\partial}{\partial Z} + Y \cdot Y^{-1} {}^t V \cdot Y \cdot \frac{\partial}{\partial W} \cdot \frac{\partial}{\partial W} \\ & + V^t \cdot Y \cdot \frac{\partial}{\partial Z} \cdot \frac{\partial}{\partial W} + {}^t V \cdot Y \cdot \frac{\partial}{\partial W} \cdot \frac{\partial}{\partial Z} \\ & + \frac{4}{B} \cdot Y \cdot \frac{\partial}{\partial W} \cdot \frac{\partial}{\partial W} : \end{aligned}$$

The following differential form

$$dv = (\det Y)^{-(n+m+1)} [dX] \wedge [dY] \wedge [dU] \wedge [dV]$$

is a G^J -invariant volume element on $H_{n,m}$, where

$$[dX] = \wedge dx; \quad [dY] = \wedge dy; \quad [dU] = \wedge_{k,l} du_{kl} \quad \text{and} \quad [dV] = \wedge_{k,l} dv_{kl}.$$

The point is that the invariant metric $ds_{n,m;A,B}^2$ and its Laplacian $\Delta_{n,m;A,B}$ are expressed in terms of the trace form.

For the case $n = m = 1$ and $A = B = 1$, Berndt proved in [1] (cf. [19]) that the metric $ds_{1,1}^2$ on $H \subset \mathbb{C}$ defined by

$$\begin{aligned} ds_{1,1}^2 = ds_{1,1;1,1} = & \frac{y+v^2}{y^3} (dx^2 + dy^2) + \frac{1}{y} (du^2 + dv^2) \\ & - \frac{2v}{y^2} (dx du + dy dv) \end{aligned}$$

is a Riemannian metric on $H \subset \mathbb{C}$ invariant under the action (1.2) of the Jacobi group and its Laplacian $\Delta_{1,1}$ is given by

$$\begin{aligned} \Delta_{1,1} = \Delta_{1,1;1,1} = & y^2 \left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} \right) + (y + v^2) \left(\frac{\partial^2}{\partial u^2} + \frac{\partial^2}{\partial v^2} \right) \\ & + 2yv \left(\frac{\partial^2}{\partial x \partial u} + \frac{\partial^2}{\partial y \partial v} \right) : \end{aligned}$$

It is a pleasure to thank Eberhard Freitag for his helpful advice and letting me know the paper [8] of H. Maass.

Notations: We denote by $\mathbb{R}$ and $\mathbb{C}$ the field of real numbers, and the field of complex numbers respectively. The symbol $\stackrel{\text{def}}{=}$ means that the expression on the right is the definition of that on the left. For two positive integers k and l , $F^{(k;l)}$ denotes the set of all $k \times l$ matrices with entries in a commutative ring F . For a square matrix $A \in F^{(k;k)}$ of degree k , $\text{tr}(A)$ denotes the trace of A . For any $M \in F^{(k;l)}$; M^t denotes the transpose matrix of M . E_n denotes the identity matrix of degree n . For $A \in F^{(k;l)}$ and $B \in F^{(l;k)}$, we set $B[A] = {}^tABA$. For a complex matrix A , $\overline{A}$ denotes the complex conjugate of A . For $A \in \mathbb{C}^{(k;l)}$ and $B \in \mathbb{C}^{(l;k)}$, we use the abbreviation $BfAg = {}^t\overline{A}BA$.

2. Proof of Theorem 1.1

Let $g = \begin{pmatrix} M & & & \\ & A & B & \\ & C & D & \\ & & & \end{pmatrix}$ be an element of G^J with $M = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in \text{Sp}(n; \mathbb{R})$ and $(Z; W) \in H_{n,m}$ with $Z \in H_n$ and $W \in \mathbb{C}^{(m;n)}$. If we put $(Z; W) = g \cdot (Z; W)$; then we have

$$\begin{aligned} Z &= M \cdot Z = (AZ + B)(CZ + D)^{-1}; \\ W &= (W + Z +) (CZ + D)^{-1}; \end{aligned}$$

Thus we obtain

$$(2.1) \quad dZ = dZ [(CZ + D)^{-1}] = {}^t(CZ + D)^{-1} dZ (CZ + D)^{-1}$$

and

$$(2.2) \quad dW = dW (CZ + D)^{-1} + f \cdot (W + Z +) (CZ + D)^{-1} C g dZ (CZ + D)^{-1};$$

Here we used the following facts that

$$d(CZ + D)^{-1} = -(CZ + D)^{-1} C dZ (CZ + D)^{-1}$$

and that $(CZ + D)^{-1} C$ is symmetric.

We put

$$Z = X + iY; \quad W = U + iV; \quad X, Y, U, V \text{ real};$$

From [9], p.33 or [13], p.128, we know that

$$(2.3) \quad Y = Y f(CZ + D)^{-1} g = {}^t(C\overline{Z} + D)^{-1} Y (CZ + D)^{-1};$$

First of all, we recall that the following matrices

$$\begin{aligned} t(b) &= \begin{pmatrix} E_n & b \\ 0 & E_n \end{pmatrix}; \quad b = {}^tb \text{ real}; \\ g_0(h) &= \begin{pmatrix} {}^th & 0 \\ 0 & h^{-1} \end{pmatrix}; \quad h \in \text{GL}(n; \mathbb{R}); \\ J_n &= \begin{pmatrix} 0 & E_n \\ E_n & 0 \end{pmatrix} \end{aligned}$$

generate the symplectic group $\text{Sp}(n; \mathbb{R})$ (cf. [4], [5]). Therefore the following elements

$t(b; \begin{smallmatrix} & & & \\ & & & \\ & & & \\ & & & \end{smallmatrix}); g(h)$ and $\begin{smallmatrix} & & & \\ & & & \\ & & & \\ & & & \end{smallmatrix}$ of G^J defined by

$$t(b; \begin{smallmatrix} & & & \\ & & & \\ & & & \\ & & & \end{smallmatrix}) = \begin{pmatrix} E_n & b \\ 0 & E_n \end{pmatrix}; (\begin{smallmatrix} & & & \\ & & & \\ & & & \\ & & & \end{smallmatrix}); b = {}^t b \text{ real}; (\begin{smallmatrix} & & & \\ & & & \\ & & & \\ & & & \end{smallmatrix}) \in H_R^{(n,m)};$$

$$g(h) = \begin{pmatrix} {}^t h & 0 \\ 0 & h^{-1} \end{pmatrix}; (0; 0; 0); h \in GL(n; R);$$

$$\begin{smallmatrix} & & & \\ & & & \\ & & & \\ & & & \end{smallmatrix} = \begin{pmatrix} 0 & E_n \\ E_n & 0 \end{pmatrix}; (0; 0; 0)$$

generate the Jacobi group G^J : So it suffices to prove the invariance of the metric $ds_{n,m}^2$ under the action of the generators $t(b; \begin{smallmatrix} & & & \\ & & & \\ & & & \\ & & & \end{smallmatrix}); g(h)$ and $\begin{smallmatrix} & & & \\ & & & \\ & & & \\ & & & \end{smallmatrix}$: For brevity, we write

$$(a) = Y^{-1} dZ Y^{-1} d\bar{Z};$$

$$(b) = Y^{-1} {}^t V V Y^{-1} dZ Y^{-1} d\bar{Z};$$

$$(c) = Y^{-1} {}^t (dW) d\bar{W};$$

$$(d) = V Y^{-1} dZ Y^{-1} {}^t (d\bar{W}) + V Y^{-1} d\bar{Z} Y^{-1} {}^t (dW)$$

and

$$(a) = Y^{-1} dZ Y^{-1} d\bar{Z};$$

$$(b) = Y^{-1} {}^t V V Y^{-1} dZ Y^{-1} d\bar{Z};$$

$$(c) = Y^{-1} {}^t (dW) d\bar{W};$$

$$(d) = V Y^{-1} dZ Y^{-1} {}^t (d\bar{W}) + V Y^{-1} d\bar{Z} Y^{-1} {}^t (dW)$$

Case I. $g = t(b; \begin{smallmatrix} & & & \\ & & & \\ & & & \\ & & & \end{smallmatrix})$ with $b = {}^t b \text{ real}$ and $(\begin{smallmatrix} & & & \\ & & & \\ & & & \\ & & & \end{smallmatrix}) \in H_R^{(n,m)}$:

In this case, we have

$$\begin{pmatrix} Z \\ Y \\ W \end{pmatrix} = \begin{pmatrix} Z + b \\ Y \\ W \end{pmatrix}; \begin{pmatrix} V \\ Y \\ W \end{pmatrix} = \begin{pmatrix} V \\ Y \\ W \end{pmatrix} + \begin{pmatrix} Z + b \\ 0 \\ 0 \end{pmatrix};$$

and

$$dZ = dZ; dW = dW + dZ;$$

Therefore

$$(a) = Y^{-1} dZ Y^{-1} d\bar{Z} = Y^{-1} dZ Y^{-1} d\bar{Z} = (a);$$

$$(b) = Y^{-1} {}^t V V Y^{-1} dZ Y^{-1} d\bar{Z} + Y^{-1} {}^t V dZ Y^{-1} d\bar{Z} \\ + {}^t V Y^{-1} dZ Y^{-1} d\bar{Z} + {}^t dZ Y^{-1} d\bar{Z};$$

$$(c) = Y^{-1} {}^t (dW) d\bar{W} + Y^{-1} {}^t (dW) d\bar{Z} \\ + Y^{-1} dZ {}^t d\bar{W} + Y^{-1} dZ {}^t d\bar{Z}$$

and

$$\begin{aligned}
 (d) = & \begin{pmatrix} V Y^{-1} dZ Y^{-1t} (\overline{dW}) & dZ Y^{-1t} (\overline{dW}) \\ V Y^{-1} dZ Y^{-1} \overline{dZ}^t & dZ Y^{-1} \overline{dZ}^t \\ V Y^{-1} \overline{dZ} Y^{-1t} (dW) & \overline{dZ} Y^{-1t} (dW) \\ V Y^{-1} \overline{dZ} Y^{-1} dZ^t & \overline{dZ} Y^{-1} dZ^t \end{pmatrix} :
 \end{aligned}$$

Thus we see that

$$(a) = (a) \quad \text{and} \quad (b) + (c) + (d) = (b) + (c) + (d) :$$

Hence

$$ds_{n,m;A,B}^2 = A \begin{pmatrix} n \\ 0 \end{pmatrix} (a) + B \begin{pmatrix} 0 \\ n \end{pmatrix} (b) + (c) + (d)$$

is invariant under the action of $t(B; ; ;)$:

Case II. $g = g(h)$ with $h \in GL(n; R)$:

In this case, we have

$$Z = {}^t h Z h; \quad Y = {}^t h Y h; \quad W = W h; \quad V = V h$$

and

$$dZ = {}^t h dZ h; \quad dW = dW h;$$

Therefore by an easy computation, we see that each of (a); (b); (c) and (d) is invariant under the action of all $g(h)$ with $h \in GL(n; R)$: Hence the metric $ds_{n,m;A,B}^2$ is invariant under the action of all $g(h)$ with $h \in GL(n; R)$:

$$\text{Case III. } g = \begin{pmatrix} 0 & E_n \\ E_n & 0 \end{pmatrix}; (0; 0; 0) :$$

In this case, we have

$$(2.4) \quad Z = Z^{-1} \quad \text{and} \quad W = W Z^{-1}:$$

We set

$${}_1 := \operatorname{Re} Z^{-1} \quad \text{and} \quad {}_2 := \operatorname{Im} Z^{-1}:$$

Then ${}_1$ and ${}_2$ are symmetric matrices and we have

$$(2.5) \quad Y = {}_2 \quad \text{and} \quad V := \operatorname{Im} W = V {}_1 + U {}_2:$$

It is easy to see that

$$(2.6) \quad Y = Z {}_2 \overline{Z} = \overline{Z} {}_2 Z;$$

$$(2.7) \quad {}_1 Y + {}_2 X = 0$$

and

$$(2.8) \quad {}_1 X - {}_2 Y = E_n:$$

According to (2.6) and (2.7), we obtain

$$(2.9) \quad X = ({}_2)^{-1} {}_1 Y \quad \text{and} \quad Y^{-1} = {}_1 ({}_2)^{-1} {}_1 - {}_2:$$

From (2.1) and (2.2), we have

$$(2.10) \quad dZ = Z^{-1} dZ Z^{-1}$$

and

$$(2.11) \quad dW = dW Z^{-1} - W Z^{-1} dZ Z^{-1} = dW - W Z^{-1} dZ Z^{-1};$$

Therefore we have, according to (2.6) and (2.10),

$$\begin{aligned} (a) &= (\gamma_2)^{-1} Z^{-1} dZ Z^{-1} (\gamma_2)^{-1} \overline{Z}^{-1} d\overline{Z} \overline{Z}^{-1} \\ &= Y^{-1} dZ Y^{-1} d\overline{Z} = (a); \end{aligned}$$

According to (2.5)-(2.10), we have

$$\begin{aligned} (b) &= (\gamma_2)^{-1} (\gamma_1^t V + \gamma_2^t U) (V_{-1} + U_{-2}) (\gamma_2)^{-1} Z^{-1} dZ Y^{-1} d\overline{Z} \overline{Z}^{-1} \\ &= f^t U (\gamma_2)^{-1} \gamma_1^t V g f U - V_{-1} (\gamma_2)^{-1} g Z^{-1} dZ Y^{-1} d\overline{Z} \overline{Z}^{-1} \\ &= f^t \overline{W} + (iE_n - (\gamma_2)^{-1} \gamma_1^t) V g f W - V (iE_n + \gamma_1 (\gamma_2)^{-1}) g \\ &\quad Z^{-1} dZ Y^{-1} d\overline{Z} \overline{Z}^{-1}; \end{aligned}$$

$$\begin{aligned} (c) &= (\gamma_2)^{-1} Z^{-1} {}^t(dW) Z^{-1} dZ Z^{-1} {}^tW - d\overline{W} \overline{Z}^{-1} \overline{W} \overline{Z}^{-1} d\overline{Z} \overline{Z}^{-1} \\ &= (\gamma_2)^{-1} Z^{-1} {}^t(dW) d\overline{W} \overline{Z}^{-1} - (\gamma_2)^{-1} Z^{-1} {}^t(dW) \overline{W} \overline{Z}^{-1} d\overline{Z} \overline{Z}^{-1} \\ &\quad (\gamma_2)^{-1} Z^{-1} dZ Z^{-1} {}^tW d\overline{W} \overline{Z}^{-1} \\ &\quad + (\gamma_2)^{-1} Z^{-1} dZ Z^{-1} {}^tW \overline{W} \overline{Z}^{-1} d\overline{Z} \overline{Z}^{-1} \\ &= Y^{-1} {}^t(dW) d\overline{W} - Y^{-1} {}^t(dW) \overline{W} \overline{Z}^{-1} d\overline{Z} \\ &\quad Y^{-1} dZ Z^{-1} {}^tW d\overline{W} + Y^{-1} dZ Z^{-1} {}^tW \overline{W} \overline{Z}^{-1} d\overline{Z} \end{aligned}$$

and

$$\begin{aligned}
(d) &= (V_1 + U_2)(\partial_2)^{-1} Z^{-1} dZ Z^{-1} (\partial_2)^{-1} \overline{Z}^{-1} {}^t(d\overline{W}) - \overline{Z}^{-1} d\overline{Z} \overline{Z}^{-1} {}^t\overline{W}^{\circ} \\
&\quad (V_1 + U_2)(\partial_2)^{-1} \overline{Z}^{-1} d\overline{Z} \overline{Z}^{-1} (\partial_2)^{-1} Z^{-1} {}^t(dW) - Z^{-1} dZ Z^{-1} {}^tW \\
&= (V_1 + U_2)(\partial_2)^{-1} Z^{-1} dZ Y^{-1} {}^t(d\overline{W}) \\
&\quad + (V_1 + U_2)(\partial_2)^{-1} Z^{-1} dZ Y^{-1} d\overline{Z} \overline{Z}^{-1} {}^t\overline{W} \\
&\quad (V_1 + U_2)(\partial_2)^{-1} \overline{Z}^{-1} d\overline{Z} Y^{-1} {}^t(dW) \\
&\quad + (V_1 + U_2)(\partial_2)^{-1} \overline{Z}^{-1} d\overline{Z} Y^{-1} dZ Z^{-1} {}^tW :
\end{aligned}$$

Taking the $(dZ; d\overline{W})$ -part $(Z; \overline{W})$ in (b) + (c) + (d); we have

$$\begin{aligned}
(Z; \overline{W}) &= V Y^{-1} dZ Y^{-1} {}^t(d\overline{W}) + Y^{-1} dZ ({}^tW - Z^{-1} {}^tW) d\overline{W} \\
&= V Y^{-1} dZ Y^{-1} {}^t(d\overline{W}) \quad \text{because } W = W Z^{-1} \text{ (cf. (2.4))}:
\end{aligned}$$

Similarly, if we take the $(d\overline{Z}; dW)$ -part $(\overline{Z}; W)$ in (b) + (c) + (d); we have

$$\begin{aligned}
(\overline{Z}; W) &= V Y^{-1} d\overline{Z} Y^{-1} {}^t(dW) + d\overline{Z} Y^{-1} {}^t(dW) (\overline{W} - \overline{W} \overline{Z}^{-1}) \\
&= V Y^{-1} d\overline{Z} Y^{-1} {}^t(dW) \quad \text{because } W = W Z^{-1}:
\end{aligned}$$

If we take the $(dW; d\overline{W})$ -part $(W; \overline{W})$ in (b) + (c) + (d); we have

$$(W; \overline{W}) = Y^{-1} {}^t(dW) d\overline{W} :$$

Finally, if we take the $(dZ; d\overline{Z})$ -part $(Z; \overline{Z})$ in (b) + (c) + (d); we have

$$\begin{aligned}
(Z; \overline{Z}) &= {}^t\overline{W} + (iE_n - (\partial_2)^{-1} {}_1) {}^tV - W - V (iE_n + {}_1 (\partial_2)^{-1}) \\
&\quad Z^{-1} dZ Y^{-1} d\overline{Z} \overline{Z}^{-1} \\
&\quad + Z^{-1} {}^tW \overline{W} \overline{Z}^{-1} d\overline{Z} Y^{-1} dZ \\
&\quad + {}^t\overline{W} (V_1 + U_2)(\partial_2)^{-1} Z^{-1} dZ Y^{-1} d\overline{Z} \overline{Z}^{-1} \\
&\quad + {}^tW (V_1 + U_2)(\partial_2)^{-1} \overline{Z}^{-1} d\overline{Z} Y^{-1} dZ Z^{-1} :
\end{aligned}$$

Since

$$\begin{aligned}
(V_1 + U_2)(\partial_2)^{-1} &= U + V {}_1 (\partial_2)^{-1} \\
&= W + V f iE_n + {}_1 (\partial_2)^{-1} g \\
&= \overline{W} - V f iE_n - {}_1 (\partial_2)^{-1} g;
\end{aligned}$$

we have

$$\begin{aligned}
 (Z; \bar{Z}) &= \bar{Z}^{-1} iE_n \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}^{-1} {}^t V W - V (iE_n + \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}^{-1}) \\
 &\quad Z^{-1} dZ - Y^{-1} d\bar{Z} \\
 &= \bar{Z}^{-1} iE_n \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}^{-1} {}^t V W Z^{-1} dZ - Y^{-1} d\bar{Z} \\
 &= \bar{Z}^{-1} iE_n \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}^{-1} {}^t V V iE_n + \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}^{-1} \\
 &\quad Z^{-1} dZ - Y^{-1} d\bar{Z} :
 \end{aligned}$$

By the way, according to (2.9), we obtain

$$\begin{aligned}
 \bar{Z}^{-1} iE_n \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}^{-1} &= \begin{pmatrix} 1 & i \\ 0 & 1 \end{pmatrix} iE_n \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}^{-1} \\
 &= \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}^{-1} = Y^{-1}
 \end{aligned}$$

and

$$iE_n + \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}^{-1} Z^{-1} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}^{-1} = Y^{-1}.$$

Therefore

$$(Z; \bar{Z}) = Y^{-1} {}^t V V Y^{-1} dZ - Y^{-1} d\bar{Z} :$$

Hence (a) = (a) and

$$\begin{aligned}
 (b) + (c) + (d) &= (Z; \bar{W}) + (\bar{Z}; W) + (W; \bar{W}) + (Z; \bar{Z}) \\
 &= (b) + (c) + (d):
 \end{aligned}$$

This implies that the metric

$$ds_{n,m;A,B}^2 = A(a) + B(b) + (c) + (d)$$

is invariant under the action (1.2) of G^n :

Consequently $ds_{n,m;A,B}^2$ is invariant under the action (1.2) of the Jacobi group G^J : In particular, for $(Z; W) = (iE_n; 0)$; we have

$$\begin{aligned}
 ds_{n,m;A,B}^2 &= A \left(dZ d\bar{Z} + B {}^t(dW) d\bar{W} \right) \\
 &= A \left(\sum_{k=1}^n (dx_k^2 + dy_k^2) + 2 \sum_{1 \leq k < l \leq n} (dx_k^2 + dy_k^2) \right) \\
 &\quad + B \sum_{1 \leq k \leq m; 1 \leq l \leq n} (du_{kl}^2 + dv_{kl}^2);
 \end{aligned}$$

which is clearly positive definite. Since G^J acts on $H_{n,m}$ transitively, $ds_{n,m;A,B}^2$ is positive definite everywhere in $H_{n,m}$: This completes the proof of Theorem 1.1.

Remark 2.1. The scalar curvature of the Siegel-Jacobi space $(H_{n,m}; ds_{n,m;A,B}^2)$ is constant because of the transitive group action of G^J on $H_{n,m}$: In the special case $n = m = 1$ and

$A = B = 1$, by a direct computation, we see that the scalar curvature of $(H_{1,1}; ds_{1,1;1}^2)$ is 3:

3. Proof of Theorem 1.2

If $(Z; W) = g(Z; W)$ with $g = \begin{pmatrix} A & B \\ C & D \end{pmatrix}; (\cdot; \cdot) = 2G^J$; we can see easily that

$$(3.1) \quad \frac{\partial}{\partial Z} = (CZ + D)^t (CZ + D) \frac{\partial}{\partial Z} + (CZ + D)^t C^t W + C^t D^t \frac{\partial}{\partial W}$$

and

$$(3.2) \quad \frac{\partial}{\partial W} = (CZ + D) \frac{\partial}{\partial W} :$$

For brevity, we put

$$\begin{aligned} (\cdot) &:= 4 \quad Y^t Y \frac{\partial}{\partial Z} \frac{\partial}{\partial Z} ; \\ (\cdot) &:= 4 \quad Y \frac{\partial}{\partial W}^t \frac{\partial}{\partial W} ; \\ (\cdot) &:= 4 \quad V Y^{-1} V^t Y \frac{\partial}{\partial W} \frac{\partial}{\partial W} ; \\ (\cdot) &:= 4 \quad V^t Y \frac{\partial}{\partial Z} \frac{\partial}{\partial W} \end{aligned}$$

and

$$(\cdot) := 4 \quad V^t V^t Y \frac{\partial}{\partial W} \frac{\partial}{\partial Z} :$$

We also set

$$\begin{aligned} (\cdot) &:= 4 \quad Y^t Y \frac{\partial}{\partial Z} \frac{\partial}{\partial Z} ; \\ (\cdot) &:= 4 \quad Y \frac{\partial}{\partial W}^t \frac{\partial}{\partial W} ; \\ (\cdot) &:= 4 \quad V Y^{-1} V^t Y \frac{\partial}{\partial W} \frac{\partial}{\partial W} ; \\ (\cdot) &:= 4 \quad V^t Y \frac{\partial}{\partial Z} \frac{\partial}{\partial W} \end{aligned}$$

and

$$(\cdot) := 4 \quad V^t V^t Y \frac{\partial}{\partial W} \frac{\partial}{\partial Z} :$$

We need the following lemma for the proof of Theorem 1.2. H. Maass [8] observed the following useful fact.

Lemma 3.1. (a) Let A be an $n \times k$ matrix and let B be a $k \times n$ matrix. Assume that the entries of A commute with the entries of B . Then $(AB)^t = (BA)^t$:

(b) Let A be an $m \times n$ matrix and B an $n \times l$ matrix. Assume that the entries of A commute with the entries of B . Then ${}^t(AB) = {}^tB {}^tA$:

(c) Let A, B and C be a $k \times l$, an $n \times m$ and an $m \times l$ matrix respectively. Assume that the entries of A commute with the entries of B . Then

$${}^t(A {}^t(BC)) = B {}^t(A {}^tC):$$

Proof. The proof follows immediately from the direct computation.

Now we are ready to prove Theorem 1.2. First of all, we shall prove that ${}_{n,m} \mathcal{A}_B$ is invariant under the action of the generators $t(b; \cdot; \cdot; \cdot); g(h)$ and ${}_{n,m}$:

Case I. $g = t(b; \cdot; \cdot; \cdot) = \begin{pmatrix} E_n & b \\ 0 & E_n \end{pmatrix}; (\cdot; \cdot; \cdot)$ with $b = {}^t b$ real.

In this case, we have

$$Y = Y; \quad V = V + Y$$

and

$$\frac{\partial}{\partial Z} = \frac{\partial}{\partial Z} - {}^t t - {}^t \frac{\partial}{\partial W} \quad \text{and} \quad \frac{\partial}{\partial W} = \frac{\partial}{\partial W}:$$

Using Lemma 3.1, we obtain

$$\begin{aligned} () &= () - Y {}^t Y \frac{\partial}{\partial Z} - \frac{\partial}{\partial W} \\ &\quad - Y {}^t t Y \frac{\partial}{\partial W} - \frac{\partial}{\partial Z} + Y {}^t t Y \frac{\partial}{\partial W} - \frac{\partial}{\partial W} \\ () &= Y \frac{\partial}{\partial W} - \frac{\partial}{\partial W} = (); \\ () &= () + {}^t V {}^t Y \frac{\partial}{\partial W} - \frac{\partial}{\partial W} \\ &\quad + V {}^t t Y \frac{\partial}{\partial W} - \frac{\partial}{\partial W} + Y {}^t t Y \frac{\partial}{\partial W} - \frac{\partial}{\partial W}; \\ () &= () + Y {}^t Y \frac{\partial}{\partial Z} - \frac{\partial}{\partial W} \\ &\quad - V {}^t t Y \frac{\partial}{\partial W} - \frac{\partial}{\partial W} + Y {}^t t Y \frac{\partial}{\partial W} - \frac{\partial}{\partial W} \end{aligned}$$

and

$$\begin{aligned} () &= () + Y {}^t t Y \frac{\partial}{\partial W} - \frac{\partial}{\partial Z} \\ &\quad - {}^t V {}^t Y \frac{\partial}{\partial W} - \frac{\partial}{\partial W} - Y {}^t t Y \frac{\partial}{\partial W} - \frac{\partial}{\partial W}: \end{aligned}$$

Thus $() = ()$ and

$$() + () + () + () = () + () + () + ():$$

Hence

$${}_{n,m} \mu_{A,B} = \frac{4}{B} () + \frac{4}{A} () + () + () + ()$$

is invariant under the action of all $t(b; ; ;)$:

$$\text{Case II. } g = g(h) = \begin{pmatrix} t_h & 0 \\ 0 & h^{-1} \end{pmatrix}; (0; 0; 0) \quad \text{with } h \in GL(n; R).$$

In this case, we have

$$Y = t_h Y h^{-1} \quad V = V h$$

and

$$\frac{\partial}{\partial Z} = h^{-1} t_h^{-1} \frac{\partial}{\partial Z}; \quad \frac{\partial}{\partial W} = h^{-1} \frac{\partial}{\partial W}:$$

According to Lemma 3.1, we see that each of $(); (); (); ()$ and $()$ is invariant under the action of all $g(h)$ with $h \in GL(n; R)$: Therefore ${}_{n,m} \mu_{A,B}$ is invariant under the action of all $g(h)$ with $h \in GL(n; R)$:

$$\text{Case III. } g = g_n = \begin{pmatrix} 0 & E_n \\ E_n & 0 \end{pmatrix}; (0; 0; 0) :$$

In this case, we have

$$Z = Z^{-1} \quad \text{and} \quad W = W Z^{-1}:$$

We set

$$_1 := \operatorname{Re} Z^{-1} \quad \text{and} \quad _2 := \operatorname{Im} Z^{-1}:$$

Then we obtain the relations (2.5)–(2.9). From (2.6), we have the relation

$$(3.3) \quad \overline{Z} = Z^{-1} Y:$$

It follows from the relation (2.3) that

$$(3.4) \quad Y = \overline{Z}^{-1} Y Z^{-1} = Z^{-1} Y \overline{Z}^{-1} = _2:$$

From (2.9), we obtain

$$(3.5) \quad _1 _2^{-1} _1 = Y^{-1} _2:$$

According to (3.1) and (3.2), we have

$$(3.6) \quad \frac{\partial}{\partial Z} = Z^{-t} Z \frac{\partial}{\partial Z} + Z^{-t} t_W^{-t} \frac{\partial}{\partial W}$$

and

$$(3.7) \quad \frac{\partial}{\partial W} = Z \frac{\partial}{\partial W}:$$

From (2.6), (3.3) and Lemma 3.1, we obtain

$$\begin{aligned} () &= () \quad U _2 \overline{Z}^{-t} Y \frac{\partial}{\partial Z} \frac{\partial}{\partial W} + i \quad V _2 \overline{Z}^{-t} Y \frac{\partial}{\partial Z} \frac{\partial}{\partial W} \\ &\quad Z _2 t_U^{-t} Y \frac{\partial}{\partial W} \frac{\partial}{\partial Z} + i \quad Z _2 t_V^{-t} Y \frac{\partial}{\partial W} \frac{\partial}{\partial Z} \\ &\quad W _2 t_U^{-t} Y \frac{\partial}{\partial W} \frac{\partial}{\partial W} + \quad W _2 t_V^{-t} Y \frac{\partial}{\partial W} \frac{\partial}{\partial W} : \end{aligned}$$

From the relation (3.4), we see $() = ()$. According to (3.3), (3.5) and Lemma 3.1, we obtain

$$() = () + (V_2 {}^t V - V_1 {}^t U - U_1 {}^t V - U_2 {}^t U) {}^t Y \frac{\partial}{\partial \bar{W}} - \frac{\partial}{\partial W} :$$

Using the relation (3.3) and Lemma 3.1, we finally obtain

$$\begin{aligned} () = & V_1 \bar{Z} {}^t Y \frac{\partial}{\partial \bar{Z}} - \frac{\partial}{\partial W} + U_2 \bar{Z} {}^t Y \frac{\partial}{\partial \bar{Z}} - \frac{\partial}{\partial W} \\ & + V_1 {}^t \bar{W} {}^t Y \frac{\partial}{\partial \bar{W}} - \frac{\partial}{\partial W} + U_2 {}^t \bar{W} {}^t Y \frac{\partial}{\partial \bar{W}} - \frac{\partial}{\partial W} \end{aligned}$$

and

$$\begin{aligned} () = & Z_1 {}^t V {}^t Y \frac{\partial}{\partial \bar{W}} - \frac{\partial}{\partial Z} + Z_2 {}^t U {}^t Y \frac{\partial}{\partial \bar{W}} - \frac{\partial}{\partial Z} \\ & + W_1 {}^t V {}^t Y \frac{\partial}{\partial \bar{W}} - \frac{\partial}{\partial W} + W_2 {}^t U {}^t Y \frac{\partial}{\partial \bar{W}} - \frac{\partial}{\partial W} : \end{aligned}$$

Using the fact $Z^{-1} = Z_1 + i Z_2$, we can show that

$$() + () + () + () = () + () + () + () :$$

Hence

$$\Delta_{n,m;A,B} = \frac{4}{B} () + \frac{4}{A} {}^n () + () + () + ()^0$$

is invariant under the action of Δ_n .

Consequently $\Delta_{n,m;A,B}$ is invariant under the action (1.2) of G^J . In particular, for $(Z; W) = (iE_n; 0)$; the differential operator $\Delta_{n,m;A,B}$ coincides with the Laplacian for the metric $ds_{n,m;A,B}^2$. It follows from the invariance of $\Delta_{n,m;A,B}$ under the action (1.2) and the transitivity of the action of G^J on $H_{n,m}$ that $\Delta_{n,m;A,B}$ is the Laplacian of $(H_{n,m}; ds_{n,m;A,B}^2)$: The invariance of the differential form dv follows from the fact that the following differential form

$$(\det Y)^{(n+1)} [dX] \wedge [dY]$$

is invariant under the action (1.1) of $Sp(n; R)$ (cf. [13], p.130).

4. Remark on Spectral Theory of $\Delta_{n,m;A,B}$ on Siegel-Jacobi Space

Before we describe a fundamental domain for the Siegel-Jacobi space, we review the Siegel's fundamental domain for the Siegel upper half plane.

We let

$$P_n = \{ Y \in R^{(n,m)} \mid Y = {}^t Y > 0 \}$$

be an open cone in $R^{n(n+1)/2}$. The general linear group $GL(n; R)$ acts on P_n transitively by

$$h \cdot Y = h Y {}^t h; \quad h \in GL(n; R); \quad Y \in P_n :$$

Thus P_n is a symmetric space diffeomorphic to $GL(n; R)/O(n)$: We let

$$GL(n; Z) = \{ h \in GL(n; R) \mid h \text{ is integral} \}$$

be the discrete subgroup of $GL(n; \mathbb{R})$.

The fundamental domain R_n for $GL(n; \mathbb{Z}) \backslash P_n$ which was found by H. Minkowski [10] is defined as a subset of P_n consisting of $Y = (y_{ij}) \in P_n$ satisfying the following conditions (M.1)–(M.2) (cf. [9] p.123):

(M.1) $aY^t a = y_{kk}$ for every $a = (a_i) \in \mathbb{Z}^n$ in which a_k and a_n are relatively prime for $k = 1, 2, \dots, n$.

(M.2) $y_{k, k+1} = 0$ for $k = 1, \dots, n-1$.

We say that a point of R_n is Minkowski reduced or simply M-reduced.

Siegel [12] determined a fundamental domain F_n for $\Gamma_n \backslash H_n$; where $\Gamma_n = Sp(n; \mathbb{Z})$ is the Siegel modular group of degree n . We say that $Y = X + iY \in H_n$ with X, Y real is Siegel reduced or S-reduced if it has the following three properties:

(S.1) $\det(\operatorname{Im}(Y)) = \det(\operatorname{Im}(Y'))$ for all $Y' \in \Gamma_n Y$;

(S.2) $Y = \operatorname{Im}(Y)$ is M-reduced, that is, $Y \in R_n$;

(S.3) $|x_{ij}| \leq \frac{1}{2}$ for $1 \leq i, j \leq n$; where $X = (x_{ij})$.

F_n is defined as the set of all Siegel reduced points in H_n : Using the highest point method, Siegel [12] proved the following (F.1)–(F.3) (cf. [9], p.169):

(F.1) $\Gamma_n \backslash F_n = H_n$; i.e., $H_n = \bigcup_{\gamma \in \Gamma_n} \gamma F_n$;

(F.2) F_n is closed in H_n ;

(F.3) F_n is connected and the boundary of F_n consists of a finite number of hyperplanes.

The metric ds_n^2 given by (1.3) induces a metric $ds_{F_n}^2$ on F_n . Siegel [12] computed the volume of F_n

$$\operatorname{vol}(F_n) = 2^{-\sum_{k=1}^n k} \prod_{k=1}^n \frac{\Gamma(k)}{(2k)};$$

where $\Gamma(s)$ denotes the Gamma function and $\zeta(s)$ denotes the Riemann zeta function. For instance,

$$\operatorname{vol}(F_1) = \frac{1}{3}; \quad \operatorname{vol}(F_2) = \frac{1}{270}; \quad \operatorname{vol}(F_3) = \frac{1}{127575}; \quad \operatorname{vol}(F_4) = \frac{1}{200930625};$$

Let f_{kl} ($1 \leq k \leq m; 1 \leq l \leq n$) be the $m \times n$ matrix with entry 1 where the k -th row and the l -th column meet, and all other entries 0. For an element $Y \in H_n$, we set for brevity

$$h_{kl}(Y) := f_{kl}; \quad 1 \leq k \leq m; 1 \leq l \leq n;$$

For each $Y \in F_n$; we define a subset P of $\mathbb{C}^{(m, n)}$ by

$$P = \left\{ \begin{pmatrix} X^m & X^n \\ \vdots & \vdots \end{pmatrix} : \begin{matrix} \sum_{k=1}^m \sum_{j=1}^n f_{kj} X^k X^j + \sum_{k=1}^m \sum_{j=1}^n h_{kj}(Y) X^k X^j = 0 \\ \vdots \\ \sum_{k=1}^m \sum_{j=1}^n h_{kj}(Y) X^k X^j = 0 \end{matrix} \right\};$$

For each $Y \in F_n$; we define the subset D of $H_n \times \mathbb{C}^{(m, n)}$ by

$$D := \left\{ (Y; Z) \in H_n \times \mathbb{C}^{(m, n)} : Z \in P \right\};$$

We define

$$F_{n, m} := \bigcup_{Y \in F_n} D;$$

Theorem 4.1. Let

$$\Gamma_{n,m} = \text{Sp}(n; \mathbb{Z}) \cap H_{\mathbb{Z}}^{(n,m)}$$

be the discrete subgroup of G_n^J , where

$$H_{\mathbb{Z}}^{(n,m)} = \left\{ \begin{pmatrix} x & y \\ z & w \end{pmatrix} \in H_{\mathbb{R}}^{(n,m)} \mid x, y, z, w \text{ are integral} \right\} :$$

Then $F_{\Gamma_{n,m}}$ is a fundamental domain for $\Gamma_{n,m} \backslash H_{\mathbb{R}}^{(n,m)}$:

Proof. The proof can be found in [20].

In the case $n = m = 1$, R. Berndt [2] introduced the notion of Maass-Jacobi forms. Now we generalize this notion to the general case.

Definition 4.1. For brevity, we set $\Gamma_{n,m} = \Gamma_{n,m;1,1}$ (cf. Theorem 1.2). Let

$$\Gamma_{n,m} = \text{Sp}(n; \mathbb{Z}) \cap H_{\mathbb{Z}}^{(n,m)}$$

be the discrete subgroup of G_n^J , where

$$H_{\mathbb{Z}}^{(n,m)} = \left\{ \begin{pmatrix} x & y \\ z & w \end{pmatrix} \in H_{\mathbb{R}}^{(n,m)} \mid x, y, z, w \text{ are integral} \right\} :$$

A smooth function $f : H_{\Gamma_{n,m}} \rightarrow \mathbb{C}$ is called a Maass-Jacobi form on $H_{\Gamma_{n,m}}$ if f satisfies the following conditions (MJ1)–(MJ3):

- (MJ1) f is invariant under $\Gamma_{n,m}$;
- (MJ2) f is an eigenfunction of the Laplacian $\Delta_{\Gamma_{n,m}}$.
- (MJ3) f has a polynomial growth, that is, there exist a constant $C > 0$ and a positive integer N such that

$$|f(X + iY; Z)| \leq C |p(Y)|^N \quad \text{as } \det Y \rightarrow 1;$$

where $p(Y)$ is a polynomial in $Y = (Y_{ij})$:

It is natural to propose the following problems.

Problem A : Construct Maass-Jacobi forms.

Problem B : Find all the eigenfunctions of $\Delta_{\Gamma_{n,m}}$:

We consider the simple case $n = m = 1$: A metric $ds_{1,1}^2$ on $H_1 \rightarrow \mathbb{C}$ given by

$$ds_{1,1}^2 = \frac{y + v^2}{y^3} (dx^2 + dy^2) + \frac{1}{y} (du^2 + dv^2) + \frac{2v}{y^2} (dx du + dy dv)$$

is a G_1^J -invariant Kähler metric on $H_1 \rightarrow \mathbb{C}$. Its Laplacian $\Delta_{1,1}$ is given by

$$\begin{aligned} \Delta_{1,1} = & y^2 \left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} \right) \\ & + (y + v^2) \left(\frac{\partial^2}{\partial u^2} + \frac{\partial^2}{\partial v^2} \right) \\ & + 2yv \left(\frac{\partial^2}{\partial x \partial u} + \frac{\partial^2}{\partial y \partial v} \right) : \end{aligned}$$

We provide some examples of eigenfunctions of $\Delta_{1,1}$.

(1) $h(x; y) = y^{\frac{1}{2}} K_{s-\frac{1}{2}}(2\sqrt{xy}) e^{2iax}$ ($s \in \mathbb{C}$; $a \in \mathbb{R}$) with eigenvalue $s(s-1)$: Here

$$K_s(z) := \frac{1}{2} \int_0^z t^{-1} \exp\left(-\frac{z}{2}(t+t^{-1})\right) t^{s-1} dt;$$

where $\operatorname{Re} z > 0$:

(2) y^s ; $y^s x$; $y^s u$ ($s \in \mathbb{C}$) with eigenvalue $s(s-1)$:

(3) $y^s v$; $y^s uv$; $y^s xv$ with eigenvalue $s(s+1)$:

(4) x ; y ; u ; v ; xv ; uv with eigenvalue 0.

(5) All mass wave forms.

We fix two positive integers m and n throughout this section.

For an element $\gamma \in H_n$, we set

$$L_\gamma := Z^{(m,n)} + \gamma Z^{(m,n)}$$

It follows from the positivity of $\operatorname{Im} \gamma$ that the elements $f_{kl}; h_{kl}(\gamma)$ ($1 \leq k \leq m$; $1 \leq l \leq n$) of L_γ are linearly independent over $\mathbb{R}$. Therefore L_γ is a lattice in $\mathbb{C}^{(m,n)}$ and the set $\{f_{kl}; h_{kl}(\gamma)\}_{1 \leq k \leq m; 1 \leq l \leq n}$ forms an integral basis of L_γ . We see easily that if γ is an element of H_n , the period matrix $\gamma = (I_n; \gamma)$ satisfies the Riemann conditions (RC.1) and (RC.2):

$$(RC.1) \quad J_n^t = 0;$$

$$(RC.2) \quad \frac{1}{i} J_n^t \gamma^t > 0.$$

Thus the complex torus $A_\gamma := \mathbb{C}^{(m,n)} / L_\gamma$ is an abelian variety.

It might be interesting to investigate the spectral theory of the Laplacian $\Delta_{n,m}$ on a fundamental domain $F_{n,m}$. But this work is very complicated and difficult at this moment. It may be that the first step is to develop the spectral theory of the Laplacian Δ_n on the abelian variety A : The second step will be to study the spectral theory of the Laplacian Δ_n (see (1.4)) on the moduli space $\mathcal{M}_n$ of principally polarized abelian varieties of dimension n . The final step would be to combine the above steps and more works to develop the spectral theory of the Laplacian $\Delta_{n,m}$ on $F_{n,m}$: Mass-Jacobi forms play an important role in the spectral theory of $\Delta_{n,m}$ on $F_{n,m}$: Here we deal only with the spectral theory on $L^2(A_\gamma)$:

We fix an element $\gamma = X + iY$ of H_n with $X = \operatorname{Re} \gamma$ and $Y = \operatorname{Im} \gamma$: For a pair $(A; B)$ with $A; B \in \mathbb{Z}^{(m,n)}$, we define the function $E_{A,B} : \mathbb{C}^{(m,n)} \rightarrow \mathbb{C}$ by

$$E_{A,B}(Z) = e^{2i((tAU) + (B-AX)Y^{-1}tV)};$$

where $Z = U + iV$ is a variable in $\mathbb{C}^{(m,n)}$ with real $U; V$.

Lemma 4.1. For any $A; B \in \mathbb{Z}^{(m,n)}$, the function $E_{A,B}$ satisfies the following functional equation

$$E_{A,B}(Z + \gamma + \gamma') = E_{A,B}(Z); \quad Z \in \mathbb{C}^{(m,n)}$$

for all $\gamma; \gamma' \in \mathbb{Z}^{(m,n)}$: Thus $E_{A,B}$ can be regarded as a function on A_γ :

Proof. We write $Z = X + iY$ with real X, Y : For any $(U, V) \in \mathbb{Z}^{(m, n)}$, we have

$$\begin{aligned} E_{A, B}(Z + U + i(V + Y)) &= E_{A, B}((U + X + iY) + i(V + Y)) \\ &= e^{2\pi i \text{tr} (A(U + X + iY) + (B - AX)Y^{-1} \text{tr}(V + Y))} g \\ &= e^{2\pi i \text{tr} (AU + AX + AY + (B - AX)Y^{-1} \text{tr}(V + B^{-1}AX))} g \\ &= e^{2\pi i \text{tr} (AU) + (B - AX)Y^{-1} \text{tr} V} g \\ &= E_{A, B}(Z): \end{aligned}$$

Here we used the fact that A and B^{-1} are integral.

Lemma 4.2. The metric

$$ds^2 = (\text{Im } Y)^{-1} \text{tr}(dZ) d\bar{Z}$$

is a Kähler metric on A invariant under the action (5.15) of $J = \text{Sp}(n; \mathbb{Z}) \ltimes H_{\mathbb{Z}}^{(m, n)}$ on (Z) with fixed. Its Laplacian of ds^2 is given by

$$= (\text{Im } Y)^{-1} \frac{\partial}{\partial Z} \text{tr} \frac{\partial}{\partial \bar{Z}} :$$

Proof. The proof can be found [20].

We let $L^2(A)$ be the space of all functions $f: A \rightarrow \mathbb{C}$ such that

$$\|f\|^2 = \int_A |f(Z)|^2 dv ;$$

where dv is the volume element on A normalized so that $\int_A dv = 1$: The inner product (f, g) on the Hilbert space $L^2(A)$ is given by

$$(f, g) = \int_A f(Z) \overline{g(Z)} dv ; \quad f, g \in L^2(A) :$$

Theorem 4.2. The set $\{E_{A, B} \mid A, B \in \mathbb{Z}^{(m, n)}\}$ is a complete orthonormal basis for $L^2(A)$. Moreover we have the following spectral decomposition of Δ :

$$L^2(A) = \bigoplus_{A, B \in \mathbb{Z}^{(m, n)}} \mathbb{C} E_{A, B} :$$

Proof. The complete proof can be found in [20]

References

- [1] R. Berndt, Sur l'arithmétique du corps des fonctions elliptiques de niveau N , Séminaire de Théorie des nombres de Paris 1982-83, Birkhäuser, Boston (1984), 21-32.
- [2] R. Berndt and R. Schmidt, Elements of the Representation Theory of the Jacobi Group, Birkhäuser, 1998.
- [3] M. Eichler and D. Zagier, The Theory of Jacobi Forms, Progress in Math., 55, Birkhäuser, Boston, Basel and Stuttgart, 1985.
- [4] E. Freitag, Siegel'sche Modulformen, Grundlehren der Math. Wissenschaften, Springer-Verlag, 1983.
- [5] M. Kashiwara and M. Vergne, On the Segal-Shale-Weil Representations and Harmonic Polynomials, Invent. Math. 44 (1978), 1-47.

- [6] J. Kramer, A geometrical approach to the theory of Jacobi forms, *Compositio Math.* 79 (1991), 1{19.
- [7] J. Kramer, An arithmetical theory of Jacobi forms in higher dimensions, *J. reine angew. Math.* 458 (1995), 157{182.
- [8] H. Maass, Die Differentialgleichungen in der Theorie der Siegelschen Modulfunktionen, *Math. Ann.* 126 (1953), 44{68.
- [9] H. Maass, Siegel modular functions and Dirichlet series, *Lecture Notes in Math.* 216, Springer-Verlag, Berlin and New York, 1971.
- [10] Minkowski, H.: *Gesammelte Abhandlungen*. Chelsea, New York (1967).
- [11] B. Runge, Theta functions and Siegel-Jacobi functions, *Acta Math.* 175 (1995), 165{196.
- [12] C. L. Siegel, Symplectic Geometry, *Gesammelte Abhandlungen*, Band II, Springer-Verlag (1966), 274{359.
- [13] C. L. Siegel, *Topics in Complex Function Theory : Abelian Functions and Modular Functions of Several Variables*, vol. III, Wiley-Interscience, 1973.
- [14] J.-H. Yang, The Siegel-Jacobi Operator, *Abh. Math. Sem. Univ. Hamburg* 63 (1993), 135{146.
- [15] J.-H. Yang, Remarks on Jacobi forms of higher degree, *Proc. of the 1993 Workshop on Automorphic Forms and Related Topics*, the Pyungsan Institute for Mathematical Sciences, Seoul (1993), 33{58.
- [16] J.-H. Yang, Singular Jacobi Forms, *Trans. of American Math. Soc.* 347, No. 6 (1995), 2041{2049.
- [17] J.-H. Yang, Construction of Modular Forms from Jacobi Forms, *Canadian J. of Math.* 47 (1995), 1329{1339.
- [18] J.-H. Yang, A geometrical theory of Jacobi forms of higher degree, *Proceedings of Symposium on Hodge Theory and Algebraic Geometry* (edited by Tadao Oda), Sendai, Japan (1996), 125-147 or *Kyungpook Math. J.* 40 (2), 209{237 (2000) or *arXiv math.NT/0602267*.
- [19] J.-H. Yang, A Note on Maass-Jacobi Forms, *Kyungpook Math. J.* 43 (2003), 547{566.
- [20] J.-H. Yang, A note on a fundamental domain for Siegel-Jacobi space, *Houston Journal of Mathematics*, Vol. 32, No. 3 (2006), 701{712.
- [21] C. Ziegler, Jacobi Forms of Higher Degree, *Abh. Math. Sem. Hamburg* 59 (1989), 191{224.

Department of Mathematics, Inha University, Incheon 402-751, Korea

E-mail address: jhyang@inha.ac.kr