

# Lattices generated by skeletons of reflexive polytopes

Christian Haase

Mathematics Department, Duke University,

Durham, NC 27708, USA

email: haase@math.duke.edu

Benjamin Nill

Mathematisches Institut, Universität Tübingen

Auf der Morgenstelle 10, 72076 Tübingen, Germany

email: nill@algebra.mathematik.uni-tuebingen.de

## Abstract

Lattices generated by lattice points in skeletons of reflexive polytopes are essential in determining the fundamental group and integral cohomology of Calabi-Yau hypersurfaces. Here we prove that the lattice generated by all lattice points in a reflexive polytope is already generated by lattice points in codimension two faces. This answers a question of J. Morgan.

## 1 Introduction and main result

Since its introduction by Batyrev in [Bat94] dual pairs of reflexive polytopes have been used to successfully construct mirror pairs of smooth Calabi-Yau varieties as resolutions of non-degenerate anti-canonical hypersurfaces in Gorenstein toric Fano varieties, see, e.g., [CK99].

Recall that a reflexive polytope is an  $n$ -dimensional lattice polytope  $P \subset M_{\mathbb{R}} = M \otimes_{\mathbb{Z}} \mathbb{R}$  for a lattice  $M = \mathbb{Z}^n$  such that  $P$  contains the origin in its interior and the dual polytope  $P^{\vee}$  is also a lattice polytope with respect to the dual lattice  $N = M^{\vee} = \text{Hom}_{\mathbb{Z}}(M; \mathbb{Z})$ . There is an associated pair  $P \in \mathcal{P}_M, P^{\vee} \in \mathcal{P}_N$  of Gorenstein toric Fano

varieties, where  $\mathcal{P}$  is just the fan of cones over the faces of  $P$ , see, e.g., [Bat94, Nil05].

For a given reflexive polytope  $P \in \mathcal{M}_R$  there exist infinitely many choices of sublattices of  $M$  such that  $P$  is reflexive with respect to this lattice. Obviously the minimal one of these is the lattice  $\Lambda_0$  generated by the vertices of  $P$ . More generally we define

$\Lambda_k :=$  lattice generated by all lattice points in the  $k$ -skeleton of  $P$ ,

where the  $k$ -skeleton is the union of  $k$ -dimensional faces.

Due to [Nil05, Lemma 1.17] and [Ful93, 3.2] we see that for  $k = 0, 1, 2$  the quotient group  $M/\Lambda_k$  equals the fundamental group  $\pi_1$  of the union of all  $k+1$ -codimensional torus orbits in  $P_{P, M}$ .

Returning to the relevance of these lattices in mirror symmetry Batyrev and Kreuzer [BK05, Cor.1.9] recently proved that for  $n = 3, 4$  the quotient group  $M/\Lambda_{n-2}$  is precisely the fundamental group  $\pi_1(X)$  of a projective crepant resolution  $X$  of an  $(n-1)$ -dimensional non-degenerate anticanonical Calabi-Yau hypersurface in  $P_{P, M}$ . Moreover for  $n = 4$  they showed that the torsion group of  $H^2(X; \mathbb{Z})$  equals  $\text{Hom}(M/\Lambda_{n-2}; \mathbb{Q}/\mathbb{Z})$  and the torsion group of  $H^3(X; \mathbb{Z})$  is isomorphic to  $\text{Hom}(\wedge^2 M/\Lambda_{n-3}; \mathbb{Q}/\mathbb{Z})$ , cf. [BK05, Cor.3.9]. Mirror symmetry should exchange these torsion parts of integral cohomology of  $X$  and a mirror  $X^\vee$ . Hence, this yields the surprising isomorphism  $M/\Lambda_2 = \wedge^2 N/N^\vee_{\Lambda_1}$ , where  $\Lambda_1$  is the sublattice of  $N$  generated by lattice points in edges of  $P$ . Using the classification of Kreuzer and Skarke [KS00] they confirmed this conjecture.

These results and open questions motivate further investigation of the lattices  $\Lambda_k$  for an  $n$ -dimensional reflexive polytope  $P$ . Here we first note the following two observations:

Since there are no non-zero lattice points in the interior of  $P$ , we have  $\Lambda_{n-1} = \Lambda_n$ .

If there exists a crepant toric resolution of  $P_{P, M}$ , then the boundary  $\partial P$  contains a lattice basis, hence  $\Lambda_{n-1} = M$ . This holds for  $n = 2, 3$ .

For  $n = 2$  there are precisely three isomorphism classes of reflexive polytopes with  $\Lambda_{n-2} = \Lambda_0 \notin M$ , here  $\Lambda_0$  has index 2, 2, and 3:




Figure 1: Reflexive polygons with  $\Lambda_{n-2} \notin M$

As noted in [BK 05, Remark 1.10] for  $n = 3$  a non-degenerate Calabi-Yau hypersurface  $X$  arising from a crepant resolution of  $P_{\mathbb{P}^3, M}$  is a smooth K3-surface, hence simply-connected. So the fundamental group  $\pi_1(X) = M = \pi_2$  is trivial, i.e.,  $\pi_2 = \pi_1$ . The authors did not know of a combinatorial proof of this result in the literature.

At PCM I 2004, John Morgan (for a mirror-symmetric conjecture concerning integral variations of Hodge structure see Conjecture 2.9 in [DM 05]) asked the first author for a combinatorial proof that  $\pi_2 = \pi_1$  would also hold for  $n = 4$ . This was confirmed by the classification of all four-dimensional reflexive polytopes [KS00]. John Morgan, and independently Batyrev and Kreuzer [BK 05], have listed all 16 isomorphism classes of four-dimensional reflexive polytopes with  $\pi_2 \in M$ . However in contrast to  $n = 3$  not even an algebraic geometric proof seemed to be known.

The goal of this article is to provide a purely convex-geometric proof valid in arbitrary dimension  $n > 2$ . (Figure 1 shows all "counter-examples" for  $n = 2$ ).

**Theorem 1.1.** If  $n \geq 3$ , then  $\pi_2 = \pi_1$ .

The main idea of the proof is to show that any lattice point inside a facet can be obtained from lattice points on lower dimensional faces. For this we use a partial addition on the set of lattice points in  $P$ , observed by the second author in [Nil05, Prop. 4.1]. In [Nil05] that method has been applied to investigate the set of lattice points inside the facets, since this is precisely the set of Demazure roots for the automorphism group of the corresponding Gorenstein toric Fano variety  $P_{\mathbb{P}^n, N}$ .

Acknowledgments: Work of the first author was partially supported by NSF grant DMS-0200740. The second author would like to thank Duke University for hospitality and support.

## 2 Proof of theorem

Let  $P \subset M_{\mathbb{R}}$  be an  $n$ -dimensional reflexive polytope. We denote by  $h; i$  the non-degenerate symmetric pairing of the dual lattices  $M; N$ .

For the proof we will use the following notions.

**Definition 2.1.**

We denote by  $R$  the set of Demazure roots of  $P$ , i.e., the set of lattice points in the interior of facets.

For  $x \in R$  we denote by  $F_x$  the unique facet of  $P$  containing  $x$ , and by  $\nu_x$  the unique inner normal of  $F_x$ , i.e.,  $\nu_x = \nu_{F_x} \in N$  s.t.  $\langle \nu_x, x \rangle = 1$ .

A pair of roots  $x, y \in R$  is called orthogonal, if  $\langle \nu_x, y \rangle = 0 = \langle \nu_y, x \rangle$ . In particular  $x \notin F_y \iff y \notin F_x$ . In this case  $x + y \in M \setminus F_x \setminus F_y$  (see the next result).

The following lemma is the engine running our proof.

**Lemma 2.2** ([Nil04, Lemma 3.5 & Corollary 3.6]). For  $x, y \in P \setminus M$  exactly one of the following is true. Either  $x$  and  $y$  belong to a common facet, or  $x + y = 0$ , or  $x + y \in P$ . In the last case, there is a unique  $z(x, y) = ax + by \in P$  with  $a, b \in \mathbb{Z}_{>0}$ . We have  $a = 1$  or  $b = 1$ , and  $x$  and  $z(x, y)$ , respectively  $y$  and  $z(x, y)$ , belong to a common facet. Furthermore, if  $x \in R$ , then  $a = \langle \nu_x, y \rangle + 1$ .

The custom tailored version for our purposes reads as follows.<sup>1</sup> We abbreviate  $\nu_x := \nu_{F_x}$ .

**Corollary 2.3.** Let  $x \in R^n$ ,  $y \in P \setminus M$  with  $y \notin F_x$ . Then  $\langle \nu_x, y \rangle \geq 1$ , and  $p(x, y) := \langle \nu_x, y \rangle x + y \in R^n$  is a root orthogonal to  $x$ .




Figure 2:  $p(x, y)$

**Proof of Theorem 1.1.** Let  $n \geq 3$ , and let  $x_1 \in R$  arbitrary. We have to show  $x_1 \in M$ . Assume not, that is,  $x_1 \notin M$ .

Let  $y_0 \in V(P) \cap F_{x_1}$  be a vertex of  $P$  outside  $F_{x_1}$ . Corollary 2.3 provides  $x_2 := p(x_1, y_0) \in R^n$ . The points  $x_1$  and  $x_2$  span a 2-dimensional linear space  $L$ . Either all vertices of  $P$  belong to  $(F_{x_1} \setminus F_{x_2}) \cup L$  or not.

$V(P) \cap (F_{x_1} \setminus F_{x_2}) \cup L$ : The intersection  $Q = P \cap L$  is a reflexive lattice polygon since any vertex of  $Q$  that is not on  $F_{x_1} \setminus F_{x_2}$  is a vertex of  $P$ , and the other vertex  $x_1 + x_2$  is integral, too. The part of  $\partial Q$  outside  $F_{x_1} \cup F_{x_2}$  belongs to the 1-skeleton of  $P$ . The endpoints of a primitive segment form a lattice basis for  $L \cap M \subseteq \mathbb{Z}x_1$ . So  $x_1 \in M$ , a contradiction. (This is where we use  $n \geq 3$ !)

<sup>1</sup>Of course, this corollary talks about the empty set as we aim to show.




Figure 3:  $V(P) \cap (F_{x_1} \setminus F_{x_2}) \cap L$

$V(P) \cap (F_{x_1} \setminus F_{x_2}) \cap L$  : There are two subcases.

$V(P) \cap F_{x_1} \cap F_{x_2} \cap L$  : Since  $y_0 \in V(F_{x_2}) \setminus L$ , there has to exist a pair of vertices  $y_1 \in F_{x_1} \cap F_{x_2}$  and  $y_2 \in F_{x_2} \cap F_{x_1}$  with  $y_1 + y_2 \notin 0$ , such that one of the two sets  $\{x_1, y_1, y_2\}$  or  $\{x_2, y_1, y_2\}$  is linearly independent, say the latter.

Choose  $u \in \mathbb{N}_R$  so that  $h_u; x_2 = h_u; y_2 = 0$  and  $h_u; y_1 = 1$ . Set  $A := \{r \in F_{x_1} \setminus \mathbb{R}^n : h_{x_2}; r = 0\}$ . Since  $x_1 \in A$ ,  $A \neq \emptyset$ , so there exists  $r \in A$  with  $h_u; r$  maximal (in particular  $> 0$ ).

By Corollary 2.3 we have  $k_1 := h_{x_2}; y_1 > 0$  and  $k_2 := h_{x_1}; y_2 > 0$ .

1. Furthermore  $q := p(r; y_2) = k_2 r + y_2 \in \mathbb{R}^n$ ,  $h_{x_1}; q = 0$ . Since  $h_{x_2}; r = 0$  and  $y_2 \in F_{x_2}$ , we get  $q \in F_{x_2}$ . Again by Corollary 2.3  $r^0 := p(q; y_1) = k_1 q + y_1 \in \mathbb{R}^n$ ,  $h_{x_2}; r^0 = 0$ . Since  $h_{x_1}; q = 0$  and  $y_1 \in F_{x_1}$ , we get  $r^0 \in F_{x_1}$ . Hence  $r^0 \in A$ . Since  $r^0 = k_1(k_2 r + y_2) + y_1$ , we have  $h_u; r^0 = k_1 k_2 h_u; r + k_1 > h_u; r$ , a contradiction.




Figure 4: Ping-pong

$V(P) \cap F_{x_1} \cap F_{x_2} \cap L$  : Let  $y_1 \in V(P)$  outside  $F_{x_1} \cap F_{x_2} \cap L$ . Now  $x_3 := p(x_1; y_1) \in \mathbb{R}^n$  does not belong to  $F_{x_2}$  or  $L$ . By Lemma 2.2,  $z := z(x_2; x_3) = ax_2 + bx_3 \in P \setminus L$  with  $h_{x_1}; z = b h_{x_1}; x_3 = 0$ , a contradiction to Corollary 2.3.

□




Figure 5: Finish

## References

- [Bat94] Batyrev, V.V.: Dual polyhedra and mirror symmetry for Calabi-Yau hypersurfaces in toric varieties. *J. Algebr. Geom.* 3, 493-535 (1994)
- [BK05] Batyrev, V. V.; Kreuzer, M.: Integral cohomology and mirror symmetry for Calabi-Yau 3-folds. Preprint, math.AG/0505432 (2005)
- [CK99] Cox, D.A.; Katz, S.: Mirror symmetry and algebraic geometry. *Mathematical surveys and monographs* 68. Providence, RI: Amer. Math. Soc. 1999
- [DM05] Doran, C.F.; Morgan, J.W.: Mirror symmetry and integral variations of Hodge structure underlying one parameter families of Calabi-Yau threefolds. Preprint, math.AG/0505272 (2005)
- [Ful93] Fulton, W.: Introduction to toric varieties. *Annals of Mathematics Studies* 131. Princeton, NJ: Princeton University Press 1993
- [KS00] Kreuzer, M.; Skarke, H.: Complete classification of reflexive polyhedra in four dimensions. *Adv. Theor. Math. Phys.* 4, 1209-1230 (2000)
- [Nil04] Nil, B.: Complete toric varieties with reductive automorphism group. Preprint, math.AG/0407491 (2004)
- [Nil05] Nil, B.: Gorenstein toric Fano varieties. *Manuscr. Math.* 116, 183-210 (2005)