Gam e theoretic derivation of discrete distributions and discrete pricing formulas

A kim ichi Takem ura and Taiji Suzuki
Departm ent of Mathematical Informatics
Graduate School of Information Science and Technology
University of Tokyo

September, 2005

A bstract

In this expository paper we illustrate the generality of game theoretic probability protocols of Shafer and Vovk (2001) in nite-horizon discrete games. By restricting ourselves to nite-horizon discrete games, we can explicitly describe how discrete distributions with nite support and the discrete pricing formulas, such as the Cox-Ross-Rubinstein formula, are naturally derived from game-theoretic probability protocols. Corresponding to any discrete distribution with nite support, we construct a nite-horizon discrete game, a replicating strategy of Skeptic, and a neutral forecasting strategy of Forecaster, such that the discrete distribution is derived from the game. Construction of a replicating strategy is the same as in the standard arbitrage arguments of pricing European options in the binomial tree models. However the game theoretic framework is advantageous because no a priori probabilistic assumption is needed.

Keywords and phrases: binom ial distribution, Cox-Ross-Rubinstein formula, hypergeometric distribution, lower price, Polya's distribution, probability protocol, replicating strategy, upper price

1 Introduction

In the gam e theoretic probability of Shafer and Vovk (2001), probability distributions and probability models are not assumed a priori but derived as logical consequences of certain protocol of a gam e between two players \Skeptic" and \Reality". In this game Skeptic tries to become rich by exploiting patterns in the moves of Reality. In order to prevent Skeptic from becoming rich, Reality is in a sense forced to behave probabilistically. Therefore probability distributions are determined by the protocol of the game. This feature of the game theoretic probability is well illustrated by Shafer and Vovk (2001) in

their derivation of Skeptic's strategy forcing the strong law of large numbers (Chapter 3) and the derivation of Black-Scholes formula (Chapter 9). Also in Takeuchi's exposition of the game theoretic probability and nance (Takeuchi (2004)) this point is discussed with many interesting examples. Recently Kumon and Takemura (2005) gave a very simple strategy forcing the strong law of large numbers.

In the standard stochastic derivation of option pricing form ulas, empirical probability is assumed rst, but then by arbitrage arguments, the empirical probability is replaced by the risk neutral probability and the price of an option is given as the expected value with respect to the risk neutral probability. The risk neutral probability is often explained as a purely operational device useful in expressing the option price in a convenient form. On the other hand in the game theoretic probability the risk neutral probability is more substantial, in the sense that Reality is forced to behave according to the risk neutral probability to avoid arbitrage by Skeptic. We should mention here that in Shafer and Vovk (2001) \forcing" is used only for in nite-horizon games. In this paper we somewhat informally use the word to mean that Reality should avoid arbitrage by Skeptic in the setting of nite-horizon games.

Additional exibility of game theoretic probability is gained by introducing the third player \Forecaster" into the game. At the beginning of each round Forecaster sets the price for Reality's move. By appropriately specifying the strategy of Forecaster, Reality's moves can be forced to follow any prespecied distribution.

In this paper we demonstrate the above features of the game theoretic probability in the setting of nite-horizon discrete games. For expository purposes we start with the simplest setting of the coin-tossing game and derive binomial distribution in Section 2 and give an analogous derivation of the Cox-Ross-Rubinstein formula in Section 3. We discuss derivation of hypergeometric distribution and Polya's distribution in Section 4 in order to illustrate the role of Forecaster. Then in Section 5 we discuss derivation of an arbitrary discrete distribution with nite support. Multivariate extension is given in Section 6. Some preliminary material on game theoretic probability is given in Appendix.

2 Derivation of binomial distribution

Consider the nite-horizon fair-coin game in Section 6.1 of Shafer and Vovk (2001). Its protocol is given as follows.

```
Fair-Coin Game
Protocol:

K_0 = :given

FOR n = 1; :::;N

Skeptic announces M_n \ge R.

Reality announces x_n \ge f = 1;1g.

K_n := K_{n-1} + M_n x_n.

END FOR
```

In this protocola gam e theoretic version of Chebyshev inequality is proved in (6.9) of

S&V in the following form:

$$\underline{P} = \frac{S_N}{N}$$
 1 $\frac{1}{N^2}$;

where $S_N = x_1 + \frac{1}{N}$ and \underline{P} denotes the lower probability. A ctually the equality of the upper probability and the lower probability

$$\underline{P} \quad \frac{S_{N}}{N} \qquad = P \quad \frac{S_{N}}{N} \tag{1}$$

holds here and this probability is given by binom ial distribution. Although this fact is contained in a more general statement of Proposition 8.5 of S&V, we give a full proof of this fact employing standard arbitrage arguments.

In order to treat success probability p = 1 = 2, let us consider the following biased-coin gam e.

Biased-Coin Game

P rotocol:

 $K_0 = 2 R; a;b > 0:given$

FOR n = 1; :::; N

Skeptic announces M_n 2 R.

Reality announces x_n 2 fa; bg.

 $K_n := K_{n-1} + M_n x_n$.

END FOR

As above we write $S_n = x_1 + \frac{1}{\pi}$.xLet $S_0 = 0$. Consider a random variable $x() = (S_N)$ which depends only on S_N (European option). Then we have the following basic result.

Theorem 2.1 The upper and the lower expected values of (S_N) coincide and given by

$$E((S_N)) = \underline{E}((S_N)) = \frac{X^N}{m=0} \quad N \quad p^m (1 \quad p)^{N-m} \quad (ma \quad (N \quad m)b); \tag{2}$$

where p = b = (a + b) is the risk neutral probability.

Proof: The rst step of our proof consists of de ning a \candidate" price of the European option. In the second step we verify that the candidate price is actually the precise price, by constructing a replicating strategy.

Let $(n; S_n)$, $S_n = nb$; $(n \ 1)b + a$;:::na, denote the price of (S_N) at time n. We require $(n; S_n)$ to satisfy the following \partial di erence equation"

$$(n;S_n) = p (n + 1;S_n + a) + q (n + 1;S_n b); 0 n < N;$$
 (3)

where q = 1 p. Note that (3) with p = 1=2 is a discrete version of the heat equation. The term inal condition for $(n; S_n)$ is given by

$$(N; S_N) = (S_N); S_N = ma (N m)b; m = 0; ...; N:$$
 (4)

Starting with the term in al condition (4) we can solve for $(n; S_n)$ in (3) by backward induction n = N 1; N 2; ...; 0. Then the initial value (0; 0) is easily calculated as

$$(0;0) = p (1;a) + q (1; b)$$

$$= p(p (2;2a) + q (2;a b)) + q(p (2;a b) + q (2; 2b))$$

$$= p^{2} (2;2a) + 2pq (2;a b) + q^{2} (2; 2b)$$

$$= :::$$

$$= \sum_{m=0}^{X^{N}} \sum_{m=0}^{N} p^{m} (1 p)^{N-m} (ma (N m)b):$$

Now we describe a replicating strategy for (S_N) with the the replicating initial capital (0;0). For n=1;:::;N, let

$$M_{n} = \frac{(n; S_{n-1} + a) \quad (n; S_{n-1} \quad b)}{a + b};$$
 (5)

Note that a + b can be written as

$$a + b = (S_{n-1} + a) (S_{n-1} b)$$
:

Therefore M_n is the ratio of the increments of $(n;S_n)$ and S_n and is called the \delta hedge". We now check that this M_n gives a replicating strategy P. This can be consimed by forward induction. At the end of the strough n=1,

$$(0;0) + K_{1}^{P} = (0;0) + M_{1}x_{1}$$

$$= (0;0) + \frac{(1;a) \quad (1; b)}{a+b} x_{1}$$

$$= (0;0) + q((1;a) \quad (1; b)); \text{ if } x_{1} = a;$$

$$(0;0) \quad p((1;a) \quad (1; b)); \text{ if } x_{1} = b;$$

$$= (1;a); \text{ if } x_{1} = a;$$

$$= (1;b); \text{ if } x_{1} = b;$$

$$= (1;S_{1}):$$

Sim ilarly at the end of round n = 2, we have

$$(0;0) + K_2^P =$$

$$(2;S_1 + a); \text{ if } x_2 = a;$$

$$(2;S_1 + b); \text{ if } x_2 = b;$$

$$= (2;S_2):$$

Now by induction we arrive at

$$(0;0) + K_N^P = (N;S_N) = (S_N)$$
:

We have con rmed that M $_n$ in (5) with the replicating initial capital (2) gives a replicating strategy for (S_N) . Hence the theorem holds by Proposition A.1 in Appendix.

In particular if we take

$$(S_N) = \begin{cases} 1; & \text{if } S_N \neq N \\ 0; & \text{otherw ise;} \end{cases}$$

we see that the equality holds in (1) and the probability is given by binom ial distribution.

In this section we took Reality's move space as fa; bg. This is convenient in comparing Theorem 2.1 with the Cox-Ross-Rubinstein formula in the next section. However for generalization of binomial distribution to hypergeometric distribution in Section 4, it is more convenient to rescale Reality's move space to f0;1g. Then we need to introduce the price p for the \ticket" x_n . The rescaled protocol is written as follows.

Rescaled Biased-Coin Game Protocol:

$$K_0 = ;0 : given FOR $n = 1; :::;N$ Skeptic announces $M_n \ge R$. Reality announces $x_n \ge f0;1g$. $K_n := K_{n-1} + M_n (x_n - p)$. END FOR$$

It is clear that the biased-coin gam e and the rescaled biased-coin gam e is equivalent by the a ne correspondence $x_n \$ (a + b) $(x_n \ p)$; p = b = (a + b). In the rescaled version the expected value in (2) is $\sin p \ln w$ written as

$$E ((S_N)) = \underline{E} ((S_N)) = \begin{pmatrix} X^N & N \\ m = 0 & m \end{pmatrix} p^m (1 p)^{N-m} (m):$$
 (6)

Furtherm ore, since the increm ent of S_n is norm alized to be 1, the replicating strategy in (5) is $\sin p \ln w$ written as

$$M_n = (n; S_{n-1} + 1) \quad (n; S_{n-1}):$$
 (7)

It is also conceptually very important to consider the single step game i.e. the game with N=1. Note that each round n of the N step biased-coin game can be viewed as a single step game. In the single step game binomial distribution reduces to a Bernoulli trial. This implies that given the price p, Reality's move x_n for each round n is exactly the same as a single Bernoulli trial with success probability p. Furtherm ore this behavior of Reality is dictated solely by the value of p, independently from the past moves $x_1; \ldots; x_{n-1}$ of Reality. Therefore in the Rescaled Biased-Coin Game, Reality's moves $x_1; \ldots; x_N$ are independent Bernoulli trials.

3 Derivation of the Cox-Ross-Rubinstein formula

Here we present a game theoretic formulation and derivation of the Cox-Ross-Rubinstein formula (Cox, Ross and Rubinstein (1979)), which is fully discussed in many introductory textbooks on option pricing (e.g., Shreve (2004), Chapter 2 of Baxter and Rennie (1996), Chapter 8 of Capinski and Zastawniak (2003)). Once an appropriate game is formulated, the rest of the argument is the same as in the previous section.

Our protocol for Cox-Ross-Rubinstein game is as follows.

Cox-Ross-Rubinstein Game Protocol:

```
\begin{split} &S_0 > 0; u > r > d > 0: \text{given} \\ &FOR \ n = 1; :::; N \\ &\quad \text{Skeptic announces M}_n \ 2 \ R. \\ &\quad \text{Reality announces x}_n \ 2 \ \text{fu}; \text{dg.} \\ &\quad S_n \ \coloneqq \ S_{n-1} \quad x_n. \\ &\quad K_n \ \coloneqq \ K_{n-1} + M_n \ (S_n \quad S_{n-1}) + \ (r-1) \ (K_{n-1} \quad M_n S_{n-1}). \\ &\quad \text{END FOR} \end{split}
```

Here K $_{\rm n-1}$ M $_{\rm n}$ S $_{\rm n-1}$ is the am ount of risk less bond held by Skeptic for the round n and r 1 is the xed risk less interest rate. A lthough by appropriate discounting we may put r = 1 w ithout essential loss of generality (Section 12.1 of S & V), here we leave the interest rate ras in standard derivation of the C ox-R oss-R ubinstein formula. p = (r d)=(u d) is called the risk neutral probability.

Let (S_N) denote a payo function of a European option depending on S_N . Corresponding to Theorem 2.1 we have the following result.

Theorem 3.1 (the Cox-Ross-Rubinstein formula) The upper and the lower expected values of (S_N) coincide and given by

$$E ((S_N)) = \underline{E} ((S_N)) = \frac{1}{r^N} \sum_{m=0}^{X^N} \sum_{m=0}^{N} p^m (1 - p)^{N-m} (u^m d^{N-m} S_0);$$
 (8)

where p = (r d) = (u d) is the risk neutral probability.

Proof: As in the previous section we de ne $(n;S_n)$ by backward induction. Put $(N;S_n) = (S_n)$ and for n = N 1;:::;0, de ne

$$(n;S_n) = \frac{1}{r} p (n + 1;uS_n) + (1 p) (n + 1;dS_n)$$

Then the initial value $(0; S_0)$ is easily calculated as

$$(0;S_0) = \frac{1}{r^N} X^N \qquad N \qquad p^m (1 p)^{N m} (u^m d^{N m} S_0)$$
:

This becomes the replicating initial capital of the following replicating strategy:

$$M_n = \frac{(n; uS_{n-1}) (n; dS_{n-1})}{(u d)S_{n-1}}$$
:

Since the gam e is coherent by the requirement u > r > d, the theorem follows from Proposition A 1.

4 Hypergeom etric distribution and Polya's distribution

In the rescaled biased-coin game of Section 2, the price p of the ticket x_n was a constant. Therefore the third player \Forecaster" did not enter the protocol. Now we introduce Forecaster, who sets the price of the ticket at the beginning of each round in the rescaled biased-coin game. We illustrate the role of Forecaster below by deriving the hypergeom etric distribution. We also derive Polya's distribution.

Rescaled Biased-Coin Game W ith Forecaster Protocol:

```
K_0 = :given

FOR n = 1; :::;N

Forecaster announces p_n \ge R.

Skeptic announces M_n \ge R.

Reality announces x_n \ge f0;1g.

K_n = K_{n-1} + M_n (x_n - p_n).

END FOR
```

Note that if Forecaster announces $p_n > 1$ or $p_n < 0$, then Skeptic can become in nitely rich im mediately by taking M_n jarbitrarily large. Therefore we can restrict them ove space of Forecaster to be [0;1]. Furtherm ore if $p_n = 0$, Skeptic can still take M_n arbitrarily large, which forces Reality to choose $x_n = 0$. Similarly if $p_n = 1$, then Reality is forced to choose $x_n = 1$.

Now consider a strategy of Forecaster. A strategy of Forecaster is called neutral (Section 8.2 of S& V) if p_n is determined by the past moves of Reality $x_1:::x_{n-1}$. From now on we only consider neutral strategies for Forecaster. Furthermore for simplicity we consider neutral strategy depending on $S_{n-1}=x_1+\frac{1}{n}$ and write $p_n=p_n$ (S_{n-1}), which we may call \M arkovian neutral strategy". In M arkovian neutral strategy Forecaster only needs to keep S_{n-1} in memory to choose his move.

Consider an um with $\ _1$ red balls and $\ _2$ black balls, where $\ _1+\ _2$ N. Let $x_n=1$ correspond to drawing a red ball and let $x_n=0$ correspond to drawing a black ball from the um by Reality. Let p_n be the ratio of red balls in the um at the n-th round. Then

$$p_n = p_n (S_{n-1}) = \frac{m ax (0; 1 S_{n-1})}{1 + 2 (n-1)}$$
:

A ctually here we do need to take the positive part of $_1$ S_{n-1} , because as remarked above once the boundary $S_{n-1} = _1$ is attained, then $0 = p_n = p_{n+1} = :::$ and Reality is forced to choose $0 = x_n = x_{n+1} = :::$, which results in $_1 = S_n = S_{n+1} = :::$. Now we write out a game of sampling without replacement from an um.

Game of Sampling W ithout Replacement From An Urn Protocol:

```
\begin{split} &K_0 = \text{ ; }_1 > 0\text{; }_2 > 0\text{; }_1 + \text{ }_2 \quad \text{N , } S_0 = 0\text{ : given} \\ &FOR \text{ } n = 1\text{; :::;} \text{N} \\ &Forecaster announces } p_n = \left( \text{ }_1 \quad S_{n-1} \right) = \left( \text{ }_1 + \text{ }_2 \quad n+1 \right) \\ &Skeptic announces M_n 2 R \text{ .} \\ &Reality announces x_n 2 f0\text{; } 1g\text{.} \\ &K_n \coloneqq K_{n-1} + M_n \left( x_n \quad p_n \right) \text{.} \\ &S_n \coloneqq S_{n-1} + x_n \text{.} \end{split} END FOR
```

For this gam e the upper and the lower values of the payo (S_N) coincide and are given by the expected value with respect to the hypergeom etric distribution:

$$E ((S_N)) = \underline{E} ((S_N)) = \frac{m \operatorname{in} X_{1,N}}{m = m \operatorname{ax}(0,N - 2)} (m) \frac{\frac{1}{m} \frac{2}{N - m}}{\frac{1+2}{N}} :$$
 (9)

This result is actually almost obvious from the discussion at the end of Section 2, namely, at each round n Reality's move \mathbf{x}_n is like drawing a ball from an unw ith $_1$ S_{n 1} red balls and $_2$ (n 1 S_{n 1}) black balls. However it is instructive to look at a formal proof of (9).

De ne a candidate price of (S_N) at time n by backward induction:

$$(n; S_n) = p_{n+1}(S_n) (n+1; S_n+1) + (1 p_{n+1}(S_n)) (n+1; S_n); n=N 1; :::; 0;$$

with the term inal condition $(N;S_N) = (S_N)$. Then by fully expanding the recurrence relation we have

$$(0;0) = \begin{array}{c} X & Y^{N} \\ p_{n} (S_{n-1})^{x_{n}} (1 - p_{n} (S_{n-1}))^{1 - x_{n}} (S_{N}) : \end{array} (10)$$

$$\underset{\text{max}(0;N-2)}{(x_{1}; \dots; x_{N})^{2} f_{0}; l_{g}^{N}} n = 1} p_{n} (S_{n-1})^{x_{n}} (1 - p_{n} (S_{n-1}))^{1 - x_{n}} (S_{N}) : \tag{10}$$

A ctually we do not need the restriction max(0;N $_2$) S_N m in($_1$;N) in the sum mation, because $\sum_{n=1}^N p_n (S_{n-1})^{x_n} (1 - p_n (S_{n-1}))^{1-x_n} = 0$ for S_N outside of this range. Now

it is easily seen that

$$\begin{split} & \overset{Y^{N}}{\underset{n=1}{\bigvee}} p_{n} \left(S_{n-1}\right)^{x_{n}} \left(1 - p_{n} \left(S_{n-1}\right)\right)^{1 - x_{n}} \\ & = \frac{1 \left(1 - 1\right) - 1 \left(S_{N} + 1\right) - 2 \left(2 - 1\right) - 2 \left(N + S_{N} + 1\right)}{\left(1 + 2\right) \left(1 + 2 - 1\right) - 1 \cdot \left(2 - N + 1\right)} \\ & = \frac{\frac{1!}{\left(1 - S_{N}\right)!} \frac{2!}{\left(2 - N - S_{N}\right)!}}{\frac{\left(1 + 2\right)!}{\left(1 + 2 - N\right)!}} ; \end{split}$$

T herefore

$$(0;0) = X \frac{\frac{1!}{(1 S_N)!} \frac{2!}{(2 (N S_N))!}}{\frac{(1+2)!}{(1+2 N)!}} (S_N):$$

$$\frac{(S_N)!}{(1+2)!}$$

Now for a given value of S_N , the sum mation just counts the number of ways of choosing S_N 1's among x_1 ;:::; x_N . It follows that

$$(0;0) = \begin{array}{c} X \\ \max(0;N) = \\ \max(0;N) = 2 \\ \end{array} \xrightarrow{S_{N} \text{ m in } (1;N)} \frac{N!}{S_{N}!(N)} \frac{\frac{1!}{(1 + S_{N})!} \frac{2!}{(2 + N)!}}{\frac{(1+2)!}{(1+2 + N)!}} (S_{N});$$

which proves (9).

The above argument can be immediately applied to Polya's ummodel (Section V 2 of Feller (1968)). In this scheme, when a ball is drawn from an um, it is replaced and, moreover, c balls of the same color are added. Then the game corresponding to Polya's ummodeldiers from the game of sampling without replacement only in the speciation of Forecaster's neutral strategy. In Polya's ummodel

$$p_n = p_n (S_{n-1}) = \frac{1 + cS_{n-1}}{1 + 2 + (n-1)c}$$
:

Then as in (2.4) of Section V 2 of Feller (1968), the expected value of (S_N) in this game is written as

$$E ((S_N)) = \underline{E} ((S_N)) = \frac{X^N}{m = 0} (m) \frac{\frac{1-C}{m} \frac{2-C}{N}}{(\frac{1+2}{2}) = C};$$

$$(11)$$

where the binomial coecient ${r\atop n}$ for a real r and nonnegative integer k denotes

$$r = \frac{r(r-1)}{k} = \frac{(r-k+1)}{(12)}$$

Note that (6) and (9) are special cases of (11) with c = 0 and c = 1, respectively. In (9) the range of sum m ation can be taken as m = 0; ...; N, with the convention (12).

5 Arbitrary discrete distribution with nite support

So far we have discussed how to derive some classical distributions. We now show that given any distribution on f0;:::;N g, we can specify a neutral strategy of Forecaster in a game with N rounds such that Reality follows the distribution.

Let $q_n = 0$; m = 0; m =

$$q_n = \frac{q_n + q_n + q_n}{q_{n-1} + q_n + q_n}; m = 1; ...; N :$$

Then

$$q_{n} = q_{1}$$
 $_{m}$ (q_{n+1}); $m = 1; :::; N; q_{N+1} = 0:$ (13)
$$q_{n} = q_{n}; \text{ if } S_{n-1} = n - 1;$$

$$p_{n} = q_{n}; \text{ otherwise:}$$

Let

The idea here is to let Reality increase S_{n-1} by 1 with probability q_{n-1} if $S_{n-1}=n-1$ or otherwise let him stop at the current level for the rest of the rounds. Note that $p_n=p_n\left(S_{n-1}\right)$ is indeed a function of S_{n-1} , because it is written as

$$p_n = q_n$$
 I($S_{n-1} = n - 1$);

where I () is the indicator function.

Biased-Coin Game W ith Forecaster For Arbitrary Distribution Protocol:

The tree of this game is illustrated in Figure 1. For this game we have the following result.

Theorem 5.1 The upper and the lower expected values of (S_N) coincide and given by

$$E ((S_N)) = \underline{E} ((S_N)) = \sum_{m=0}^{X^N} (m) q_m :$$
 (14)




Figure 1: Tree of the gam e for arbitrary distribution

Proof: As in the case of hypergeom etric distribution

$$(0;0) = \begin{array}{c} X & Y^{N} \\ & p_{n} (S_{n-1})^{x_{n}} (1 & p_{n} (S_{n-1}))^{1 \times x_{n}} (S_{N}) : \\ & (x_{1}; :::; x_{N}) 2 f_{0}; 1 g^{N} = 1 \end{array}$$

In this gam e, the path leading to S_{m} is uniquely determ ined as

$$x_1 ::: x_N = 1 ::: 10 ::: 0$$

with m initial 1's. By (13), for this path $\sum_{n=1}^{Q} p_n (S_{n-1})^{x_n} (1 - p_n (S_{n-1}))^{1-x_n} = q_n$. Therefore (0;0) = $\sum_{m=0}^{N} (m) q_m$. The replicating strategy con rm ing this candidate price is given as

$$M_{n} = \begin{pmatrix} (P_{N} & (m)q_{n} \\ \frac{m}{n} = n & (m)q_{n} \\ \vdots & \vdots \\ m = n & q_{n} \end{pmatrix}$$
 (S_n 1); if S_n 1 = n 1; otherwise:

In Theorem 5.1 we have considered a discrete distribution with the support f0;:::;N g. We can dealwith the support of the form fa;a+1;:::;bg, by letting N=b a and setting the initial value $S_0=a$.

In Section 8.3 of their book, Shafer and Vovk discuss \adding tickets" to make the upper expected value and the lower expected value to coincide. Note that if Reality's move space has more than two elements in a single step game, then the upper expected value is generally larger than the lower expected value. Theorem 5.1 shows that if we add so cient number of steps to a single step game, the equality of the upper and the lower prices is achieved.

6 Multivariate extension

In the previous sections we have considered univariate random variable S_N . In this section we give a straightforward multivariate extension of the results of the previous sections. We

em ploy the multi-label classication protocoldiscussed in Vovk, Nouretdinov, Takemura and Shafer (2005).

Our extension corresponds to generalizing B inom ial distribution to multinom ial distribution. Let $S_N = (S_N^1; :::; S_N^d)$ be a d-dim ensional vector. As in multinom ial distribution, for the sake of sym metry, we leave one-dim ensional redundancy $N = S_N^1 + \frac{d}{N}$ In the components of S_N . Therefore x_n in Rescaled B iased-C oin G ame now corresponds to a 2-dim ensional vector $(x_n; 1 \quad x_n)$. For the general d-dim ensional case the m ove space of Reality

$$X = fe_1; ...; e_d q = f(1;0; ...;0); (0;1;0; ...;0); ...; (0; ...;0;1)q$$

consists of d standard coordinate vectors.

Multilabel Classification Game With Neutral Forecasting Strategy Protocol:

```
K_0 = , S_0 = 0 : given

FOR n = 1; :::; N

Forecaster announces p_n = p_n (S_{n-1}) 2 R^d.

Skeptic announces M_n 2 R^d.

Reality announces x_n 2 X.

K_n = K_{n-1} + M_n (x_n p_n).

S_n = S_{n-1} + x_n.

END FOR
```

Here $\$ "denotes the standard inner product of \mathbb{R}^{1} .

In the above protocol we took the whole Rd as the move space of Forecaster. Let

$$(X) = f(p^1; ...; p^d) jp^i = 0;$$
 $p^i = 1g$

denote the probability sim plex spanned by the standard coordinate vectors. If Forecaster announces p_n (%), then by the hyperplane separation theorem. Skeptic can choose M $_n$ 2 R d such that he becomes in nitely rich immediately, nomeatter what move Reality chooses. See Vovk, Nouretdinov, Takemura and Shafer (2005) for a discussion of this point. Therefore we can restrict Forecaster's move space to the probability simplex (%). Also if $p_n^i = 0$ for some i, Skeptic can choose M $_n^i$ arbitrarily large and Reality is forced to choose $x_n^i = 0$.

We also note that there is a redundancy in the move space of Skeptic, once p_n is restricted to lie in (X). M $_n$ + c(1;:::;1) for any c 2 R leads to the same increment of the capital process K_n . However it is often convenient to ignore this redundancy in specifying Reality's move M $_n$.

For notational sim plicity write

$$p_n^{x_n} = (p_n^1)^{x_n^1}$$
 $\binom{d}{n}(p^0) = p_n^1$ for $x_n = e_i$:

As a straightforward generalization of results in the previous sections we have the following theorem .

Theorem 6.1 The upper and the lower expected values of (S_N) coincide and given by

$$E ((S_N)) = \underline{E} ((S_N)) = \begin{array}{c} X & Y^N \\ p_n (S_{n-1})^{x_n} (S_N) : \end{array}$$

$$(15)$$

The line of the proof is the same as in the previous theorem s and we om it the details. The price $(n;S_n)$ at time n is de ned recursively by

$$(n;S_n) = \sum_{i=1}^{X^d} p_{n+1}^i (S_n) (n+1;S_n + e_i)$$

and the replicating strategy M $_{\rm n}^{\rm i}$ is \sin ply given by

$$M_{n}^{i} = (n; S_{n+1} + e_{i})$$
:

From Theorem 6.1 we can easily derive multinomial distribution, multivariate hypergeometric distribution as well as multivariate Polya's distribution.

A generalization of Theorem 5.1 to an arbitrary (d 1)-dimensional discrete distribution of $(S_N^1; :::: S_N^{d-1})$ with nite support can be explained as follows. We rst use Theorem 5.1 on the rst component S_n^1 to derive the one-dimensional marginal distribution of S_N^1 . Then, given the realization of the rst component, we derive the conditional distribution of S_N^2 given S_N^1 by another application of Theorem 5.1 to the second component. We can continue this process up to the (d 1)th component. The last component S_n^d is used as a slack variable.

Finally as an illustration of Theorem 6.1 we show how the Cox-Ross-Rubinstein formula of Section 3 is reduced to our multivariate fram ework. De ne

$$x_n = \frac{S_n - rS_{n-1}}{r^n}$$
:

Furtherm ore by discounting de ne

$$K_n = \frac{K_n}{r^n}$$
:

Then the recurrence relation of the Cox-Ross-Rubinstein game

$$K_n = K_{n-1} + M_n (S_n - S_{n-1}) + (r - 1) (K_{n-1} - M_n S_{n-1})$$

= $rK_{n-1} + (S_n - rS_{n-1})M_n$

is written as

$$K_{n} = K_{n-1} + M_{n} x_{n}$$
:

 $H \ ere \ x_n \ can \ take \ two \ values$

$$(x_n^1; x_n^2) = \frac{S_{n-1}(u-r)}{r^n}; \frac{S_{n-1}(d-r)}{r^n}$$
:

Rescaling the values we de ne d = 2, x_n 2 fe₁; e₂g = f(1;0); (0;1)g and

$$M_{n} = (x_{n}^{1}; x_{n}^{2});$$

$$p_{n} = \frac{x_{n}^{2}}{x_{n}^{1} x_{n}^{2}}; \frac{x_{n}^{1}}{x_{n}^{1} x_{n}^{2}} :$$

Then $M_n x_n$ is written as

$$M_n x_n = M_n \quad (x_n \quad p_n)$$
:

Therefore the iteration part of the the Cox-Ross-Rubinstein game is written as

FOR
$$n = 1; :::; N$$

Skeptic announces $M_n \ge R^2$.
Reality announces $x_n \ge fe_1; e_2g$.
 $K_n := K_{n-1} + M_n$ ($x_n = p_n$).
END FOR

This shows that the Cox-Ross-Rubinstein formula is also a special case of our multi-variate extension.

A P relim inaries on game theoretic probability

Here we sum marize prelim inary material (Chapter 1 of S&V) of the game theoretic probability. We also state a basic proposition on the existence of a replicating strategy and the existence of the game theoretic expectation in a coherent game.

In this paper all the gam es are nite-horizon gam es with N rounds. Therefore a path of the gam e is a nite sequence = $x_1:::x_N$ of Reality's moves. A random variable x () denotes a payo to Skeptic, when Reality chooses the path . G iven a strategy P of Skeptic, K^P denotes the capital process of P with zero initial capital. Furtherm ore in this paper we only consider symmetric games, in the sense that if P is a strategy of Skeptic, then P is also a strategy of Skeptic and

$$K^{P} = K^{P}$$
:

The upper expected value Ex and the lower expected value $\underline{E}x$ of x is de ned as

$$Ex = \inf j9P;8;K_N^P() x() g;$$

 $\underline{E}x = \sup j9P;8;K_N^P() x()g:$

In a sym m etric gam e \underline{E} x can also be written as

$$\underline{E}x = \text{supf j9P;8;} + K_N^P () x ()g:$$
 (16)

A gam e is coherent if Skeptic is not allowed to make money for certain, i.e.,

8P; 9;
$$K_N^P$$
() < 0:

If a game is coherent, then Ex \underline{E} x for every random variable x (Proposition 72 of S&V).

We call Pareplicating strategy for x with the replicating initial capital 2R if

$$+ K_{N}^{P} () = x(); 8 :$$

We now state the following basic fact.

Proposition A.1 In a coherent symmetric game, suppose that P is a replicating strategy for x with the replicating initial capital . Then

$$Ex = \underline{E}x =$$
:

Proof: By de nition of Ex we have Ex . Furtherm ore in a symmetric game $\underline{E}x$ follows from (16). Therefore Ex $\underline{E}x$. Combining this with the inequality Ex Ex we obtain the proposition.

We should note that the proof of the inequality $Ex = \underline{E}x$ in S&V and the above proof are standard arbitrage arguments.

A cknow ledgm ent. We are grateful for insightful comments by Vladim ir Vovk, Glenn Shafer, Kei Takeuchi and Masayuki Kumon on earlier drafts of this paper.

References

- [1] Martin Baxter and Andrew Rennie. Financial Calculus, An Introduction to Derivative Pricing. Cambridge University Press, Cambridge, 1996.
- [2] Marek Capinski and Tomasz Zastawniak. Mathematics for Finance, An Introduction to Financial Engineering. Springer, London, 2003.
- [3] John C. Cox, Stephen A. Ross, and Mark Rubinstein. Option pricing: A simplied approach. Journal of Financial Economics, 3, 229 (263, 1979.
- [4] William Feller. An Introduction to Probability Theory and Its Applications. third edition, volume 1, Wiley, New York, 1968.
- [5] Masayuki Kumon and Akim ichi Takemura. On a simple strategy weakly forcing the strong law of large numbers in the bounded forecasting game. Technical Report METR 05-20, Department of Mathematical Informatics, University of Tokyo. 2005.
- [6] Glenn Shafer and Vladim ir Vovk. Probability and Finance: It's Only a Game! Wiley, New York, 2001.
- [7] Steven E. Shreve. Stochastic Calculus for Finance I. The Binomial Asset Pricing Model. Springer, New York. 2004.

- [8] Kei Takeuchi. Kake no suuri to kinyu kogaku (Mathematics of betting and nancial engineering). Saiensusha, Tokyo, 2004. (in Japanese)
- [9] V ladim ir Vovk, Ilia Nouretdinov, Akim ichi Takemura, and Glenn Shafer. Defensive forecasting for linear protocols. The Game-Theoretic Probability and Finance project, http://probabilityandfinance.com, Working paper # 10, 2005. To appear in Proceedings of the Sixteenth International Conference on Algorithm ic Learning Theory (ed. by Sanjay Jain, Hans Ulrich Simon, and Etsuji Tomita), Lecture Notes in Articial Intelligence, Springer, Berlin.