On phylogenetic trees { a geom eter's view

W eronika Buczynska and Jaroslaw A.W isniewski Instytut M atem atyki UW, Banacha 2,02-097 W arszawa, Poland

M arch 8, 2022

A bstract: In the present note we investigate projective varieties which are geom etric m odels of binary symmetric phylogenetic 3-valent trees. We prove that these varieties have G orenstein term inal singularities (with small resolution) and they are Fano varieties of index 4. Moreover any two such varieties associated to trees with the same number of leaves are deformation equivalent, that is, they are in the same connected component of the Hilbert scheme of the projective space. As an application we provide a simple formula for computing their Hilbert-Ehrhard polynom ial.

C ontents

0	Intr	oduction	2
	0.1	Notation	5
1	P re	lim inaries: phylogenetic trees.	5
	1.1	Trees and linear algebra	5
	12	Binary symmetric trees.	8
2	Tor	ic geom etry.	11
	2.1	Lattice of a tree and the action of the torus	11
	22	Polytopemodelofatree	14
	2.3	Geometricmodelofatree	18
	2.4	1-param eter group action, quotients	20

Deticated to Andrzej Bialynicki-Birula

 $^{\mathrm{y}}$ Partially supported by Polish KBN (grant 1P03A03027) and MPIM, Bonn, Germany

· · · 2	22 25
2	25
~	
••• 3	30
3	34
3	38
3	38
4	11
4	12
	3

0 Introduction

A lgebraic geom etry, a classical, alm ost ancient, branch of pure m athem atics, is constantly stim ulated by questions arising in applicable m athem atics and other sciences. String theory and m intor conjecture from m athem atical physics, coding theory and im age recognition from computer sciences | to m ention just a few of the big areas of sciences which had enorm ous im pact on the developm ent of algebraic geom etry in the past decade. Now the m odern biology with its computational aspects and relations to statistics seems to be m aking its way into this branch of m athem atics.

A lthough the roots of questions which we tackle are beyond the area of our professional interest and we do not claim any thorough understanding of them still, the questions form ulated in the language of our trade seem to be extrem ely interesting for its own, m athem atical m eaning. In fact, we believe that m ost of the important things in m athem atics are related to real phenom ena of N ature. The interpretation of this profound feature of M athem atics is left for the reader and it will de nitely depend on the reader's attitude tow ards fundam ental C reation vs. Evolution problem, cf. [Shafarevich] and [R eid '87].

K now ing our limitations as laymen in computational biology and statistics we try to stay within borders of the branch of mathematics which we believe we understand. That is why we take a relatively simplemodel, rede neit in purely algebraic language and examine it using methods of algebraic geometry. The result exceeds our original expectations, we nd the object appearing in this process very interesting for its own, pure geometric aspects, with properties which we have not expected originally.

O ur original task was com puting H ilbert-Ehrhard polynom ials for varieties arising as geometrical models of binary symmetric 3-valent phylogenetic trees. The question is consistent with the attitude of computational algebraic geometry and algebraic statistics where the point is to compute and understand the ideal of the variety in question in the ambient projective space. Then the Hilbert-Ehrhard polynom ial provides a fundam ental invariant of such an ideal, the dimensions of hom ogeneous parts of it. To our surprise the polynom ial does not depend on the shape of the tree but merely on its size, the number of leaves or, equivalently the dimension of its geometric model. The strive to understanding this phenomenon lead us to proving one of the main results of the present paper, 3.26, which asserts that models of trees with the same number of leaves are deformation equivalent, that is they are in the same connected component of the Hilbert scheme of the projective space in question (hence they have the same Hilbert polynom ial).

The fact that the geom etric m odels of trees m odelling som e processes | the discreet objects | live in a connected continuous family of geom etric objects probably deserves its explanation in terms of algebraic statistic or even biology. For the algebraic geom etry part we have a natural question arising about irreducibility of the component of the H ilbert scheme e containing these m odels and (if the irreducibility is con m ed) about varieties which arise as general deform ations (that is, over a general point of the component of the H ilbert scheme in question). The question about a general deform ation of the m odel is related to the other m ain result of the present paper, 3.17, which is that these m odels are index 4 Fano varieties with G orenstein term inal singularities. Thus one would expect that their general deform ation is a sm ooth Fano variety of index 4, c.f. [N am ikawa].

The present paper is organized as follows. We deal with varieties de ned over complex numbers. In the 1st section we de ne phylogenetic trees and their geom etric models. We do it in pure algebraic way and with many simplications: we deal with unrooted symmetric trees which are then assumed to be binary and eventually 3-valent. From the algebraic geometer point of view studying geometric models in this case can be reduced to understanding special linear subsystems of the Segre linear system on a product of P^{1} 's, 1.9. Eventually, the question boils down to studying xed points of the Segre system with respect to an action of a group of involutions, 1.12. Since the action can be diagonalized this brings us down to toric geometry.

In the second section we de ne a geom etric model of a tree in terms of toric geom etry, via a polytope in the space of characters of a complex torus, which we call a polytope model of the tree and to which we subsequently associate a projective variety. The main results of this part are 2.12 and 2.24 which assert that the models de ned in the rst part are the same as these de ned the toric way. In this part we also prove results which are of the fundam ental technical in portance: this is a ber product form ula for polytopes of trees, 2.20, and its counterpart for varieties,

a quotient form ula 2.26. The latter asserts that the geometric model of a tree obtained by gluing two smaller trees is a Mum ford's GIT (Geometric Invariant Theory) quotient of the product of their respective models.

The third section of the present paper contains its main results. A fler a brief discussion of equations de ning a geom etric model of a tree, with special consideration to a tree with two inner nodes and four leaves, we exam ine fans of geom etric models and resolution of their singularities. We prove that geom etric models of 3-valent binary symmetric trees are index 4 Fano varieties with G orenstein term inal singularities which admit small resolution, 3.17. Next we consider deform ations of models of trees. The approach is, roughly, as follows: we know how to deform equations of a small tree with four leaves and one inner edge, the result of the deform ation is another tree with the inner edge $\$ opped":



Applying the G IT quotient formula, 2.26, we are able to use this elementary deformation associated to four leaves trees to get a similar deformation for every inner edge of any tree, 3.24. This implies the result about deforming one geometric model to another, 3.26.

In the last part of section 3, we discuss H ilbert-Ehrhard polynom ial of m odels (both polytopes and varieties) of trees. We de near elative version of the polynom ial and then a product of such polynom ials which is related to gluing respective trees. The elementary deformation procedure in plies associativity of the product which not only im plies the invariance of the H ilbert-Ehrhard polynom ial for trees with the same number of leaves but also provides a simple formula for computing it, 3.38.

The appendix contains som e com putations. Firstly we prove that the polytopes of the 3-valent trees are norm alwhich is needed to ensure the proper de nition of their geom etrical models. Next, using the [polymake] software we verify a sim ple (yet 9-dim ensional) example to check that the polytope models of di erent trees in this case are di erent. The question if the polytope (or geom etric) models of nonisom orphic trees are non-isom orphic is open, c.f. [A llm an, Rhodes, '05]. Finally, we make num erical experiments (using [maxima] and [gnuplot]) to book at the behaviour of the relative volume distribution which measures the (norm alized) volume of the model with respect to a xed leaf of a tree.

The paper uses consistently the language of algebraic geom etry, including toric geom etry. We ignore, or barely mention, relations to algebraic statistic and biology, suggesting the reader to look into Pachter, Sturm fels] (or into ERSS] for a concise version of exposition), to get an idea about the background of the problem s that we

deal with. It was our primary intention to make the present paper self-contained so that it can be read as it is by an algebraic geometer with no know ledge of its possible applications outside algebraic geometry. On the other hand, a reader who is not familiar with algebraic geometry but is interested in acquiring ideas which are important in our approach (regarding quotients and deform ations) is advised to bok into [Reid '92] and [Altman] for a short exposition to these matters.

We would like to thank Jaroslaw Buczynski for his remarks and Piotr Zwiernik for bringing this subject to our attention.

0.1 Notation

A jdenotes cardinality of a nite set A.

A lattice is a nitely generated free abelian group.

Depending on the context a subscript denotes the extension of the basic ring or a ber of a morphism, e.g. $M_R = M_Z R$.

Given a nite dimensional vector space (or a lattice) V with a basis $fv_1; :::; v_n g$, by $fv_1; :::; v_n g$ we will denote the dual base of V, that is $v_i (v_i) = 1$ and $v_i (v_j) = 0$ if $i \in j$.

1 Prelim inaries: phylogenetic trees.

Sum m ary: (for algebraic geom eters) phylogenetic trees are a clever way of describing linear subsystems of Segre system on the product of projective spaces. In case of binary symmetric trees the question is to nd subsystems of sections of Segre system on a product of P^1 's invariant with respect to some $Z_2^{\frac{N}{2}}$ action.

1.1 Trees and linear algebra

N otation 1.1. A tree T is a simply connected graph (1-dimensional CW complex) with a set of edges E = E(T) and vertices V = V(T) and the (unordered) boundary map $@:E ! V^{2}$, where V^{2} denotes the set of unordered pairs of distinct elements in V. The number $\pm j$ 1 is, by de nition, the number of edges of T, then number of vertices $J_{V} j_{1} = j_{1} + 1$. We write $@(e) = f@_{1}(e);@_{2}(e)g$ and say v is a vertex of e, or e contains v if v 2 f@_{1}(e);@_{2}(e)g, we simply write v 2 e. The valency of a vertex v is the number of edges which contain v (the valency is positive since T is connected

and we assume it has at least one edge). A vertex v is called a leaf if its valency is 1, otherwise it is called an inner vertex or a node. If the valency of each inner node is m then the tree will be called m -valent. The set of leaves and nodes will be denoted L and N, respectively, V = L [N. An edge which contains a leaf is called a petiole, an edge which is not a petiole is called an inner edge (or branch), and the set of inner edges will be denoted by E° .

E xam ple 1.2. An caterpillar of length n is a 3-valent tree with n inner edges and n + 1 inner nodes whose defoliation (i.e. after removing all leaves and petioles) is just a string of edges. That is, there are exactly two inner nodes to which of them there are attached two petioles (we call them heads or tails), any other inner node has exactly one petiole (called a leg) attached.



N otation 1.3. Let W be a (complex, nite dimensional) vector space with a distinguished basis, sometimes called letters: $f_0;_1;_2:p_3$. We consider the map :W ! C, such that (i) = 1 for every i, that is = i.

P Let \widehat{W} be a subspace of the second tensor product W W. An element $_{i;j}a_{ij}(_{i}_{j})$ of \widehat{W} can be represented as a matrix (a_{ij}) . Through the present paper we will assume that these matrices are symmetric, that is \widehat{W} is contained in $S^2(W)$.

Given a tree T and a vector space W, and a subspace \widehat{W} W W we associate to any vertex v of V(T) a copy of W denoted by W_v and for any edge e 2 E(T) we associate a copy of \widehat{W} understood as the subspace in the tensor product $\widehat{W} \stackrel{e}{=} W_{\mathfrak{G}_1(e)}$ W_{$\mathfrak{G}_2(e)$}. Note that although the pair f $\mathfrak{G}_1(e)$; $\mathfrak{G}_2(e)$ g is unordered, this de nition makes sense since \widehat{W} consists of symmetric tensors. Elements of $\widehat{W} \stackrel{e}{=}$ will be written as (symmetric) matrices ($a^{e}_{\mathfrak{G}_1(e)}$).

De nition 1.4. The triple (T;W; \Re) together with the above association is called a (symmetric, unrooted) phylogenetic tree.

Construction 1.5. Let us consider a linear map of tensor products

$$b: \widehat{W}^{E} = \bigcup_{e \ge E}^{O} \widehat{W}^{e} ! W_{V} = \bigcup_{v \ge V}^{O} W_{v}$$

de ned by setting its dual as follows

$$D (v_2 v_v) = e^{2E} (e_1 (e) e^{2E})_{\text{H} e}$$

where v_v stands for an element of the chosen basis f ig of the space W v_v . The complete a negeometric model of the phylogenetic tree (T;W; \widehat{W}) is the image of the associated multi-linear map

$$e : \bigvee_{e^{2E}} W_{e^{e}} = W_{v} = \bigvee_{v^{2V}} W_{v}$$

The induced rationalm ap of projective varieties will be denoted by :

$$\begin{array}{ccc} Y & & O \\ \vdots & P(\mathcal{W}^{e}) & ! & P(\mathcal{W}_{V}) = P(& \mathcal{W}_{V}) \\ e^{2E} & & v^{2V} \end{array}$$

and the closure of the image of is called the complete projective geometric model, or just the complete model of (T; W; \Re). The maps ^e and are called the parameterization of the respective model.

 G_{p} iven a set of vertices of the tree we can \hide" them by applying the map = _____i to their tensor factors. In what follows will hide inner nodes and project to leaves. That is, we consider the map

$${}_{\mathrm{L}} : \mathbb{W}_{\mathrm{V}} = {}^{\mathrm{N}}_{\mathrm{v2v}} \mathbb{W}_{\mathrm{v}} ! \quad \mathbb{W}_{\mathrm{L}} = {}^{\mathrm{N}}_{\mathrm{v2L}} \mathbb{W}_{\mathrm{v}}$$
$${}_{\mathrm{L}} = ({}_{\mathrm{v2L}} \operatorname{id}_{\mathbb{W}_{\mathrm{v}}}) \quad ({}_{\mathrm{v2N}} {}_{\mathbb{W}_{\mathrm{v}}})$$

Denition 1.6. The ane geometrical model of a phylogenetic tree (T;W; \hat{W}) is an an esubvariety of W_L = $_{v^{2}L}$ W_v which is the image of the composition = _ Respectively, the projective geometrical model, or just a model, denoted by X (T) is the underlying projective variety in P(W_L). For X = X (T) by O_X (1) we will denote the the hyperplane section bundle coming from the embedding in the projective space P(W_L).

Note that X (T) is the closure of the image of the respective rational map

which is defined by a special linear subsystem in the Segre linear system $j_{e^{2E}} p_{p(\mathcal{R}^{e})} O_{P(\mathcal{R}^{e})}(1) j$ where $p_{p(\mathcal{R}^{e})}$ is the projection from the product to the respective component. We will call this map a rational parametrization of the model.

The above de nition of param etrization is an unrooted and algebracized version of what is commonly considered in the literature, see e.g. [Allman, Rhodes '03], [Sturm fels, Sullivant] or [CGS].

1.2 Binary symmetric trees.

Depending on the choice of \Re W W we get di erent phylogenetic træes and their models. A natural assumption is that in the matrix representation the elements of \Re the sum of the numbers in each row and each column is the same (in applications, these numbers would stand for the probability distribution so their sum should be equal to 1). If W is of dimension 2 this is equivalent to saying that the respective matrix is of the form

for some a and b in C .

From now on we will consider binary symmetric phylogenetic trees, that is, we assume that dimension of W is 2 and \widehat{W} consists of matrices (tensors) satisfying the above symmetric condition. The elements of the distinguished basis of W will be denoted and . Note that \widehat{W} has dimension 2 as well. We will call them binary symmetric trees or just trees when the context is obvious. Our task is to understand geometric models of these trees.

E xam ple 1.7. Let T be a tree which has one inner node v_0 , three leaves v_1 ; v_2 ; v_3 whose petioles we denote, respectively, by e_1 ; e_2 ; e_3 . We denote the basis of W v_1 by

 $_i$; , while ${\bf \widehat{W}}^{~e_i}$ consists of m atrices $\begin{array}{cc} a_i & b_i \\ b_i & a_i \end{array}$. Then the parameterization m ap

 $e: \mathfrak{F}^{e_1} \quad \mathfrak{F}^{e_1} \quad \mathfrak{F}^{e_3} \quad ! \quad W_{v_1} \quad W_{v_2} \quad W_{v_3}$

is as follow s:

$e_{(a_1;b_1;a_2;b_2;a_3;b_3)} =$								
$(a_1a_2a_3 + b_1b_2b_3)$	6	2	₃ +	1	2	3) +		
$(b_1a_2a_3 + a_1b_2b_3)$	(2	₃ +	1	2	3) +		
$(a_1b_2a_3 + b_1a_2b_3)$	6	2	₃ +	1	2	3) +		
$(a_1a_2b_3 + b_1b_2a_3)$	6	2	₃ +	1	2	3)		

Notation 1.8. Let :W ! W be a linear involution () = , () = , the map

is relection with respect to the linear space W spanned by +. We note that on \widehat{W} the right and left action of coincide, i.e. $(id_{W})_{\widehat{M}} = (id_{W})_{\widehat{M}}$, and the resulting involution will be denoted by b, note that

 $b \quad \begin{array}{c} a & b \\ b & a \end{array} = \begin{array}{c} b & a \\ a & b \end{array}$

In particular, is identity on \Re .

G iven a binary symmetric tree (T;W; \Re) we de ne respective involutions:

$$V = V_{2V} V : W_{v} ! W_{v}$$

$$V_{2V} V_{2V} V_{2V}$$

$$V_{2V} V_{2V}$$

$$V_{2V} V_{2V}$$

$$V_{2V} V_{2V}$$

$$V_{2V} V_{2V}$$

$$W_{v} V_{v}$$

$$W_{v} V_{v}$$

$$V_{v} V_{v}$$

$$V_{v} V_{v}$$

Let $W_{v} = \begin{pmatrix} N \\ v_{2v} & W_{v} \end{pmatrix}^{v}$ and $W_{L} = \begin{pmatrix} N \\ v_{2L} & W_{v} \end{pmatrix}^{L}$ be their xed points, that is the maximal subspace on which v_{v} and, respectively, L acts trivially.

Lem m a 1.9. The image of is contained in P (W $_{\rm v}$) and the induced m ap

is Segre em bedding.

$$\begin{array}{c} b \quad (\begin{array}{c} & & \\ & & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & & \\ & & \\ &$$

(where, again, as in 1.5 v denotes either or in the space W_v) so that b v = bwhich implies v = b, hence im (b) W_v .

Next, let us note that dim $W_v = 2^{y j l}$ so that it is equal to dim \widehat{W}^{E} because $\underline{F} j = \underline{j} y j$ 1. The proof (e.g. by induction with respect to $\underline{j} y \underline{j}$) is instantaneous if one observes that the basis of W_v can be made of tensor products of (+ 1) and (1) eigenvectors of each $_v$ and thus W_v splits into the sum of (+ 1) and (1) eigenspaces of $_v$, each of the same dimension.

Now, to conclude the proof we have to show that ^b is in jective which is equivalent to ^b being surjective. Note that \widehat{W} is spanned by two form s:

 $\begin{array}{ccc} a & b \\ 0 & b & a \end{array} = a; \quad \begin{array}{ccc} a & b \\ 1 & b & a \end{array} = b$

and 1 = 0 b.

Now given an element $_{e2E}$ $_{i(e)}$ 2 $_{e2E}^{N} \Re^{e}$ we de ne inductively a sequence $_{v}$, indexed by vertices of T such that $b(_{v2V} _{v}) = _{e2E} _{i(e)}$. We choose a vertex v_{0} and set $_{v_{0}}$ to be either or . Now suppose that $_{v}$ are de ned for v in a subtree T⁰ of T. Suppose that v^{0} is not in T⁰ but is joined to a vertex v^{0} in T⁰ by and edge e^{0} . Then we set $_{v^{0}} = _{v^{0}}$ if $_{i(e^{0})} = _{0}$ or $_{v^{0}} = (_{v^{0}})$ if $_{i(e^{0})} = _{1}$.

N otation 1.10. Let us choose a node v 2 N . We consider an involution b_v^E on the space $\hat{W}^E = \bigcup_{e \ge E} \hat{W}^e$. First, for any e 2 E we set $b_v^e = id_{N_e}^e$ if v is not a vertex of e and $b_v^e = b$ if v is a vertex of e. Next, we de ne $b_v^E = \bigcup_{e \ge E} b_v^e$. Let G_N be the group of autom orphism s of \hat{W}^E generated by involutions $\sum_{v=1}^{E} b_v^e$, for v 2 N.

The following observation about a convenient choice of coordinates is sometimes referred to as a Fourier transform, see e.g. [Sturm fels, Sullivant].

Lem m a 1.11. $G_N = Z_2^{\mathcal{N} j}$, the action of G_N restricts to $Q_{e^2 E} \widehat{W}^e$ and it is is diagonalizable.

Proof. This follows immediately from the denition of the action of b_v^e on \mathfrak{A}^e . Namely, in basis of \mathfrak{A}^e consisting of a + b and a - b the action of \mathfrak{B}^e is diagonal. \Box

Lem m a 1.12. The m ap b _L m aps (W_L) invectively into the space ($(\Re ^{E})$)^{G_N} on which G_N acts trivially.

Proof. The proof is similar to that of 1.9. Firstly, the map L_{L} is injective on (W_{L}) because of 1.9 and injectivity of L_{L} . Next, we note that the action of G_{N} is trivial on its image. Indeed, we de ne $V_{V} = V_{W2V} V_{V}$, where, for any w 2 V we set $b_{V}^{W} = id_{W_{N}}$ if $W \in V$ and $V_{V} = V$. Then for $V \ge N$ we have

$$(b_v^E)$$
 b = b (v_v^V) and $v_v^V = v_v^V$

where the rst equality follows directly from the de nition of the map b, 1.5. This in plies (b_v^E) $b_L = b_L$ which is what we want.

W ewillprove that, in fact, ^b $_{L}$ is an isom orphism , 2.12, so that the geom etric m odel of the tree is de ned by G^{N} invariant sections of the Segre linear system. Because of 1.11 G^{N} can be treated as a subgroup of a complex torus and thus we can use toric geom etry.

2 Toric geom etry.

Sum m ary: W e study invariants of an action of Z_2^N on $(\mathbb{P}^1)^{\frac{p}{2}}$ and a related polytope in the cube $[0;1]^{\frac{p}{2}}$ which we call a polytope m odel of the tree. The polytope m odels are used to de ne geom etric m odels in terms of toric geom etry. These polytopes turn out to be ber products of elementary ones. This leads to interpreting the geom etrical m odel of a tree as a quotient of products.

2.1 Lattice of a tree and the action of the torus

Given a tree T we encode it in terms of dual lattices.

Denition 2.1. Let T_L be a tree with the set of vertices V and the set of edges E. We set M = M (T) = $_{e^{2E}}Z$ e to be a lattice, or free abelian group, spanned on the set E. Let N = N (T) = H om (M; Z) be the dual lattice. We represent elements of V as elements of N. Namely, for $v \ge V$ we set v(e) = 1 if e contains the vertex v and v(e) = 0 otherwise. The pair (M; N) together with the choice of the basis E of M and set V N is called the lattice pair of the tree T.

From this point on we identify the edges and the vertices of T with the respective elements in M (T) and N (T). The elements of the basis of N dual to fe 2 Eg will be denoted by e. Then for any v 2 V we have, by de nition, $v = \begin{bmatrix} e^{3v} e & :N \end{bmatrix}$ Z. In particular, v is a leaf if and only if v = e for som e e, which is a petiole for v.

Let us recall that $jV j = j \pm j + 1$ so the set of vertices has to be linearly dependent in N. The set of vertices of T can be divided into two disjoint classes, say $V = V [V^+, each class consisting of vertices which can be reached one from another by passing through an even number of edges.$

Lem m a 2.2. The equality $P_{v2V} v = P_{v2V^+} v$ is, up to multiplication by a constant, the only linear relation in N between vectors v from V. In particular, any proper subset of V consists of linearly independent vectors in N.

Proof. Suppose that
$$a_v = 0$$
, for som $e_va^2 k$. For any $e^2 E$ we have
 X
 $(a_v = 0) (e) = e_{1(e)}^2 + a_{e_2(e)}$

and therefore $a_{\theta_1(e)} = a_{\theta_2(e)}$. Thus we get the desired relation.

The operations on trees can be translated to lattices, here is an example.

C on struction 2.3. Let v_0 be a 2-valent inner node of T which belongs to exactly two edges e_1 and e_2 . Let T_{v_0} be a tree obtained by removing the node v_0 from T and replacing the edges e_1 and e_2 by a single edge e_0 . Let (M; N) be the lattice pair of T. We set M_{v_0} M to be the kernel of q_0 e and $N_{v_0} = N = Z$ (e q), clearly M_{v_0} and N_{v_0} are dual. We den $e_0^0 = e_1 + e_2$. Note that $E_{v_0} = E n fe_1; e_2g [fe_0^0g$ is a basis of M_{v_0} . For $v \ge V n fv_0 g$ by v^0 we denote the in age of a vertex v under the projection N ! N_{v_0} and set of all v^0 we denote by V_{v_0} . One can verify easily the follow ing.

Lem m a 2.4. The above de ned pair $(M_{v_0}; N_{v_0})$ together with the above choice of E_{v_0} and V_{v_0} is the lattice pair of the tree T_{v_0} obtained from T by removing the 2-valent inner node v_0 .

Now we set up the toric environment.

Construction 2.5. We deal with a binary symmetric tree (T;W; \Re). Because of 1.11 for any edge e 2 E there exists an inhom ogeneous coordinate z_e on $P_e^1 = P(\Re^e)$ such that for v = 1 there exists a follows $\sum_{v=1}^{E} (z_e) = -z_e$.

Let $T = \sum_{e^{2E}} C$ be a torus with coordinates $fz_e \ 2 \ C : e \ 2 \ Eg$ and with the natural action Y = Y

$$\begin{array}{cccccccc} & Y & Y \\ T & P_e^1 & ! & P_e^1 \\ & e^{2E} & e^{2E} \end{array}$$

which is the multiplication of z_e 's coordinate wise. We consider an injective map : G_N ! T = ${}_{e^{2E}}C$ which is dened as follows. For any ${}_v^E \ 2 \ G_N$ we take ${}_v^E \ 2 \ T$ such that $z_e(\ {}_v^E \)) = 1$ if $v \ 2 \ e$ and $z_e \ {}_e^{(E_v)} \) = 1$ if $v \ B \ e$. Then extends to a homomorphism of groups : G_N ! T = ${}_{e^{2E}}C$ and the action of G_N on ${}_{e^{2E}}P_e^1$ factors through $\ .$

We explain this situation using the lattices of the tree T and toric geom etry formalism. Our notation is consistent with this of standard toric geom etry textbooks, e.g. [O da] or [Fulton]. We take the torus $T = T_N = N_Z C$ with coordinates $z_e = e^{e}$, where e 2 E M is the distinguished basis. Recall that the elements of M can be identified with monomials in coordinates z_e , that is, each u 2 M such that $u = a_i e_i$ represents a monomial $u = z_{e_i}^{a_i}$. Forw 2 N and t 2 C the z_e -th coordinate of the respective point on $T_N = N_Z C$ is as follows $z_e (w = t) = t^{e} (e^{e})$. Moreover, recall that every element w of N can be identified with algebraic 1-parameter

subgroups $_{w}$ of T_{N} . That is, for $w \ge N$ and $t \ge C$ we set $_{w}$ (t) (z_{e}) = t^{w (e)} $_{e}z$ In short, $N = Hom_{alg}(C; T_{N})$ and $M = Hom_{alg}(T_{N}; C)$, Fulton, Sect 2.3]

The com plexi ed lattice N_C = N_ZC can interpreted as the tangent space to the unit element in the torus T_N and we have the natural exponential map N_C ! T_N. In particular, $z_e (exp (2 i(w))) = exp (2 i(w (e))$. The image of the real vector space N_R N_C under this exponential map is the maxim alcom pact real subgroup S^1 of T_N. Using the exponential map we can relate the vertices v 2 N viewed as elements of the lattice N to their respective automorphism s $\frac{E}{v} 2 G^N$. The following lemma is not used directly in our arguments so we skip its proof.

Lem m a 2.6. For every v 2 V we have $\exp(2 i(v=2)) = \binom{E}{v}$. If $\overset{P}{\mathbb{N}}$ is the lattice spanned in N_R by N and N = 2 = fv=2 : v 2 N g then the inclusion N ,! $\overset{P}{\mathbb{N}}$ yields an exact sequence of groups

$$O ! (G^{M}) ! T_{N} = N _{Z} C ! M^{P} _{Z} C ! O$$

For our purposes we need the following lemma which provides a clear description of functions on the torus $T_{\rm N}\,$ which are invariant with respect to the action of G $_{\rm N}$.

Lem m a 2.7. A monom ial function ^u on T_N with u 2 M is invariant with respect to the action of $\frac{E}{V}$ if and only if v(u) 2 2Z.

Proof. First, we note that, by de nition, $_{v}^{E}(e) = _{v}^{E}(z_{e}) = (1)^{v(e)} e^{z} = (1)^{v(e)} e^{z}$. Next, we write the exponent of the monom ial u in terms of the distinguished coordinates: $u = e^{2E} e(u)e$. Then, since $_{v}^{E}$ is a homomorphism, we get

$${}^{E}_{v}({}^{u}) = (1)^{P}_{e}e^{(u)v(e)} {}^{u} = (1)^{v(u)} {}^{u}$$

which concludes the proof.

De nition 2.8. Given a tree T with the lattice pair (M; N) = (M(T); N(T)) we de ne its normalized lattice pair (M; N) = (M(T); N(T)) as follows: M =fu 2 M : 8v 2 V v(u) 2 2Zg and N(T) is a dual of M(T) which contains N and the set N = 2 = fv=2 : v 2 N g.

In view of 2.7 the lattice \mathbb{M} contains m onom ials which are $\mathbb{G}^{\mathbb{N}}$ invariant.

2.2 Polytope model of a tree

The complete Segre linear system on ${}^{Q}_{e2E} P_{e}^{1}$ is spanned on monomials ${}^{Q}_{e2E} z_{e}^{e}$ where ${}_{e} 2$ f0; 1g. Equivalently, once the big torus action T_{N} on ${}^{P}_{e}^{1}$ is chosen, the complete Segre system is represented by vertices of the unit cube ${}_{M} = fu 2 M_{R}$: $8_{i} 0 = q(u)$ 1g in the space of characters M_{R} , or by zero-one sequences indexed by E.

Because of 1.9 and 1.12 we are interested in subsystems of the Segre linear systems or, equivalently, subsets of of vertices of $_{\rm M}$. If is a polytope in M $_{\rm R}$ whose vertices are contained in the set of vertices of $_{\rm M}$ then we call it a subcube.

Denition 2.9. Given a binary tree T with its lattice pair (M;N) its pplytope model (T) is a polytope in the lattice M which is the convex hull of $fu = a_i e_i 2 M$: $a_i = 0; 1 \text{ and } v(u) 2 2Z$ for every v 2 N g.

We note that the vertices of are precisely these among vertices of $_{M}$ which are in the sublattice M M and because of 2.7 they are exactly these monomials in the complete Segre system which are invariant with respect to the action of G_{N} .

Since the cube $_{\rm M}$ is the fundam ental dom ain in dividing M modulo 2 we can interpret the elements of the complete Segre system as points in the linear space M $_{\rm Z_2}$ = M $_{\rm Z}$ Z₂.

Lem m a 2.10. If the vertices of the cube $_{M}$ are identified with the points in linear space M Z_{2} then vertices of (T) form the linear subspace N $^{?}$ M Z_{2} of zeros of form s v 2 N Z_{2} , where v 2 N (T).

Proof. This is a restatem ent of 2.7.

Corollary 2.11. The polytope (T) has $2^{j j l}$ vertices.

Proof. We use 2.10: by 2.2 the elements v's are linearly independent in N Z_2 so dimension of the space of their zeroes in M Z_2 is $\pm j$ $N = j_1 j 1$.

Using the above information we can conclude identifying the linear subsystem in the Segre system which de nes the projective model of a binary symmetric tree.

Theorem 2.12. In the situation of section 1.2 the map b_{L} maps (W_{L}) isomorphically to ((M^{E}))^{G_N}. In particular, in terms of the toric coordinates on $Q_{e^{2E}} P(M^{e})$ introduced in section 2.1, the rational parametrization map

is dened by elements of the Segre linear system on ${}^{Q}_{e^{2}E} P (\Re^{e})$ which are associated to vertices of (T).

Proof. By the construction the vertices of (T) are these monomial in the Segre system which are invariant with respect to the action of G^N . In other words they form a basis for $((\Re^{E}))^{G_N}$. In 1.12 have proved that the parametrization map injects (M_L) into the space $((\Re^{E}))^{G_N}$ and now by 2.11 they are of the same dimension so this is an isomorphism.

Thus we have determ ined that studying projective geometric models of binary symmetric trees is essentially equivalent to understanding their polytopes. We start with the simplest, in fact trivial, example.

Example 2.13. Let T be a tree consisting of two leaves, two petioles e_1 and e_2 , and one inner node v_0 . Then (T) is spanned on 0 and $e_1 + e_2$.

M ore generally we have the following result which extends 2.4.

Lem m a 2.14. Suppose that T is a tree with a 2-valent node v_0 . Let T_{v_0} be a tree obtained from T by removing v_0 , as in the situation of lem m a 2.4. Then, under the natural inclusion M (T_{v_0}) M (T) we have (T_{v_0}) = (T) and M (T_{v_0}) = M (T) \ M (T_{v_0})_R

Proof. We use the notation of 2.4, in particular e_1 and e_2 denote the edges containing v_0 and M $(T_{v_0}) = M_{v_0} = \ker(e_2 \quad e_1)$. Note that the parity of the node $v_0 = e_1 + e_2$ is equivalent to that of $e_2 \quad e_1$, in particular for $u \ 2 \ M (T_{v_0}) = \ker(e_2 \quad e_1)$ we have v_0 (u) 2 2Z. Since N (T) = N (T_{v_0}) [fv_0 g the conditions defined by N (T) and N (T_{v_0}) on M (T_{v_0})_R are the same. Similarly, the conditions defining (T) and (T v_0) inside M $v_0 \quad R \ M = M_{v_0}$ are the same.

By 2.23 we have that rem oving the 2-valent node does not change the model of the tree. Thus, from now on we consider trees with no 2-valent nodes.

A star tree is a tree which has exactly one inner node, a star tree with d leaves will be denoted by T d .

Lem m a 2.15. If d 3 then vertices of (T^{a}) generate \mathbb{N} (T^{a}) , in particular dim $(T^{a}) = d$.

Proof. If fe_ig is the set of edges then (T^d) contains sum $se_i + e_j$ for all possible pairs $i \in j$ and for d 3 they span M (T^d) which is of index 2 in M (T^d).

E xam ple 2.16. The vertices of the polytope (T^{3}) with edges e_{0} ; e_{1} ; e_{2} are as follows: $0, e_{1} + e_{2}, e_{2} + e_{0}$ and $e_{0} + e_{1}$ so that (T^{3}) is a 3-dimensional tetrahedron. If M is the sublattice spanned by the vertices of (T^{3}) then $M = M = Z_{2}$.

The inequalities de ning (T^{-3}) are as follows

(v=2)() 1; $(v=2_{0})(e)$ 0; $(v=2_{1})e()$ 0; $(v=2_{2})e()$ 0: where, recall, $v = e_{0} + e_{1} + e_{2}$.

C on struction 2.17. A pointed tree (T; ') is a pair consisting of a tree T and a leaf '2 L (T). Given two pointed trees $(T_1; '_1)$ and $(T_2; '_2)$ we de ne their graft as follows: $T = T_1 \xrightarrow{i_1-i_2} T_2$ is a tree obtained by removing from each T_1 the leaf 'i and identifying their respective petioles which becomes an inner edge of the resulting tree T.

For example, a graft of two trees of type T 3 with distinguished leaves denoted by is the following operation

$$\succ$$
 _ _ < = \succ

Let us take two lattices M₁ and M₂ with distinguished bases $fe_0^i; :::; e_{m_1}^i g$, for i = 1; 2 and respective subcube polytopes $_i$, each of them having the set of vertices A_i. Let $i = (e_0^i) : M_i ! Z$ be the projection to the zeroth coordinate and, by abuse of notation, by the same letter we will denote its composition with the projection M₁ M₂ ! M_i ! Z. Now we can take ber product of each of these objects, relative over the projection, e.g. M₁ $i_1 i_2 M_2 M_1 M_2$ consists of pairs $(u_1; u_2)$ such that $i_1(u_1) = i_2(u_2)$. In other words M₁ $i_1 i_2 M_2 = ker(i_1 i_2)$ and $1 i_1 i_2 i_2 = (i_1 i_2) \setminus ker(i_1 i_2)$

Lem m a 2.18. In the above situation = $_{1 i_{1} i_{2} 2}$ is a subcube polytope in $M = M_{1 i_{1} i_{2}}M_{2}$ with the set of vertices $A = A_{1 i_{1} i_{2}}A_{2}$. In general, if $_{i} (M_{i})_{R}$ and $i_{i}: M_{i} !$ Z are lattice hom on orphism s such that $i_{i}(i)$ [0;1] then the set of vertices of $= _{1 i_{1} i_{2} 2}$ is the ber product of the vertices of $_{i}$'s.

Proof. The only non-trivial thing is to show that all vertices of are in the ber product of vertices of $_1$ and $_2$. Since = $(_1 _ 2) \setminus \ker(1 _ 2)$ is a codimension 1 linear section of $_1 _ 2$ its vertices are either vertices of $_1 _ 2$ (which is what we want) or are obtained by intersecting the hyperplane ker $(1 _ 2)$ with an edge of $_1 _ 2$. To this end, let us take two pairs of vertices $(u_1^1; u_2^1); (u_1^2; u_2^2),$ where u_1^i is a

vertex of j. Suppose that for som et 2 (0;1) the point $u = t(u_1^1; u_2^1) + (1 t)(u_1^2; u_2^2)$ is in ker('1 '2), that is, we have

$$'_{1}(tu_{1}^{1} + (1 t)u_{1}^{2}) = '_{2}(tu_{2}^{1} + (1 t)u_{2}^{2})$$

and moreover $\mathbf{1}_1(\mathbf{u}_1^1) \notin \mathbf{1}_2(\mathbf{u}_2^1)$. Thus, we may assume that $\mathbf{1}_1(\mathbf{u}_1^1) = 0$ and $\mathbf{1}_2(\mathbf{u}_2^1) = 1$. Hence, because of the above equality and since $t \notin 0$; 1, we get $\mathbf{1}_1(\mathbf{u}_1^2) = 1$, $\mathbf{1}_2(\mathbf{u}_2^2) = 0$ and t = 1=2. So $u = t(\mathbf{u}_1^1; \mathbf{u}_2^1) + (1 t)(\mathbf{u}_1^2; \mathbf{u}_2^2) = t(\mathbf{u}_1^1; \mathbf{u}_2^2) + (1 t)(\mathbf{u}_1^2; \mathbf{u}_2^1)$ and $(\mathbf{u}_1^1; \mathbf{u}_2^2)$; $(\mathbf{u}_1^2; \mathbf{u}_2^1)$ are vertices of so u lies in the interior of an edge of . \Box

Example 2.19. Let us consider two copies of a tetrahedron, as in example 2.16. That is, for i = 1;2, in a lattice $M^{i} = Ze_{0}^{i}$ Ze_{1}^{i} Ze_{2}^{i} , we consider a tetrahedron $_{i}^{3}$ spanned on vertices 0, $e_{0}^{i} + e_{1}^{i}$, $e_{1}^{i} + e_{2}^{i}$ and $e_{2}^{i} + e_{0}^{i}$. We take the projections $(e_{0}^{i}) : M^{i} ! Z$ and in the berproduct

$$M = M_1 \cdot V_2 M_2 = \ker (e_0^2 \cdot e_0^2) \qquad M_1 \cdot M_2$$

by e_0 we denote the element $e_0^1 + e_0^2$. The resulting ber product of tetrahedra

$$= \begin{pmatrix} 3 & 3 \\ 1 & 2 \end{pmatrix} \setminus M_{R}$$

has the following vertices: $0;e_1^1 + e_2^1;e_1^2 + e_2^2;e_1^1 + e_2^1 + e_2^2;e_0 + e_1^1 + e_2^2;e_0 + e_1^1 + e_2^2;e_0 + e_1^1 + e_2^2;e_0 + e_2^1 + e_2^2;e_0 + e_2^2;$

P roposition 2.20. Let $(T_1; '_1)$ and $(T_2; '_2)$ be two pointed trees. Then

 $M (T_1 \xrightarrow{1} T_2) = M (T_1) \xrightarrow{1} T_2 M (T_2)$ $(T_1 \xrightarrow{1} T_2) = (T_1) \xrightarrow{1} T_2 (T_2)$

Proof. Let $M_1 = M_1(T_1)$, $M_2 = M_1(T_2)$, and similarly for N's, M'' and M''s. We set $T = T_1 \xrightarrow{i_1-i_2} T_2$. Then, by construction 2.17, $M = M_1(T) = M_1 \xrightarrow{i_1-i_2} M_2$ and $M = M_1 \xrightarrow{i_1-i_2} M_2$. The two projections $p_1 : M_1 M_1$ yield respective injections of Hom (;Z)-spaces: $i_1 : N_1 I N_1$ (in fact $N = (N_1 N_2) = Z(i_1 I_2)$). If N_1 and N_1 denote, respectively, inner nodes of T_1 and T then $N = I_1(N_1) [I_2(N_2)]$. Since M_1 , M_1 , M_2 are dened by extending $N_1 N_1$, N_2 by N = 2, $N_1 = 2$ and $N_2 = 2$, respectively, it follows that $M = M_1 + M_2$ in N_R . This in plies the rst equality of the lemma. Similarly, since the set N_1 determines vertices of M_1 which span (T), see 2.9, we get the second equality.

The above result can be expressed as follows: the polygon of a tree T is a ber product of polygons of star trees associated to inner nodes of the tree, bered over the relations encoded in the inner branches of the tree. Since (T^{-3}) is a 3-dimensional tetrahedron this is especially straightforward in case of 3-valent trees.

For any inner node v 2 N of a 3-valent tree T we consider the lattice M_v = $Ze_1^v \quad Ze_2^v \quad Ze_3^v$, where e_1 , e_2 and e_3 are the three edges stem m ing from v. Inside M_v we have the tetrahedron v with vertices 0, $e_1^v + e_2^v$, $e_2^v + e_3^v$ and $e_3^v + e_1^v$. We consider the big lattice $M = {}_{v2N} M_v$ and M_R contains the product ${}_{v2N} v$. Now for each inner edge e we have a form e : M ! Z such that $e(e_1^v)_T = (1)$ if e(e) = v and $e_1^v = e$, and $e(e_1^v) = 0$ otherwise. Then the intersection ${}_{e2E^\circ}$ ker(e) can be identied with the lattice M, that is, we map $e 2 E^\circ$ to $e_1^v + e_{1^0}^v$ where $e(e) = (v; v^0)$, and $e_1^v = e_{1^0}^{v^0} = e$, while a petiole e is just mapped to its unique representation in M. Then, by 2.20 we get



2.3 Geometric model of a tree

First, let us recall the construction of a projective torivariety from a lattice polytope of characters. Let \mathbb{M} and \mathbb{N} be dual lattices of characters and 1 parameter subgroups for a torus $T_{\mathbb{N}} = \mathbb{N}_{2} \mathbb{C}$.

De nition 2.21. A lattice polytope M_R is called norm al if

the sublatice of \mathbb{M} spanned by the di erences of points in $\setminus \mathbb{M}$ is equal to \mathbb{M}

for every integer d 0 any point in d $\$ is equal to a sum of d points in $\$ M .

Equivalently, the second condition in the above de nition can be restated as follows. Let $\mathbb{M}^0 = \mathbb{M}$ Z and take an a nem ap $i_1 : \mathbb{M}^1 : \mathbb{M}^0$ such that $i_1 (u) = (u; 1)$. Then is normal in \mathbb{M}^1 if and only if the sem igroup spanned in \mathbb{M}^0 by $i_1 (\setminus \mathbb{M})$ is equal to the sem igroup of lattice points in cone spanned in \mathbb{M}^0_R by $i_1 ()$, that is the sem igroup R $_0 (i_1 ()) \setminus \mathbb{M}^0$.

Denition 2.22. Suppose that is a normal polytope in \mathbb{M} . Let A^d be a C-linear space with the basis f^u: u 2 d \ \mathbb{M} g. We consider a graded C-algebra A() = $_{d 0} A^d$, with multiplication $^{u_1 u_2} = ^{u_1+u_2}$. Then X() = ProjA() is called the projective model of .

We note that in the above situation A () is a norm along, that is, it integrally closed in its eld of fractions. This, by de nition, is equivalent to saying that a ne spectrum Spec(A()) is a norm alow a variety. In fact, in such a case A() is the sem igroup algebra of R₀(i₁()) \ M⁰ so Spec(A()) is an a netoric variety with the big torus T_{N^0-Z} . In the projective case we have the following general result which summarizes properties of the projective model of a norm algolytope, see [0 da, Sect. 2.1{2.4}, [Sturm fels] or [Fulton].

P roposition 2.23. Suppose that is a normal polytope in the lattice M of characters of a torus T_{vb} . Then the following holds:

- 1. X () is a toric variety on which T $_{\rm wb}$ acts e ectively,
- 2. X () is embedded in P $^{j \ M} j \,^{1}$ as a projectively normal variety such that H 0 (X ();O $_{X}$ (d)) = A d ,
- 3. Characters from $\ \ M$ de ne a diagonal action of T_p on $P^{j \ M j l}$ which restricts to the torus action on X (),
- 4. The induced action of $T_{_{\rm ff}}\,$ on H 0 (X ();O $_{\rm X}\,$ (d)) is linearizable with weights in d $\,\setminus\,$ IF .
- 5.X () $P^{j \setminus M j l}$ is the closure of the image of the map T_{M} ! $P^{j \setminus M j l}$ de ned by the characters from $\setminus M$

Because of A.5 the polytope model (T) of a 3-valent tree T is normal so we can consider its projective model. The following is the key result of the paper which allows us to study projective models of binary symmetric trees in purely toric way.

Theorem 2.24. Let (T;W; \Re) be a binary symmetric 3-valent tree. Then the varieties X (T) and X ((T)) are projectively equivalent in P (W_L) = P^{2^{jL j 1} 1}.

Proof. By 2.12 the parametrization of X (T) is defined as a rational map from $Q_{e^{2E}} P(\mathbf{M}^{e})$ defined by characters of torus T_N which are vertices of (T). Thus, X (T) is the closure of the respective map $T_N \ ! \ P^{2^{|\mathcal{I}|\mathcal{I}|} \ 1}$. Since \mathbf{M} is the sublattice spanned by vertices of (T) this factors to the map $T_{\mathbb{N}}^{e}$! $P^{2^{|\mathcal{I}|\mathcal{I}|} \ 1}$ the image of which defines X (), 2.23.

2.4 1-param eter group action, quotients.

In this section we consider quotients of projective varieties as in M um ford's G IT [M um ford]. For a comprehensive exposition of the theory, including a relevant de – nition of good quotient we refer to [B ialynickiB irula]. In the present section as well as in section 3.3 we consider an algebraic action of a torus T on a projective variety X , P^{m} which is given by a choice of weights hence it extends to the a ne cone over X and thus it determ ines its linearization, its set of sem i-stable points X ^{ss} and its good quotient X ^{ss} ! X ^{ss}=T, see [B ialynickiB irula, Ch.6].

C on struction 2.25. Let $_{i}$ $(\mathbf{F}_{i})_{R}$, for i = 1; 2 be two lattice polytopes adm itting unimodular covers hence normal, see A.1, and X ($_{i}$) $P^{n_{i}-1}$, where $n_{i} = \mathbf{F}_{i} \setminus_{i} \mathbf{F}_{i}$ their associated toric varieties. In $M = \mathbf{F}_{1} \quad \mathbf{F}_{2}$ we take the product polytope $= 1 \dots 2$ which is also normal, referoduct-unimodular. Then the associated toric variety X = X ($_{i}$) $P^{n_{1}n_{2}-1}$ is the Segre in age of X ($_{1}$) X ($_{2}$).

Suppose that 'i : M i ! Z are lattice hom on orphism s such that ('i)_R (i) [0;1]. We pull 'i to the product of lattices and on M 1 M 2 we dene the form ('1 '2). The form deness a diagonal action '1 '2 of C on X $P^{n_1n_2 - 1}$ which on the coordinate associated to ${}^{(u_1,u_2)}$, where $u_i 2$ i \ M i, has the weight '1 (u_1) '2 $(u_2) 2$ f 1;0;1g. A coordingly, we regroup the coordinates of $P^{n_1n_2 - 1}$ and write them as $[z_i; z_j^0; z_k^+]$ depending on whether they are of weight 1, 0 and 1, respectively. That is

$$x_1 x_2$$
 (t) $[z_1; z_j^0; z_k^+] = [t^1 z_1; z_j^0; tz_k^+]$

The above formula de nes the action of $\frac{1}{1} \frac{1}{2}$ on the cone over X and thus a C -linearization of the bundle O_X (1) in the sense of G IT . By X⁰ let us denote the intersection of X with the complement of the space spanned on the eigenvectors of $\frac{1}{1} \frac{1}{2}$ of weight \neq 0, that is X⁰ = X nf[z₁; z₁⁰; z_k⁺]: 8j z₁⁰ = 0g

We set $\mathbb{M} = \ker(\mathbf{1} \ \mathbf{2})$ and $= \ \ \ker(\mathbf{1} \ \mathbf{2}) = \ \mathbf{1} \ \mathbf{1} \ \mathbf{2} = 2$. By A.4 is a norm alpolytope and by X () we denote its associated toric variety.

P roposition 2.26. In the above situation the set X 0 is equal to the set of the sem istable points of the action of i_1 i_2 . The projection to the weight 0 eigenspace $[z_{i}^{0};z_{j}^{0};z_{k}^{+}]$? $[z_{j}^{0}]$ de ness a regular map of X 0 to X () and X () is a good quotient for the action of i_1 i_2 .

Proof. The sections of O_x (m) form > 0 m ake a vector space spanned on ^u, where u 2 m $\setminus M$. Am ong them, these which are invariant with respect to the action of ^{'1} ^{'2} are associated to u's in the intersection with ker '1 ² thus in m $\setminus M$. By the norm ality of , see A.4, the algebra of invariant sections is generated by these from O_X (1). Thus the set of sem istable points of the action of $\frac{1}{1} \frac{1}{2}$ is where at least one of the coordinates z_j^0 is non-zero and the quotient m ap is the projection to the weight zero eigenspace.

C orollary 2.27. Let $(T_1; `_1)$ and $(T_2; `_2)$ be two pointed trees. Then X $(T_1 :_{!-`2} T_2)$ is a good quotient of X (T_1) X (T_2) with respect to an action of $`^1 :_2$.

E xam ple 2.28. Consider the C action on the product $P_1^3 = P_2^3$ given by the form ula:

(t)
$$([z_0^1; z_1^1; z_2^1; z_3^1]; [z_0^2; z_1^2; z_2^2; z_3^2]) = ([z_0^1; tz_1^1; tz_2^1; z_3^1]; [z_0^2; t^{-1}z_1^2; t^{-1}z_2^2; z_3^2])$$

where the superscripts of the coordinates indicate the factor in the product $P_1^3 = P_2^3$. The following rational map $P_1^3 = P_2^3 = P^7$ is equivariant and regular outside the set $fz_0^1 = z_3^1 = z_1^2 = z_2^2 = 0$ g [$fz_1^1 = z_1^2 = z_0^2 = z_3^2 = 0$ g, each component of this set is a quadric $P^1 = P^1$:

If $[x_0; ::: x_7]$ are coordinates in P⁷ then the intersection of two quadrics $fx_0x_7 = x_1x_6g \setminus fx_2x_5 = x_3x_4g$.

The above claim will be clear if we write functions $z_i^1 z_j^2$ in terms of characters of the respective torus, which we denote by e_i^1 and e_j^2 , respectively. Namely, dividing the right hand side of the above displayed formula by $z_0^1 z_0^2$ we get the following sequence of rational functions:

$$\begin{bmatrix} 1 \\ e_1^2 + e_2^2 \\ e_0^1 + e_1^1 \\ e_0^2 + e_1^2 \\ e_0^1 + e_1^1 \\ e_0^2 + e_1^2 \\ e_0^2 + e_2^2 \\ e_0^1 + e_1^1 \\ e_0^2 + e_1^2 \\ e_0^2 + e_1^2 \\ e_0^1 + e_1^1 \\ e_0^2 + e_2^2 \\ e_0^1 + e_1^1 \\ e_0^2 + e_2^2 \\ e_0^1 + e_1^2 \\ e_0^2 + e_1^2 \\ e_0^2$$

If we write the sum s of the exponents of the above rational functions in M₁ M₂ and call $e_0 = e_0^1 + e_0^2$ then we get the vertices of (T ³_T T ³) which we computed in example 2.19. From the above formula we can read the weights with which 1-parameter groups (e_j^i) , for i; j = 1; 2, associated to leaves, act on the quotient variety in P⁷.

3 3-valent binary trees.

Sum m ary: From this point on we concentrate on understanding varieties associated to 3-valent binary trees and we prove m ain results of the present note which are as

follows: (1) such varieties have only G orenstein term inal singularities and are Fano of index 4, (2) any two such varieties associated to trees with the same number of leaves are in the same connected component of the H ilbert scheme of the projective space, (3) their H ilbert-Ehrhard polynom ial can be computed e ectively..

3.1 Paths, networks and sockets.

Let T be a 3-valent binary symmetric tree. In section 2.2 we identified the variety X (T) in $P^{2^{j \cdot j \cdot 1} - 1}$ with the closure of the image of a torus map dened by a polytope

(T). We recall that the linear coordinates on the ambient projective space can be idential with the vertices of (T) which are among the vertices of the cube $_{\rm M}$ satisfying parity relation with respect to the form s v 2 N N,2.9. For 3-valent trees we have a convenient interpretation of these points.

Denition 3.1. Let T be a 3-valent tree. A path on T of length m 1 is a choice of m + 1 distinct vertices v_0 ;:::; v_m such that v_0 and v_m are leaves (called the ending points of) and there exists m edges, e_i ;:::; e_m such that for i = 1;:::;m it holds $(e_i) = fv_{i-1}$; v_i g.

A network of paths (or just a network) on T is a set of paths (possibly an empty set), each two of them have no common vertex (neither edge). For any network of paths on T we de ne the socket () L to be the set of leaves which are ending points of paths in .

A tree T is labeled if its leaves are numbered by $1; :::; j_j j_i$ A subset L is represented by a characteristic sequence $(1); :::; (j_j)$ in which (i) = 1 or 0, depending on whether the leaf numbered by i is in or not.

Sockets of networks will identied by their characteristic binary sequences. We note that, clearly, every socket consists of even number of elements in L .

E xam ple 3.2. Let us consider a labeled 3-valent tree with four leaves. In the following diagram, in the upper row we draw all possible networks on this tree, where paths are denoted by solid line segments. In the lower row we write down the respective sockets in terms of characteristic sequences of length four

Lem m a 3.3. Let T be a 3-valent tree. A ssociating to a network a point u() = e (e) e 2 M (T), where (e) = 1;0 depending on whether e is on or not, de nes a bijection between networks and vertices of (T)

Proof. First note that u() 2 (T). To de ne the inverse of ! u(), for any vertex $u = e^{2E} e = 2$ (T) we de ne the support of u consisting of edges of T whose contribution to u is nonzero, i.e. fe 2 E : e (u) = 1g. The parity condition 8_{v2N} either v(u) = 0 or v(u) = 2 yields that these edges de ne a network on T.

We note that, because of 2.11, there are $2^{j_{L}j}$ ¹ networks. On the other hand, the association of the socket to a network gives a map from the set of networks to the subsets of leaves. This map is surjective, that is, every subset of L with even umber of elements is a socket of a network. Indeed, this follows by a straightforward induction with respect to the number of leaves of the tree: in the induction step we write a tree T_{n+1} with n + 1 leaves as a graft of a tree T_n with n leaves and a star tree with 3 leaves and consider three cases depending on how many of the two new leaves replacing one old are in the set L.

Finally, because the number of all subsets of L with even number of elements equals to $2^{j \cdot j - 1}$ we get the following.

Lem m a 3.4. Let T be a 3-valent tree. Then associating to a network its socket de nes a bijection between the set of networks of paths on T and the set of subsets of L which have even number of elements.

We note that the sockets of a tree T form a convenient basis in the space W_L, which was introduced in section 1.2. Indeed, in order to use toric arguments we have diagonalized the action of the involution on W with a basis ₀, ₁ such that $(_i) = (1)^{i}_{i}$. Now any socket (or, equivalently, a subset of L with even number of elements) whose characteristic binary function is :L ! f0; lg, de nes an element $_{v^{2}L}_{v^{2}}$ in (W_L). Similarly, to any network on T we associate a vector $_{e^{2}E}$! (e) in $\mathfrak{M} \stackrel{E}{=}$, where $!_i$ is such b($!_i$) = $(1)^{i}!_i$ and (e) = 1;0 depending on whether e is in or not. Now associating to a network its sockets de nes an isomorphism ($\mathfrak{M} \stackrel{E}{=}$)^{G_N} ! (W_L) which one can compare to what we discuss in 1.12.

We have a convenient description of the action of one-parameter groups associated to leaves of T in terms of socket coordinates of P (W_L). Namely, given a leaf 'the 1-parameter group 'acts on the coordinate with the weight (').

C on struction 3.5. Using networks and sockets, and the toric form alism, one can explain the inclusion X (T) $P(W_L)$ as follows. Let $M = \int_{0}^{\infty} Z$ be a lattice, a free abelian group generated by non-empty sockets of a tree T. The empty socket = 0 we interpret as the zero of the lattice. Then $P(W_L)$ is a toric variety X (e^0) associated to a unit simplex e^0 in M spanned on the vectors of the distinguished basis.

Now the bijective map sockets \$ networks gives rise to a homomorphism of lattices M ! M, where, recall, the latter lattice is spanned in M by the points associated to networks. This gives a surjective map from the symmetric graded algebra spanned by all the sockets, which is just algebra of polynomials C [], to the algebra A (), hence we get the inclusion X (T) P ($M_{\rm L}$), c.f. 2.3 and 2.23.

De nition 3.6. Let be a norm al lattice polytope in a lattice M. Let us choose two collection of points u_1 ;:::; u_r and w_1 ;:::; w_s in M and positive integers a_1 ;:::; a_r and b_1 ;:::; b_s . This data de nes a relation of degree d for if $a_1 + r + a = b_1 + s + b_1$ and

$$a_1u_1 + rut_r = b_1w_1 + swt_s b_1w_1$$

The relation is called primitive if fu_1 ;:::; $u_rg \setminus fw_1$;:::; $w_sg =$;.

Let us recall that given the projective variety $X = P^r$ with graded coordinate ring $S(X) = \int_{m=0}^{m} S^m(X)$ its ideal I(X) is the kernel of evaluation map $Symm(S^1(X)) \leq S(X)$. The following result is known as binomial generation of a toric ideal, see [Eisenbud, Sturm fels], [Sturm fels].

Lem m a 3.7. Suppose that we are in the situation of 2.23. Then the ideal I (X ()) is generated by polynom ials

 $\begin{pmatrix} u_1 \end{pmatrix}^{a_1} \quad u_r \end{pmatrix}^{a_r} \begin{pmatrix} w_1 \end{pmatrix}^{b_1} \quad w_s \end{pmatrix}^{b_s}$

where u_1 ;:::; u_r and w_1 ;:::; w_s , together with a_1 ;:::; a_r and b_1 ;:::; b_s de ne a prim - itive relation for .

E xam ple 3.8. The following are primitive relations and respective equations for the polytope coming from a 3-valent tree with four leaves, c.f. example 32. First, we describe them in terms of networks; they are as follows:

$$\frac{1}{2} \underbrace{\overset{3}{\overset{}}_{4}}_{2} + \underbrace{\overset{1}{\overset{}}_{2}}_{2} \underbrace{\overset{3}{\overset{}}_{4}}_{4} = \underbrace{\overset{1}{\overset{}}_{2}}_{2} \underbrace{\overset{3}{\overset{}}_{4}}_{4} + \underbrace{\overset{1}{\overset{}}_{2}}_{2} \underbrace{\overset{3}{\overset{}}_{4}}_{4}$$

and

$$1 - 3 + 1 - 3 = 1 - 3 + 1 - 3$$

 $On P(W_L)$ we introduce coordinates $x_{(1)}$ (indexed by characteristic sequences for the sockets in L. Then the respective equations dening X (T) are as follows:

x ₀₀₀₀	$1X_{11}$	=	x_{1100}	0011
x_{1001}	0 X 10	=	x_{1010}	0 X 01

F inally, let us note that renum bering the leaves or, equivalently, changing the shape of a 3-valent tree connecting the four num bered leaf vertices, produces the following respective equations

We note that all the above equations involve only four quadratic monom ials: $x_{0000}x_{1111}$, $x_{1100}x_{0011}$, $x_{1010}x_{0101}$, $x_{1001}x_{0110}$. Moreover, given any leaf ', the 1-parameter group `acts with weight 1 on each of them.

3.2 Dualpolytopes, fans, resolutions and Fano varieties.

In the situation of 2.22 the description of the fan of the variety X () in \mathbb{N}^{1} is given in term s of its support functions [O da, Thm . 2.22] or dual polytopes [Fulton].

E xam ple 3.9. By boking at the exam ple 2.16 and the inequalities which appear there we see that the fan of X (T³) in \mathbb{N}^{0} N has rays generated by the following elements: v=2 = ($e_{1} + e_{2}$)=2, v=2 $e_{2} = (e_{1} + e_{2} - e_{3})=2$, v=2 $e_{1} = (e_{0} + e_{2} - e_{3})=2$, v=2 $e_{2} = (e_{0} + e_{1} - e_{2})=2$.

The formula from 2.20 can be used to get the description of the polytope dual to (T), hence to describing the fan of X (T) for 3-valent trees.

Lem m a 3.10. Let T be a 3-valent binary symmetric tree with n inner nodes. Then the polytope (T) is de ned in M_R by 4n inequalities, which are as follows: for any inner node v 2 N, such that $v = e_{0:1} + e_{v:1} + e_{v:2}$ we take

$$(v=2)()$$
 1; $(v=2_{v,0})e()$ 0; $(v=2_{v,1}e)()$ 0; $(v=2_{v,2}e)()$ 0:

Proof. Let $(T_1; `_1)$ and $(T_2; `_2)$ be pointed trees. If $_i = (T_i) (M_i)_R$ is defined by inequalities with respect to some form $s w_j^i$ in $(N_i)_R$ then $_1 _2$ is defined by form $s (w_j^1; 0)$ and $(0; w_j^2)$ in $(N_1)_R (N_2)_R$. Then the classes of these form s in $N = (N_1 N_2)_R = R(`_1 `_2)$ define the berproduct of $_i$'s.

Denition 3.11. For a binary symmetric 3-valent tree T we dene a polytope - (T) in N_R which is the convex hull of $v=2 = (e_{0} + e_{v:1} + e_{v:2})=2, v=2$ $e_{0} = (e_{v:1} + e_{v:2} - e_{0})=2, v=2$ $e_{1} = (e_{v:0} + e_{v:2} - e_{0})=2, v=2$ $e_{2} = (e_{v:0} + e_{v:1} - e_{0})=2, v=2$ for v 2 N and $e_{v:0}, e_{v:1}, e_{v:2}$ edges containing v.

Let us note that the listed above points are in fact vertices of -(T). Indeed, take v 2 N and $e_{v:0}$, $e_{v:1}$, $e_{v:2}$ the edges containing v. Then by boking at the points which span -(T) we see that $(e_{v:0} + e_{v:1} + e_{v:2})(-(T))$ 3=2 with the equality only for the point $(e_{v:0} + e_{v:1} + e_{v:2})=2$ which therefore is a vertex. Similarly, $(e_{v:0} + e_{v:1} - e_{v:1}) = 2$ with the equality only for $(e_{v:0} + e_{v:1} - e_{v:2})=2$. Lem m a 3.12. Let b = P_{e2E} e then 4 (T) 2 and -(T) are dual, or polar, one to another in the sense that

> $-(T) = fw 2 N_R : w (4 (T) 2b)$ 1g 4 (T) 2b = fu 2 M_R : u (- (T)) 1g

Proof. The rst equality is a restatement of 3.10, the second equality follows because the polar polytope of the polar is the original polytope, Fulton, Sect 1.5].

N otation 3.13. For a vertex of (T) we de ne its dual face $u^{?} = -(T) \setminus fw$: w (4u 2b) = 1g. By \hat{e} we will understand the polytope which is the convex hull of u[?] and 0 2 N_R while by **b**[?] we will understand the cone spanned in N_R by u[?].

Let u be a vertex of (T) which we can represent as a network of paths, (u). Then v(u) is either 0 or 2, depending on whether (u) contains v and, similarly e (u) is, respectively 0 or 1. Thus (v=2)(4u 2b) = 1 if v is in (u) and (v=2)(4u 2b) = 3 otherwise. On the other hand (v=2 $_{v_{0}}$)(4u 2b) = 1 if either v is not in (u) or if both v and $e_{v_{10}}$ are in (u). Finally, (v=2 $e_{v_{10}}$)(4u 2b) = 3 if v is in (u) but $e_{v_{10}}$ is not.

Therefore, for any vertex u of (T) and any node v 2 N exactly three of the following four points v=2 = $(e_{0} + e_{v:1} + e_{v:2})=2$, v=2 $e_{0} = (e_{v:1} + e_{v:2} + e_{0})=2$, v=2 $e_{1} = (e_{v:0} + e_{v:2} + e_{v:2})=2$, v=2 $e_{1} = (e_{v:0} + e_{v:2} + e_{v:2})=2$ are in u? which therefore has 3n vertices.

E xam ple 3.14. We will visualize the points of \mathbb{N}^{0} on the graph of the tree in the following way. Given a 3-valent node v with edges $e_{v:0}$, $e_{v:1}$, $e_{v:2}$, which for simplicity we denote just by numbers on the graph, the point v=2 will be denoted by the dot at the vertex, while the point v=2 $e_{v:0}$ by the secant opposing the edge $e_{v:0}$, that is



Using this notation we can put on the same picture both, the system of paths associated to a vertex u of (T) as well as the respective points in u[?]. We put only four out of eight systems of paths from 3.2 since the other ones are obtained by renum bering of leaves.



In each of these cases the polytope $\mathbf{e}^{?}$ can be divided into two simplexes, each of them having edges which make a basis of the lattice \mathbb{N}° . For example:



The rst equality means that $\mathbf{e}^{?}$ in this case is a union of a simplex with edges $(\mathbf{e}_{1} + \mathbf{e}_{2} - \mathbf{e}_{3})=2$, $(\mathbf{e}_{0} + \mathbf{e}_{2} - \mathbf{e}_{1})=2$, $(\mathbf{e}_{0} + \mathbf{e}_{1} - \mathbf{e}_{2})=2$, $(\mathbf{e}_{3} + \mathbf{e}_{4} - \mathbf{e}_{3})=2$, $(\mathbf{e}_{0} + \mathbf{e}_{3} - \mathbf{e}_{4})=2$, and another one with edges $(\mathbf{e}_{1} + \mathbf{e}_{2} - \mathbf{e}_{3})=2$, $(\mathbf{e}_{0} + \mathbf{e}_{2} - \mathbf{e}_{3})=2$, $(\mathbf{e}_{0} + \mathbf{e}_{1} - \mathbf{e}_{2})=2$, $(\mathbf{e}_{3} + \mathbf{e}_{4} - \mathbf{e}_{3})=2$, $(\mathbf{e}_{0} + \mathbf{e}_{4} - \mathbf{e}_{3})=2$. The common part of these two simplexes is a simplex with edges $(\mathbf{e}_{1} + \mathbf{e}_{2} - \mathbf{e}_{3})=2$, $(\mathbf{e}_{0} + \mathbf{e}_{2} - \mathbf{e}_{3})=2$, $(\mathbf{e}_{0} + \mathbf{e}_{1} - \mathbf{e}_{2})=2$, $(\mathbf{e}_{3} + \mathbf{e}_{4} - \mathbf{e}_{3})=2$, which contains $\mathbf{e}_{0}=2 = ((\mathbf{e}_{0} + \mathbf{e}_{3} - \mathbf{e}_{3})=2 + (\mathbf{e}_{0} + \mathbf{e}_{4} - \mathbf{e}_{3})=2)=2$.

This example is even more transparent when we write NP as a sum of a rank 2 lattice spanned by $(e_1 + e_2 - e_3)=2$ and $(e_3 + e_4 - e_3)=2$, and of rank 3 lattice spanned by $(e_0 + e_2 - e_3)=2$, $(e_0 + e_1 - e_3)=2$, $(e_0 + e_3 - e_4)=2$ which contains also $(e_0 + e_4 - e_3)=2$. Then our division of the cone b[?] comes by multiplying by the cone R $_0(e_1 + e_2 - e_3)+R _0(e_3 + e_4 - e_3)$ the standard division of the 3-dimensional cone generated by $(e_0 + e_2 - e_3)=2$, $(e_0 + e_1 - e_2)=2$, $(e_0 + e_3 - e_4)=2$ and $(e_0 + e_4 - e_3)=2$, see Fulton, p. 49], which in geometric terms is a small resolution of a 3-dimensional quadric cone singularity giving rise to so-called A tiyah op.

The same argument works whenever (u) does not contain e_0 . Then u^2 contains $(e_0 + e_2 - e_1)=2$, $(e_0 + e_1 - e_2)=2$, $(e_0 + e_4 - e_3)=2$, $(e_0 + e_3 - e_4)=2$ and we can make a similar division of e^2 using the equality

$$(e_0 + e_2 - e_1)=2 + (e_0 + e_1 - e_2)=2 = (e_0 + e_4 - e_3)=2 + (e_0 + e_3 - e_4)=2$$

If (u) contains e_0 then we use the identity

 $(e_{0} + e_{1} + e_{2})=2 + (e_{1} + e_{2} - e_{0})=2 = (e_{0} + e_{3} + e_{4})=2 + (e_{3} + e_{4} - e_{0})=2$

which presents $e_{r}=22$ u[?] as an average of two di erent pairs of vertices to make a similar decomposition



Now we shall show that the above discussion can be generalized to the case of trees with more inner nodes.

Lem m a 3.15. Suppose that T is a binary symmetric 3-valent tree with n inner nodes. For any u, a vertex of (T) there exists a division of u[?] (or, equivalently of $\mathbf{e}^{?}$) into a union of 2^{n-1} (normalized) volume 1 simplexes. Equivalently, the cone $\mathbf{e}^{?}$ can be divided into a union of simplicial cones which are regular (i.e. their generators form bases of \mathbb{N}^{2}).

Proof. The construction of the division will proceed along an ascending sequence of subtrees of T, starting from an inner node of T. That is we have an ascending sequence of 3-valent trees

$$T_1 \quad T_2 \quad n \quad 1 \quad T \quad T_n = T$$

where T_i has i inner nodes and T_{i+1} is obtained from T_i as a graft with a star 3-valent tree. Forgetting of edges which are not in T_i gives a sequence of surjective m aps $M(T) ! M(T) ! M(T) ! M(T) = M(T) M(T_i)$ $M(T_i) = M(T_i) M(T_i) = M(T_i) =$

Now we will de ne the division of u_i^2 inductively. The polytope u_1^2 is just a simplex so let us assume that $u_i^2 = \begin{bmatrix} j \\ i \end{bmatrix}$ where $j = 1; :::; 2^{i-1}$ and the normalized volume of $\begin{bmatrix} j \\ i \end{bmatrix}$ with respect to the lattice $\mathbb{N}(T_i)$ is 1. Let v^i be an inner node of T_{i+1} which was a leaf of T_i , let e_0^i be a petiole of T_i which become an inner edge of T_{i+1} and let e_1^i and e_2^i are the two new petioles of T_{i+1} which contain v^i .

Now we make argument as in 3.14. If e_0^i is in u then $(e_0^i) = 22 u_i^2$ and we may assume that e_1^i is in u and e_2^i is not. Now from any simplex i_i^j from the original division of u_i^2 we produce two simplexes by adding a new vertex at $((e_0^i) + (e_2^i))$

 $(e_1^i) = 2$ and another one at either $((e_0^i) + (e_1^i) + (e_2^i)) = 2$ or at $((e_1^i) + (e_2^i)) = 2$. Because

$$((e_1^{i}) + (e_1^{i}) + (e_2^{i})) = 2 + ((e_1^{i}) + (e_2^{i})) = 2 = (e_1^{i})$$

and $(e_{0}^{i}) = 22 u_{1}^{?}$ this denes a good division of $u_{1+1}^{?}$.

If e_0^i is not in u then $(e_0^i) = 22 u_i^2$ and we make a similar construction but now we have to consider two cases: either none of e_1^i , e_2^i is in u or both are in u. At either case the discussion is similar to that we encountered in 3.14.

In terms of toric geometry the division process in plies the following.

C orollary 3.16. The a netoric variety associated to the cone \mathbf{b}^2 has G orenstein term in al singularities which adm it a sm all resolution.

P roof. The toric singularities are C ohen-M acaulay and since all the generators of the rays of \mathbf{b}^2 lie on the hyperplane (4u 2)() = 1 the singularities in question are G orenstein. The division into regular simplicial cones involves adding no extra ray so the respective resolution is small which also in plies that the original singularity is term inal.

W e note that the construction of the division certainly depends on the choice of the root of the tree and changing the root gives a op.

Let be a fan in $\mathbb{N}_{\mathbb{R}}$ consisting of cones \mathbf{b}^2 , where u is a vertex of (T), and their faces. In other words, contains cones spanned by the proper faces of -(T)(including the empty face, whose cone is the zero cone). Let us recall that equivariant line bundles on toric varieties are in a standard way described by piecew ise linear functions on its fan, see [D da, Sect. 2.1]. Setting $\mathbf{p}^2 = \mathbf{u}$ we de ne a continuous piecew ise linear function on the fan in N $_{\mathbb{R}}$ such that for every v 2 N and \mathbf{e}_{v} 2 E containing v we have (v=2) = 1 and (v=2 \mathbf{e}) = 0. The sections of the bundle related to , see [O da, Prop. 2.1], are in $\mathbb{M} \setminus (T)$. Therefore the toric variety X () given by the fan can be identied with the original variety X ((T)) and the line bundle associated to is O_X (1). On the other hand the function 4 2 assum es value 1 on the primitive vectors in rays of which allow s us to identify the canonical divisor of X (), see [O da, Sect 2.1]. The result is the follow ing.

Theorem 3.17. Let T be a 3-valent binary symmetric tree. Then the variety X (T) G orenstein and Fano with term inal singularities. Moreover it is of index 4, that is the canonical divisor $K_{X(T)}$ is linearly equivalent to $O_{X(T)}$ (4).

W e note the follow ing consequence of K odaira-K aw am ata-V iehw eg vanishing, see e.g. [K ollar, M ori, Sect 2.5]

C orollary 3.18. In the above situation $H^{i}(X (T); O (d)) = 0$ for i > 0 and d 3. In particular for d 0 we have $\dim_{C} H^{0}(X (T); O (d)) = h_{X (T)}(d)$ where the latter is Poincare-Hilbert polynom ial of (X (T); O (1)).

3.3 M utation of a tree, deform ation of a model.

In example 3.8 we noted that a four-leaf 3-valent tree can be labeled in three nonequivalent ways. We can revert it to say that given four numbered leaves we have three 3-valent labeled trees connecting these leaves. By grouping in pairs the leaves whose petioles are attached to common inner nodes we can list these as follows: (1;2) (3;4), (1;3) (2;4), and (1;4) (2;3).

Now, given four pointed trees T_i , where i = 1; :::; 4 we can produce a tree T by grafting the tree T_i along the i-th leaf of a labeled 3-valent 4-leaf tree T_0 . Here are possible con gurations, e_0 denotes the inner edge of the tree T_0



D e nition 3.19. In the above situation we say that there exists an elementary mutation along e_0 from one of the above trees to the other two. (W e note that a mutation m ay actually yield an equivalent tree.) W e say that two trees are mutation equivalent if there exists a sequence of elementary mutations from one to the other.

Lem m a 3.20. Any two 3-valent trees with the same number of leaves are mutation equivalent.

P roof. We prove, by induction, that any 3-valent tree is mutation equivalent to a caterpillar. To get the induction step it is enough to note that the graft of a caterpillar tree pointed at one of its legs with a star 3 valent tree contains a distinguished inner edge the mutation of which gives a caterpillar.

Now, let us recall the basics regarding deforming subvarieties in the projective space. Let B be an irreducible variety (possibly non-complete). Consider the product P^m B with the respective projections p_{B} and p_{B} . Suppose that X P^m B is a subscheme such that the induced projection $p_{B,K}$: X ! B is proper and at. Suppose that for two points a; b 2 B the respective scheme - theoretic bers X_a = X_a and X_b = X_b are reduced and irreducible. Then we say that the subvariety X_a in P^m

can be deformed to X $_{\rm b}$ over the base B. This gives rise to a notion of deformation equivalent subvarieties of P^m.

D e nition 3.21. Given two subvarieties X_1 ; X_2 in P^m we say that they are deformation equivalent if their classes are in the sam e connected component of the H ibert scheme of P^m .

Complete intersections of the same type are deformation equivalent. Let us consider an fundamental example, understanding of which is essential for the proof of the main result of this section.

E xam ple 3.22. Let us consider P^7 with hom ogeneous coordinates indexed by sockets of a 4-leaf tree T_0 , as in example 3.8. In P^7 we consider a family of intersections of 2 quadrics parametrized by an open subset B of P^2 with coordinates $t_{(12)(34)}$; $t_{(13)(24)}$; $t_{(14)(23)}$]. We set $B = P^2 n f[1;";"^2]$: "³ = 1g and over B we consider X⁰ given in B P^7 by equations

t _{(12) (34)}	₁¥00×0011	+ $t_{(13)(24)}$ + $t_{(14)(23)}$	₁ ¥310 ×0101 1¥301 ×0110	=	$t_{(12)(34)} + t_{(13)(24)} + t_{(14)(23)}$	$x_{0000}x_{1111}$
+ +	t _{(13) (24)} t _{(14) (23)} t _{(12) (34)}	t _{14) (23)} t _{12) (34)} t _{13) (24)}	₩00×0011 ₩10×0101 ₩01×0110	=	0	

Three special bers of the projection X⁰! B, namely X $^{0}_{[1;0,0]}$, X $^{0}_{[0;1;0]}$ and X $^{0}_{[0;0;1]}$, are varieties associated to three 4-leaf trees labeled by (12) (34), (13) (24) and (14) (23), respectively. On the other hand X⁰ is a complete intersection of two quadrics and the m ap X⁰! B is equidimensional. The latter statement follows because over B the matrix

is of rank 2 hence any berover B is a complete intersection of two non-proportional quadrics. Hence X^0 ! B is at because of [E isenbud, Thm . 18.16].

By T_0 T_N denote the 4-dimensional subtorus associated to the lattice spanned by leaves, that is a subtorus of T_N with coordinates v_i , where v_i , $i = 1; \dots; 4$ are leaves of T_0 . Torus T_0 acts on P^7 B via the rst coordinate, that is, for a leaf v_i of T_0 and a socket we have v_i (t) (x; $t_{(i)(i)}$) = $t^{(v_i)}x$; $t_{(i)(i)}$ Then by looking at the equations de ning X⁰ wee see that the inclusion X⁰, P⁷ B is equivariant with respect to this action. We also note that a rationalm ap P^7 ! P^3 , regular outside 16 linear P^3 's, which is given by four quadrics:

[x]! [$x_{0000}x_{1111}$; $x_{0011}x_{1100}$; $x_{0101}x_{1010}$; $x_{0110}x_{1001}$]

de nes a good quotient with respect to the action of T_0 on P^7 , c.f. B ialynicki B irula, 7.1.1]. If we take a subvariety Z^0 in the product P^3 B de ned by the equations

In what follows we construct an ambient variety which contains as locally com - p lete intersections a at family of varieties containing a geometric model of tree as well as models of the tree's elementary mutations.

C on struction 3.23. Let T be a tree with an inner edge e_0 which contains two 3-valent inner vertices. We can write T as a graft of ve trees: a labeled tree T_0 with four leaves v_i , i = 1; :::4, containing e_0 as an inner edge and four pointed trees $(T_i; `_i)$, with i = 1; :::4 which are attached to T_0 along the respectively labeled leaves. The edges in T which have common nodes with e_0 we denote, respectively, by e_i , each e_i comes from a petiole of `_i (or v_i). Recall, see 2.20, that M (T) and

(T) can be expressed as berproduct of M(T_i) and (T_i), respectively. That is, M(T) = $\begin{pmatrix} Q & \\ i=0 \end{pmatrix}$ M(T_i) $\setminus \begin{pmatrix} T & \\ i=1 \end{pmatrix}$ ker('_i \vee _i); (T) = $\begin{pmatrix} Q & \\ i=0 \end{pmatrix}$ (T_i) $\setminus \begin{pmatrix} T & \\ i=1 \end{pmatrix}$ ker('_i \vee _i)

Now, as in 3.5, we consider the lattice M_0 spanned on the non-trivial sockets of the tree T_0 together with the unit simplex e_0 M_0 R and the maps M_0 ! M $_0$ and e_0 ! $_0$ which give the inclusion X (T_0) P^7 as a complete intersection of two quadrics. Form s v_i , i = 1; :::4 pullback to M_0 and we denote them by v_i , respectively. Now we denote

$$\overline{\mathbf{M}} = \mathbf{M}_{0} \qquad \begin{array}{c} \mathbf{Q}_{4} \\ i=1 \\ \mathbf{M}_{i=1} \\$$

As in 2.3 we de ne the toric variety $Y = X(\overline{T})$. We note that, by A.4 the polytope is normal in the lattice $M_0 = \begin{pmatrix} 4 \\ i=1 \end{pmatrix} (T_i) \setminus \begin{pmatrix} T_4 \\ i=1 \end{pmatrix} (r_i \cdot \mathbf{y})$, which is spanned by its vertices. Also, by the construction we have the embeddings X(T), Y, $P(W_1)$. Lem m a 3.24. The inclusions

$$\overline{M}$$
, M_0 $\overset{Q_4}{\underset{i=1}{}}M$ (T_i) and $\overset{Q_4}{\underset{i=1}{}}$ e $\overset{Q_4}{\underset{i=1}{}}$ (T_i)

induce a rationalm ap

$$P^{7} \qquad \begin{array}{c} Y^{4} \\ X (T_{i}) & ! \\ Y \\ i=1 \end{array}$$

which is a good quotient map (of the set over which it is de ned) with respect to the action of the 4-dimensional torus T_0 generated by 1-parameter groups $v_i \rightarrow v_i$, where i = 1; :::4. The subvariety

$$X^{D} = X^{0}$$
 X^{4} $(T_{i}) , B$ P^{7} X^{4} (T_{i})

is T_0 equivariant and its quotient X is locally complete intersection in B Y.

Proof. The rst (quotient) part is the same as what we claim in 2.26, this time how – ever we repeat the argument for all four ber products in question. The invariance of the variety $\overset{1}{X}$ follows by the invariance of X^0 , $\overset{1}{B}$ $P^7_{Q^4}$ which we discussed in 3.22. Finally, since $\overset{1}{X}$ is a complete intersection in B $P^7 \qquad \overset{4}{\overset{1}{i=1}} X$ (T_i) its image X is a locally complete intersection in the quotient which is B Y, this follows from the denition of good quotient which locally is an a nequotient, [Bialynicki:Birula, Ch. 5], hence functions dening $\overset{1}{X}$ locally descend to functions dening X.

Lem m a 3.25.0 ver an open set $B^0 P^2$ containing points [1;0;0], [0;1;0], [0;0;1] the projection morphism X ! B^0 is at. The bers over points [1;0;0], [0;1;0], [0;0;1] are reduced and isomorphic to, respectively, the geometric model of T and of its elementary mutations along the edge e_0 .

Proof. First we note that the bers in question, $X_{[;;i]}$ of X ! B are geometric models as we claim. Indeed this follows from the universal properties of good quotients, c.f. $B_{ialynicki:B}$ inula], as they are quotients of the respective products $X_{[;;i]}^{0}$, $A_{i=1}^{4} X$ (T_{i}), which are located, as three invariant subvarieties, in $X^{b} = X^{0}$, $A_{i=1}^{4} X$ (T_{i}). This, in particular, implies that the respective bers of X ! B are of the expected dimension, hence they are contained in a set $B^{0} = P^{2}$ over which the map in question is equidimensional. Since Y is toric it is C ohen-M acaulay too [E isenbud, Prop. 18.13]. Finally, the map X ! B^{0} is equidimensional hence it is at, because B^{0} is smooth, see [E isenbud, Thm . 18.16]

Theorem 3.26. Geometric models of 3-valent trees with the same number of baves are deformation equivalent in $P(W_{T})$.

Proof. This is a combination of 3.20 and of 3.25.

3.4 Hilbert-Ehrhard polynom ial.

Denition 3.27. Given two pointed trees $(T_1; `_1)$ and $(T_2; `_2)$ we dene a pointed graft which is a pointed tree $(T; o) = (T_1; `_1)? (T_2; `_2)$ where $T = T_1 \ _{+o_1} T^{-3} \ _{o_2-`_2} T_2$, and o, o_1 and o_2 are the bayes of T^{-3} .

Exam ple 3.28. Pointed graft of two 3-valent stars

$$\rightarrow$$
 ? \prec = \rightarrow

By arguments used in the proof of 2.20 we also get.

P roposition 3.29. Let $(T_1; '_1)$ and $(T_2; '_2)$ be two pointed trees. Then

$$(T_1 \cdot T_2, T_2) = (T_1) \cdot T_0 (T^3) \cdot T_2$$

Let us consider a 3-dimensional lattice $M = Ze_0$ Ze Ze with a xed tetrahedron ⁰ with vertices 0; $e_0 + e_1$; $e_0 + e_2$; $e_1 + e_2$. By M we denote the index 2 sublattice spanned on the vertices of ⁰.

De nition 3.30. Let n be a positive integer and let $f_1^n = f_1$; $f_2^n = f_2$ be two functions de ned on the set f0; ::: ng with values in Z or, more generally, in an arbitrary ring or algebra (we use the superscript ⁿ to indicate the domain of f's). For any k 2 f0; ::: ; ng we de ne

$$(f_{1}?f_{2})(k) = \begin{pmatrix} X \\ (f_{1}(e_{1}(u)) & _{2}f(e_{2}(u))) \\ e_{0}(u) = k \end{pmatrix}$$

We note that ? is commutative, that is $f_1^n ? f_2^n = f_2^n ? f_1^n$, but possibly not associative. By $(f^n)^{?m}$ we denote the ? product of m copies of a chain of f^n , that is $f^n ? (f^n ? (::: (f^n ? f^n) :::).$ By 1^n we denote the constant function f0;:::;ng ! flg Z.

A function $f^n : f0; :::; ng ! Z will be called symmetric if <math>f^n (k) = f^n (n k)$.

Lem m a 3.31. If $f_1 = f_1^n$; $f_2 = f_2^n : f_0; \dots; ng !$ Z are symmetric functions then $f_1 ? f_2$ is a symmetric function as well and moreover for k n=2 we have

$$(f_{1}?f_{2})(k) = 2 \qquad \begin{array}{c} X \ ^{1}X^{1} \\ f_{1}(i)f_{2}(k+i) \\ i = 0 \ j = 0 \end{array} \qquad \begin{array}{c} X \ ^{k}X^{k} \\ f_{1}(i)f_{2}(k+i) \\ f_{2}(k+i) \\ i = k \ j = 0 \end{array}$$

In particular, for k n=2

$$(f_1 \quad 1) (k) = 2 \quad (i+1) f_1 (i) + X^{k} (k+1) f_1 (i)$$

Proof. Let us look at the sections of the tetrahedron n ⁰ with hyperplanes (e_0) ¹ (k). We picture the situation for n = 6 and k = 0; :::6, the dotted square is the section of the cube with the lower left corner satisfying relation $e_1 = e_2 = 0$, the section of the tetrahedron denoted with solid line and points inside the (closed) tetrahedron denoted by .



The denition of f_1^n ? f_2^n is sum of the product of f_i^n 's over the lattice points of such a section. The sections over k and n k are obtained by a relection with respect to either $e_1 = 1=2$ or $e_2 = 1=2$. Thus if one of f_i^n 's is symmetric then the f_1^n ? f_2^n is symmetric as well.

On the other hand for 0 k n k the tetrahedron's section is a rectangle with vertices (k;0); (0;k); (n k;n); (n;n k) which we divide into two triangles and a parallelogram, the division is indicated by dotted vertical line segments for boxes labeled by k = 1;2 in the above diagram. Because functions f_i^n are symmetric the values of the product f_1^n $_2^n$ fare the same for the points which are central symmetric with respect to the center of the square. Thus in the form ula of the lemma we take the value $f_1(a)f_2(b)$ for all integral pairs (a;b) in the left hand side triangle and multiply it by 2 (that is the rst summand in the form ula) and add the sum over the parallelogram.

Example 3.32. We note that $(1^n)^{2}(k) = (k+1)(n + 1)$ is the number of lattice points in the rectangle used in the argument in the above proof of 3.31. On the

other hand by using the form ula from 3.31 one gets

$$(1^n)^{23}$$
 (k) = $\frac{1}{6}$ (k + 1) (n k + 1) (r² + kn k² + 5n + 6)

Let us recall that given a lattice polytope M_R for any positive integer n we de ne Ehrhard function h as follows:

h (n) = j(n
$$\setminus M$$
) j

If satisfies the assumptions of 2.3 then $h = h_{X()}$ where the latter is the Poincare-Hilbert polynomial of (X ();0 (1)) which, by denition, is equal to $\dim_{C} H^{0}(X();0(m))$ form 0.

Denition 3.33. Let M_R be a lattice polytope which is not contained in any hyperplane and let v 2 N be a non-zero form on M. Suppose that v() [0;1]. We de ne its relative Ehrhard function f_{rv}^{*} :f0;:::ng! Z by setting

$$f_{v}^{n}(k) = jv^{1}(k) \setminus n \qquad \setminus M j$$

We note that, clearly, $\int_{k=0}^{P} f_{v}^{n}(k) = h$ (n) is the usual Ehrhard function. Thus, in case of 2.3 the above de nition can be restated in purely geometric fashion.

Lem m a 3.34. Suppose that satis es assumptions of 2.3 and v is as in 3.33. Let us consider a linearization of the action of the 1-parameter group $_{v}$ on H 0 (X ();0 (n)) which has non-negative weights and the eigenspace of the zero weight is nontrivial. Then f_{v}^{n} (k) is equal to the dimension of the eigenspace of the action of $_{v}$ of weight k.

Proof. This is a consequence of the standard properties of X (), 223.4. $\hfill \Box$

Lem m a 3.35. Let $(T_1; `_1)$ and $(T_2; `_2)$ be two pointed trees and let f_1^n and f_2^n be two relative Ehrhard functions associated to (T_1) and (T_2) , respectively. If $(T; o) = (T_1; `_1) ? (T_2; `_2)$ and f_o^n is the relative Ehrhard function associated to (T) then $f_o^n = f_1^n ? f_2^n$

Proof. The de nitions of? are made accordingly.

Example 3.36. By using 3.32 we nd out that

$$X^{n} (1^{n})^{2} (k) = \frac{(n+1)(n+2)(n+3)}{6}$$

which is the Poincare-H ilbert polynom ial of (P³;O (1)) while

$$X^{n} (1^{n})^{23} (k) = \frac{(n+1)(n+2)(n+3)(n^{2}+4n+5)}{30}$$

which is Poincare H ibert polynom ial of intersection of two quadrics in P^7 .

Theorem 3.37. Let us consider three pointed trees $(T_i; '_i)$, with i = 1;2;3 with relative Ehrhard functions $f_i^n = f_{i_i}^n$ associated to polytopes (T_i) , respectively. Then

$$(f_1^n ? f_2^n) ? f_3^n = f_1^n ? (f_2^n ? f_3^n)$$

Proof. Let 'denote the distinguished leaf of the result of the ? operation on the trees. Then the relative Ehrhard function $(f_1^n ? f_2^n) ? f_3^n$ and, respectively, $f_1^n ? (f_2^n ? f_3^n)$ is related to one of the following trees, each of them is obtained by an elementary mutation from the other:

Now we repeat the construction 3.23, with obvious modi cations. Namely, we de ne a polytope

$$= e_0 \qquad (T_i) \setminus ker('_i v_i)$$

$$= 1 \qquad i = 1$$

where e_0 is the unit simplex as in 3.5. We de ne a toric variety Y = X() with the embedding in $P(W_E)$ and the action of the group \cdot .

Next, as in 3.24 we de ne a subvariety X В Y such that the projection p_B : X ! B is at and its two bers are varieties associated to the above two pointed trees, see 3.25. Because of the atness the sheaf (p_B) $(p_Y (0 (n)))$ is locally 0, see #artshome, III.9.9, III.12.9] and 3.18. M oreover, by the free for each n construction, the action of the group \neg on Y leaves X В Y, as we noted in 3.22. Finally, the decomposition into eigenspaces of the action of $\sqrt{}$ on H 0 (Y; O(n)) restricts into a respective eigenspace decomposition of the action of , on bers of (p_B) $(p_v (0 (n)))$, which are equal to H⁰ $(X_b; 0 (n))$, for b 2 B. This implies that the dimension of the respective eigenspaces is locally constant, with respect to the parameter b 2 B hence the relative Ehrhard function of bers of p_B is constant which concludes the argum ent.

Let us underline the fact that although the invariance of the H ilbert polynom ial is a standard property of a at family the above result is about the invariance of the family with respect to an action of a 1-parameter group, the group \cdot in our case.

The above theorem 3.37 in plies that the operation ? on relative Ehrhard functions of polytopes of 3-valent trees is not only commutative (which is obvious from its de nition) but also associative. This im plies that the function does not depend on either the shape nor the location of the leaf. M ore precisely we have the follow ing form ula which allows to compute the H ilbert-Ehrhard polynom ial very e ciently.

C orollary 3.38. If (T; ') is a pointed 3-valent tree with r + 1 leaves then

$$f_{(T)}^{n} = (1^{n})^{2r}$$

A Appendix

A.1 Norm alpolytopes, unim odular covers

A lattice simplex 0 M_{R} with vertices v_{0} ;::: v_{r} is called unimodular if vectors v_{1} v_{3} ;:::; v_{r} v_{3} span M. We say at a lattice polytope M_{R} has a unimodular covering if $= {}^{0}$ where 0 are unimodular simplexes. This de nition is taken from [BGT] where we also have the following result.

Lem m a A .1. If a lattice polytope M_R has a unimodular covering then it is normal.

The following observation is probably known but we include its proof because of the proof of the subsequent lemma.

Lemma A 2. Let $_1$ (M₁)_R and $_2$ (M₂)_R be two unimodular simplexes. Then $_1$ $_2$ has a unimodular covering in M₁ M₂.

Proof. We can assume that $_1$ has vertices $0;e_1;:::;e_r$ and $_2$ has vertices $0;f_1;:::;f_s$. Suppose that x 2 (M₁)_R (M₂)_R is as follow s:

$$x = \begin{matrix} X^{r} & X^{s} \\ a_{i}e_{i} + \begin{matrix} X^{s} \\ b_{j}f_{j} \end{matrix}$$

where a_i ; b_j 0 and a_i b_j 1

The union of unimodular simplexes contained in 1 = 2 is a closed subset. Therefore if x is not contained in any modular subsimplex of 1 = 2 then any sm all perturbation of x has this property as well. Thus we are free to assume that all a_i 's and b_j 's are nonzero and any two non-empty subsets of a_i 's and b_j 's have di erent sum, in particular $a_1 + \ldots + a_p \in b_1 + \ldots + b_q$ for any reasonable (p;q). Let m be such $b_1 + \ldots + b_m$ $_1 < a_1 + \ldots + a_r < b_1 + \ldots + b_m$. We set $b_m^0 = (b_1 + \ldots + b_m)$ ($a_i + \ldots + a_r$).

In order to prove the km m a we will nd r + m 1 positive num bers q_{j} indexed by some pairs (i; j) 2 f1; ...; rg f1; ...; m g such that

$$x = \sum_{ij}^{X} c_{i;j} (e_i + f_j) + b_m^0 f_m + b_{m+1} f_m + \dots + b_s f_s$$

and the set of respective vectors $(e_i + f_j)$ together with f_m ;::: f_s can be modied via addition or subtraction of pairs among them to the standard basis e_1 ;::: e_r ; f_1 ;:::; f_s .

The coe cients c_{i;j} are de ned inductively according to the follow ing rules. The rst coe cient is c_{1;1} = m infa₁; b₁g. Suppose that the last de ned coe cient is c_{i0;j0}. If $(i_0; j_0) = (r; m)$ then we are done so assume that it is not the case. Then, because of our assumption that the sequences (a_i) and (b_j) have no equal partial sum s, either $a_1 + \ldots + a_{i_0} > b_1 + \ldots + b_{j_0}$, or $a_1 + \ldots + a_{i_0} < b_1 + \ldots + b_{j_0}$. In the form er case we set

$$C_{i_0;i_0+1} = m \inf b_{i_0+1}; (a_1 + :::+ a_{i_0}) (b_1 + :::+ b_{i_0})g$$

whereas in the latter case we de ne

$$c_{i_0+1;j_0} = m \inf_{a_{i_0+1}} (b_1 + \dots + b_{j_0}) (a_1 + \dots + a_{i_0})g$$

The veri cation that $P_{i=1}^{r} c_{i;j} = b_j$ for j = 1:::m 1 and $P_{j=1}^{m} c_{i;j} = a_i$ for i = 1;::r is easy and left for the reader. Sim ilarly, a sim ple backtracking allows to modify the set of the respective vectors $e_i + f_j$ with f_m ; ::: f_s to the standard basis for $M_1 = M_2$.

Lem m a A .3. Let $_{1}$ $(M_{1})_{R}$, $_{2}$ $(M_{2})_{R}$ be two unimodular simplexes. We consider two homomorphisms $i_{i}: M_{i} ! Z$ such that $(i_{i})_{R} (i_{i})$ [0;1]. The the ber product = $(1)_{i_{1}} i_{2} (i_{2})$ has a unimodular covering with respect to the ber product lattice $M = (M_{1})_{i_{1}} i_{2} (M_{2})$.

Proof. The argument is a variation of the one used in the previous km m a. We can assume that $_1$ has vertices $0;e_1^0;\ldots;e_{r_1}^0;e_1^1;\ldots;e_{r_1}^1$ and $_2$ has vertices

 $0; f_1^0; \dots; f_{s_0}^0; f_1^1; \dots; f_{s_1}^1 \text{ where } '_1(e_i^0) = '_2(f_j^0) = 0 \text{ and } '_1(e_i^1) = '_2(f_j^1) = 1 \text{ for suitable is and j's. Suppose that x 2 (M₁)_R (M₂)_R is as follows:$

$$\mathbf{x} = \begin{bmatrix} X^{r_0} \\ a_1^0 e_1^0 + \begin{bmatrix} X^{r_1} \\ a_1^1 e_1^1 + \begin{bmatrix} X^{s_0} \\ b_j^0 f_j^0 + \end{bmatrix} \begin{bmatrix} X^{s_1} \\ b_j^1 f_j^1 \end{bmatrix}$$

where $a_i^0; a_i^1; b_j^0; b_j^1 = 0$, $a_i^0 + P = a_i^1 = 1$, $b_j^0 + P = b_j^1 = b_j^1$. The latter conditions ensures that $a_i^1 = b_j^1$. The latter conditions ensures that $a_i^1 = b_j^1$ and moreover $a_i^1 = b_j^1$. The latter conditions ensures that $a_i^1 = b_j^1$ and $a_j^1 = b_j^1$. The latter conditions ensures that $a_j^1 = b_j^1 = b_j^1$. The latter conditions ensures that $a_j^1 = b_j^1 = b_j^1$. The latter conditions ensures that $a_j^1 = b_j^1 = b_j^1$. The latter conditions ensures that $a_j^1 = b_j^1 = b_j^1$. The latter conditions ensures that $a_j^1 = b_j^1 = b_j^1$. The latter conditions ensures that $a_j^1 = b_j^1 = b_j^1$. The latter conditions ensures that $a_j^1 = b_j^1 = b_j^1 = b_j^1$. The latter conditions ensures that $a_j^1 = b_j^1 = b_$

We write $x = x_0 + x_1$ where $x_0 = \begin{bmatrix} a_i^0 e_i^0 + b_j^0 f_j^0 & \text{and } x_1 = \begin{bmatrix} a_i^1 e_i^1 + b_j^1 f_j^1 \\ \text{and we repeat the proof of } A_p^2 & \text{for } x_{q_p} & \text{and } x_1 & \text{separately. The only difference is that, because of the equality <math>a_i^1 = b_j^1$, the construction will give $r_1 + s_1 = 1$ are class of the equality $a_i^1 = b_j^1$, the construction will enable to write $x_1 = \begin{bmatrix} c_{i,j}^1 (e_i^1 + f_j^1) \\ 0 & \text{not constitute} \end{bmatrix}$ abasis of vectors $e_i^1 + f_j^1$ which will enable to write a basis of the lattice spanned by $e_1^1; \ldots; e_{r_1}^1; f_1^1; \ldots; f_{s_1}^1$ but of this lattice intersected with ker($\begin{bmatrix} r_1 \\ 0 \end{bmatrix}$). That is, among the chosen $r_1 + s_1 = 1$ vectors $e_i^1 + f_j^1$ we have $e_1^1 + f_1^1$ and $e_{r_1}^1 + f_{s_1}^1$ and if $e_i^1 + f_j^1$ is among them then either $e_{i+1}^1 + f_j^1$ or $e_i^1 + f_{j+1}^1$ is among them as well (but not both). We are to prove that any $e_i^1 + f_j^1$ can be obtained as a sum of them . But this follows because

$$e_i^1 + f_j^1 + e_{i+1}^1 + f_{j+1}^1 = e_{i+1}^1 + f_j^1 + e_i^1 + f_{j+1}^1$$

so any one of the above above four vectors is a combination of the other three and this observation can be used repeteadly to complete our claim . $\hfill \Box$

Corollary A.4. Let $_{1}$ $(M_{1})_{R}$, $_{2}$ $(M_{2})_{R}$ be two polytopes which have covering by unim odular simplexes. We consider two hom on orphism s $i : M_{i} ! Z$ such that $(i)_{R}(i)$ [0;1]. Then the ber product = $(1)_{1} i : 2$ (2) has a unim odular covering with respect to the ber product lattice $M = (M_{1})_{1} i : 2$ (M₂)

Proof. The ber product of $_1$ and $_2$ is covered by ber products of simplexes from the unim odular cover of each of them. Thus the result follows by A.3.

Since the polytope of the star 3-valent tree is a unit tetrahedron we get the following.

P roposition A .5. If T is a binary symmetric 3-valent tree then its polytope in \mathbb{M} (T) has unimodular covering hence it is normal.

A.2 Two 3-valent trees with 6 leaves

O ne of the fundam ental questions regarding the phylogenetic trees is the follow ing. G iven two (3-valent binary symmetric) trees T_1 and T_2 suppose that $(T_1) = (T_2)$ as lattice polytopes, or the projective models X (T_1) and X (T_2) are projectively equivalent. Does it in ply that the trees are equivalent (as CW complexes) as well?

We tackled the problem by understanding the di erence of models of the two simplest non-equivalent trees. These are 6-leaf trees pictured below, respectively, a 3-caterpillar tree and a tree which we call a snow ake, [Sturm fels, Sullivant].



The snow ake tree is obtained from the 3-caterpillar tree by elementary mutation along its middle inner edge. Therefore their Hilbert-Ehrhard polynom ials are equal and computed with [maxima] to be as follows.

h (n) =
$$\frac{1}{22680}$$
 (n + 1) (n + 2) (n + 3)
(31 n⁶ + 372 n⁵ + 1942 n⁴ + 5616 n³ + 9511 n² + 8988 n + 3780)

On the other hand we can distinguish their polytopes in term s of som e com binatorial invariants.

Example A .6. Given a polytope we de ne its incidence matrix (a_{ij}) as follows: (a_{ij}) is a symmetric matrix with integral entries such that for i j the number a_{ij} is equal to the number of i-dimensional faces contained in j-dimensional faces of . In particular a_{ii} is the number of i-dimensional faces. The following is the incidence matrix of a polytope of the snow ake tree.

32	480	2400	6144	9312	8832	5280	1920	384
480	240	2400	9456	19920	24960	19200	8880	2256
2400	2400	760	5944	19008	32552	32408	18792	5872
6144	9456	5944	1316	8400	21744	29308	21720	8388
9312	19920	19008	8400	1392	7200	14640	14640	7200
8832	24960	32552	21744	7200	940	3820	5760	3820
5280	19200	32408	29308	14640	3820	406	1224	1224
1920	8880	18792	21720	14640	5760	1224	108	216
384	2256	5872	8388	7200	3820	1224	216	16

And this is the incidence matrix of the polytope of a 3-caterpillar tree.

32	480	2400	6144	9312	8832	5280	1920	384
480	240	2400	9456	19904	24896	19104	8816	2240
2400	2400	760	5944	18976	32408	32168	18616	5824
6144	9456	5944	1316	8384	21648	29112	21552	8336
9312	19904	18976	8384	1392	7184	14584	14576	7176
8832	24896	32408	21648	7184	940	3816	5752	3816
5280	19104	32168	29112	14584	3816	406	1224	1224
1920	8816	18616	21552	14576	5752	1224	108	216
384	2240	5824	8336	7176	3816	1224	216	16

Both m atrices were computed by [polymake]. We note that although both polytopes have the same number of faces of respective dimension their incidences are di erent (indicated in boldface).

A.3 Volume distribution

The leading coe cient in the Ehrhard polynom ial of a lattice polytope can be identified as the volume of (with respect to the lattice in question, whose unit cube is assumed to have volume 1). Similarly, we can de ne a relative volum e function which will measure the distribution of the volume of (T) with respect to a leaf ' of T. Because of 3.38 this function does not depend either on the shape of the tree nor on the choice of the leaf '. Moreover we will normalize it so that its integral over the unit segment is 1.

If r:[0;1]! R is the normalized volume distribution with respect to a leaf of a 3-valent tree with r leaves then because of 3.31 we have n(t) = n(1 - t) and for t2 (0;1=2) we get the following recursive formula

$$r^{r+1}(t) = d_{n+1}$$
 2 s $r(s)ds + t$ t $r(s)ds$

where d_{n+1} is a constant such that $\int_{0}^{n+1} r^{r+1}(s) ds = 1$. From this it follows that r is a polynom ial of degree 2r. However, the numerical experiments which we have made seem to indicate that for r > 3 the actual values of r do not depend too much on r, see Fig. 1. It seems that this function does not see the shape of the tree (which is because it comes from the relative H ilbert-Ehrhard polynom ial) but also alm ost disregards its size (or dimension of the model)



Figure 1: Polynom ials 2 and 100 at the same diagram, by [gnuplot].

References

- [A llm an, Rhodes '03] E.S.A llm an, J.A.Rhodes, Phylogenetic ideals and varieties for the general M arkov m odel, M ath. Biosciences 186 (2003), 133-144, arX ive m ath AG /0410604.
- [A llm an, Rhodes, '05] E.S.A llm an, J.A.Rhodes, The identi ability of tree topology for phylogenetic models, including covarion and mixture models, arX ive q-bio PE/0511009.
- [Altman] K. Altmann, Deformation theory, 1995, notes at http://pagemifu-berlin.de/ altmann/PAPER/dmvps
- Bialynicki-Birula] A.Bialynicki-Birula, Quotients by algebraic groups, Encyclopedia of Mathematical Sciences vol 131, Springer 2003.
- [BGT] W. Bruns, J. Gubeladze, N.V. Trung, Norm al polytopes, triangulations and Koszul algebras, J. reine u angew. M ath. 485 (1997), 123-160.

- [CGS] M. Casanellas, L. Garcia, S. Sullivant, Catalog of small trees, in A lgebraic Statistics for Computational Biology, ed. Pachter, Sturm fels, Cambridge 2005.
- [E isenbud] D.E isenbud, Commutative algebra with a view toward algebraic geometry, Springer 1994.
- [Eisenbud, Sturm fels] D. Eisenbud, B. Sturm fels, Binom ial ideals. Duke Math. J. 84 (1996), 1{45.
- [ERSS] N.Eriksson, K.Ranestad, B.Sturm fels, S.Sullivant, Phylogenetic Algebraic Geometry, arX ive m ath AG /0407033.
- [Fulton] W. Fulton, Introduction to toric varieties, Princeton University Press 1993.
- [Hartshome] R.Hartshome, Algberaic Geometry, GTM 52, Springer, 1997.
- [M um ford] D.M um ford, J.Fogarty, G eom etric Invariant Theory, 2nd ed. Springer 1982.
- [Kollar, Mori] J. Kollar, Sh. Mori, Birational geometry of algebraic varieties, Cambridge Tracts in Math 134, 1998.
- [Namikawa] Y. Namikawa, Smoothing Fano 3-folds. J. Algebraic Geom. 6 (1997), 307{324.
- [0 da] T.O da, Convex bodies and algebraic geometry, Springer 1987.
- [Pachter, Sturm fels] L. Pachter, B. Sturm fels et al, A lgebraic Statistics for C om putational Biology, C am bridge 2005.
- [Reid '87] M. Reid, Tendencious survey of 3-folds. A lgebraic geometry, Bowdoin, 1985 333{344, Proc. Sympos. Pure Math. 46, AMS 1987.
- [Reid '92] M.Reid, What is a ip?, UnivUtahColloquium talk notes, 1992.
- [Shafarevich] I. R. Shafarevich, On certain tendencies in the development of mathematics, Math. Intelligencer 3, (1980), 182–184.

[Stum fels] B. Stum fels, Equations de ning toric varieties. A lgebraic geom etry | Santa C nuz 1995, 437{449, P roc. Sym pos. Pure M ath., 62, AM S 1997.
[Stum fels, Sullivant] B. Stum fels, S. Sullivant, Toric ideals of phylogenetic invariants, arX ive q-bio PE/0402015.
[gnuplot] G nuplot 3.7, T. W illiam s, C. K elley et al, www.gnuplot.info
[maxima] M axim a 5.4, W. Schelter et al, http://m axim a sourceforge.net/
[polymake] Polym ake 2.0.1, E. G avrilow, M. Joswig et al, www.math.tu-

Authors' address: Instytut M atem atykiUW , Banacha 2,02–097 W arszawa, Poland wkrych@mimuw.edu.pl jarekw@mimuw.edu.pl

berlin de/polym ake