

ON THE AVERAGE NUMBER OF SHARP CROSSINGS OF CERTAIN GAUSSIAN RANDOM POLYNOMIALS

S. Shemehsavar, S. Rezakhah*

Abstract

Let $Q_n(x) = \sum_{i=0}^n A_i x^i$ be a random algebraic polynomial where the coefficients $A_0, A_1, \dots$ form a sequence of centered Gaussian random variables. Moreover, assume that the increments $\Delta_j = A_j - A_{j-1}$, $j = 0, 1, 2, \dots$ are independent, assuming $A_{-1} = 0$. The coefficients can be considered as n consecutive observations of a Brownian motion. We obtain the asymptotic behaviour of the expected number of u -sharp crossings of polynomial $Q_n(x)$. We refer to u -sharp crossings as those zero up-crossings with slope greater than u , or those down-crossings with slope smaller than $-u$. We consider the cases where u is unbounded and is increasing with n , where $u = o(n^{5/4})$, and $u = o(n^{3/2})$ separately.

Keywords and Phrases: random algebraic polynomial, number of real zeros, sharp crossings, expected density, Brownian motion.

AMS(2000) subject classifications. Primary 60H42, Secondary 60G99.

*Faculty of Mathematics and Computer Science, Amirkabir University of Technology, 424 Hafez Avenue, Tehran Iran. Email: Rezakhah@aut.ac.ir, Shemehsavar@aut.ac.ir.

1 Preliminaries

The theory of the expected number of real zeros of random algebraic polynomials was addressed in the fundamental work of M. Kac[6]. The works of Logan and Shepp [7, 8], Ibragimov and Maslova [5], Wilkins [14], and Farahmand [3] and Sambandham [12, 13] are other fundamental contributions to the subject. For various aspects on random polynomials see Bharucha-Reid and Sambandham [1], and Farahmand [4]. There has been recent interest in cases where the coefficients form certain random processes, Rezakhah and Soltani [10, 11], Rezakhah and Shemehsavar [9].

Let

$$Q_n(x) = \sum_{i=0}^n A_i x^i, \quad -\infty < x < \infty, \quad (1.1)$$

where $A_0, A_1, \dots$ are mean zero Gaussian random variables for which the increments $\Delta_i = A_i - A_{i-1}$, $i = 1, 2, \dots$ are independent, $A_{-1} = 0$. The sequence $A_0, A_1, \dots$ may be considered as successive Brownian points, i.e., $A_j = W(t_j)$, $j = 0, 1, \dots$, where $t_0 < t_1 < \dots$ and $\{W(t), t \geq 0\}$ is the standard Brownian motion. In this physical interpretation, $\text{Var}(\Delta_j)$ is the distance between successive times t_{j-1} , t_j . Thus for $j = 1, 2, \dots$, we have that $A_j = \Delta_0 + \Delta_1 + \dots + \Delta_j$, where $\Delta_i \sim N(0, \sigma_i^2)$ and are independent. Thus $Q_n(x) = \sum_{k=0}^n (\sum_{j=k}^n x^j) \Delta_k = \sum_{k=0}^n a_k(x) \Delta_k$, and $Q'_n(x) = \sum_{k=0}^n (\sum_{j=k}^n j x^{j-1}) \Delta_k = \sum_{k=0}^n b_k(x) \Delta_k$, where

$$a_k(x) = \sum_{j=k}^n x^j, \quad b_k(x) = \sum_{j=k}^n j x^{j-1}, \quad k = 0, \dots, n. \quad (1.2)$$

We say that $Q_n(x)$ has a zero up-crossing at t_0 if there exists $\varepsilon > 0$ such that $Q_n(x) \leq 0$ in $(t_0 - \varepsilon, t_0)$ and $Q_n(x) > 0$ in $(t_0, t_0 + \varepsilon)$. Similarly $Q_n(x)$ is said to have a zero down crossing at t_0 , if $Q_n(x) > 0$ in $(t_0 - \varepsilon, t_0)$ and $Q_n(x) \leq 0$ in $(t_0, t_0 + \varepsilon)$. We study the asymptotic behavior of the expected

number of, u -sharp zero crossings, those zero up-crossings with slope greater than $u > 0$, or those down-crossings with slope smaller than $-u$.

Cramer and Leadbetter [1967 p 287] have shown that the expected number of total zeros of any Gaussian non stationary process, say $Q_n(x)$, is calculated via the following formula

$$EN(a, b) = \int_a^b dt \int_{-\infty}^{\infty} |y| p_t(0, y) dy$$

where $p_t(z, y)$ denotes the joint density of $Q_n(x) = z$ and its derivative $Q'_n(x) = y$, and

$$p_t(0, y) = [2\pi\gamma\sigma(1 - \mu^2)^{1/2}]^{-1} \times \exp \left\{ - \frac{\gamma^2 m^2 + 2\mu\gamma\sigma m(y - m') + \sigma^2(y - m')^2}{2\gamma^2\sigma^2(1 - \mu^2)} \right\}$$

in which $m = E(Q_n(x))$, $m' = E(Q'_n(x))$, $\sigma^2 = \text{Var}(Q_n(x))$, $\gamma^2 = \text{Var}(Q'_n(x))$, and $\mu = \text{Cov}(Q_n(x), Q'_n(x))/(\gamma\sigma)$.

By a similar method as in Farahmand [4] we find that $ES_u(a, b)$, the expected number of u -sharp zero crossings of $Q_n(x)$ in any interval (a, b) , satisfies

$$ES_u(a, b) = \int_a^b dt \left\{ \int_{-\infty}^{-u} + \int_u^{\infty} \right\} |y| p_t(0, y) dy = \int_a^b f_n(x) dx \quad (1.3)$$

where

$$f_n(x) = \frac{1}{\pi} g_{1,n}(x) \exp(g_{2,n}(x)) \quad (1.4)$$

and

$$g_{1,n}(x) = EA^{-2}, \quad g_{2,n}(x) = -\frac{A^2 u^2}{2E^2}, \quad (1.5)$$

in which

$$A^2 = \text{Var}(Q_n(x)) = \sum_{k=0}^n a_k^2(x) \sigma_k^2, \quad B^2 = \text{Var}(Q'_n(x)) = \sum_{k=0}^n b_k^2(x) \sigma_k^2,$$

$$D = \text{Cov}(Q_n(x), Q'_n(x)) = \sum_{k=0}^n a_k(x) b_k(x) \sigma_k^2, \quad \text{and} \quad E^2 = A^2 B^2 - D^2,$$

and $a_k(x)$, $b_k(x)$ is defined by (1.2).

2 Asymptotic behaviour of ES_u

In this section we obtain the asymptotic behaviour of the expected number of u -sharp crossings of $Q_n(x) = 0$ given by (1.1). We prove the following theorem for the case that the increments $\Delta_1 \cdots \Delta_n$ are independent and have the same distribution. Also we assume that $\sigma_k^2 = 1$, for $k = 1 \cdots n$.

Theorem(2.1): Let $Q_n(x)$ be the random algebraic polynomial given by (1.1) for which $A_j = \Delta_1 + \dots + \Delta_j$ where Δ_k , $k = 1, \dots, n$ are independent and $\Delta_j \sim N(0, \sigma_j^2)$ then the expected number of u -sharp crossings of $Q_n(x) = 0$ satisfies

(i)- for $u = o(n^{5/4})$

$$\begin{aligned} ES_u(-\infty, \infty) = & \frac{1}{\pi} \log(2n+1) + \frac{1}{\pi}(1.920134502) \\ & + \frac{1}{\pi\sqrt{2n}} \left(-\pi + 2 \arctan\left(\frac{1}{2\sqrt{2n}}\right) \right) + \frac{C1}{n\pi} \\ & + \frac{u^2}{n^3\pi} \left(19.05803659 - \frac{8}{3} \ln(n^3 + 1) \right) \\ & + \frac{u^4}{n^6\pi} \left(-34989.96324 + \frac{32}{3} \ln(n^6 + 1) \right) + o(n^{-1}) \end{aligned}$$

where $C1 = -0.7190843756$ for n even and $C1 = 1.716159410$ for n odd.

(ii)- for $u = o(n^{3/2})$

$$ES_u(-\infty, \infty) = \frac{1}{\pi} \log(2n+1) + \frac{1}{\pi}(1.920134502) + o(1)$$

proof: Due to the behaviour of $Q_n(x)$, the asymptotic behaviour is treated separately on the intervals $1 < x < \infty$, $-\infty < x < -1$, $0 < x < 1$ and $-1 < x < 0$. For $1 < x < \infty$, using (1.3), the change of variable $x = 1 + \frac{t}{n}$ and the equality $\left(1 + \frac{t}{n}\right)^n = e^t \left(1 - \frac{t^2}{n}\right) + O\left(\frac{1}{n^2}\right)$, we find that

$$ES_u(1, \infty) = \frac{1}{n} \int_0^\infty f_n\left(1 + \frac{t}{n}\right) dt,$$

where $f_n(\cdot)$ is defined by (1.4). Using (1.5), and by tedious manipulation we have that

$$g_{2,n} \left(1 + \frac{t}{n}\right) = o(n^{-2})$$

and

$$n^{-1}g_{1,n} \left(1 + \frac{t}{n}\right) = \left(R_1(t) + \frac{S_1(t)}{n} + O\left(\frac{1}{n^2}\right)\right), \quad (2.1)$$

where

$$R_1(t) = \frac{\sqrt{(4t-15)e^{4t} + (24t+32)e^{3t} - e^{2t}(8t^3+12t^2+36t+18) + 8e^t t + 1}}{2t(-1 - 3e^{2t} + 4e^t + 2te^{2t})}$$

and $S_1(t) = S_{11}(t)/S_{12}(t)$ in which

$$\begin{aligned} S_{11}(t) = & -0.25 \left((4t^2 - 6t - 27)e^{6t} + (156 - 84t + 116t^2 - 24t^3)e^{5t} \right. \\ & + (16t^5 - 72t^4 + 96t^3 - 212t^2 + 220t - 331)e^{4t} + (328 - 168t + 128t^2 - 104t^3)e^{3t} \\ & \left. + (8t^4 + 8t^3 - 32t^2 + 42t - 153)e^{2t} + (28 - 4t - 4t^2)e^t - 1 \right) \end{aligned}$$

$$\begin{aligned} S_{12}(t) = & \left(2e^{2t}t - 1 + 4e^t - 3e^{2t} \right)^2 \\ & \times (1 - 8t^3e^{2t} - 12e^{2t}t^2 + 8e^t t - 18e^{2t} - 36e^{2t}t - 15e^{4t} + 32e^{3t} + 24e^{3t}t + 4te^{4t})^{1/2} \end{aligned}$$

One can easily verify that as $t \rightarrow \infty$,

$$R_1(t) = \frac{1}{2t^{3/2}} + O(t^{-2}), \quad S_1(t) = -\frac{1}{8t^{1/2}} + O(t^{-3/2}).$$

As (2.1) can not be integrated term by term, we use the equality

$$\frac{I_{[t>1]}}{8n\sqrt{t}} = \frac{I_{[t>1]}}{8n\sqrt{t} + t\sqrt{t}} + O\left(\frac{1}{n^2}\right), \quad (2.2)$$

where

$$I_{[t>1]} = \begin{cases} 1 & \text{if } t \geq 1 \\ 0 & \text{if } t < 1 \end{cases},$$

Thus by (2.2) we have that

$$\frac{1}{n}f_n\left(1 + \frac{t}{n}\right) = -\frac{I_{[t>1]}}{\pi(8n\sqrt{t} + t\sqrt{t})} + \frac{R_1(t)}{\pi} + \frac{1}{\pi} \left(\frac{S_1(t)}{n} + \frac{I_{[t>1]}}{8n\sqrt{t}} \right) + O\left(\frac{1}{n^2}\right).$$

This expression is term by term integrable, and provides that

$$ES_u(1, \infty) = \frac{1}{n} \int_0^\infty f_n \left(1 + \frac{t}{n}\right) dt = \frac{1}{2\pi\sqrt{2n}} \left(-\pi + 2 \arctan\left(\frac{1}{2\sqrt{2n}}\right)\right) \\ + \frac{1}{\pi} \int_0^\infty R_1(t) dt + \frac{1}{\pi n} \int_0^\infty \left(S_1(t) + \frac{I_{[t>1]}}{8\sqrt{t}}\right) dt + O\left(\frac{1}{n^2}\right),$$

where $\int_0^\infty R_1(t) dt = 0.7348742023$, and $\int_0^\infty \left(S_1(t) + I_{[t>1]}(8\sqrt{t})^{-1}\right) dt = -0.2496371198$.

For $-\infty < x < -1$, using $x = -1 - \frac{t}{n}$, we have $ES_u(-\infty, -1) = \frac{1}{n} \int_0^\infty f_n \left(-1 - \frac{t}{n}\right) dt$, where $f_n(\cdot)$ is defined by (1.4), and by (1.5)

$$n^{-1} g_{1,n} \left(-1 - \frac{t}{n}\right) = \left(R_2(t) + \frac{S_2(t)}{n} + O\left(\frac{1}{n^2}\right)\right) \quad (2.3)$$

where

$$R_2(t) = 1/2 \sqrt{\frac{-2e^{2t} + e^{4t} + 1 - 12e^{2t}t^2 - 8t^3e^{2t} + 4te^{4t} - 4te^{2t}}{t^2(e^{2t} - 1 + 2te^{2t})^2}};$$

$$\text{for } n \text{ even} \quad S_2(t) = \frac{S_{21}(t) + S_{22}(t)}{4S_{23}(t)}, \quad \text{and for } n \text{ odd} \quad S_2(t) = \frac{S_{21}(t) - S_{22}(t)}{4S_{23}(t)}$$

in which

$$S_{21}(t) = 1 + (-8t^4 + 30t - 8t^3 + 48t^2 - 3)e^{2t} \\ + (3 - 12t + 52t^2 + 96t^3 + 40t^4 - 16t^5)e^{4t} - (18t + 4t^2 + 1)e^{6t}, \\ S_{22}(t) = (8t + 32t^3 + 40t^2)e^{3t} + (-8t^2 - 12t)e^{5t} + 4e^{7t}, \\ S_{23}(t) = (e^{4t}(4t + 1) - 2e^{2t}(1 + 2t + 6t^2 + 4t^3) + 1)^{1/2} (e^{2t}(2t + 1) - 1)^2.$$

and

$$g_{2,n} \left(-1 - \frac{t}{n}\right) = \left(\frac{u^2}{n^3}\right) G_{2,1}(t) + o\left(\frac{1}{n}\right) \quad (2.4)$$

where

$$G_{2,1}(t) = \frac{16(2e^{2t}t + e^{2t} - 1)t^3}{2e^{2t} - e^{4t} + 12e^{2t}t^2 - 1 + 8e^{2t}t^3 - 4e^{4t}t + 4e^{2t}t}$$

It can be seen that as $t \rightarrow \infty$,

$$R_2(t) = \frac{1}{2t^{3/2}} + O(t^{-2}), \quad S_2(t) = \frac{-1}{8t^{1/2}} + O(t^{-3/2}), \quad G_{2,1}(t) = o(e^{-t}).$$

Now by using the equality (2.2) we have that

$$\begin{aligned} \frac{1}{n} f_n(-1 - \frac{t}{n}) &= -\frac{I_{[t>1]}}{\pi(8n\sqrt{t} + t\sqrt{t})} + \frac{1}{\pi} \left(R_2(t) + \frac{1}{n} (S_2(t) + \frac{I_{[t>1]}}{8\sqrt{t}}) \right. \\ &\quad \left. + \frac{u^2}{n^3} R_2(t) G_{2,1}(t) + \frac{u^4}{2n^6} R_2(t) G_{2,1}^2(t) \right) + o(n^{-1}). \end{aligned}$$

Thus

$$\begin{aligned} ES_u(-\infty, -1) &= \frac{1}{n} \int_0^\infty f_n(-1 - \frac{t}{n}) dt = \frac{1}{2\pi\sqrt{2n}} \left(-\pi + 2 \arctan(\frac{1}{2\sqrt{2n}}) \right) \\ &\quad + \frac{1}{\pi} \int_0^\infty R_2(t) dt + \frac{1}{n\pi} \int_0^\infty \left(S_2(t) + \frac{I_{[t>1]}}{8\sqrt{t}} \right) dt \\ &\quad + \frac{u^2}{n^3\pi} \int_0^\infty R_2(t) G_{2,1}(t) dt + \frac{u^4}{n^6\pi} \int_0^\infty \frac{1}{2} R_2(t) G_{2,1}^2(t) dt + o(n^{-1}) \end{aligned}$$

where $\int_0^\infty R_2(t) dt = 1.095640061$, and

$$\int_0^\infty R_2(t) G_{2,1}(t) dt = -2.418589510, \quad \int_0^\infty \frac{1}{2} R_2(t) G_{2,1}^2(t) dt = 7.057233216,$$

and for n odd $\int_0^\infty \left(S_2(t) + I_{[t>1]}(8\sqrt{t})^{-1} \right) dt = -0.0322863105$, and for n even $\int_0^\infty \left(S_2(t) + I_{[t>1]}(8\sqrt{t})^{-1} \right) dt = -0.4677136959$.

For $0 < x < 1$, let $x = 1 - \frac{t}{n+t}$, then $ES_u(0, 1) = \left(\frac{n}{(n+t)^2} \right) \int_0^\infty f_n \left(1 - \frac{t}{n+t} \right) dt$, where $f_n(\cdot)$ is defined by (1.4), and by (1.5)

$$g_{2,n} \left(1 - \frac{t}{n+t} \right) = o(n^{-2}), \quad (2.5)$$

and

$$\begin{aligned} \left(\frac{n}{(n+t)^2} \right) g_{1,n} \left(1 - \frac{t}{n+t} \right) &= \left(1 - \frac{2t}{n} + O\left(\frac{1}{n^2}\right) \right) \left(R_3(t) + \frac{S_3(t)}{n} + O\left(\frac{1}{n^2}\right) \right) \\ &= \left(R_3(t) + \frac{S_3(t) - 2tR_3(t)}{n} \right) + O\left(\frac{1}{n^2}\right), \quad (2.6) \end{aligned}$$

where we observe that $R_3(t) \equiv R_1(-t)$ and $S_3(t) = S_{31}(t)/S_{32}(t)$, in which

$$\begin{aligned} S_{31}(t) &= \left(\left(-7t^2 - \frac{69}{2}t - \frac{63}{4} \right) e^{-6t} + \left(6t^3 + 35t - 55t^2 + 39 \right) e^{-5t} \right. \\ &\quad \left. + \left(49t - 4t^5 + 22t^4 + 91t^2 - \frac{63}{4} - 12t^3 \right) e^{-4t} + \left(-6t^3 - 30 - 44t^2 - 66t \right) e^{-3t} \right. \\ &\quad \left. + \left(\frac{35}{2}t + 2t^4 - 6t^3 + 16t^2 + \frac{123}{4} \right) e^{-2t} + \left(-9 - t - t^2 \right) e^{-t} + 3/4 \right), \end{aligned}$$

and $S_{32}(t) \equiv S_{12}(-t)$. Now as $t \rightarrow \infty$, $R_3(t) = \frac{1}{2t} + O(t^{-1/2}e^{-t/2})$, and $S_3(t) = \frac{3}{4} + O(t^2e^{-t})$.

Since the relation (2.6) is not term by term integrable we use the equality

$$\frac{I_{[t>1]}}{2t} - \frac{I_{[t>1]}}{4n+2t} = \frac{I_{[t>1]}}{2t} - \frac{I_{[t>1]}}{4n} + O\left(\frac{1}{n^2}\right), \quad (2.7)$$

can be written as

$$\begin{aligned} \frac{n}{(n+t)^2} f_n\left(1 - \frac{t}{n+t}\right) &= \frac{1}{\pi} \left(R_3(t) - \frac{I_{[t>1]}}{2t}\right) + \frac{1}{\pi} \left(\frac{I_{[t>1]}}{2t} - \frac{I_{[t>1]}}{4n+2t}\right) \\ &\quad + \frac{1}{n\pi} \left(S_3(t) - 2tR_3(t) + \frac{I_{[t>1]}}{4}\right) + O\left(\frac{1}{n^2}\right) \end{aligned}$$

Thus we have that

$$\begin{aligned} ES_u(0,1) &= \frac{n}{(n+t)^2} \int_0^\infty f_n\left(1 - \frac{t}{n+t}\right) dt \\ &= \frac{1}{\pi} \int_0^\infty \left(R_3(t) - \frac{I_{[t>1]}}{2t}\right) dt + \frac{1}{2\pi} (\log(4n+2) - \log(2)) \\ &\quad + \frac{1}{n\pi} \int_0^\infty \left(S_3(t) - 2tR_3(t) + \frac{I_{[t>1]}}{4}\right) dt + O\left(\frac{1}{n^2}\right) \end{aligned} \quad (2.8)$$

where $\int_0^\infty \left(R_3(t) - \frac{I_{[t>1]}}{2t}\right) dt = -0.2897712456$ and $\int_0^\infty \left(S_3(t) - 2tR_3(t) + \frac{I_{[t>1]}}{4}\right) dt = 0.498174649$.

For $-1 < x < 0$, let $x = -1 + \frac{t}{n+t}$, then $ES_u(-1,0) = \left(\frac{n}{(n+t)^2}\right) \int_0^\infty f_n\left(-1 + \frac{t}{n+t}\right) dt$, again by (1.4) and (1.5) we have

$$\left(\frac{n}{(n+t)^2}\right) g_{1,n}(t) = \left(\frac{n^2}{(n+t)^2}\right) \left(R_4(t) + \frac{S_4(t)}{n} + O\left(\frac{1}{n^2}\right)\right), \quad (2.9)$$

in which $R_4(t) \equiv R_2(-t)$.

for n even $S_4(t) = \frac{S_{41}(t) + S_{42}(t)}{4S_{43}(t)}$, and for n odd $S_4(t) = \frac{S_{41}(t) - S_{42}(t)}{4S_{43}(t)}$

In which

$$\begin{aligned} S_{41}(t) &= 8 \left(-9/4 t + 6 t^2 - 3 t^3 - \frac{9}{8} + t^4\right) e^{-2t} \\ &\quad + 8 \left(15 t^4 - 3/2 t - 22 t^3 + \frac{9}{8} + 19/2 t^2 - 2 t^5\right) e^{-4t} \\ &\quad + 8 \left(\frac{15}{4} t - 7/2 t^2 - 3/8\right) e^{-6t} + 3, \end{aligned}$$

$S_{42}(t) \equiv S_{22}(-t)$, and $S_{43}(t) \equiv S_{23}(-t)$. Finally we have

$$g_{2,n} \left(-1 + \frac{t}{n+t} \right) = \left(\frac{u^2}{n^3} \right) G_{2,1}(t) + o(n^{-1}) \quad (2.10)$$

where

$$G_{2,1}(t) = -\frac{16(-e^{-2t} + 1 + 2e^{-2t}t)t^3}{(-4e^{-4t}t + e^{-4t} - 12e^{-2t}t^2 + 4e^{-2t}t + 1 + 8e^{-2t}t^3 - 2e^{-2t})}$$

As $t \rightarrow \infty$ we have

$$R_4(t) = \frac{1}{2t} + O(t^{1/2}e^{-t}), \quad S_4(t) = \frac{3}{4} + O(te^{-t}), \quad G_{2,1}(t) = -16t^3 + O(t^4e^{-2t}).$$

We have that

$$\begin{aligned} \frac{n^2}{(n+t)^2} f_n \left(-1 + \frac{t}{n+t} \right) &= \frac{1}{\pi} \left(1 - \frac{2t}{n} + O\left(\frac{1}{n^2}\right) \right) \left(R_4(t) + \frac{S_4(t)}{n} + O\left(\frac{1}{n^2}\right) \right) \\ &\times \left(1 + \frac{u^2}{n^3} G_{2,1}(t) + \frac{u^4}{2n^6} G_{2,1}^2(t) + o(n^{-1}) \right) \\ &= \frac{1}{\pi} \left\{ R_4(t) + \frac{S_4(t) - 2tR_4(t)}{n} + \frac{u^2}{n^3} R_4(t) G_{2,1}(t) \right. \\ &\quad \left. + \frac{u^4}{2n^6} R_4(t) G_{2,1}^2(t) \right\} + o(n^{-1}) \end{aligned}$$

Since this is not term by term integrable we use (2.7) and the following equalities to solve this

$$\frac{8u^2t^2}{n^3} = \frac{8u^2t^2}{n^3 + \exp(t^3)} + o(n^{-2}), \quad \frac{64u^4t^5}{n^6} = \frac{64u^4t^5}{n^6 + \exp(t^6)} + o(n^{-2})$$

Thus we have that

$$\begin{aligned} ES_u(-1, 0) &= \frac{n^2}{(n+t)^2} \int_0^\infty f_n \left(-1 + \frac{t}{n+t} \right) dt \\ &= \frac{1}{2\pi} (\log(2n+1)) + \frac{1}{\pi} \left\{ \int_0^\infty \left(R_4(t) - \frac{I_{[t>1]}}{2t} \right) dt(2) \right\} \\ &+ \frac{1}{n} \int_0^\infty \left(S_4(t) - 2tR_4(t) + \frac{I_{[t>1]}}{4} \right) dt \\ &+ \frac{u^2}{n^3} \int_0^\infty (R_4(t)G_{2,1}(t) + 8t^2) dt - u^2 \int_0^\infty \frac{8t^2}{n^3 + \exp(t^3)} dt \\ &+ \frac{u^4}{n^6} \int_0^\infty \left(\frac{1}{2} R_4(t)G_{2,1}^2(t) - 64t^5 \right) dt + u^4 \int_0^\infty \frac{64t^5}{n^6 + \exp(t^6)} dt \Big\} \\ &+ o(n^{-1}) \end{aligned} \quad (2.11)$$

where $\int_0^\infty (R_4(t) - \frac{I_{[t>1]}}{2t})dt = .3793914851$, and

$$\int_0^\infty (R_4(t)G_{2,1}(t) + 8t^2)dt = 21.47662610, \quad \int_0^\infty (\frac{1}{2}R_4(t)G_{2,1}^2(t) - 64t^5)dt = -34997.02047$$

and we have for n even $\int_0^\infty (S_4(t) - 2tR_4(t) + \frac{I_{[t>1]}}{4})dt = -0.4999081999$, and for n odd $\int_0^\infty (S_4(t) - 2tR_4(t) + \frac{I_{[t>1]}}{4})dt = 1.499908200$. Also we have that

$$\int_0^\infty \frac{8t^2}{n^3 + \exp(t^3)}dt = \frac{8\ln(n^3 + 1)}{3n^3}, \quad \int_0^\infty \frac{64t^5}{n^6 + \exp(t^6)}dt = \frac{32\ln(n^6 + 1)}{3n^6}.$$

So we arrive at the first assertion of the theorem.

Now for the proof of the second part of the theorem, that is for the case $k = o(n^{3/2})$ we study the asymptotic behavior of $ES_u(a, b)$ for different intervals $(-\infty, -1)$, $(-1, 0)$, $(0, 1)$ and $(1, \infty)$ separately.

For $1 < x < \infty$, using the change of variable $x = 1 + \frac{t}{n}$, and by (1.4), (1.5), and (2.1) we find that

$$\frac{1}{n} \int_0^\infty f_n(1 + \frac{t}{n})dt = \frac{1}{\pi} \int_0^\infty R_1(t)dt + o(1)$$

where $\int_0^\infty R_1(t)dt = 0.734874192$.

For $-\infty < x < -1$ using the change of variable $x = -1 - \frac{t}{n}$, and by the fact that $u^2/n^3 = o(1)$, as $n \rightarrow \infty$, we have that

$$\exp\{u^2 G_{2,1}(-1 - t/n)/n^3\} = 1 + o(1)$$

Therefore the relations (2.4) implies that $\exp\{g_2(-1 - t/n)\} = 1 + o(1)$. Thus by the relations (1.4), (1.5), (2.3), (2.4) we have that

$$\frac{1}{n} \int_0^\infty f_n(-1 - \frac{t}{n})dt = \frac{1}{\pi} \int_0^\infty R_2(t)dt + o(1)$$

where $\int_0^\infty R_2(t)dt = 1.095640061$.

For $0 < x < 1$ using the change of variable $x = 1 - \frac{t}{n+t}$, and relations (1.4), (1.5), (2.5), (2.6), and by using the equality (2.7) we have that

$$\frac{n}{(n+t)^2} \int_0^\infty f_n(-1 - \frac{t}{n+t})dt = \frac{1}{\pi} \int_0^\infty (R_3(t) - \frac{I_{[t>1]}}{2t})dt + \frac{1}{2\pi} \log(2n+1) + o(1)$$

where $\int_0^\infty (R_3(t) - \frac{I[t>1]}{2t})dt = -.28977126$.

For $-1 < x < 0$, using the change of variable $x = -1 + \frac{t}{n+t}$, (2.9), and by the same reasoning as above, the case $-\infty < x < -1$, we have that $\exp\{g_2(-1 + \frac{t}{n+t})\} = 1 + o(1)$. Thus by using the relations (2.9),(2.10), and by using the equality (2.7) we have that

$$\frac{n}{(n+t)^2} \int_0^\infty f_n(-1 + \frac{t}{n+t})dt = \frac{1}{\pi} \int_0^\infty (R_4(t) - \frac{I[t > 1]}{2t})dt + \frac{1}{2\pi} \log(2n+1) + o(1)$$

where $\int_0^\infty (R_4(t) - \frac{I[t>1]}{2t})dt = .3793914850$. This complete the proof of the theorem.

References

- [1] A T Bharucha-Ried and M Sambandham, *Random Polynomials*. Academic Press, New York (1986).
- [2] H Cramér and M R Leadbetter, *Stationary and Related Stochastic Processes*, John Wiley New York (1967).
- [3] K Farahmand, *Real zeros of random algebraic polynomials*, Proc. Amer. Math. Soc., 113(1991), 1077-1084.
- [4] K Farahmand, (1998). *Topics in Random Polynomials*, Chapman & Hall
- [5] I A Ibragimov and N B Maslova, *On the expected number of real zeros of random polynomials, coefficients with zero means*. Theory Probab. Appl., 16(1971, 228-248.
- [6] M Kac, *On the average number of real roots of a random algebraic equation*, Bull. Amer. Math. Soc., 49(1943), 314-320.
- [7] B F Logan and L A Shepp, *Real zeros of random polynomials*, Proc. London Math. Soc., Ser. 3, 18(1968), 29-35.
- [8] B F Logan and L A Shepp, *Real zeros of random polynomials*, Proc. London Math. Soc., Ser. 3, 18(1968), 308-314.

- [9] S Rezakhah and S Shemehsavar, *On the average Number of Level Crossings of Certain Gaussian Random Polynomials*. Nonlinear. Anal., 63, (2005), pp. e555-567.
- [10] S Rezakhah, and A R Soltani, *On the Levy and Harmonizable random polynomials*. Georgian. Math. Journal, vol. 7, No. 2,(2000) pp. 379-386.
- [11] S Rezakhah, A R Soltani, *On the Expected Number of Real Zeros of Certain Gaussian Random Polynomials*. Stoc. Anal, Appl, vol. 21, No. 1, pp. 223-234.
- [12] M Sambandham, *On the real roots of the random algebraic polynomial*, Indian J. Pure Appl. Math., 7(1976), 1062-1070.
- [13] M Sambandham, *On a random algebraic equation*, J. Indian Math. Soc., 41(1977), 83-97.
- [14] J E Wilkins, *An asymptotic expansion for the expected number of real zeros of a random polynomial*. Proc. Amer. Math. Soc. 103(1988), 1249-1258.