

# Profile-Kernel Likelihood Inference With Diverging Number of Parameters \*

By Clifford Lam and Jianqing Fan

Department of Operations Research and Financial Engineering  
Princeton University, Princeton, NJ, 08544

The generalized varying coefficient partially linear model with growing number of predictors arises in many contemporary scientific endeavor. In this paper we set foot on both theoretical and practical sides of profile likelihood estimation and inference. When the number of parameters grows with sample size, the existence and asymptotic normality of the profile likelihood estimator are established under some regularity conditions. Profile likelihood ratio inference for the growing number of parameters is proposed and Wilk's phenomenon is demonstrated. A new algorithm, called the accelerated profile-kernel algorithm, for computing profile-kernel estimator is proposed and investigated. Simulation studies show that the resulting estimates are as efficient as the fully iterative profile-kernel estimates. For moderate sample sizes, our proposed procedure saves much computational time over the fully iterative profile-kernel one and gives stabler estimates. A set of real data is analyzed using our proposed algorithm.

*Short Title:* High-dimensional profile likelihood.

*AMS 2000 subject classifications.* Primary 62G08; secondary 62J12, 62F12.

*Key words and phrases.* Generalized linear models, varying coefficients, high dimensionality, asymptotic normality, profile likelihood, generalized likelihood ratio tests.

---

\*Clifford Lam is PhD student, Department of Operation Research and Financial Engineering, Princeton University, Princeton, NJ 08544 (email: wlam@princeton.edu); Jianqing Fan is Professor, Department of Operation Research and Financial Engineering, Princeton University, Princeton, NJ 08544 (email: jqfan@princeton.edu). Financial support from the NSF grant DMS-0354223, DMS-0704337 and NIH grant R01-GM072611 is gratefully acknowledged.

# 1 Introduction

Semiparametric models with large number of predictors arise frequently in many contemporary statistical studies. Large data set and high-dimensionality characterize many contemporary scientific endeavors ([6]; [8]). Statistical models with many predictors are frequently employed to enhance the explanatory and predictive powers. At the same time, semiparametric modeling is frequently incorporated to balance between modeling biases and “curse of dimensionality”. Profile likelihood techniques ([23]) are frequently applied to this kind of semiparametric models. When the number of predictors is large, it is more realistic to regard it growing with the sample size. Yet, few results are available for semiparametric profile inferences when the number of parameters diverges with sample size. This paper focuses on profile likelihood inferences with diverging number of parameters in the context of the generalized varying coefficient partially linear model (GVCPLM).

GVCPLM is an extension the generalized linear model ([20]) and the generalized varying-coefficient model ([12]; [4]). It allows some coefficient functions to vary with certain covariates  $U$  such as age ([9]), toxic exposure level or time variable in a longitudinal data or survival analysis ([22]). Therefore, general interactions, not just the linear interaction as in parametric models, between the variable  $U$  and these covariates are explored nonparametrically.

If  $Y$  is a response variable and  $(U, \mathbf{X}, \mathbf{Z})$  is the associated covariates, then by letting  $\mu(u, \mathbf{x}, \mathbf{z}) = E\{Y|(U, \mathbf{X}, \mathbf{Z}) = (u, \mathbf{x}, \mathbf{z})\}$ , the GVCPLM takes the form

$$g\{\mu(u, \mathbf{x}, \mathbf{z})\} = \mathbf{x}^T \boldsymbol{\alpha}(u) + \mathbf{z}^T \boldsymbol{\beta}, \quad (1.1)$$

where  $g(\cdot)$  is a known link function,  $\boldsymbol{\beta}$  a vector of unknown regression coefficients and  $\boldsymbol{\alpha}(\cdot)$  a vector of unknown regression functions. One of the advantages over the varying coefficient model is that GVCPLM allows more efficient estimation when some coefficient functions are not really varying with  $U$ , after adjustment of other genuine varying effects. It also allows more interpretable model, where primary interest is focused on the parametric component.

## 1.1 A motivating example

We use a real data example to demonstrate the need for GVCPLM. The Fifth National Bank of Springfield faced a gender discrimination suit in which female received substan-

Table 1: Proportions of employees having job grade higher than 4

	Covariate <b>TotalYrsExp</b>		
	0-7	8-16	$\geq 17$
Age $\leq 35$	1/11	1/9	0
Age $> 35$	0	2/11	8/21

tially smaller salaries than male employees. This example is based on a real case with data dated 1995. Only the bank’s name is changed. See Example 11.3 of [2]. Among 208 employees, eight variables are collected. They include employee’s salary; age; year hired; number of years of working experience at another bank; gender; PC Job, a dummy variable with value 1 if the employee’s job is computer related; educational level, a categorical variable with categories 1 (finished school), 2 (finished some college courses), 3 (obtained a bachelor’s degree), 4 (took some graduate courses), 5 (obtained a graduate degree); job grade, a categorical variable indicating the current job level, the possible levels being 1–6 (6 the highest).

[9] has conducted such a salary analysis using an additive model with quadratic spline and does not find a significant evidence of gender difference. However, salary is directly related to the job grade. With the adjustment for the job grade, the salary discrimination can not easily be seen. An important question then arises if female employees have lower probability getting promoted. In analyzing such probability, a common tool will be the logistic regression, a class of the generalized linear model (for example, see [20]).

To this end, we create a binary response variable **HighGrade4**, indicating if Job Grade is greater than 4. The associated covariates are **Female**(1 for female employee and 0 otherwise), **Age**, **TotalYrsExp**(total years of working experience), **PCJob**, **Edu**(level of education). If the covariate **Female** has a significantly negative coefficient, then it would suggest that female employees are harder to promote to higher grade jobs.

However, in a simple logistic regression, the effect of a covariate cannot change with another covariate nonparametrically. Table 1 shows the proportion of employees having a job grade higher than 4, categorized by **Age** and **TotalYrsExp**. Clearly interactions between **Age** and **TotalYrsExp** have to be considered.

This can be done by creating categorical variables over the covariate **Age**. However this would increase the number of predictors considerably if we create many categories of **Age**. More importantly, we do not know where to draw the borders of each **Age** category and how many categories should be produced. This problem is nicely overcome if we allow

the coefficient of **TotalYrsExp** to vary with **Age**, so that we obtain a coefficient function of **Age** for **TotalYrsExp**. See section 4.3 for a detail analysis of the data.

If interactions between different variables are considered, then the number of predictors will be large compare with the sample size  $n = 208$ . This motivates us to consider the setting  $p_n \rightarrow \infty$  as  $n \rightarrow \infty$  and present general theories in section 2, where such a setting will be faced by many modern statistical applications.

## 1.2 Goals of the paper

When the number of parameters  $\beta$  is fixed and the link  $g$  is identity, the model (1.1) has been considered by [33], [17] and [31], and [1]. [7] propose a profile-kernel inference for such a varying coefficient partial linear model (VCPLM) and [18] considered a backfitting-based procedure for model selection in VCPLM. All of these papers rely critically on the explicit form of the estimation procedures and the techniques can not easily be applied to the GVCPLM.

Modern statistical applications often involve estimation of large number of parameters. It is of interest to derive asymptotic properties for the profile likelihood estimators under model (1.1) when number of parameters diverges. The fundamental questions arise naturally whether the profile likelihood estimator ([23]) still possesses efficient sampling properties, whether the profile likelihood ratio test for the parametric component possesses Wilks type of phenomenon, namely whether the asymptotic null distributions are independent of nuisance functions and parameters, and whether the usual sandwich formula provides a consistent estimator of the covariance matrix of the profile likelihood estimator. These questions are poorly understood and will be thoroughly investigated in Section 2. Pioneering work on statistical inference with diverging number of parameters include [14] which gave related results on M-estimators, and [25] which analyzed a regular exponential family under the same setting. [9] studied the penalized likelihood approach under such setting, whereas [10] investigated a semiparametric model with growing number of nuisance parameters.

Another goal of this paper is to provide an efficient algorithm for computing profile likelihood estimates under the model (1.1). To this end, we propose a new algorithm, called the accelerated profile-kernel algorithm, based on an important modification of the Newton-Raphson iterations. Computational difficulties ([19]) of the profile-kernel approach is significantly reduced, while nice sampling properties of such approach over

the backfitting algorithm (e.g. [13]) are retained. This will be convincingly demonstrated in Section 4, where the Poisson and Logistic specifications are considered for simulations. A new difference-based estimate for the parametric component is proposed as an initial estimate of our proposed profile-kernel procedure. Our method expands significantly the idea used in [32] and [7] for the partial linear model.

The outline of the paper is as follows. In Section 2 we briefly introduce the profile likelihood estimation with local polynomial modeling and present our main asymptotic results. Section 3 turns to the computational aspect, discussing the elements of computing in the accelerated profile-kernel algorithm. Simulation studies and an analysis of real data set are given Section 4. The proofs of our results are given in Section 5, and technical details in the appendix.

## 2 Properties of profile likelihood inference

Let  $(Y_{ni}; \mathbf{X}_i, \mathbf{Z}_{ni}, U_i)$ , where  $1 \leq i \leq n$  be a random sample where  $Y_{ni}$  is a scalar response variable,  $U_i, \mathbf{X}_i \in \mathbb{R}^q$  and  $\mathbf{Z}_{ni} \in \mathbb{R}^{p_n}$  are vectors of explanatory variables. We consider model (1.1) with  $\beta_n$  and  $\mathbf{Z}_n$  having dimensions  $p_n \rightarrow \infty$  as  $n \rightarrow \infty$ . Like the distributions in the exponential family, we assume that the conditional variance depends on the conditional mean so that  $\text{Var}(Y|U, \mathbf{X}, \mathbf{Z}_n) = V(\mu(u, \mathbf{X}, \mathbf{Z}_n))$  for a given function  $V$  (Our result is applicable even when  $V$  is multiplied by an unknown scale). Then, the conditional quasi-likelihood function is given by

$$Q(\mu, y) = \int_{\mu}^y \frac{s - y}{V(s)} ds.$$

As in [28], we denote by  $\alpha_{\beta_n}(u)$  the ‘least favorable curve’ of the nonparametric function  $\alpha(u)$ , which is defined as the one that maximizes

$$E_0\{Q(g^{-1}(\boldsymbol{\eta}^T \mathbf{X} + \beta_n^T \mathbf{Z}_n), Y_n)|U = u\} \quad (2.1)$$

with respect to  $\boldsymbol{\eta}$ , where  $E_0$  is the expectation taken under the true parameters  $\alpha_0(u)$  and  $\beta_{n0}$ . As will be discussed in section 2.1, through the use of least favorable curve, no undersmoothing of the nonparametric component is required to achieve asymptotic normality when  $p_n$  is diverging with  $n$ . Note that  $\alpha_{\beta_{n0}}(u) = \alpha_0(u)$ . Under some mild conditions, it satisfies

$$\frac{\partial}{\partial \boldsymbol{\eta}} E_0\{Q(g^{-1}(\boldsymbol{\eta}^T \mathbf{X} + \beta_n^T \mathbf{Z}_n), Y_n)|U = u\}|_{\boldsymbol{\eta}=\alpha_{\beta_n}(u)} = 0. \quad (2.2)$$

The profile-likelihood function for  $\beta_n$  is then

$$Q_n(\beta_n) = \sum_{i=1}^n Q\{g^{-1}(\alpha_{\beta_n}(U_i)^T \mathbf{X}_i + \beta_n^T \mathbf{Z}_{ni}), Y_{ni}\}, \quad (2.3)$$

if the least-favorable curve  $\alpha_{\beta_n}(\cdot)$  is known.

The least-favorable curve defined by (2.1) can be estimated by its sample version through a local polynomial regression approximation. For  $U$  in a neighborhood of  $u$ , approximate the  $j^{th}$  component of  $\alpha_{\beta_n}(\cdot)$  as

$$\begin{aligned} \alpha_j(U) &\approx \alpha_j(u) + \frac{\partial \alpha_j(u)}{\partial u}(U - u) + \cdots + \frac{\partial^p \alpha_j(u)}{\partial u^p}(U - u)^p/p! \\ &\equiv a_{0j} + a_{1j}(U - u) + \cdots + a_{pj}(U - u)^p/p!. \end{aligned}$$

Denoting  $\mathbf{a}_r = (a_{r1}, \dots, a_{rq})^T$  for  $r = 0, \dots, p$ , for each given  $\beta_n$ , we then maximize the local likelihood

$$\sum_{i=1}^n Q\{g^{-1}(\sum_{r=0}^p \mathbf{a}_r^T \mathbf{X}_i (U_i - u)^r / r! + \beta_n^T \mathbf{Z}_{ni}), Y_{ni}\} K_h(U_i - u) \quad (2.4)$$

with respect to  $\mathbf{a}_0, \dots, \mathbf{a}_p$ , where  $K(\cdot)$  is a kernel function and  $K_h(t) = K(t/h)/h$  is a re-scaling of  $K$  with bandwidth  $h$ . Thus, we get estimate  $\hat{\alpha}_{\beta_n}(u) = \hat{\mathbf{a}}_0(u)$ .

Plugging our estimates into the profile-kernel likelihood function (2.3), we have

$$\hat{Q}_n(\beta_n) = \sum_{i=1}^n Q\{g^{-1}(\hat{\alpha}_{\beta_n}(U_i)^T \mathbf{X}_i + \beta_n^T \mathbf{Z}_{ni}), Y_{ni}\}. \quad (2.5)$$

Maximizing  $\hat{Q}_n(\beta_n)$  with respect to  $\beta_n$  to get  $\hat{\beta}_n$ . With  $\hat{\beta}_n$ , the varying coefficient functions are estimated as  $\hat{\alpha}_{\hat{\beta}_n}(u)$ .

One property of the profile quasi-likelihood is that the first and second order Bartlett's identities continue to hold. In particular, with the definition given by (2.3), then for any  $\beta_n$ , we have

$$\mathbf{E}_{\beta_n} \left( \frac{\partial Q_n}{\partial \beta_n} \right) = 0, \quad \mathbf{E}_{\beta_n} \left( \frac{\partial Q_n}{\partial \beta_n} \frac{\partial Q_n}{\partial \beta_n^T} \right) = -\mathbf{E}_{\beta_n} \left( \frac{\partial^2 Q_n}{\partial \beta_n \partial \beta_n^T} \right). \quad (2.6)$$

See [28] for more details. These properties give rise to the asymptotic efficiency of the profile likelihood estimator.

## 2.1 Consistency and asymptotic normality of $\hat{\beta}_n$

We need Regularity Conditions (A) - (G) in Section 5 for the following results.

**Theorem 1** (Existence of profile likelihood estimator). *Assume that Conditions (A)-(G) are satisfied. If  $p_n^4/n \rightarrow 0$  as  $n \rightarrow \infty$  and  $h = O(n^{-a})$  with  $(4(p+1))^{-1} < a < 1/2$ , then there is a local maximizer  $\hat{\beta}_n \in \Omega_n$  of  $\hat{Q}_n(\beta_n)$  such that  $\|\hat{\beta}_n - \beta_{n0}\| = O_P(\sqrt{p_n/n})$ .*

The above rate is the same as the one established by [14] for the M-estimator.

Note that the optimal bandwidth  $h = O(n^{-1/(2p+3)})$  is included in Theorem 1. Hence  $\sqrt{n/p_n}$ -consistency is achieved without the need of undersmoothing of the nonparametric component. In particular, when  $p_n$  is fixed, the result is in line with those, for instance, by [27] in a different context.

Define  $I_n(\beta_n) = n^{-1} \mathbf{E}_{\beta_n}(\frac{\partial Q_n}{\partial \beta_n} \frac{\partial Q_n}{\partial \beta_n^T})$ , which is an extension of the Fisher matrix. Since the dimensionality grows with sample size, we need to consider the arbitrary linear combination of the profile kernel estimator  $\hat{\beta}_n$  as stated in the following theorem.

**Theorem 2** (Asymptotic normality). *Under Conditions (A) - (G), if  $p_n^5/n = o(1)$  and  $h = O(n^{-a})$  for  $3/(10(p+1)) < a < 2/5$ , then the consistent estimator  $\hat{\beta}_n$  in Theorem 1 satisfies*

$$\sqrt{n} A_n I_n^{1/2}(\beta_{n0})(\hat{\beta}_n - \beta_{n0}) \xrightarrow{D} N(0, G),$$

where  $A_n$  is an  $l \times p_n$  matrix such that  $A_n A_n^T \rightarrow G$ , and  $G$  is an  $l \times l$  nonnegative symmetric matrix.

A remarkable technical achievement of our result is that it does not require undersmoothing of the nonparametric component, as in Theorem 1, thanks to the profile likelihood approach. The key lies in a special orthogonality property of the least favorable curve (see equation (2.2) and Lemma 2). Asymptotic normality without undersmoothing is also proved in [30] for both backfitting and profiling methods.

Theorem 2 shows that profile likelihood produces a semi-parametric efficient estimate even when the number of parameters diverges. To see this more explicitly, let  $p_n = r$  be a constant. Then, by taking  $A_n = I_r$ , we obtain

$$\sqrt{n}(\hat{\beta}_n - \beta_{n0}) \xrightarrow{D} N(0, I^{-1}(\beta_{n0})).$$

The asymptotic variance of  $\hat{\beta}_n$  achieves the efficient lower bound given, for example, in [28].

## 2.2 Profile likelihood ratio test

After estimation of parameters, it is of interest to test the statistical significance of certain variables in the parametric component. Consider the problem of testing linear hypotheses:

$$H_0 : A_n \beta_{n0} = 0 \longleftrightarrow H_1 : A_n \beta_{n0} \neq 0,$$

where  $A_n$  is an  $l \times p_n$  matrix and  $A_n A_n^T = I_l$  for a fixed  $l$ . Note that both the null and the alternative hypotheses are semi-parametric, with nuisance functions  $\alpha(\cdot)$ . The generalized likelihood ratio test (GLRT) is defined by

$$T_n = 2 \left\{ \sup_{\Omega_n} \hat{Q}_n(\beta_n) - \sup_{\Omega_n; A_n \beta_n = 0} \hat{Q}_n(\beta_n) \right\}.$$

Note that the testing procedure does not depend explicitly on the estimated asymptotic covariance matrix. The following theorem shows that, even when the number of parameters diverges with sample size,  $T_n$  still follows a chi-square distribution asymptotically, without reference to any nuisance parameters and functions. This reveals the Wilk's phenomenon, as termed in [11].

**Theorem 3** *Assuming Conditions (A) - (G), under  $H_0$ , we have*

$$T_n \xrightarrow{D} \chi_l^2,$$

*provided that  $p_n^5/n = o(1)$  and  $h = O(n^{-a})$  for  $3/(10(p+1)) < a < 2/5$ .*

## 2.3 Consistency of the sandwich covariance formula

The estimated covariance matrix for  $\hat{\beta}_n$  can be obtained by the sandwich formula

$$\hat{\Sigma}_n = n^2 \{ \nabla^2 \hat{Q}_n(\hat{\beta}_n) \}^{-1} \widehat{\text{cov}} \{ \nabla \hat{Q}_n(\hat{\beta}_n) \} \{ \nabla^2 \hat{Q}_n(\hat{\beta}_n) \}^{-1},$$

where the middle matrix has  $(j, k)$  entry given by

$$\begin{aligned} (\widehat{\text{cov}} \{ \nabla \hat{Q}_n(\hat{\beta}_n) \})_{jk} &= \left\{ \frac{1}{n} \sum_{i=1}^n \frac{\partial \hat{Q}_{ni}(\hat{\beta}_n)}{\partial \beta_{nj}} \frac{\partial \hat{Q}_{ni}(\hat{\beta}_n)}{\partial \beta_{nk}} \right\} \\ &\quad - \left\{ \frac{1}{n} \sum_{i=1}^n \frac{\partial \hat{Q}_{ni}(\hat{\beta}_n)}{\partial \beta_{nj}} \frac{1}{n} \sum_{i=1}^n \frac{\partial \hat{Q}_{ni}(\hat{\beta}_n)}{\partial \beta_{nk}} \right\}. \end{aligned}$$

With the notation  $\Sigma_n = I_n^{-1}(\beta_{n0})$ , we have the following consistency result for the sandwich formula.



**Theorem 4** *Assuming Conditions (A) - (G). If  $p_n^4/n = o(1)$  and  $h = O(n^{-a})$  with  $(4(p+1))^{-1} < a < 1/2$ , we have*

$$A_n \hat{\Sigma}_n A_n^T - A_n \Sigma_n A_n^T \xrightarrow{\mathbb{P}} 0 \text{ as } n \rightarrow \infty$$

for any  $l \times p_n$  matrix  $A_n$  such that  $A_n A_n^T = G$ .

This result provides a simple way to construct confidence intervals for  $\beta_n$ . Simulation results show that this formula indeed provides a good estimate of the covariance of  $\hat{\beta}_n$  for a variety of practical sample sizes.

### 3 Computation of the estimates

Finding  $\hat{\beta}_n$  to maximize the profile likelihood (2.5) poses some interesting challenges, as the function  $\hat{\alpha}_{\beta_n}(u)$  in (2.5) depends on  $\beta_n$  implicitly (except the least-square case). The full profile-kernel estimate is to directly employ the Newton-Raphson iterations:

$$\beta_n^{(k+1)} = \beta_n^{(k)} - \{\nabla^2 \hat{Q}_n(\beta_n^{(k)})\}^{-1} \nabla \hat{Q}_n(\beta_n^{(k)}), \quad (3.1)$$

starting from the initial estimate  $\beta^{(0)}$ . We will call the estimate  $\beta_n^{(k)}$  and  $\hat{\alpha}_{\beta_n^{(k)}}(u)$  the  $k$ -step estimate ([3]; [26]). The initial estimate for  $\beta_n$  is critically important for the computational speed. We will propose a new and fast initial estimate in Section 3.1.

The first two derivatives of  $\nabla \hat{Q}_n(\beta_n)$  is given by

$$\begin{aligned} \nabla \hat{Q}_n(\beta_n) &= \sum_{i=1}^n q_{1i}(\beta_n) (\mathbf{Z}_{ni} + \hat{\alpha}'_{\beta_n}(U_i) \mathbf{X}_i), \\ \nabla^2 \hat{Q}_n(\beta_n) &= \sum_{i=1}^n q_{2i}(\beta_n) (\mathbf{Z}_{ni} + \hat{\alpha}'_{\beta_n}(U_i) \mathbf{X}_i) (\mathbf{Z}_{ni} + \hat{\alpha}'_{\beta_n}(U_i) \mathbf{X}_i)^T \\ &\quad + \sum_{i=1}^n \left\{ q_{1i}(\beta_n) \sum_{r=1}^q \frac{\partial^2 \hat{\alpha}_{\beta_n}^{(r)}(U_i)}{\partial \beta_n \partial \beta_n^T} X_{ir} \right\}, \end{aligned} \quad (3.2)$$

where  $q_l(x, y) = \frac{\partial^l}{\partial x^l} Q(g^{-1}(x), y)$ ,  $q_{ki}(\beta_n) = q_k(\hat{m}_{ni}(\beta_n), Y_{ni})$  ( $k = 1, 2$ ) with  $\hat{m}_{ni}(\beta_n) = \hat{\alpha}_{\beta_n}(U_i)^T \mathbf{X}_i + \mathbf{Z}_{ni}^T \beta_n$ . In the above formulae,  $\hat{\alpha}'_{\beta_n}(u) = \frac{\partial \hat{\alpha}_{\beta_n}(u)}{\partial \beta_n}$  is a  $p_n$  by  $q$  matrix and  $\alpha_{\beta_n}^{(r)}(u)$  is the  $r^{\text{th}}$  component of  $\alpha_{\beta_n}(u)$ .

As the first two derivatives of  $\hat{\alpha}_{\beta_n}(u)$  are hard to compute in (3.2), one can employ the backfitting algorithm, which iterates between (2.4) and (2.3). This is really the same as the fully iterated algorithm (3.1) but ignores the functional dependence of  $\hat{\alpha}_{\beta_n}(u)$

in (2.5) on  $\beta_n$ ; it uses the value of  $\beta_n$  in the previous step of the iteration as a proxy. More precisely, the backfitting algorithm treats the terms  $\hat{\alpha}'_{\beta_n}(u)$  and  $\hat{\alpha}''_{\beta_n}(u)$  in (3.2) as zero and computes  $\hat{m}_{ni}(\beta_n)$  using the value of  $\beta_n$  from the previous iteration. The maximization is thus much easier to carry out, but the convergence speed can be reduced. See [13] and [19] for more descriptions of the two methods and some closed-form solutions proposed for the partially linear models.

Between these two extreme choices is our modified algorithm, which ignores the computation of the second derivative of  $\hat{\alpha}_{\beta_n}(u)$  in (3.1), but keeps its first derivative in the iteration. Namely, the second term in (3.2) is treated as zero. Details will be given in Section 3.2. It turns out that this algorithm improves significantly the computation with achieved accuracy. At the same time, it enhances dramatically the stability of the algorithm. We will term the algorithm as the accelerated profile-kernel algorithm.

When the quasi-likelihood becomes a square loss, the accelerated profile-kernel algorithm is exactly the same as that used to compute the full profile likelihood estimate, since  $\hat{\alpha}_{\beta_n}(\cdot)$  is linear in  $\beta_n$ .

### 3.1 Difference-based estimation

We generalize the difference-based idea to obtain an initial estimate  $\beta_n^{(0)}$ . The idea has been used in [32] and [7] to remove the nonparametric component in the partially linear model.

We first consider the specific case of the GVCPLM:

$$Y = \alpha(U)^T \mathbf{X} + \beta_n^T \mathbf{Z}_n + \varepsilon. \quad (3.3)$$

This is the varying-coefficient partially linear model studied by [33] and [31]. Let the random sample  $\{(U_i, \mathbf{X}_i^T, \mathbf{Z}_{ni}^T, Y_i)\}_{i=1}^n$  be from the model (3.3), with the data ordered according to the  $U_i$ 's. Under mild conditions, the spacing  $U_{i+j} - U_i$  is  $O_P(1/n)$ , so that

$$\alpha(U_{i+j}) - \alpha(U_i) \approx \gamma_0 + \gamma_1(U_{i+j} - U_i), \quad j = 1, \dots, q. \quad (3.4)$$

Indeed, it can be approximately zero; the linear term is used to reduce the approximation errors.

For given weights  $w_j$  (its dependence on  $i$  is suppressed for simplicity), define

$$Y_i^* = \sum_{j=1}^{q+1} w_j Y_{i+j-1}, \quad \mathbf{Z}_{ni}^* = \sum_{j=1}^{q+1} w_j \mathbf{Z}_{n(i+j-1)}, \quad \varepsilon_i^* = \sum_{j=1}^{q+1} w_j \varepsilon_{i+j-1}.$$

If we choose the weights to satisfy  $\sum_{j=1}^{q+1} w_j \mathbf{X}_{i+j-1} = \mathbf{0}$ , then using (3.3) and (3.4), we have

$$Y_i^* \approx \boldsymbol{\gamma}_0^T \mathbf{X}_i w_1 + \boldsymbol{\gamma}_1^T \sum_{j=1}^{q+1} w_j U_{i+j-1} \mathbf{X}_{i+j-1} + \boldsymbol{\beta}_n^T \mathbf{Z}_{ni}^* + \varepsilon_i^*,$$

Ignoring the approximation, which is of order  $O_P(n^{-1})$ , the above is a multiple regression model with parameters  $(\boldsymbol{\gamma}_0, \boldsymbol{\gamma}_1, \boldsymbol{\beta}_n)$ . The parameters can be found by a weighted least square fit to the  $(n - q)$  starred data. This yields a root-n consistent estimate of  $\boldsymbol{\beta}_n$ , as the above approximation for the finite  $q$  is of order  $O_P(n^{-1})$ .

To solve  $\sum_{j=1}^{q+1} w_j \mathbf{X}_{i+j-1} = \mathbf{0}$ , we need to find the rank of the matrix  $(\mathbf{X}_i, \dots, \mathbf{X}_{i+q})$ , denoted it by  $r$ . Fix  $q + 1 - r$  of the  $w_j$ 's and the rest can be determined uniquely by solving the system of linear equations for  $\{w_j, j = 1, \dots, q + 1\}$ . For random designs, with probability 1,  $r = q$ . Hence, the direction of the weights  $\{w_j, j = 1, \dots, q + 1\}$  is uniquely determined. For example, in the partial linear model,  $q = 1$  and  $\mathbf{X}_i = 1$ . Hence,  $(w_1, w_2) = c(1, -1)$  and the constant  $c$  can be taken to have a norm one. This results in the difference based estimator in [32] and [7].

To use the differencing idea to obtain an initial estimate of  $\boldsymbol{\beta}_n$  for the GVCPLM, we apply the transformation of the data. If  $g$  is the link function, we use  $g(Y_i)$  as the transformed data and proceed with the difference-based method as for the VCPLM. Note that for some models like the logistic regression with logit link and Poisson log-linear model, we need to make adjustments in transforming the data. We use  $g(y) = \log(\frac{y+\delta}{1-y+\delta})$  for the logistic regression and  $g(y) = \log(y + \delta)$  for the Poisson regression. Here, the parameter  $\delta$  is treated as a smoothing parameter like  $h$ , and its choice will be discussed in Section 3.4.

### 3.2 Accelerated profile-kernel algorithm

As mentioned before, the accelerated profile-kernel algorithm needs to compute  $\boldsymbol{\alpha}'_{\boldsymbol{\beta}_n}(u)$ , which will be replaced by its consistent estimate given in the following theorem. The proof is in section 5.

**Theorem 5** *Under Regularity Conditions (A)-(G), provided  $\sqrt{p_n}(h + c_n \log^{1/2}(1/h)) = o(1)$  where  $c_n = (nh)^{-1/2}$ , we have for each  $\boldsymbol{\beta}_n \in \Omega_n$ ,*

$$\hat{\boldsymbol{\alpha}}'_{\boldsymbol{\beta}_n}(u) = - \left\{ \sum_{i=1}^n q_{2i}(\boldsymbol{\beta}_n) \mathbf{Z}_{ni} \mathbf{X}_i^T K_h(U_i - u) \right\} \cdot \left\{ \sum_{i=1}^n q_{2i}(\boldsymbol{\beta}_n) \mathbf{X}_i \mathbf{X}_i^T K_h(U_i - u) \right\}^{-1}$$

*being a consistent estimator of  $\boldsymbol{\alpha}'_{\boldsymbol{\beta}_n}(u)$  which holds uniformly in  $u \in \Omega$ .*

Since the function  $q_2(\cdot, \cdot) < 0$  by Regularity Condition (D), by ignoring the second term in (3.2), the modified  $\nabla^2 \hat{Q}_n(\beta_n)$  in equation (3.2) is still negative-definite. This ensures the Newton-Raphson update of the profile-kernel procedure can be carried out smoothly. The intuition behind the modification is that, for a neighborhood around the true parameter  $\beta_{n0}$ , the least favorable curve  $\alpha_{\beta_n}(u)$  should be approximately linear in  $\beta_n$ .

### 3.3 One-step estimation for the nonparametric component

Given  $\beta_n = \beta_n^{(k)}$ , we need to compute  $\alpha_{\beta_n}(u)$  in order to compute  $\hat{m}_{ni}(\beta_n)$  and hence the modified gradient vector and Hessian matrix in (3.1). This is the same as estimating the varying coefficient functions under model (1.1) with known  $\beta_n$ . [4] propose a one-step local MLE, which is shown to be as efficient as the fully iterated one. They also propose an efficient algorithm to compute these varying coefficient functions. Their algorithm can be directly adapted here. Details can be found in [4].

### 3.4 Choice of bandwidth

As mentioned at the end of Section 3.1, in addition to choosing the bandwidth  $h$ , we have an extra smoothing parameter  $\delta$  to be determined due to the adjustments to the transformation of the response  $Y_{ni}$ . This two dimensional smoothing parameters  $(\delta, h)$  can be selected by a  $K$ -fold cross-validation, using the quasi-likelihood as a criterion function. As demonstrated in Section 4, the practical accuracy can be achieved in several iterations using the accelerated profile-kernel algorithm. Hence, the profile-kernel estimate can be computed rapidly. As a result, the  $K$ -fold cross-validation is not too computationally intensive, as long as  $K$  is not too large (e.g.  $K=5$  or  $10$ ).

## 4 Numerical properties

To evaluate the performance of estimator  $\hat{\alpha}(\cdot)$ , we use the square-root of average errors (RASE)

$$\text{RASE} = \left\{ n_{\text{grid}}^{-1} \sum_{k=1}^{n_{\text{grid}}} \|\hat{\alpha}(u_k) - \alpha(u_k)\|^2 \right\}^{1/2},$$

over  $n_{\text{grid}} = 200$  grid points  $\{u_k\}$ . The performance of the estimator  $\hat{\beta}_n$  is assessed by the generalized mean square error (GMSE)

$$\text{GMSE} = (\hat{\beta}_n - \beta_{n0})^T \mathbf{B} (\hat{\beta}_n - \beta_{n0}),$$

where  $\mathbf{B} = E\mathbf{Z}_n\mathbf{Z}_n^T$ .

Throughout our simulation studies, the dimensionality of parametric component is taken as  $p_n = \lfloor 1.8n^{1/3} \rfloor$  and the nonparametric component as  $q = 2$  in which  $X_1 = 1$  and  $X_2 \sim N(0, 1)$ . The rate  $p_n = O_P(n^{1/3})$  is not the same as presented in the theorems in section 2, but we use this to show the capability of handling a higher rate of parameters growth for the accelerated profile-kernel method. In addition, the covariates  $(\mathbf{Z}_n^T, X_2)^T$  is a  $(p_n + 1)$ -dimensional normal random vector with mean zero and covariance matrix  $(\sigma_{ij})$ , where  $\sigma_{ij} = 0.5^{|i-j|}$ . Furthermore, we always take  $U \sim U(0, 1)$  independent of the other covariates. Finally, we use  $\text{SD}_{\text{mad}}$  to denote the robust estimate of standard deviation, which is defined as interquartile range divided by 1.349. The number of simulations is 400 except that in Table 1 (which is 50) due to the intensive computation of the fully iterated profile-kernel estimate.

**Poisson model.** The response  $Y$ , given  $(U, \mathbf{X}, \mathbf{Z}_n)$ , has a Poisson distribution with the mean function  $\mu(U, \mathbf{X}, \mathbf{Z}_n)$  where

$$\log(\mu(U, \mathbf{X}, \mathbf{Z}_n)) = \mathbf{X}^T \boldsymbol{\alpha}(U) + \mathbf{Z}_n^T \boldsymbol{\beta}_n.$$

We have  $\beta_{n0} = (0.5, 0.3, -0.5, 1, 0.1, -0.25, 0, \dots, 0)^T$ , the  $p_n$ -dimensional parameters. The coefficient functions are given by

$$\alpha_1(u) = 4 + \sin(2\pi u), \quad \text{and} \quad \alpha_2(u) = 2u(1 - u).$$

**Bernoulli model.** The response  $Y$ , given  $(U, \mathbf{X}, \mathbf{Z}_n)$ , has a Bernoulli distribution with the success probability given by

$$p(U, \mathbf{X}, \mathbf{Z}_n) = \exp\{\mathbf{X}^T \boldsymbol{\alpha}(U) + \mathbf{Z}_n^T \boldsymbol{\beta}_n\} / [1 + \exp\{\mathbf{X}^T \boldsymbol{\alpha}(U) + \mathbf{Z}_n^T \boldsymbol{\beta}_n\}].$$

The  $p_n$ -dimensional parameters are  $\beta_{n0} = (3, 1, -2, 0.5, 2, -2, 0, \dots, 0)^T$  and the varying coefficient functions is given by

$$\alpha_1(u) = 2(u^3 + 2u^2 - 2u), \quad \text{and} \quad \alpha_2(u) = 2\cos(2\pi u).$$

Throughout our numerical studies, we use the Epanechnikov kernel  $K(u) = 0.75(1 - u^2)_+$  and the 5-fold cross-validation to choose a bandwidth  $h$  and  $\delta$ . With the assistance of

the 5-fold cross-validation, we chose  $\delta = 0.1$  and  $h = 0.1, 0.08, 0.075$  and  $0.06$  respectively for  $n = 200, 400, 800$  and  $1500$  for the Poisson model. For the Bernoulli model,  $\delta = 0.005$  and  $h = 0.45, 0.4, 0.25$  and  $0.18$  were chosen respectively for  $n = 200, 400, 800$  and  $1500$ .

Note that  $X_2$  and the  $Z_{ni}$ 's are not bounded r.v.s as needed in condition (A) in section 5. However, these still satisfy the moment conditions needed in the proofs, and condition (A) is imposed to merely simplify these proofs. Condition (B) is satisfied mainly because the correlations between further  $Z_{ni}$ 's are weak, and condition (C) is satisfied because it involves products of standard normal r.v.s which are bounded in the first two moments.

## 4.1 Comparisons of algorithms

Table 2: Computation time and accuracy for different computing algorithms

$n$	$p_n$	backfitting	accelerated profile-kernel	full profile-kernel
<i>Median and <math>SD_{mad}</math> (in parentheses) of computing times in seconds</i>				
200	10	.6(.0)	.7(.0)	77.2(.2)
400	13	.8(.0)	1.4(.0)	463.2(.9)
<i>Median and <math>SD_{mad}</math> (in parentheses) of GMSE (multiplied by <math>10^4</math>)</i>				
200	10	10.72(6.47)	5.45(2.71)	9.74(14.67)
400	13	5.63(4.39)	2.78(1.19)	5.26(9.46)
<i>Median RASE relative to the oracle estimate</i>				
200	10	.848	.970	.895
400	13	.856	.986	.882

We first compare the computing times and the accuracies among three algorithms: 3-step backfitting, 3-step accelerated profile-kernel and fully-iterated profile-kernel algorithms. All of them use the difference-based estimate as the initial estimate. Table 2 summarizes the results based on the Poisson model with 50 samples.

With the same initial values, the backfitting algorithm is slightly faster than the accelerated profile-kernel algorithm, which in turn by far faster than the full profile-kernel algorithm. Our experience shows that the backfitting algorithm needs more than 20 iterations to converge without improving too much the GMSE. In terms of the accuracy of estimating the parametric component, the accelerated profile-kernel algorithm is about twice as accurate as the backfitting algorithm and the full profile-kernel one. This demonstrates the advantage of keeping the curvature of the least-favorable function in the Newton-Raphson algorithm. For the nonparametric component, we compare RASEs of three algorithms with those based on the oracle estimator, which uses the true value of

$\beta_n$ . The ratios of the RASEs based on the oracle estimator and those based on the three algorithms are reported in Table 1. It is clear that the accelerated profile-kernel estimate performs very well in estimating the nonparametric components, mimicking very well the oracle estimator. The second best is the backfitting algorithm.

We have also compared the three algorithms using the Bernoulli model. Our proposed accelerated profile-kernel estimate still performs the best in terms of accuracy, though the improvement is not as dramatic as those for the Poisson model. We speculate that the poor performance of the full profile-kernel estimate is due to its unstable implementation that is related to computing the second derivatives of the least-favorable curve.

Table 3: Medians of the percentages of GMSE based on the accelerated profile-kernel estimates

$n$	$p_n$	<i>Poisson</i>		<i>Bernoulli</i>	
		AF/DBE	AF/3S	AF/DBE	AF/3S
200	10	8.2	99.9	64.1	101.7
400	13	6.0	100.2	52.7	104.7
800	16	5.0	100.1	50.9	102.6
1500	20	4.2	100.0	46.4	100.5

We next demonstrate the accuracy of the three-step accelerated profile-kernel estimate (3S), compared with the fully-iterated accelerated profile-kernel estimate (AF) (iterating until convergence), and the difference-based estimate (DBE), which is our initial estimate. Table 3 reports the ratios of GMSE based on 400 simulations. It demonstrates convincingly that with the DBE as the initial estimate, three iterations achieve the accuracy that is comparably with the fully iterated algorithm. In fact, the one-step accelerated profile-kernel estimates improve dramatically (not shown here) our initial estimate (DBE). On the other hand, the DBE itself is not accurate enough for GCVPLM.

The effect of bandwidth choice on the estimation of parametric component is summarized in Table 4. Denote by  $h_{CV}$  the bandwidth chosen by the cross-validation. We scaled the bandwidth up and down by using a factor of 1.5. For illustration, we use the one-step accelerated profile-kernel estimate. The results for three-step profile-kernel estimate are similar. We evaluate the performance for all components using GMSE and for the specific component  $\beta_5$  using MSE (the results for other components are similar). We do not report all the results here to save the space. It is clear that the GMSE does not sensitively depends on the bandwidth, as long as it is reasonably close to  $h_{CV}$ . This is consistent with our asymptotic results.

Table 4: One-step estimate of parametric components with different bandwidths

		<i>Poisson</i>				<i>Bernoulli</i>	
		<i>Median and <math>SD_{mad}</math> of <math>GMSE \times 10^5</math></i>		<i>Mean and SD of <math>MSE \times 10^4</math> for <math>\beta_5</math></i>		<i>Median and <math>SD_{mad}</math> of <math>GMSE \times 10</math></i>	
$n$	$p_n$	$h_{CV}$	$1.5h_{CV}$	$0.66h_{CV}$	$h_{CV}$	$0.66h_{CV}$	$h_{CV}$
200	10	5.9(3.0)	6.4(3.3)	993(112)	995(105)	8.2(4.4)	8.4(5.1)
400	13	3.1(1.4)	3.0(1.4)	1004(67)	1001(65)	4.8(2.2)	5.4(2.5)
800	16	1.7(0.7)	1.7(0.6)	999(47)	999(46)	2.7(1.0)	2.7(1.1)
1500	20	1.1(0.3)	1.1(0.4)	1000(32)	1000(32)	1.8(0.7)	1.8(0.6)

SD and  $SD_{mad}$  are shown in parentheses.

## 4.2 Accuracy of profile-likelihood inferences

Table 5: Standard deviations and estimated standard errors

		<i>Poisson, values <math>\times 1000</math></i>				<i>Bernoulli, values <math>\times 10</math></i>			
		$\hat{\beta}_1$		$\hat{\beta}_3$		$\hat{\beta}_2$		$\hat{\beta}_4$	
$n$	$p_n$	SD	$SD_m$	SD	$SD_m$	SD	$SD_m$	SD	$SD_m$
200	10	9.1	8.5(1.3)	9.9	9.4(1.3)	3.6	2.9(.4)	3.2	2.8(.4)
400	13	6.0	5.6(0.7)	6.5	6.1(0.7)	2.3	2.1(.2)	2.2	2.0(.2)
800	16	3.7	3.8(0.3)	4.1	4.2(0.4)	1.7	1.6(.1)	1.5	1.5(.1)
1500	20	2.8	2.7(0.2)	3.1	3.0(0.2)	1.2	1.2(.1)	1.1	1.1(.1)

$SD_{mad}$  are shown in parentheses.

To test the accuracy of the sandwich formula for estimating standard errors, the standard deviations of the estimated coefficients (using the one-step accelerated profile-kernel estimate) are computed from the 400 simulations using  $h_{CV}$ . These can be regarded as the true standard errors (columns labeled SD). The 400 estimated standard errors are summarized by their median (columns  $SD_m$ ) and its associated  $SD_{mad}$ . Table 4 summarizes the results. Clearly, the sandwich formula does a good job, and accuracy gets better as  $n$  increases.

We now study the performance of GLRT in Section 2.2. To this end, we consider the following null hypothesis:

$$H_0 : \beta_7 = \beta_8 = \dots = \beta_{p_n} = 0.$$

We examine the power of the test under a sequence of the alternative hypotheses indexed by a parameter  $\gamma$  as follows:

$$H_1 : \beta_7 = \beta_8 = \gamma, \beta_j = 0 \text{ for } j > 8.$$

When  $\gamma = 0$ , the alternative hypothesis becomes the null hypothesis.






Figure 1: (a) Asymptotic null distribution (solid) and estimated true null distribution (dotted) for the Poisson model. (b) The power function at significant level  $\alpha = 0.01, 0.05$  and  $0.1$ . The captions for (c) and (d) are the same as those in (a) and (b) except that the Bernoulli model is now used.

Under the null hypothesis, the GLRT statistics are computed for each of 400 simulations, using the one-step accelerated profile-kernel estimates. Their distribution is summarized by a kernel density estimate and can be regarded as the true null distribution. This is compared with the asymptotic null distribution  $\chi^2_{p_n-6}$ . Figures 1(a) and (c) show the results when  $n = 400$ . The finite sample null density is seen to be reasonably close to the asymptotic one, except for the Monte Carlo error.

The power of the GLR test is studied under a sequence of alternative models, progressively deviating from the null hypothesis, namely, as  $\gamma$  increases. Again, the one-step accelerated profile-kernel algorithm is employed. The power functions are calculated at three significance levels: 0.1, 0.05 and 0.01, using the asymptotic distribution. They are the proportion of rejection among the 400 simulations and are depicted in Figures 1(b) and (d). The power curves increase rapidly with  $\gamma$ , which shows the GLR test is powerful. The powers at  $\gamma = 0$  are approximately the same as the significance level except the Monte Carlo error. This shows that the size of the test is reasonably accurate.

### 4.3 A real data example.

This is the analysis of the data in section 1.1 in where details of data and variables are given.

To examine the nonlinear effect of age and its nonlinear interaction with the experience, we appeal to the following GVCPLM (interactions between age and covariates other than **TotalYrsExp** are considered but found to be insignificant):

$$\log\left(\frac{p_H}{1-p_H}\right) = \alpha_1(\text{Age}) + \alpha_2(\text{Age})\text{TotalYrsExp} + \beta_1\text{Female} + \beta_2\text{PCJob} + \sum_{i=1}^4 \beta_{2+i}\text{Edu}_i \quad (4.1)$$

where  $p_H$  is the probability of having a high grade job. Formally, we are testing

$$H_0 : \beta_1 = 0 \longleftrightarrow H_1 : \beta_1 < 0. \quad (4.2)$$

Table 6: Fitted coefficients (sandwich SD) for model (4.1)

<i>Response</i>	Female	PCJob	Edu <sub>1</sub>	Edu <sub>2</sub>	Edu <sub>3</sub>	Edu <sub>4</sub>
HighGrade4	-1.96(.57)	-0.02(.76)	-5.14(.85)	-4.77(.98)	-2.72(.52)	-2.85(.96)
HighGrade5	-2.22(.59)	-1.96(.61)	-5.69(.67)	-5.95(.97)	-3.09(.72)	-1.26(1.10)

A 20-fold CV is employed to select the bandwidth  $h$  and the parameter  $\delta$  in the transformation of the data. This yields  $h_{CV} = 24.2$ ,  $\delta_{CV} = 0.1$ . Table 6 shows the results of the fit using the three-step accelerated profile-kernel estimate. The coefficient for **Female** is significantly negative. The education plays also an important role in getting high grade job. All coefficients are negative, as they are contrasted with the highest education level. The **PCJob** does not seem to play any significant role in getting promotion. Figures 2(a) and (b) depict the estimated coefficient functions. They show that as age increases one has a better chance of being in a higher job grade, and then the marginal effect of working experience is large when age is around 30 or less, but start to fall as one gets older. However, the second result should be interpreted with caution, as the variables **Age** and **TotalYrsExp** are highly correlated (Figure 2(c)). The standardized residuals  $(y - \hat{p}_H)/\sqrt{\hat{p}_H(1 - \hat{p}_H)}$  against **Age** is plotted in Figure 2(d). It shows that the fit seems reasonable. Other diagnostic plots also look reasonable, but they are not shown here.




Figure 2: (a) Fitted coefficient function  $\alpha_1(\cdot)$  (b) Fitted coefficient function  $\alpha_2(\cdot)$ . (c) The scatter plot ‘TotalYrsExp’ Against ‘Age’. (d) Standardized residuals against the variable ‘Age’.

We have conducted another fit using a binary variable **HighGrade5**, which is 0 only when job grade is less than 5. The coefficients are shown in Table 6 and the **Female** coefficient is close to the first fit.

We now employ the generalized likelihood ratio test to the problem (4.2). The GLR test statistic is 14.47 with one degree of freedom, resulting in a P-value of 0.0001. We have also conduct the same analysis using **HighGrade5** as the binary response. The GLR test statistic is now 13.76 and the associated P-value is 0.0002. The fitted coefficients are summarized in Table 5. The result provides stark evidence that even after adjusting for other confounding factors and variables, female employees of the Fifth National Bank is harder to get promoted to a high grade job.

Not shown in this paper, we have conducted the analysis again after deleting 6 data points corresponding to 5 male executives and 1 female employee having many years of working experience and high salaries. The test results are still similar.

## 5 Technical proofs

In this section the proofs of Theorems 1-4 will be given. We introduce some notations and regularity conditions for our results. In the following and thereafter, the symbol  $\otimes$  represents the Kronecker product between matrices, and  $\lambda_{\min}(A)$  and  $\lambda_{\max}(A)$  denotes respectively the minimum and maximum eigenvalues of a symmetric matrix  $A$ . We let  $Q_{ni}(\boldsymbol{\beta}_n)$  be the  $i$ -th summand of (2.3).

Denote the true linear parameter by  $\boldsymbol{\beta}_{n0}$ , with parameter space  $\Omega_n \subset \mathbb{R}^{p_n}$ . Let  $\mu_k = \int_{-\infty}^{\infty} u^k K(u) du$  and  $A_p(\mathbf{X}) = (\mu_{i+j})_{0 \leq i, j \leq p} \otimes \mathbf{X}\mathbf{X}^T$ . Set

$$\begin{aligned} \rho_l(t) &= (dg^{-1}(t)/dt)^l / V(g^{-1}(t)), & m_{ni}(\boldsymbol{\beta}_n) &= \boldsymbol{\alpha}_{\boldsymbol{\beta}_n}(U_i)^T \mathbf{X}_i + \boldsymbol{\beta}_n^T \mathbf{Z}_{ni}, \\ \boldsymbol{\alpha}'_{\boldsymbol{\beta}_n}(u) &= \frac{\partial \boldsymbol{\alpha}_{\boldsymbol{\beta}_n}(u)}{\partial \boldsymbol{\beta}_n}, & \boldsymbol{\alpha}_{\boldsymbol{\beta}_n}^{(r)''}(u) &= \frac{\partial^2 \boldsymbol{\alpha}_{\boldsymbol{\beta}_n}^{(r)}(u)}{\partial \boldsymbol{\beta}_n \partial \boldsymbol{\beta}_n^T}. \end{aligned}$$

### Regularity Conditions:

- (A) The covariates  $\mathbf{Z}_n$  and  $\mathbf{X}$  are bounded random variables.
- (B) The smallest and the largest eigenvalues of the matrix  $I_n(\boldsymbol{\beta}_{n0})$  is bounded away from zero and infinity for all  $n$ . In addition,  $\mathbf{E}_0[\nabla^T Q_{n1}(\boldsymbol{\beta}_{n0}) \nabla Q_{n1}(\boldsymbol{\beta}_{n0})]^4 = O(p_n^4)$ .
- (C)  $\mathbf{E}_{\boldsymbol{\beta}_n} \left| \frac{\partial^{l+j} Q_{n1}(\boldsymbol{\beta}_n)}{\partial^j \boldsymbol{\alpha} \partial \beta_{nk_1} \cdots \partial \beta_{nk_l}} \right|$  and  $\mathbf{E}_{\boldsymbol{\beta}_n} \left| \frac{\partial^{l+j} Q_{n1}(\boldsymbol{\beta}_n)}{\partial^j \boldsymbol{\alpha} \partial \beta_{nk_1} \cdots \partial \beta_{nk_l}} \right|^2$  are bounded for all  $n$ , with  $l = 1, \dots, 4$  and  $j = 0, 1$ .
- (D) The function  $q_2(x, y) < 0$  for  $x \in \mathbb{R}$  and  $y$  in the range of the response variable, and  $\mathbf{E}_0\{q_2(m_{n1}(\boldsymbol{\beta}_n), Y_{n1}) A_p(\mathbf{X}_1) | U = u\}$  is invertible.
- (E) The functions  $V''(\cdot)$  and  $g'''(\cdot)$  are continuous. The least-favorable curve  $\boldsymbol{\alpha}_{\boldsymbol{\beta}_n}(u)$  is three times continuously differentiable in  $\boldsymbol{\beta}_n$  and  $u$ .
- (F) The random variable  $U$  has a compact support  $\Omega$ . The density function  $f_U(u)$  of  $U$  has a continuous second derivative and is uniformly bounded away from zero.
- (G) The kernel  $K$  is a bounded symmetric density function with bounded support.

Note the above conditions are assumed to hold uniformly in  $u \in \Omega$ . Condition (A) is imposed just for the simplicity of proofs. The boundedness of covariates is imposed to ensure various products involving  $q_l(\cdot, \cdot)$ ,  $\mathbf{X}$  and  $\mathbf{Z}_n$  have bounded first and second moments. Conditions (B) and (C) are uniformity conditions on higher-order moments of

the likelihood functions. They are stronger than those of the usual asymptotic likelihood theory, but they facilitate technical proofs. Condition (G) is also imposed for simplicity of technical arguments. All of these conditions can be relaxed at the expense of longer proofs.

Before proving Theorem 1, we need two important lemmas. Lemma 1 concerns the order approximations to the least-favorable curve  $\alpha_{\beta_n}(\cdot)$ , while Lemma 2 holds the key to showing why undersmoothing is not needed in Theorems 1 and 2. Let  $c_n = (nh)^{-1/2}$ ,  $\hat{\alpha}_{0\beta_n}, \dots$ , and  $\hat{\alpha}_{p\beta_n}$  maximize (2.4), and  $\alpha_{u\beta_n}^{(p)}(u) = \frac{\partial^p \alpha_{\beta_n}(u)}{\partial u^p}$ . Set

$$\begin{aligned}\bar{\alpha}_{ni}(u) &= \mathbf{X}_i^T \left( \sum_{k=0}^p \frac{(U_i - u)^k}{k!} \alpha_{u\beta_n}^{(k)}(u) \right) + \beta_n^T \mathbf{Z}_{ni}, \\ \hat{\beta}^* &= c_n^{-1} \left( (\hat{\alpha}_{0\beta_n} - \alpha_{\beta_n}(u))^T, \dots, \frac{h^p}{p!} (\hat{\alpha}_{p\beta_n} - \alpha_{u\beta_n}^{(p)}(u))^T \right)^T, \\ \mathbf{X}_i^* &= \left( 1, \frac{U_i - u}{h}, \dots, \left( \frac{U_i - u}{h} \right)^p \right)^T \otimes \mathbf{X}_i.\end{aligned}$$

**Lemma 1** *Under Regularity Conditions (A) - (G), for each  $\beta_n \in \Omega_n$ , the following holds uniformly in  $u \in \Omega$ :*

$$\|\hat{\alpha}_{0\beta_n}(u) - \alpha_{\beta_n}(u)\| = O_P(h^{p+1} + c_n \log^{1/2}(1/h)).$$

*Likewise, the norm of the  $k^{\text{th}}$  derivative of the above with respect to any  $\beta_{nj}$ 's, for  $k = 1, \dots, 4$ , all have the same order uniformly in  $u \in \Omega$ .*

**Proof of Lemma 1.** Our first step is to show that, uniform in  $u \in \Omega$ ,

$$\hat{\beta}^* = \tilde{\mathbf{A}}_n^{-1} \mathbf{W}_n + O_P(h^{p+1} + c_n \log^{1/2}(1/h)),$$

where

$$\begin{aligned}\tilde{\mathbf{A}}_n &= f_U(u) E_0 \{ \rho_2(\alpha_{\beta_n}(U))^T \mathbf{X} + \mathbf{Z}_n^T \beta_n \} A_p(\mathbf{X}) | U = u\}, \\ \mathbf{W}_n &= hc_n \sum_{i=1}^n q_1(\bar{\alpha}_{ni}, Y_{ni}) \mathbf{X}_i^* K_h(U_i - u), \\ \mathbf{A}_n &= hc_n^2 \sum_{i=1}^n q_2(\bar{\alpha}_{ni}, Y_{ni}) \mathbf{X}_i^* \mathbf{X}_i^{*T} K_h(U_i - u).\end{aligned}$$

Since expression (2.4) is maximized at  $(\hat{\mathbf{a}}_{0\beta_n}, \dots, \hat{\mathbf{a}}_{p\beta_n})^T$ ,  $\hat{\boldsymbol{\beta}}^*$  maximizes

$$\begin{aligned} l_n(\boldsymbol{\beta}^*) &= h \sum_{i=1}^n \{Q(g^{-1}(c_n \mathbf{X}_i^{*T} \boldsymbol{\beta}^* + \bar{\boldsymbol{\alpha}}_{ni}), Y_{ni}) - Q(g^{-1}(\bar{\boldsymbol{\alpha}}_{ni}), Y_{ni})\} \\ &= \mathbf{W}_n^T \boldsymbol{\beta}^* + \frac{1}{2} \boldsymbol{\beta}^{*T} \mathbf{A}_n \boldsymbol{\beta}^* + \frac{hc_n^3}{6} \sum_{i=1}^n q_3(\eta_i, y_{ni}) (\mathbf{X}_i^{*T} \boldsymbol{\beta}^*)^3 K_h(U_i - u), \end{aligned}$$

where  $\eta_i$  lies between  $\bar{\boldsymbol{\alpha}}_{ni}$  and  $\bar{\boldsymbol{\alpha}}_{ni} + c_n \mathbf{X}_i^{*T} \boldsymbol{\beta}^*$ . The concavity of  $l_n(\boldsymbol{\beta}^*)$  is ensured by Condition (D). Note that  $K(\cdot)$  is bounded, we have under Conditions (A) and (C), the third term on the right hand side is bounded by

$$O_P(nhc_n^3 E|q_3(\eta_1, Y_{n1})| \|\mathbf{X}_1\|^3 K_h(U_1 - u)) = O_P(c_n) = o_P(1).$$

Direct calculation yields  $E_0 \mathbf{A}_n = -\tilde{\mathbf{A}}_n + O(h^{p+1})$  and  $\text{Var}_0((\mathbf{A}_n)_{ij}) = O((nh)^{-1})$  so that mean-variance decomposition yields

$$\mathbf{A}_n = -\tilde{\mathbf{A}}_n + O_P(h^{p+1}).$$

Hence we have

$$l_n(\boldsymbol{\beta}^*) = \mathbf{W}_n^T \boldsymbol{\beta}^* - \frac{1}{2} \boldsymbol{\beta}^{*T} \tilde{\mathbf{A}}_n \boldsymbol{\beta}^* + o_P(1). \quad (5.1)$$

Note that  $\mathbf{A}_n$  is a sum of i.i.d. random variables of kernel form, by a result of [21],

$$\mathbf{A}_n = -\tilde{\mathbf{A}}_n + O_P\{h^{p+1} + c_n \log^{1/2}(1/h)\} \quad (5.2)$$

uniformly in  $u \in \Omega$ . Hence by the Convexity Lemma ([24]), equation (5.1) also holds uniformly in  $\boldsymbol{\beta}^* \in C$  for any compact set  $C$ . Using Lemma A.1 of [5], it yields that

$$\sup_{u \in \Omega} |\hat{\boldsymbol{\beta}}^* - \tilde{\mathbf{A}}_n^{-1} \mathbf{W}_n| \xrightarrow{\mathbb{P}} 0. \quad (5.3)$$

Furthermore, by its definition,  $\hat{\boldsymbol{\beta}}^*$  solves the local likelihood equation:

$$\sum_{i=1}^n q_1(\bar{\boldsymbol{\alpha}}_{ni} + c_n \mathbf{X}_i^{*T} \hat{\boldsymbol{\beta}}^*, Y_{ni}) \mathbf{X}_i^* K_h(U_i - u) = 0. \quad (5.4)$$

Expanding  $q_1(\bar{\boldsymbol{\alpha}}_{ni} + c_n \mathbf{X}_i^{*T} \hat{\boldsymbol{\beta}}^*, \cdot)$  at  $\bar{\boldsymbol{\alpha}}_{ni}$  yields

$$\mathbf{W}_n + \mathbf{A}_n \hat{\boldsymbol{\beta}}^* + \frac{hc_n^3}{2} \sum_{i=1}^n q_3(\bar{\boldsymbol{\alpha}}_{ni} + \hat{\zeta}_i, Y_{ni}) \mathbf{X}_i^* (\mathbf{X}_i^{*T} \hat{\boldsymbol{\beta}}^*)^2 K_h(U_i - u) = 0 \quad (5.5)$$

where  $\hat{\zeta}_i$  lies between 0 and  $c_n \mathbf{X}_i^{*T} \hat{\boldsymbol{\beta}}^*$ . Using Conditions (A) and (C), the last term has order  $O_P(c_n^3 h n \|\hat{\boldsymbol{\beta}}^*\|^2) = O_P(c_n \|\hat{\boldsymbol{\beta}}^*\|^2)$ . With this, combining (5.2) and (5.5), we obtain

$$\hat{\boldsymbol{\beta}}^* = \tilde{\mathbf{A}}_n^{-1} \mathbf{W}_n + O_P(h^{p+1} + c_n \log^{1/2}(1/h)) \quad (5.6)$$

holds uniformly in  $u \in \Omega$  by (5.3). Using the result of [21] on  $\mathbf{W}_n$ , we obtain

$$\|\hat{\mathbf{a}}_{0\beta_n}(u) - \boldsymbol{\alpha}_{\beta_n}(u)\| = O_P(h^{p+1} + c_n \log^{1/2}(1/h)) \quad (5.7)$$

which holds uniformly in  $u \in \Omega$ .

Differentiate both sides of (5.4) w.r.t.  $\beta_{nj}$ ,

$$\sum_{i=1}^n q_2(\bar{\boldsymbol{\alpha}}_{ni} + c_n \mathbf{X}_i^{*T} \hat{\boldsymbol{\beta}}^*, Y_{ni}) \left\{ \frac{\partial \bar{\boldsymbol{\alpha}}_{ni}}{\partial \beta_{nj}} + c_n \left( \frac{\partial \hat{\boldsymbol{\beta}}^*}{\partial \beta_{nj}} \right)^T \mathbf{X}_i^* \right\} \mathbf{X}_i^* K_h(U_i - u) = 0, \quad (5.8)$$

which holds for all  $u \in \Omega$ . By Taylor's expansion and similar treatments to (5.5),

$$\mathbf{W}_n^1 + \mathbf{W}_n^2 + (\mathbf{A}_n + \mathbf{B}_n^1 + \mathbf{B}_n^2) \frac{\partial \hat{\boldsymbol{\beta}}^*}{\partial \beta_{nj}} + O_P(c_n \|\hat{\boldsymbol{\beta}}^*\|^2) = 0,$$

where

$$\begin{aligned} \mathbf{W}_n^1 &= h c_n \sum_{i=1}^n q_2(\bar{\boldsymbol{\alpha}}_{ni}, Y_{ni}) \frac{\partial \bar{\boldsymbol{\alpha}}_{ni}}{\partial \beta_{nj}} \mathbf{X}_i^* K_h(U_i - u), \\ \mathbf{W}_n^2 &= h c_n \sum_{i=1}^n q_3(\bar{\boldsymbol{\alpha}}_{ni}, Y_{ni}) c_n \mathbf{X}_i^{*T} \hat{\boldsymbol{\beta}}^* \frac{\partial \bar{\boldsymbol{\alpha}}_{ni}}{\partial \beta_{nj}} \mathbf{X}_i^* K_h(U_i - u), \\ \mathbf{B}_n^1 &= h c_n^2 \sum_{i=1}^n q_3(\bar{\boldsymbol{\alpha}}_{ni}, Y_{ni}) c_n \mathbf{X}_i^{*T} \hat{\boldsymbol{\beta}}^* \mathbf{X}_i^* \mathbf{X}_i^{*T} K_h(U_i - u), \\ \mathbf{B}_n^2 &= \frac{h c_n^2}{2} \sum_{i=1}^n q_4(\bar{\boldsymbol{\alpha}}_{ni} + \hat{\zeta}_i, Y_{ni}) (c_n^2 \mathbf{X}_i^{*T} \hat{\boldsymbol{\beta}}^*)^2 \mathbf{X}_i^* \mathbf{X}_i^{*T} K_h(U_i - u), \end{aligned}$$

with  $\hat{\zeta}_i$  lies between 0 and  $c_n \mathbf{X}_i^{*T} \hat{\boldsymbol{\beta}}^*$ . The above equations hold for all  $u \in \Omega$ . The order of  $\mathbf{W}_n^2$  is smaller than that of  $\mathbf{W}_n^1$ , and the orders of  $\mathbf{B}_n^1$  and  $\mathbf{B}_n^2$  are smaller than that of  $\mathbf{A}_n$ . Hence

$$\frac{\partial \hat{\boldsymbol{\beta}}^*}{\partial \beta_{nj}} = \tilde{\mathbf{A}}_n^{-1} \mathbf{W}_n^1 + O_P(\log^{1/2}(1/h) + c_n^{-1} h^{p+1})$$

uniformly in  $u \in \Omega$ . From this, for  $j = 1, \dots, p_n$ , we have

$$\left\| \frac{\partial \hat{\mathbf{a}}_{0\beta_n}(u)}{\partial \beta_{nj}} - \frac{\partial \boldsymbol{\alpha}_{\beta_n}(u)}{\partial \beta_{nj}} \right\| = O_P(h^{p+1} + c_n \log^{1/2}(1/h)) \quad (5.9)$$

uniformly in  $u \in \Omega$ . Differentiating (5.4) again w.r.t.  $\beta_{nk}$  and repeating as needed, we get the desired results for higher order derivatives by following similar arguments as above.

□

**Lemma 2** Under Regularity Conditions (A) - (G), if  $p_n^s/n \rightarrow 0$  for  $s > 5/4$ ,  $h = O(n^{-a})$  with  $(2s(p+1))^{-1} < a < 1 - s^{-1}$ , then for each  $\beta_n \in \Omega_n$ ,

$$n^{-1/2} \|\nabla \hat{Q}_n(\beta_n) - \nabla Q_n(\beta_n)\| = o_P(1).$$

**Proof of Lemma 2.** Define

$$\begin{aligned} \mathbf{K}_1 &= n^{-1/2} \sum_{i=1}^n q_2(m_{ni}(\beta_n), Y_{ni})(\mathbf{Z}_{ni} + \alpha'_{\beta_n}(U_i)\mathbf{X}_i)(\hat{\alpha}_{\beta_n}(U_i) - \alpha_{\beta_n}(U_i))^T \mathbf{X}_i, \\ \mathbf{K}_2 &= n^{-1/2} \sum_{i=1}^n q_1(m_{ni}(\beta_n), Y_{ni})(\hat{\alpha}'_{\beta_n}(U_i) - \alpha'_{\beta_n}(U_i))^T \mathbf{X}_i. \end{aligned}$$

Then by Taylor's expansion, Lemma 1 and Condition (C),

$$n^{-1/2}(\nabla \hat{Q}_n(\beta_n) - \nabla Q_n(\beta_n)) = \mathbf{K}_1 + \mathbf{K}_2 + \text{smaller order terms.}$$

Define, for  $\Omega$  as in Condition (F),

$$S = \{f \in C^2(\Omega) : \|f\|_\infty \leq 1\},$$

equipped with a metric  $\rho(f_1, f_2) = \|f_1 - f_2\|_\infty$ , where  $\|f\|_\infty = \sup_{u \in \Omega} |f(u)|$ . We also let, for  $r = 1, \dots, q$  and  $l = 1, \dots, p_n$ ,

$$\begin{aligned} A_{rl}(y, u, \mathbf{X}, \mathbf{Z}_n) &= q_2(\mathbf{X}^T \alpha_{\beta_n}(u) + \mathbf{Z}_n^T \beta_n, y) X_r \left( Z_{nl} + \mathbf{X}^T \frac{\partial \alpha_{\beta_n}(u)}{\partial \beta_{nl}} \right), \\ B_r(y, u, \mathbf{X}, \mathbf{Z}_n) &= q_1(\mathbf{X}^T \alpha_{\beta_n}(u) + \mathbf{Z}_n^T \beta_n, y) X_r. \end{aligned}$$

By Lemma 1, for any positive sequences  $(\delta_n)$  with  $\delta_n \rightarrow 0$  as  $n \rightarrow \infty$ , we have  $P_0(\lambda_r \in S) \rightarrow 1$  and  $P_0(\gamma_{rl} \in S) \rightarrow 1$ , where

$$\begin{aligned} \lambda_r &= \delta_n (h^{p+1} + c_n \log^{1/2}(1/h))^{-1} (\hat{\alpha}_{\beta_n}^{(r)} - \alpha_{\beta_n}^{(r)}), \\ \gamma_{rl} &= \delta_n (h^{p+1} + c_n \log^{1/2}(1/h))^{-1} \left( \frac{\partial \hat{\alpha}_{\beta_n}^{(r)}}{\partial \beta_{nl}} - \frac{\partial \alpha_{\beta_n}^{(r)}}{\partial \beta_{nl}} \right), \end{aligned}$$

$r = 1, \dots, q$  and  $l = 1, \dots, p_n$ . Hence for sufficiently large  $n$ , we have  $\lambda_r, \gamma_{rl} \in S$ . The following three points allow us to utilize [15] to prove our lemma.

- I. For any  $v \in S$ , we will view the map  $v \mapsto A_{rl}(y, u, \mathbf{X}, \mathbf{Z}_n)v(u)$  as an element of  $C(S)$ , the space of continuous functions on  $S$  equipped with the sup norm. For  $v_1, v_2 \in S$ , we have

$$\begin{aligned} &|A_{rl}(y, u, \mathbf{X}, \mathbf{Z}_n)v_1(u) - A_{rl}(y, u, \mathbf{X}, \mathbf{Z}_n)v_2(u)| \\ &= |A_{rl}(y, u, \mathbf{X}, \mathbf{Z}_n)(v_1 - v_2)(u)| \leq |A_{rl}(y, u, \mathbf{X}, \mathbf{Z}_n)| \|v_1 - v_2\|. \end{aligned}$$

Similar result holds for  $B_r(y, u, \mathbf{X}, \mathbf{Z}_n)$ .



II. Note that equation (2.2) is true for all  $\beta_n$ , and by differentiating w.r.t.  $\beta_n$  we get the following formulas:

$$\begin{aligned} E_0(q_1(m_n(\beta_n), Y_n) \mathbf{X} | U = u) &= \mathbf{0}, \\ E_0(q_2(m_n(\beta_n), Y_n) \mathbf{X} (\mathbf{Z}_n + \alpha'_{\beta_n}(U) \mathbf{X})^T | U = u) &= \mathbf{0}. \end{aligned}$$

Thus, we can easily see that

$$E_0(A_{rl}(Y, U, \mathbf{X}, \mathbf{Z}_n)) = 0$$

for each  $r = 1, \dots, q$  and  $l = 1, \dots, p_n$ . Also we have

$$E_0(A_{rl}(Y, U, \mathbf{X}, \mathbf{Z}_n)^2) < \infty,$$

by Regularity Conditions (A) and (C). For  $B_r(Y, U, \mathbf{X}, \mathbf{Z}_n)$ , results hold similarly.

III. Let  $H(\cdot, S)$  denote the metric entropy of the set  $S$  w.r.t. the metric  $\rho$ . Then

$$H(\epsilon, S) \leq C_0 \epsilon^{-1}$$

for some constant  $C_0$ . Hence  $\int_0^1 H^{1/2}(\epsilon, S) d\epsilon < \infty$ .

Conditions of Theorem 1 in [15] can be derived from the three notes above, so that we have

$$n^{-1/2} \sum_{i=1}^n A_{rl}(Y_i, U_i, \mathbf{X}_i, \mathbf{Z}_{ni})(\cdot),$$

where  $A_{rl}(Y_i, U_i, \mathbf{X}_i, \mathbf{Z}_{ni})(\cdot)$ ,  $i = 1, \dots, n$  being i.i.d. replicates of  $A_{rl}(Y, U, \mathbf{X}, \mathbf{Z}_n)(\cdot)$  in  $C(S)$ , converges weakly to a Gaussian measure on  $C(S)$ . Hence, since  $\lambda_r, \gamma_{rl} \in S$ ,

$$n^{-1/2} \sum_{i=1}^n A_{rl}(Y_i, U_i, \mathbf{X}_i, \mathbf{Z}_{ni})(\lambda_r) = O_P(1),$$

which implies that

$$n^{-1/2} \sum_{i=1}^n A_{rl}(Y_i, U_i, \mathbf{X}_i, \mathbf{Z}_{ni})(\hat{\alpha}_{\beta_n}^{(r)} - \alpha_{\beta_n}^{(r)}) = O_P(\delta_n^{-1}(h^{p+1} + c_n \log^{1/2}(1/h))).$$

Similarly, apply Theorem 1 of [15] again, we have

$$n^{-1/2} \sum_{i=1}^n B_r(Y_i, U_i, \mathbf{X}_i, \mathbf{Z}_{ni}) \left( \frac{\partial \hat{\alpha}_{\beta_n}^{(r)}}{\partial \beta_{nl}} - \frac{\partial \alpha_{\beta_n}^{(r)}}{\partial \beta_{nl}} \right) = O_P(\delta_n^{-1}(h^{p+1} + c_n \log^{1/2}(1/h))).$$

Then the column vector  $\mathbf{K}_1$  which is  $p_n$ -dimensional, has the  $l^{\text{th}}$  component equals

$$\sum_{r=1}^q \left\{ n^{-1/2} \sum_{i=1}^n A_{rl}(Y_i, U_i, \mathbf{X}_i, \mathbf{Z}_{ni})(\hat{\alpha}_{\beta_n}^{(r)} - \alpha_{\beta_n}^{(r)}) \right\} = O_P(\delta_n^{-1}(h^{p+1} + c_n \log^{1/2}(1/h)),$$

using the result just proved. Hence we have shown

$$\|\mathbf{K}_1\| = O_P(\sqrt{p_n} \delta_n^{-1}(h^{p+1} + c_n \log^{1/2}(1/h))) = o_P(1),$$

since  $\delta_n$  can be made arbitrarily slow in converging to 0. Similarly, we have  $\|\mathbf{K}_2\| = o_P(1)$  as well. The conclusion of the lemma follows.  $\square$

### Proof of Theorem 1.

Let  $\gamma_n = \sqrt{p_n/n}$ . Our aim is to show that, for a given  $\epsilon > 0$ ,

$$\mathbb{P} \left\{ \sup_{\|\mathbf{v}\|=C} \hat{Q}_n(\beta_{n0} + \gamma_n \mathbf{v}) < \hat{Q}_n(\beta_{n0}) \right\} \geq 1 - \epsilon, \quad (5.10)$$

so that this implies with probability tending to 1 there is a local maximum  $\hat{\beta}_n$  in the ball  $\{\beta_{n0} + \gamma_n \mathbf{v} : \|\mathbf{v}\| \leq C\}$  such that  $\|\hat{\beta}_n - \beta_{n0}\| = O_P(\gamma_n)$ .

Define the terms  $\hat{I}_1 = \gamma_n \nabla^T \hat{Q}_n(\beta_{n0}) \mathbf{v}$ ,  $\hat{I}_2 = \frac{\gamma_n^2}{2} \mathbf{v}^T \nabla^2 \hat{Q}_n(\beta_{n0}) \mathbf{v}$  and  $\hat{I}_3 = \frac{\gamma_n^3}{6} \nabla^T (\mathbf{v}^T \nabla^2 \hat{Q}_n(\beta_n^*) \mathbf{v}) \mathbf{v}$ . By Taylor's expansion,

$$\hat{Q}_n(\beta_{n0} + \gamma_n \mathbf{v}) - \hat{Q}_n(\beta_{n0}) = \hat{I}_1 + \hat{I}_2 + \hat{I}_3,$$

where  $\beta_n^*$  lies between  $\beta_{n0}$  and  $\beta_{n0} + \gamma_n \mathbf{v}$ .

We further split  $\hat{I}_1 = D_1 + D_2$ , where

$$\begin{aligned} D_1 &= \sum_{i=1}^n q_1(\hat{m}_{ni}(\beta_{n0}), Y_{ni})(\mathbf{Z}_{ni} + \alpha'_{\beta_{n0}}(U_i) \mathbf{X}_i)^T \mathbf{v} \gamma_n, \\ D_2 &= \sum_{i=1}^n q_1(\hat{m}_{ni}(\beta_{n0}), Y_{ni}) \mathbf{X}_i^T (\hat{\alpha}'_{\beta_{n0}}(U_i) - \alpha'_{\beta_{n0}}(U_i))^T \mathbf{v} \gamma_n, \end{aligned}$$

with  $\hat{m}_{ni}(\beta_n) = \hat{\alpha}_{\beta_n}(U_i)^T \mathbf{X}_i + \beta_n^T \mathbf{Z}_{ni}$ . By Condition (A) and Lemma 1,  $D_2$  has order smaller than  $D_1$ . Using Taylor's expansion, we have

$$D_1 = \gamma_n \mathbf{v}^T \left( \sum_{i=1}^n \frac{\partial Q_{ni}(\beta_{n0})}{\partial \beta_n} + \sqrt{n} \mathbf{K}_1 \right) + \text{smaller order terms},$$

where  $\mathbf{K}_1$  is as defined in Lemma 2 so that within the lemma's proof we have  $\|\mathbf{K}_1\| = o_P(1)$ . Using equation (2.6), we have by the mean-variance decomposition

$$\left\| \mathbf{v}^T \sum_{i=1}^n \frac{\partial Q_{ni}(\beta_{n0})}{\partial \beta_n} \right\| = O_P(\sqrt{n \mathbf{v}^T I_n(\beta_{n0}) \mathbf{v}}) = O_P(\sqrt{n}) \|\mathbf{v}\|,$$

where last inequality follows from Condition (B). Hence

$$|\hat{I}_1| = O_P(\sqrt{n}\gamma_n)\|\mathbf{v}\|.$$

Next, consider  $\hat{I}_2 = I_2 + (\hat{I}_2 - I_2)$ , where

$$\begin{aligned} I_2 &= \frac{1}{2}\mathbf{v}^T \nabla^2 Q_n(\boldsymbol{\beta}_{n0}) \mathbf{v} \gamma_n^2 \\ &= -\frac{n}{2}\mathbf{v}^T I_n(\boldsymbol{\beta}_{n0}) \mathbf{v} \gamma_n^2 + \frac{n}{2}\mathbf{v}^T \{n^{-1} \nabla^2 Q_n(\boldsymbol{\beta}_{n0}) + I_n(\boldsymbol{\beta}_{n0})\} \mathbf{v} \gamma_n^2 \\ &= -\frac{n}{2}\mathbf{v}^T I_n(\boldsymbol{\beta}_{n0}) \mathbf{v} \gamma_n^2 + o_P(n\gamma_n^2)\|\mathbf{v}\|^2 \end{aligned}$$

with the last line follows from Lemma 5 in the Appendix. Using Lemma 4,

$$\|\hat{I}_2 - I_2\| = o_P(n\gamma_n^2\|\mathbf{v}\|^2).$$

On the other hand, by Condition (B), we have

$$|n\gamma_n^2 \mathbf{v}^T I_n(\boldsymbol{\beta}_{n0}) \mathbf{v}| \geq O(n\gamma_n^2 \lambda_{\min}(I_n(\boldsymbol{\beta}_{n0}))\|\mathbf{v}\|^2) = O(n\gamma_n^2\|\mathbf{v}\|^2).$$

Hence,  $\hat{I}_2 - I_2$  has a smaller order than  $I_2$ .

Finally consider  $\hat{I}_3$ . We suppress the dependence of  $\boldsymbol{\alpha}_{\boldsymbol{\beta}_n}(U_i)$  and its derivatives on  $U_i$ , and denote  $q_{1i} = q_1(m_{ni}(\boldsymbol{\beta}_{n0}), Y_{ni})$ . Using Taylor's expansions, expanding  $\hat{Q}_n(\boldsymbol{\beta}_n^*)$  at  $\boldsymbol{\beta}_{n0}$  and then  $\hat{Q}_n(\boldsymbol{\beta}_{n0})$  at  $\boldsymbol{\alpha}_{\boldsymbol{\beta}_{n0}}$ , we can arrive at

$$\begin{aligned} \hat{Q}_n(\boldsymbol{\beta}_n^*) &= Q_n(\boldsymbol{\beta}_{n0}) + \sum_{i=1}^n \{q_{1i} \mathbf{X}_i^T (\hat{\boldsymbol{\alpha}}_{\boldsymbol{\beta}_{n0}} - \boldsymbol{\alpha}_{\boldsymbol{\beta}_{n0}}) \\ &\quad + q_{1i} (\mathbf{Z}_{ni} + \hat{\boldsymbol{\alpha}}'_{\boldsymbol{\beta}_{n0}} \mathbf{X}_i)^T (\boldsymbol{\beta}_n^* - \boldsymbol{\beta}_{n0})\} (1 + o_P(1)). \end{aligned}$$

Substituting  $\hat{Q}_n(\boldsymbol{\beta}_n^*)$  into  $\hat{I}_3$  with the right hand side above, by Condition (C) and Lemma 1, we have

$$\hat{I}_3 = \frac{1}{6} \sum_{i,j,k=1}^{p_n} \frac{\partial^3 Q_n(\boldsymbol{\beta}_{n0})}{\partial \boldsymbol{\beta}_{ni} \partial \boldsymbol{\beta}_{nj} \partial \boldsymbol{\beta}_{nk}} v_i v_j v_k \gamma_n^3 + \text{smaller order terms.}$$

Hence,

$$|\hat{I}_3| = O_P(np_n^{3/2} \gamma_n^3 \|\mathbf{v}\|^3) = O_P(\sqrt{p_n^4/n} \|\mathbf{v}\|) n\gamma_n^2 \|\mathbf{v}\|^2 = o_P(1) n\gamma_n^2 \|\mathbf{v}\|^2.$$

Comparing, we find the order of  $-n\gamma_n^2 \mathbf{v}^T I_n(\boldsymbol{\beta}_{n0}) \mathbf{v}$  dominates all other terms by allowing  $\|\mathbf{v}\| = C$  to be large enough. This proves (5.10).  $\square$

**Proof of Theorem 2.**

Note that by Theorem 1,  $\|\hat{\beta}_n - \beta_{n0}\| = O_P(\sqrt{p_n/n})$ . Since  $\nabla \hat{Q}_n(\hat{\beta}_n) = 0$ , by Taylor's expansion,

$$\nabla \hat{Q}_n(\beta_{n0}) + \nabla^2 \hat{Q}_n(\beta_{n0})(\hat{\beta}_n - \beta_{n0}) + \mathcal{C} = 0, \quad (5.11)$$

where  $\beta_n^*$  lies between  $\beta_{n0}$  and  $\hat{\beta}_n$  and  $\mathcal{C} = \frac{1}{2}(\hat{\beta}_n - \beta_{n0})^T \nabla^2(\nabla \hat{Q}_n(\beta_n^*))(\hat{\beta}_n - \beta_{n0})$  which is understood as a vector of quadratic components.

Using similar argument to approximating  $\hat{I}_3$  in Theorem 1, by Lemma 1 and noting  $\|\beta_n^* - \beta_{n0}\| = o_P(1)$ , we have  $\|\nabla^2 \frac{\partial \hat{Q}_n(\beta_n^*)}{\partial \beta_{nj}}\|^2 = O_P(n^2 p_n^2)$ . Hence

$$\|n^{-1} \mathcal{C}\|^2 \leq n^{-2} \|\hat{\beta}_n - \beta_{n0}\|^4 \sum_{j=1}^{p_n} \left\| \nabla^2 \frac{\partial \hat{Q}_n(\beta_n^*)}{\partial \beta_{nj}} \right\|^2 = O_P(p_n^5/n^2) = o_P(n^{-1}). \quad (5.12)$$

At the same time, by Lemma 5 and the Cauchy-Schwarz inequality,

$$\begin{aligned} & \|n^{-1} \nabla^2 \hat{Q}_n(\beta_{n0})(\hat{\beta}_n - \beta_{n0}) + I_n(\beta_{n0})(\hat{\beta}_n - \beta_{n0})\| \\ &= o_P((np_n)^{-1/2}) + O_P(\sqrt{p_n^3/n}(h^{p+1} + c_n \log^{1/2}(1/h))) = o_P(n^{-1/2}). \end{aligned} \quad (5.13)$$

Combining (5.11), (5.12) and (5.13), we have

$$\begin{aligned} I_n(\beta_{n0})(\hat{\beta}_n - \beta_{n0}) &= n^{-1} \nabla \hat{Q}_n(\beta_{n0}) + o_P(n^{-1/2}) \\ &= n^{-1} \nabla Q_n(\beta_{n0}) + o_P(n^{-1/2}), \end{aligned} \quad (5.14)$$

where the last line follows from Lemma 2. Consequently, using equation (5.14), we get

$$\begin{aligned} \sqrt{n} A_n I_n^{-1/2}(\beta_{n0})(\hat{\beta}_n - \beta_{n0}) &= n^{-1/2} A_n I_n^{-1/2}(\beta_{n0}) \nabla Q_n(\beta_{n0}) \\ &\quad + o_P(A_n I_n^{-1/2}(\beta_{n0})) \\ &= n^{-1/2} A_n I_n^{-1/2}(\beta_{n0}) \nabla Q_n(\beta_{n0}) + o_P(1), \end{aligned} \quad (5.15)$$

since  $\|A_n I_n^{-1/2}(\beta_{n0})\| = O(1)$  by conditions of Theorem 2.

We now check the Lindeberg-Feller Central Limit Theorem (see for example, [29]) for the last term in (5.15). Let  $B_{ni} = n^{-1/2} A_n I_n^{-1/2}(\beta_{n0}) \nabla Q_{ni}(\beta_{n0})$ ,  $i = 1, \dots, n$ . Given  $\epsilon > 0$ ,

$$\sum_{i=1}^n E_0 \|B_{ni}\|^2 1\{\|B_{ni}\| > \epsilon\} \leq n \sqrt{E_0 \|B_{n1}\|^4 \cdot \mathbb{P}(\|B_{n1}\| > \epsilon)}.$$

Using Chebyshev's inequality,

$$\begin{aligned} \mathbb{P}(\|B_{n1}\| > \epsilon) &\leq n^{-1} \epsilon^{-2} E \|A_n I_n^{-1/2}(\beta_{n0}) \nabla Q_{n1}(\beta_{n0})\|^2 \\ &= n^{-1} \epsilon^{-2} \text{tr}(G) = O(n^{-1}), \end{aligned} \quad (5.16)$$

where  $\text{tr}(A)$  is the trace of square matrix  $A$ . Similarly, we can show that, using Condition (B),

$$\begin{aligned} E_0 \|B_{n1}\|^4 &\leq \sqrt{l} n^{-2} \lambda_{\min}^2(A_n A_n^T) \lambda_{\max}^2(I_n(\beta_{n0})) \sqrt{E_0 (\nabla Q_{n1}(\beta_{n0})^T \nabla Q_{n1}(\beta_{n0}))^4} \\ &= O(p_n^2/n^2). \end{aligned} \quad (5.17)$$

Therefore (5.16) and (5.17) together imply

$$\sum_{i=1}^n E_0 \|B_{ni}\|^2 1\{\|B_{ni}\| > \epsilon\} = O(\sqrt{p_n^2/n}) = o(1).$$

Also,

$$\begin{aligned} \sum_{i=1}^n \text{Var}_0(B_{ni}) &= \text{Var}_0(A_n I_n^{-1/2}(\beta_{n0}) \nabla Q_{n1}(\beta_{n0})) \\ &= A_n A_n^T \rightarrow G. \end{aligned}$$

Therefore  $B_{ni}$  satisfies the conditions of the Lindeberg-Feller Central Limit Theorem. Consequently, using (5.15), it follows that

$$\sqrt{n} A_n I_n^{1/2}(\beta_{n0}) (\hat{\beta}_n - \beta_{n0}) \xrightarrow{D} N(0, G),$$

and this completes the proof.  $\square$

Referring back to Section 2.2, let  $B_n$  be a  $(p_n - l) \times p_n$  matrix satisfying  $B_n B_n^T = I_{p_n - l}$  and  $A_n B_n^T = 0$ . Since  $A_n \beta_n = 0$  under  $H_0$ , rows of  $A_n$  are perpendicular to  $\beta_n$  and the orthogonal complement of rows of  $A_n$  is spanned by rows of  $B_n$  since  $A_n B_n^T = 0$ . Hence

$$\beta_n = B_n^T \gamma$$

under  $H_0$ , where  $\gamma$  is a  $(p_n - l) \times 1$  vector. Then under  $H_0$  the profile likelihood estimator is also the local maximizer  $\hat{\gamma}_n$  of the problem

$$\hat{Q}_n(B_n^T \hat{\gamma}_n) = \max_{\gamma_n} Q_n(B_n^T \gamma_n).$$

### Proof of Theorem 3.

By Taylor's expansion, expanding  $\hat{Q}_n(B_n^T \hat{\gamma}_n)$  at  $\hat{\beta}_n$  and noting that  $\nabla^T \hat{Q}_n(\hat{\beta}_n) = 0$ , then  $\hat{Q}_n(\hat{\beta}_n) - \hat{Q}_n(B_n^T \hat{\gamma}_n) = T_1 + T_2$ , where

$$\begin{aligned} T_1 &= -\frac{1}{2} (\hat{\beta}_n - B_n^T \hat{\gamma}_n)^T \nabla^2 \hat{Q}_n(\hat{\beta}_n) (\hat{\beta}_n - B_n^T \hat{\gamma}_n), \\ T_2 &= \frac{1}{6} \nabla^T \{ (\hat{\beta}_n - B_n^T \hat{\gamma}_n)^T \nabla^2 \hat{Q}_n(\beta_n^*) (\hat{\beta}_n - B_n^T \hat{\gamma}_n) \} (\hat{\beta}_n - B_n^T \hat{\gamma}_n). \end{aligned}$$

Denote by  $\Theta_n = I_n(\beta_{n0})$  and  $\Phi_n = \frac{1}{n}\nabla Q_n(\beta_{n0})$ . Using equation (5.14) and noting that  $\Theta_n$  has eigenvalues uniformly bounded away from 0 and infinity by Condition (B), we have

$$\hat{\beta}_n - \beta_{n0} = \Theta_n^{-1}\Phi_n + o_P(n^{-1/2}).$$

Combining this with Lemma 6 in the Appendix, under the null hypothesis  $H_0$ ,

$$\begin{aligned} \hat{\beta}_n - B_n^T \hat{\gamma}_n &= \Theta_n^{-1/2} \{I_{p_n} - \Theta_n^{1/2} B_n^T (B_n \Theta_n B_n^T)^{-1} B_n \Theta_n^{1/2}\} \Theta_n^{-1/2} \Phi_n \\ &\quad + o_P(n^{-1/2}). \end{aligned} \quad (5.18)$$

Since  $S_n = I_{p_n} - \Theta_n^{1/2} B_n^T (B_n \Theta_n B_n^T)^{-1} B_n \Theta_n^{1/2}$  is a  $p_n \times p_n$  idempotent matrix with rank  $l$ , it follows by mean-variance decomposition of the term  $\|\hat{\beta}_n - B_n^T \hat{\gamma}_n\|^2$  and Condition (B) that

$$\|\hat{\beta}_n - B_n^T \hat{\gamma}_n\| = O_P(n^{-1/2}).$$

Hence, using similar argument as in the approximation of order for  $|\hat{I}_3|$  in Theorem 1, we have

$$|T_2| = O_P(np_n^{3/2}) \cdot \|\hat{\beta}_n - B_n^T \hat{\gamma}_n\|^3 = o_P(1).$$

Hence  $\hat{Q}_n(\hat{\beta}_n) - \hat{Q}_n(B_n^T \hat{\gamma}_n) = T_2 + o_P(1)$ .

By Lemma 5 and an approximation to  $n^{-1}\|\nabla^2 \hat{Q}_n(\hat{\beta}_n) - \nabla^2 \hat{Q}_n(\beta_{n0})\| = o_P(p_n^{-1/2})$  (the proof is similar to that for Lemma 3 with the proof of order for  $|\hat{I}_3|$  in Theorem 1, and is omitted), we have

$$\begin{aligned} &\left\| \frac{1}{2}(\hat{\beta}_n - B_n^T \hat{\gamma}_n)^T \{\nabla^2 \hat{Q}_n(\hat{\beta}_n) + nI_n(\beta_{n0})\}(\hat{\beta}_n - B_n^T \hat{\gamma}_n) \right\| \\ &= O_P(l/n) \cdot n\{o_P(p_n^{-1/2}) + O_P(p_n(h^{p+1} + c_n \log^{1/2}(1/h)))\} = o_P(1). \end{aligned}$$

Therefore,

$$\hat{Q}_n(\hat{\beta}_n) - \hat{Q}_n(B_n^T \hat{\gamma}_n) = \frac{n}{2}(\hat{\beta}_n - B_n^T \hat{\gamma}_n)^T I_n(\beta_{n0})(\hat{\beta}_n - B_n^T \hat{\gamma}_n) + o_P(1).$$

By (5.18), we have

$$\hat{Q}_n(\hat{\beta}_n) - \hat{Q}_n(B_n^T \hat{\gamma}_n) = \frac{n}{2}\Phi_n^T \Theta_n^{-1/2} S_n \Theta_n^{-1/2} \Phi_n + o_P(1).$$

Since  $S_n$  is idempotent, it can be written as  $S_n = D_n^T D_n$  where  $D_n$  is an  $l \times p_n$  matrix satisfying  $D_n D_n^T = I_l$ . By Theorem 2, we have already shown that  $\sqrt{n} D_n \Theta_n^{-1/2} \Phi_n \xrightarrow{D} N(\mathbf{0}, I_l)$ . Hence

$$2\{\hat{Q}_n(\hat{\beta}_n) - \hat{Q}_n(B_n^T \hat{\gamma}_n)\} = n(D_n \Theta_n^{-1/2} \Phi_n)^T (D_n \Theta_n^{-1/2} \Phi_n) \xrightarrow{D} \chi_l^2. \quad \square$$

**Proof of Theorem 4.**

Let  $\hat{\mathcal{A}}_n = -n^{-1}\nabla^2\hat{Q}_n(\hat{\beta}_n)$ ,  $\hat{\mathcal{B}}_n = \widehat{\text{cov}}\{\nabla\hat{Q}_n(\hat{\beta}_n)\}$  and  $\mathcal{C} = I_n(\beta_{n0})$ . Write

$$I_1 = \hat{\mathcal{A}}_n^{-1}(\hat{\mathcal{B}}_n - \mathcal{C})\hat{\mathcal{A}}_n^{-1}, \quad I_2 = \hat{\mathcal{A}}_n^{-1}(\mathcal{C} - \hat{\mathcal{A}}_n)\hat{\mathcal{A}}_n^{-1}, \quad I_3 = \hat{\mathcal{A}}_n^{-1}(\mathcal{C} - \hat{\mathcal{A}}_n)\mathcal{C}^{-1}.$$

Then,  $\hat{\Sigma}_n - \Sigma_n = I_1 + I_2 + I_3$ . Our aim is to show that, for all  $i = 1, \dots, p_n$ ,

$$\lambda_i(\hat{\Sigma}_n - \Sigma_n) = o_P(1),$$

so that  $A_n(\hat{\Sigma}_n - \Sigma_n)A_n^T \xrightarrow{\mathbb{P}} 0$ , where  $\lambda_i(A)$  is the  $i$ th eigenvalue of a symmetric matrix  $A$ . Using the inequalities

$$\begin{aligned} \lambda_{\min}(I_1) + \lambda_{\min}(I_2) + \lambda_{\min}(I_3) &\leq \lambda_{\min}(I_1 + I_2 + I_3) \\ \lambda_{\max}(I_1 + I_2 + I_3) &\leq \lambda_{\max}(I_1) + \lambda_{\max}(I_2) + \lambda_{\max}(I_3), \end{aligned}$$

it suffices to show that  $\lambda_i(I_j) = o_P(1)$  for  $j = 1, 2, 3$ . From the definition of  $I_1, I_2$  and  $I_3$ , it is clear that we only need to show  $\lambda_i(\mathcal{C} - \hat{\mathcal{A}}_n) = o_P(1)$  and  $\lambda_i(\hat{\mathcal{B}}_n - \mathcal{C}) = o_P(1)$ . Let  $K_1 = I_n(\beta_{n0}) + n^{-1}\nabla^2 Q_n(\beta_{n0})$ ,  $K_2 = n^{-1}(\nabla^2 Q_n(\hat{\beta}_n) - \nabla^2 Q_n(\beta_{n0}))$ , and  $K_3 = n^{-1}(\nabla^2 \hat{Q}_n(\hat{\beta}_n) - \nabla^2 Q_n(\hat{\beta}_n))$ . Then,

$$\mathcal{C} - \hat{\mathcal{A}}_n = K_1 + K_2 + K_3.$$

Applying Lemma 5 to  $K_1$ , Lemma 3 to  $K_2$ , and Lemma 4 to  $K_3$ , we have  $\|\mathcal{C} - \hat{\mathcal{A}}\| = o_P(1)$ . Thus,  $\lambda_i(\mathcal{C} - \hat{\mathcal{A}}) = o_P(1)$ . Hence the only thing left to show is  $\lambda_i(\hat{\mathcal{B}}_n - \mathcal{C}) = o_P(1)$ .

To this end, consider the decomposition

$$\hat{\mathcal{B}}_n - \mathcal{C} = K_4 + K_5$$

where

$$\begin{aligned} K_4 &= \left\{ \frac{1}{n} \sum_{i=1}^n \frac{\partial \hat{Q}_{ni}(\hat{\beta}_n)}{\partial \beta_{nj}} \frac{\partial \hat{Q}_{ni}(\hat{\beta}_n)}{\partial \beta_{nk}} \right\} - I_n(\beta_{n0}), \\ K_5 &= -\left\{ \frac{1}{n} \sum_{i=1}^n \frac{\partial \hat{Q}_{ni}(\hat{\beta}_n)}{\partial \beta_{nj}} \right\} \left\{ \frac{1}{n} \sum_{i=1}^n \frac{\partial \hat{Q}_{ni}(\hat{\beta}_n)}{\partial \beta_{nk}} \right\}. \end{aligned}$$

Our goal is to show that  $K_4$  and  $K_5$  are  $o_P(1)$ , which then implies  $\lambda_i(\hat{\mathcal{B}}_n - \mathcal{C}) = o_P(1)$ .

We consider  $K_4$  first, which can be further decomposed into  $K_4 = K_6 + K_7$ , where

$$\begin{aligned} K_6 &= \left\{ \frac{1}{n} \sum_{i=1}^n \frac{\partial \hat{Q}_{ni}(\hat{\beta}_n)}{\partial \beta_{nj}} \frac{\partial \hat{Q}_{ni}(\hat{\beta}_n)}{\partial \beta_{nk}} - \frac{1}{n} \sum_{i=1}^n \frac{\partial Q_{ni}(\beta_{n0})}{\partial \beta_{nj}} \frac{\partial Q_{ni}(\beta_{n0})}{\partial \beta_{nk}} \right\}, \\ K_7 &= \left\{ \frac{1}{n} \sum_{i=1}^n \frac{\partial Q_{ni}(\beta_{n0})}{\partial \beta_{nj}} \frac{\partial Q_{ni}(\beta_{n0})}{\partial \beta_{nk}} \right\} - I_n(\beta_{n0}). \end{aligned}$$

Observe that

$$\begin{aligned}
K_6 = & \left\{ \frac{1}{n} \sum_{i=1}^n \frac{\partial Q_{ni}(\boldsymbol{\beta}_{n0})}{\partial \beta_{nj}} \left\{ \frac{\partial \hat{Q}_{ni}(\hat{\boldsymbol{\beta}}_n)}{\partial \beta_{nk}} - \frac{\partial Q_{ni}(\boldsymbol{\beta}_{n0})}{\partial \beta_{nk}} \right\} \right. \\
& + \frac{1}{n} \sum_{i=1}^n \frac{\partial Q_{ni}(\boldsymbol{\beta}_{n0})}{\partial \beta_{nk}} \left\{ \frac{\partial \hat{Q}_{ni}(\hat{\boldsymbol{\beta}}_n)}{\partial \beta_{nj}} - \frac{\partial Q_{ni}(\boldsymbol{\beta}_{n0})}{\partial \beta_{nj}} \right\} \\
& \left. + \frac{1}{n} \sum_{i=1}^n \left\{ \frac{\partial \hat{Q}_{ni}(\hat{\boldsymbol{\beta}}_n)}{\partial \beta_{nk}} - \frac{\partial Q_{ni}(\boldsymbol{\beta}_{n0})}{\partial \beta_{nk}} \right\} \left\{ \frac{\partial \hat{Q}_{ni}(\hat{\boldsymbol{\beta}}_n)}{\partial \beta_{nj}} - \frac{\partial Q_{ni}(\boldsymbol{\beta}_{n0})}{\partial \beta_{nj}} \right\} \right\},
\end{aligned}$$

and this suggests that an approximation of the order of  $\frac{\partial}{\partial \beta_{nk}}(\hat{Q}_{ni}(\hat{\boldsymbol{\beta}}_n) - Q_{ni}(\boldsymbol{\beta}_{n0}))$  for each  $k = 1, \dots, p_n$  and  $i = 1, \dots, n$  is rewarding. Define

$$a_{ik} = \frac{\partial}{\partial \beta_{nk}}(\hat{Q}_{ni}(\hat{\boldsymbol{\beta}}_n) - Q_{ni}(\hat{\boldsymbol{\beta}}_n)), \quad \text{and} \quad b_{ik} = \frac{\partial}{\partial \beta_{nk}}(Q_{ni}(\hat{\boldsymbol{\beta}}_n) - Q_{ni}(\boldsymbol{\beta}_{n0})),$$

then  $\frac{\partial}{\partial \beta_{nk}}(\hat{Q}_{ni}(\hat{\boldsymbol{\beta}}_n) - Q_{ni}(\boldsymbol{\beta}_{n0})) = a_{ik} + b_{ik}$ . By Taylor's expansion, suppressing dependence of  $\boldsymbol{\alpha}_{\boldsymbol{\beta}_n}(U_i)$  and its derivatives on  $U_i$ ,

$$a_{ik} = \left\{ \frac{\partial^2 Q_{ni}(\hat{\boldsymbol{\beta}}_n)}{\partial \beta_{nk} \partial \boldsymbol{\alpha}_{\boldsymbol{\beta}_n}^T} (\hat{\boldsymbol{\alpha}}_{\hat{\boldsymbol{\beta}}_n} - \boldsymbol{\alpha}_{\hat{\boldsymbol{\beta}}_n}) + \frac{\partial Q_{ni}(\hat{\boldsymbol{\beta}}_n)}{\partial \boldsymbol{\alpha}_{\boldsymbol{\beta}_n}^T} \left( \frac{\partial \hat{\boldsymbol{\alpha}}_{\hat{\boldsymbol{\beta}}_n}}{\partial \beta_{nk}} - \frac{\partial \boldsymbol{\alpha}_{\hat{\boldsymbol{\beta}}_n}}{\partial \beta_{nk}} \right) \right\} (1 + o_P(1)).$$

Using Lemma 1, Condition (C), with argument similar to the proof of Lemma 4, we then have

$$a_{ik} = O_P(h^{p+1} + c_n \log^{1/2}(1/h)).$$

Similarly, Taylor's expansion gives

$$b_{ik} = \frac{\partial^2 Q_{ni}(\boldsymbol{\beta}_{n0})}{\partial \beta_{nk} \partial \boldsymbol{\beta}_n^T} (\hat{\boldsymbol{\beta}}_n - \boldsymbol{\beta}_{n0}) (1 + o_P(1)),$$

which implies that, by Theorem 1 and Regularity Condition (C),

$$|b_{ik}| = O_P(\sqrt{p_n^2/n}).$$

Using the approximations of  $a_{ik}$  and  $b_{ik}$  above, by Condition (C),

$$\begin{aligned}
& \left| \frac{1}{n} \sum_{i=1}^n \frac{\partial Q_{ni}(\boldsymbol{\beta}_{n0})}{\partial \beta_{nj}} \left\{ \frac{\partial \hat{Q}_{ni}(\hat{\boldsymbol{\beta}}_n)}{\partial \beta_{nk}} - \frac{\partial Q_{ni}(\boldsymbol{\beta}_{n0})}{\partial \beta_{nk}} \right\} \right| \\
& \leq \frac{1}{n} \sum_{i=1}^n \left| \frac{\partial Q_{ni}(\boldsymbol{\beta}_{n0})}{\partial \beta_{nj}} \right| \cdot |a_{ik} + b_{ik}| \\
& = O_P(h^{p+1} + c_n \log^{1/2}(1/h) + n^{-1/2} p_n).
\end{aligned}$$



This shows that

$$\|K_6\| = O_P(p_n(h^{p+1} + c_n \log^{1/2}(1/h)) + p_n^2 n^{-1/2}) = o_P(1)$$

by the conditions of the theorem.

For  $K_7$ , note that

$$\begin{aligned} E_0 K_7 &= n^{-2} (np_n^2) E_0 \left\{ \frac{\partial Q_{ni}(\beta_{n0})}{\partial \beta_{nj}} \frac{\partial Q_{ni}(\beta_{n0})}{\partial \beta_{nk}} - E_0 \left( \frac{\partial Q_{ni}(\beta_{n0})}{\partial \beta_{nj}} \frac{\partial Q_{ni}(\beta_{n0})}{\partial \beta_{nk}} \right) \right\}^2 \\ &= O(p_n^2/n) \end{aligned}$$

which implies that  $\|(K_7)\| = O_P(p_n^2/n) = o(1)$ . Hence using  $K_4 = K_6 + K_7$ ,

$$\|K_4\| = o_P(1) + O_P(p_n(h^{p+1} + c_n \log^{1/2}(1/h)) + \sqrt{p_n^4/n}) = o_P(1).$$

Finally consider  $K_5$ . Define  $A_j = n^{-1} \sum_{i=1}^n (a_{ij} + b_{ij}) + n^{-1} \sum_{i=1}^n \frac{\partial Q_{ni}(\beta_{n0})}{\partial \beta_{nj}}$ , where  $a_{ij}$  and  $b_{ij}$  are defined as before, we can then rewrite  $K_5 = \{A_j A_k\}$ . Now

$$\begin{aligned} |A_j| &\leq \sup_{i,j} |a_{ij} + b_{ij}| + \left| \frac{1}{n} \sum_{i=1}^n \frac{\partial Q_{ni}(\beta_{n0})}{\partial \beta_{nj}} \right| \\ &= O_P(h^{p+1} + c_n \log^{1/2}(1/h) + n^{-1/2} p_n) + O_P(n^{-1/2}), \end{aligned}$$

where the last line follows from the approximations for  $a_{ij}$  and  $b_{ij}$ , and mean-variance decomposition of the term  $n^{-1} \sum_{i=1}^n \frac{\partial Q_{ni}(\beta_{n0})}{\partial \beta_{nj}}$ . Hence

$$\|K_5\| = O_P(p_n(h^{p+1} + c_n \log^{1/2}(1/h) + n^{-1/2} p_n)^2) = o_P(1),$$

and this completes the proof.  $\square$

### Proof of Theorem 5.

In expression (2.4), we set  $p = 0$ , which effectively assumes  $\alpha_{\beta_n}(U_i) \approx \alpha_{\beta_n}(u)$  for  $U_i$  in a neighborhood of  $u$ . Using the same notation as in the proof of Lemma 1, we have  $\bar{\alpha}_{ni}(u) = \alpha_{\beta_n}(u)^T \mathbf{X}_i + \mathbf{Z}_{ni}^T \beta_n$ ,  $\hat{\beta}^* = c_n^{-1}(\hat{\alpha}_{0\beta_n}(u) - \alpha_{\beta_n}(u))$  and  $\mathbf{X}_i^* = \mathbf{X}_i$ . Following the proof of Lemma 1, we arrive at equation (5.8), which in this case is reduced to

$$\sum_{i=1}^n q_2(\mathbf{X}_i^T \hat{\alpha}_{0\beta_n}(u) + \mathbf{Z}_{ni}^T \beta_n, Y_{ni}) \left( Z_{nij} + \left( \frac{\partial \hat{\alpha}_{0\beta_n}(u)}{\partial \beta_{nj}} \right)^T \mathbf{X}_i \right) \mathbf{X}_i K_h(U_i - u) = 0.$$

Solving for  $\frac{\partial \hat{\alpha}_{0\beta_n}(u)}{\partial \beta_{nj}}$  from the above equation, which is true for  $j = 1, \dots, p_n$ , we get the same expression as given in the lemma.

Hence it remains to show that  $\frac{\partial \hat{\mathbf{a}}_{0\beta_n}(u)}{\partial \beta_n}$  is a consistent estimator of  $\boldsymbol{\alpha}'_{\beta_n}(u)$ . However this is done by the proof of Lemma 1 already, where equation (5.9) becomes

$$\left\| \frac{\partial \hat{\mathbf{a}}_{0\beta_n}(u)}{\partial \beta_n} - \hat{\boldsymbol{\alpha}}'_{\beta_n}(u) \right\| = O_P(\sqrt{p_n}(h + c_n \log^{1/2}(1/h))) = o_P(1)$$

and the proof completes.  $\square$

## APPENDIX: PROOFS OF LEMMAS 3 - 6

**Lemma 3** *Assuming Conditions (A) - (G) and  $p_n^4/n = o(1)$ , we have*

$$n^{-1} \|\nabla^2 Q_n(\hat{\beta}_n) - \nabla^2 Q_n(\beta_{n0})\| = o_P(1).$$

*Proof of Lemma 3.* Consider

$$\begin{aligned} n^{-1} \|\nabla^2 Q_n(\beta_n) - \nabla^2 Q_n(\beta_{n0})\|^2 &= \frac{1}{n^2} \sum_{i,j=1}^{p_n} \left( \frac{\partial^2 Q_n(\hat{\beta}_n)}{\partial \beta_{ni} \partial \beta_{nj}} - \frac{\partial^2 Q_n(\beta_{n0})}{\partial \beta_{ni} \partial \beta_{nj}} \right)^2 \\ &= \frac{1}{n^2} \sum_{i,j=1}^{p_n} \left( \sum_{k=1}^{p_n} \frac{\partial^3 Q_n(\beta^*)}{\partial \beta_{ni} \partial \beta_{nj} \partial \beta_{nk}} (\hat{\beta}_{nk} - \beta_{0k}) \right)^2 \\ &\leq \frac{1}{n^2} \sum_{i,j=1}^{p_n} \sum_{k=1}^{p_n} \left( \frac{\partial^3 Q_n(\beta^*)}{\partial \beta_{ni} \partial \beta_{nj} \partial \beta_{nk}} \right)^2 \|\hat{\beta}_{nk} - \beta_{0k}\|^2, \end{aligned}$$

where  $\beta^*$  lies between  $\hat{\beta}_n$  and  $\beta_{n0}$ . Similar to approximating the order of  $\hat{I}_3$  in the proof of Theorem 1, the last line of the above equation is less than or equal to

$$n^{-2} O_P(n^2 p_n^3) \|\hat{\beta}_n - \beta_{n0}\|^2 = n^{-2} O_P(n^2 p_n^3) O_P(p_n/n) = o_P(1)$$

by the conclusion of Theorem 1.  $\square$

**Lemma 4** *Assuming Regularity Conditions (A) - (G), we have for each  $\beta_n \in \Omega_n$ ,*

$$n^{-1} \|\nabla^2 \hat{Q}_n(\beta_n) - \nabla^2 Q_n(\beta_n)\| = O_P(p_n(h^{p+1} + c_n \log^{1/2}(1/h))).$$

*Proof of Lemma 4.* By Taylor's expansion and Lemma 1,

$$\begin{aligned} &n^{-1} \frac{\partial}{\partial \beta_{nk}} (\nabla \hat{Q}_n(\beta_n) - \nabla Q_n(\beta_n)) \\ &= n^{-1} \left\{ \frac{\partial^3 Q_n(\beta_n)}{\partial \beta_{nk} \partial \beta_n \partial \boldsymbol{\alpha}_{\beta_n}^T} (\hat{\boldsymbol{\alpha}}_{\beta_n} - \boldsymbol{\alpha}_{\beta_n}) + \frac{\partial^2 Q_n(\beta_n)}{\partial \beta_n \partial \boldsymbol{\alpha}_{\beta_n}^T} \left( \frac{\partial \hat{\boldsymbol{\alpha}}_{\beta_n}}{\partial \beta_{nk}} - \frac{\partial \boldsymbol{\alpha}_{\beta_n}}{\partial \beta_{nk}} \right) \right. \\ &\quad \left. + \left( \frac{\partial \hat{\boldsymbol{\alpha}}'_{\beta_n}}{\partial \beta_{nk}} - \frac{\partial \boldsymbol{\alpha}'_{\beta_n}}{\partial \beta_{nk}} \right) \frac{\partial Q_n(\beta_n)}{\partial \boldsymbol{\alpha}_{\beta_n}} + (\hat{\boldsymbol{\alpha}}'_{\beta_n} - \boldsymbol{\alpha}'_{\beta_n}) \frac{\partial^2 Q_n(\beta_n)}{\partial \boldsymbol{\alpha}_{\beta_n} \partial \beta_{nk}} \right\} (1 + o_P(1)) \end{aligned}$$

Hence, using Regularity Condition (C),

$$\begin{aligned}
& \left\| n^{-1} \frac{\partial}{\partial \beta_{nk}} (\nabla \hat{Q}_n(\boldsymbol{\beta}_n) - \nabla Q_n(\boldsymbol{\beta}_n)) \right\| \\
&= O(1) \cdot \left( \sup_i \|\hat{\boldsymbol{\alpha}}_{\boldsymbol{\beta}_n}(U_i) - \boldsymbol{\alpha}_{\boldsymbol{\beta}_n}(U_i)\| + \sup_i \left\| \frac{\partial \hat{\boldsymbol{\alpha}}_{\boldsymbol{\beta}_n}(U_i)}{\partial \beta_{nk}} - \frac{\partial \boldsymbol{\alpha}_{\boldsymbol{\beta}_n}(U_i)}{\partial \beta_{nk}} \right\| \right. \\
&\quad \left. + \sup_i \|\hat{\boldsymbol{\alpha}}'_{\boldsymbol{\beta}_n}(U_i) - \boldsymbol{\alpha}'_{\boldsymbol{\beta}_n}(U_i)\| + \sup_i \left\| \frac{\partial \hat{\boldsymbol{\alpha}}'_{\boldsymbol{\beta}_n}(U_i)}{\partial \beta_{nk}} - \frac{\partial \boldsymbol{\alpha}'_{\boldsymbol{\beta}_n}(U_i)}{\partial \beta_{nk}} \right\| \right) \\
&= O_P(\sqrt{p_n}(h^{p+1} + c_n \log^{1/2}(1/h))),
\end{aligned}$$

where the last line follows from Lemma 1. Hence

$$n^{-1} \|\nabla^2 \hat{Q}_n(\boldsymbol{\beta}_n) - \nabla^2 Q_n(\boldsymbol{\beta}_n)\| = O_P(p_n(h^{p+1} + c_n \log^{1/2}(1/h))). \quad \square$$

**Lemma 5** *Under Regularity Conditions (A) - (G) and  $p_n^4/n = o(1)$ ,*

$$\begin{aligned}
\|n^{-1} \nabla^2 Q_n(\boldsymbol{\beta}_{n0}) + I_n(\boldsymbol{\beta}_{n0})\| &= o_P(p_n^{-1}), \\
\|n^{-1} \nabla^2 \hat{Q}_n(\boldsymbol{\beta}_{n0}) + I_n(\boldsymbol{\beta}_{n0})\| &= o_P(p_n^{-1}) + O_P(p_n(h^{p+1} + c_n \log^{1/2}(1/h))).
\end{aligned}$$

*Proof of Lemma 5.* The first conclusion follows from

$$\begin{aligned}
& E_0 p_n^2 \|n^{-1} \nabla^2 Q_n(\boldsymbol{\beta}_{n0}) + I_n(\boldsymbol{\beta}_{n0})\|^2 \\
&= p_n^2 n^{-2} E_0 \sum_{i,j=1}^{p_n} \left\{ \frac{\partial^2 Q_n(\boldsymbol{\beta}_{n0})}{\partial \beta_{ni} \partial \beta_{nj}} - E_0 \frac{\partial^2 Q_n(\boldsymbol{\beta}_{n0})}{\partial \beta_{ni} \partial \beta_{nj}} \right\}^2 = O(p_n^4/n) = o(1).
\end{aligned}$$

From this, triangle inequality immediately gives

$$\|n^{-1} \nabla^2 \hat{Q}_n(\boldsymbol{\beta}_{n0}) + I_n(\boldsymbol{\beta}_{n0})\| = o_P(p_n^{-1}) + \|n^{-1} \nabla^2 (\hat{Q}_n(\boldsymbol{\beta}_{n0}) - Q_n(\boldsymbol{\beta}_{n0}))\|.$$

The second equation then follows from Lemma 4.  $\square$

**Lemma 6** *Assuming the conditions in Theorem 3 and under the null hypothesis  $H_0$  as in the theorem,*

$$B_n^T (\hat{\gamma}_n - \gamma_{n0}) = \frac{1}{n} B_n^T \{B_n I_n(\boldsymbol{\beta}_{n0}) B_n^T\}^{-1} B_n^T \nabla Q_n(\boldsymbol{\beta}_{n0}) + o_P(n^{-1/2}).$$

*Proof of Lemma 6.* Since  $B_n B_n^T = I_{p_n-l}$ , for each  $\mathbf{v} \in \mathbb{R}^{p_n-l}$ , we have

$$\|B_n^T \mathbf{v}\| \leq \|\mathbf{v}\|. \quad (5.19)$$

Following the proof of Theorem 1, we have  $\|B_n^T(\hat{\gamma}_n - \gamma_n)\| = O_P(\sqrt{p_n/n})$ . Following the proof of Theorem 2 and by Lemma 2,

$$I_n(\beta_{n0})B_n^T(\hat{\gamma}_n - \gamma_{n0}) = n^{-1}\nabla Q_n(\beta_{n0}) + o_P(n^{-1/2}).$$

Left-multiplying with  $B_n$  and using equation (5.19), the right hand side of the above equation becomes  $n^{-1}B_n\nabla Q_n(\beta_{n0}) + o_P(n^{-1/2})$ . Hence,

$$B_n^T(\hat{\gamma}_n - \gamma_{n0}) = n^{-1}B_n^T(B_nI_n(\beta_{n0})B_n^T)^{-1}B_n\nabla Q_n(\beta_{n0}) + o_P(n^{-1/2}),$$

since  $B_nI_n(\beta_{n0})B_n^T$  has eigenvalues uniformly bounded away from 0 and infinity, like  $I_n(\beta_{n0})$  does.  $\square$

## References

- [1] Ahmad, I., Leelahanon, S. and Li, Q. (2005), Efficient Estimation of a Semiparametric Partially Linear Varying Coefficient Model, *Ann. Statist.*, **33**, 258–283.
- [2] Albright, S.C., Winston, W.L. and Zappe, C.J. (1999), *Data Analysis and Decision Making with Microsoft Excel*, Pacific Grove, CA: Duxbury.
- [3] Bickel, P.J. (1975), One-step Huber estimates in linear models, *J. Amer. Statist. Assoc.*, 70, 428–433.
- [4] Cai, Z., Fan, J. and Li, R. (2000), Efficient Estimation and Inferences for Varying-Coefficient Models, *J. Amer. Statist. Assoc.*, **95**, 888–902.
- [5] Carroll, R.J., Fan, J., Gijbels, I. and Wand, M.P. (1997), Generalized Partially Linear Single-Index Models, *J. Amer. Statist. Assoc.*, **92**, 477–489.
- [6] Donoho, D.L. (2000), High-Dimensional Data Analysis: The Curses and Blessings of Dimensionality, Lecture on August 8, 2000, to the American Mathematical Society on “Math Challenges of the 21st Century”.
- [7] Fan, J. and Huang, T. (2005), Profile Likelihood Inferences on Semiparametric Varying-Coefficient Partially Linear Models, *Bernoulli*, **11**, 1031–1057.
- [8] Fan, J. and Li, R. (2006), Statistical challenges with high-dimensionality: feature selection in knowledge discovery, *Proceedings of International Congress of Mathematicians* (M. Sanz-Solé, J. Soria, J.L. Varona, J. Verdera, eds.), Vol. III, 595–622.

- [9] Fan, J. and Peng, H. (2004), Nonconcave penalized likelihood with a diverging number of parameters, *Ann. Statist.*, **32**, 928–961.
- [10] Fan, J., Peng, H. and Huang, T. (2005), Semilinear high-dimensional model for normalization of microarray data: a theoretical analysis and partial consistency, *Jour. Amer. Statist.*, (with discussion), **100**, 781 – 813.
- [11] Fan, J., Zhang, C. and Zhang, J. (2001), Generalized Likelihood Ratio Statistics and Wilks Phenomenon, *Ann. Statist.*, **29**, 153–193.
- [12] Hastie, T.J. and Tibshirani, R. (1993), Varying-coefficient models, *J. R. Statist. Soc. B*, **55**, 757–796.
- [13] Hu, Z., Wang, N. and Carroll, R.J. (2004), Profile-kernel versus backfitting in the partially linear models for longitudinal/clustered data, *Biometrika*, **91**, 251–262.
- [14] Huber, P.J. (1973), Robust Regression: Asymptotics, Conjectures and Monte Carlo, *Ann. Statist.*, **1**, 799–821.
- [15] Jain, N. and Marcus, M. (1975), Central Limit Theorems for C(S)-valued Random Variables, *J. Funct. Anal.*, **19**, 216–231.
- [16] Kauermann, G. and Carroll, R.J. (2001), A note on the efficiency of sandwich covariance matrix estimation, *J. Amer. Statist. Assoc.*, **96**, 1387–1396.
- [17] Li, Q., Huang, C.J., Li, D. and Fu, T.T. (2002), Semiparametric smooth coefficient models, *J. Bus. Econom. Statist.*, **20**, 412–422.
- [18] Li, R. and Liang, H. (2005), Variable Selection in Semiparametric Regression Modeling, *Manuscript*.
- [19] Lin, X. and Carroll, R.J. (2006), Semiparametric estimation in general repeated measures problems, *J. R. Statist. Soc. B*, **68**, Part 1, 69–88.
- [20] McCullagh, P. and Nelder, J.A. (1989), *Generalized Linear Models* (2nd ed.), London: Chapman and Hall.
- [21] Mack, Y. P., Silverman, B. W. (1982), Weak and strong uniform consistency of kernel regression estimates, *Z. Wahrscheinlichkeitstheorie verw. Gebiete*, **61**, 405–415.

- [22] Murphy, S.A. (1993), Testing for a time dependent coefficient in Cox's regression model, *Scand. J. Statist.*, **20**, 35–50.
- [23] Murphy, S.A. and van der Vaart, A.W. (2000), On Profile likelihood (with discussion), *Journal of American Statistical Association*, **95**, 449–485.
- [24] Pollard, D. (1991), Asymptotics for least absolute deviation regression estimators, *Econ. Theory*, **7**, 186–199.
- [25] Portnoy, S. (1988), Asymptotic Behavior of Likelihood Methods for Exponential Families When the Number of Parameters Tends to Infinity, *Ann. Statist.*, **16**, 356–366.
- [26] Robinson, P.M. (1988), The stochastic difference between econometric and statistics, *Econometrica*, **56**, 531–547.
- [27] Severini, T.A. and Staniswalis, J.G. (1994), Quasi-likelihood Estimation in Semi-parametric Models, *J. Amer. Statist. Assoc.*, **89**, 501–511.
- [28] Severini, T.A. and Wong, W.H. (1992), Profile Likelihood and Conditionally Parametric Models, *Ann. Statist.*, **20**, 1768–1802.
- [29] Van der Vaart, A.W. (1998), *Asymptotic Statistics*, Cambridge Univ. Press.
- [30] Van Keilegom, I. and Carroll, R.J. (2007), Backfitting versus profiling in general criterion functions, *Statist. Sinica*, **17**, 797–816.
- [31] Xia, Y., Zhang, W. and Tong, H. (2004), Efficient estimation for semivarying-coefficient models, *Biometrika*, **91**, 661–681.
- [32] Yatchew, A. (1997), An elementary estimator for the partially linear model, *Economics Letters*, **57**, 135–143.
- [33] ZHANG, W., LEE, S.Y., AND SONG, X.Y. (2002), Local Polynomial fitting in semivarying coefficient model, *J. Mult. Anal.*, **82**, 166–188.