arXiv:math/0701206v1 [math.ST] 7 Jan 2007

SOME NOTES ON IMPROVING UPON THE JAMES-STEIN ESTIMATOR

By Yuzo Maruyama

The University of Tokyo

We consider estimation of a multivariate normal mean vector under sum of squared error loss. We propose a new class of smooth estimators parameterized by α dominating the James-Stein estimator. The estimator for $\alpha = 1$ corresponds to the generalized Bayes estimator with respect to the harmonic prior. When α goes to infinity, the estimator converges to the James-Stein positive-part estimator. Thus the class of our estimators is a bridge between the admissible estimator ($\alpha = 1$) and the inadmissible estimator ($\alpha = \infty$). Although the estimators have quasi-admissibility which is a weaker optimality than admissibility, the problem of determining whether or not the estimator for $\alpha > 1$ admissible is still open.

1. Introduction. Let X be a random variable having p-variate normal distribution $N_p(\theta, I_p)$. Then we consider the problem of estimating the mean vector θ by $\delta(X)$ relative to quadratic loss. Therefore every estimator is evaluated based on the risk function

$$R(\theta, \delta) = E_\theta \left[\|\delta(X) - \theta\|^2 \right] = \int_{R^p} \frac{\|\delta(x) - \theta\|^2}{(2\pi)^{p/2}} \exp\left(-\frac{\|x - \theta\|^2}{2} \right) dx.$$

The usual estimator X, with the constant risk p, is minimax for any dimension p. It is also admissible when $p = 1$ and 2, as shown in [1] and [13], respectively. [13] showed, however, that when $p \geq 3$, there exists an estimator dominating X among a class of equivariant estimators relative to the orthogonal transformation group which have the form

$$\delta_\phi(X) = (1 - \phi(\|X\|^2)/\|X\|^2)X. \tag{1.1}$$

[7] succeeded in giving an explicit form of an estimator improving on X as

$$\delta_{JS}(X) = (1 - (p-2)/\|X\|^2)X,$$

which is called the James-Stein estimator. More generally, a large class of better estimators than X has been proposed in the literature. The strongest tool for this is [14]'s identity as follows.

AMS 2000 subject classifications: Primary 62C15; secondary 62C10
Keywords and phrases: The James-Stein estimator, quasi-admissibility

LEMMA 1.1 ([14]). *If $Y \sim N(\mu, 1)$ and $h(y)$ is any differentiable function such that $E[|h'(Y)|] < \infty$, then*

$$E[h(Y)(Y-\mu)] = E[h'(Y)]. \tag{1.2}$$

By using the identity (1.2), the risk function of the estimator of the form $\delta_g(X) = X + g(X) = (X_1 + g_1(X), \ldots, X_p + g_p(X))'$ is written as

$$\begin{aligned} R(\theta, \delta_g) &= E_\theta\left[\|\delta_g(X) - \theta\|^2\right] \\ &= E_\theta\left[\|X-\theta\|^2\right] + E_\theta\left[\|g(X)\|^2\right] + 2\sum_{i=1}^{p} E_\theta\left[(X-\theta)'g(X)\right] \\ &= p + E_\theta\left[\|g(X)\|^2\right] + 2\sum_{i=1}^{p} E_\theta\left[\frac{\partial}{\partial x_i} g_i(X)\right], \end{aligned}$$

where $g_i(x)$ is assumed to be differentiable and $E|(\partial/\partial x_i)g_i(X)| < \infty$ for $i = 1, \ldots, p$. Since a statistic $\hat{R}(\delta(X))$ given by

$$p + \|g(X)\|^2 + 2\sum_{i=1}^{p} \frac{\partial}{\partial x_i} g_i(X) \tag{1.3}$$

does not depend on the unknown parameter θ and satisfies $E(\hat{R}(\delta(X))) = R(\theta, \delta)$, it is called the unbiased estimator of risk. Clearly if

$$\|g(x)\|^2 + 2\sum_{i=1}^{p} \frac{\partial}{\partial x_i} g_i(x) \le 0, \tag{1.4}$$

for any x, then $\delta_g(X)$ dominates X.

To make the structure of estimators improving on X more comprehensible, we consider a class of orthogonally equivariant estimators of a form given in (1.1). Assigning $g(x) = -\phi(\|x\|^2)x/\|x\|^2$ in (1.4), (1.4) becomes

$$\phi(w)\left(2(p-2) - \phi(w)\right)/w + 4\phi'(w) \ge 0, \tag{1.5}$$

for $w = \|x\|^2$. The inequality (1.5) is, for example, satisfied if $\phi(w)$ is monotone nondecreasing and within $[0, 2(p-2)]$ for any $w \ge 0$.

Since $\phi(w) = c$ for $0 < c < 2(p-2)$ satisfies the inequality (1.5),

$$(1 - c/\|X\|^2)X \tag{1.6}$$

with $0 < c < 2(p-2)$ dominates X. The estimator δ_{JS}, (1.6) with $c = p-2$, is the best estimator among a class of estimators (1.6) because the risk function of (1.6) is given by

$$p + c(c - 2(p-2))E[\|X\|^{-2}]$$

and hence minimized by $c = p - 2$.

It is however noted that when $\|x\|^2 < p - 2$, the James-Stein estimator yields an over-shrinkage and changes the sign of each component of X. The James-Stein positive-part estimator

$$\delta^+_{JS}(X) = \max(0, 1 - (p-2)/\|X\|^2)X,$$

eliminates this drawback and dominates the James-Stein estimator. We notice here that the technique for proving the inadmissibility of δ_{JS} is not from the Stein identity or the unbiased estimator of risk given in (1.3). The risk difference between δ_{JS} and δ_ϕ is given by

$$R(\delta_{JS}, \theta) - R(\delta_\phi, \theta) = E\left[-\frac{(\phi(\|X\|^2) - p + 2)^2}{\|X\|^2} + 4\phi'(\|X\|^2)\right],$$

but $\phi^+_{JS}(w) = \min(w, p-2)$ which makes the James-Stein positive-part estimator does not satisfy the inequality

$$-(\phi(w) - p + 2)^2 + 4w\phi'(w) \geq 0 \tag{1.7}$$

for any $w \geq 0$. In Section 2, we will show that there is no ϕ which satisfies (1.7) for any $w \geq 0$, that is, the Stein identity itself is not useful for finding estimators dominating δ_{JS}. We call such optimality for the James-Stein estimator quasi-admissibility. We will explain the concept and give a sufficient condition for quasi-admissibility in Section 2.

In spite of such difficulty, some estimators which dominate the James-Stein estimator have been given by several authors. [11] and [6] considered the class of estimators of forms $\delta_{LK}(X) = (1 - \phi_{LK}(\|X\|^2)/\|X\|^2)X$ where

$$\phi_{LK}(w) = p - 2 - \sum_{i=1}^{n} a_i w^{-b_i},$$

where $a_i \geq 0$ for any i and $0 < b_1 < b_2 < \cdots < b_n$. For example when $n = 1$, they both showed that, $\delta_{LK}(X)$ for $0 < b_1 < 4^{-1}(p-2)$ and $a_1 = 2b_1 2^{b_1}\Gamma(p/2 - b_1 - 1)/\Gamma(p/2 - 2b_1 - 1)$ is superior to the James-Stein

estimator. [10] gave two estimators which shrink toward the ball with center 0, $\delta^i_{KT}(X) = (1 - \phi^i_{KT}(\|X\|^2)/\|X\|^2)X$ for $i = 1, 2$ where

$$\phi^1_{KT}(w) = \begin{cases} 0 & w \le r^2 \\ p - 2 - \sum_{i=1}^{p-2}(r/w^{1/2})^i & w > r^2, \end{cases}$$

$$\phi^2_{KT}(w) = \begin{cases} 0 & w \le \{(p-1)/(p-2)\}^2 r^2 \\ p - 2 - r/(w^{1/2} - r) & w > \{(p-1)/(p-2)\}^2 r^2. \end{cases}$$

They showed that when r is sufficiently small, these two estimators dominate the James-Stein estimator. However, these estimators are not appealing since they are inadmissible. In our setting, the estimation of a multivariate normal mean, [3] showed that any admissible estimator should be generalized Bayes. Since the shrinkage factor $(1-\phi_{LK}(w)/w)$ becomes negative for some w and ϕ^i_{KT} for $i = 1, 2$ fail to be analytic, neither δ_{LK} nor δ^i_{KT} can be generalized Bayes or admissible.

In general, when we propose an estimator (δ_*, say) dominating a certain inadmissible estimator, it is extremely important to find it among admissible estimators. If not, a more difficult problem (finding an estimator improving on δ_*) just occurs. To the best of our knowledge, the sole admissible estimator dominating the James-Stein estimator is [8]'s estimator $\delta_K(X) = (1 - \phi_K(\|X\|^2)/\|X\|^2)X$ where

$$\phi_K(w) = p - 2 - 2\frac{\exp(-w/2)}{\int_0^1 \lambda^{p/2-2}\exp(-w\lambda/2)d\lambda}. \tag{1.8}$$

The estimator is generalized Bayes with respect to the harmonic prior density $\|\theta\|^{2-p}$ which was originally suggested by [14]. The only fault is, however, that it does not improve upon $\delta_{JS}(X)$ at $\|\theta\| = 0$. (See Section 3 for the detail.) Shrinkage estimators like (1.1) make use of the vague prior information that $\|\theta\|$ is close to 0. It goes without saying that we would like to get the significant improvement of risk when the prior information is accurate. Though $\delta_K(X)$ is an admissible generalized Bayes estimator and thus smooth, it has no improvement on $\delta_{JS}(X)$ at the origin $\|\theta\| = 0$. On the other hand, $\delta^+_{JS}(X)$ improves on $\delta_{JS}(X)$ significantly at the origin, but it is not analytic and is thus inadmissible by [3]'s complete class theorem. Therefore a more challenging open problem is to find admissible estimators dominating the James-Stein estimator especially at $\|\theta\| = 0$. In this paper, we will consider a class of estimators

$$\delta_\alpha(X) = \frac{\int_0^1 (1-\lambda)\lambda^{\alpha(p/2-1)-1}\exp(-\|X\|^2\alpha\lambda/2)d\lambda}{\int_0^1 \lambda^{\alpha(p/2-1)-1}\exp(-\|X\|^2\alpha\lambda/2)d\lambda}X \tag{1.9}$$

for $\alpha \geq 1$. In Section 3, we show that δ_α with $\alpha \geq 1$ improves on the James-Stein estimator and that it has strictly risk improvement at $\|\theta\| = 0$. Furthermore we see that δ_α approaches δ^+_{JS} as α goes to ∞. Since δ_α with $\alpha = 1$ corresponds to δ_K, the class of δ_α with $\alpha \geq 1$ is a bridge between δ_K which is admissible and δ^+_{JS} which is inadmissible. Although we show that δ_α with $\alpha > 1$ is quasi-admissible, which is introduced in Section 2, we have no idea on its admissibility at this stage.

2. Quasi-admissibility. In this section, we introduce the concept of quasi-admissibility and give a sufficient condition for quasi-admissibility. We deal with a reasonable class of estimators which have the form

$$\begin{aligned} \delta_m &= X + \nabla \log m(\|X\|^2) \\ &= \{1 + 2m'(\|X\|^2)/m(\|X\|^2)\}X \end{aligned} \tag{2.1}$$

where ∇ is a differential operator $(\partial/\partial x_1, \ldots, \partial/\partial x_p)'$ and m is a positive function. If $m(w) = w^{-c}$ for $c > 0$, (2.1) becomes the James-Stein type estimator (1.6). Any (generalized) Bayes estimator with respect to spherical symmetric measure π should also have the form (2.1) because it is written as

$$\begin{aligned} &\frac{\int_{R^p} \theta \exp(-\|x-\theta\|^2/2)\pi(d\theta)}{\int_{R^p} \exp(-\|x-\theta\|^2/2)\pi(d\theta)} \\ &= x + \frac{\int_{R^p} (\theta - x) \exp(-\|x-\theta\|^2/2)\pi(d\theta)}{\int_{R^p} \exp(-\|x-\theta\|^2/2)\pi(d\theta)} \\ &= x + \frac{\nabla \int_{R^p} \exp(-\|x-\theta\|^2/2)\pi(d\theta)}{\int_{R^p} \exp(-\|x-\theta\|^2/2)\pi(d\theta)} \\ &= x + \nabla \log \int_{R^p} \exp(-\|x-\theta\|^2/2)\pi(d\theta) \\ &= x + \nabla \log m_\pi(\|x\|^2) \end{aligned}$$

where $m_\pi(\|x\|^2) = \int_{R^p} \exp(-\|x-\theta\|^2/2)\pi(d\theta)$. [2] and [5] called an estimator of the form (2.1) pseudo-Bayes and quasi-Bayes, respectively. If, for given m, there exists a nonnegative measure ν which satisfies

$$m(\|x\|^2) = \int_{R^p} \exp(-\|\theta - x\|^2/2)\nu(d\theta), \tag{2.2}$$

the estimator of the form (2.1) is truly generalized Bayes. However it is often difficult to determine whether or not m has such an exact integral form.

Suppose that $\delta_{m,k}(X) = \delta_m(X) + k(\|X\|^2)X$ is a competitor of δ_m. Then substituting $g(x) = \nabla \log m(\|x\|^2) = 2xm'(\|x\|^2)/m(\|x\|^2)$ and $g(x) = 2xm'(\|x\|^2)/m(\|x\|^2) + k(\|x\|^2)x$ in (1.3) respectively, we have the unbiased estimators of risk as

$$\hat{R}(\delta_m) = p - 4w\left(\frac{m'(w)}{m(w)}\right)^2 + 4p\frac{m'(w)}{m(w)} + 8w\frac{m''(w)}{m(w)} \tag{2.3}$$

for $w = \|x\|^2$ and $\hat{R}(\delta_{m,k}) = \hat{R}(\delta_m) + \Delta(m, mk)$ where

$$\Delta(m, mk) = 4w\frac{m'(w)}{m(w)}k(w) + 2pk(w) + 4wk'(w) + wk^2(w). \tag{2.4}$$

If there exists k such that $\Delta(m, mk) \leq 0$ for any $w \geq 0$ with strict inequality for some w , that implies that δ_m is inadmissible, that is, $R(\theta, \delta_{mk}) \leq R(\theta, \delta_m)$ for all θ with strict inequality for some θ. If there does not exist such k, δ_m is said to be quasi-admissible. Hence quasi-admissibility is a weaker optimality than admissibility. Now we state a sufficient condition for quasi-admissibility. The idea is originally from [4], but the paper is not so accessible. See also [12], where quasi-admissibility is called permissibility.

THEOREM 2.1. *The estimator of the form* (2.1) *is quasi-admissible if*

$$\int_0^1 w^{-p/2}m(w)^{-1}dw = \infty \quad \textit{as well as} \quad \int_1^\infty w^{-p/2}m(w)^{-1}dw = \infty.$$

PROOF. We have only to show that $\hat{R}(\delta_{m,k}) \leq \hat{R}(\delta_m)$, that is, $\Delta(m, mk) \leq 0$ implies $k(w) \equiv 0$. Let $M(w) = w^{p/2}m(w)$ and $h(w) = M(w)k(w)$. Then we have

$$\Delta(m, mk) = 4w\frac{h'(w)}{M(w)} + w\frac{h^2(w)}{M^2(w)} = \frac{4wh^2(w)}{M(w)}\left(-\frac{d}{dw}\left\{\frac{1}{h(w)}\right\} + \frac{1}{4M(w)}\right).$$

First we show that $h(w) \geq 0$ for all $w \geq 0$. Suppose to the contrary that $h(w) < 0$ for some w_0. Then $h(w) < 0$ for all $w \geq w_0$ since $h'(w)$ should be negative. For all $w > w_0$, the inequality

$$\frac{d}{dw}\left(\frac{1}{h(w)}\right) \geq \frac{1}{4M(w)} \tag{2.5}$$

should be satisfied. Integrating both sides of (2.5) from w_0 to w^* leads to

$$\frac{1}{h(w^*)} - \frac{1}{h(w_0)} \geq \frac{1}{4}\int_{w_0}^{w^*} M^{-1}(t)dt.$$

As $w^* \to \infty$, the right-hand side of above inequality tends to infinity, and this provides a contradiction since the left-hand side is less than $-1/h(w_0)$. Thus we have $g(w) \geq 0$ for all w.

Similarly we can show that $g(w) \leq 0$ for all w. It follows that $h(w)$ is zero for all w, which implies that $k(w) \equiv 0$ for all w. This completes the proof. □

Combining Theorem 2.1 and [3]'s sufficient condition for admissibility, we see that a quasi-admissible estimator of the form (2.1) is admissible if it is truly generalized Bayes, that is, there exists a nonnegative measure ν such that

$$m(\|x\|^2) = \int_{R^p} \exp(-\|x-\theta\|^2/2)\nu(d\theta). \tag{2.6}$$

However, for given m, it is often quite difficult to determine whether m has such an integral form. Furthermore, even if we find that m does not have an integral form like (2.6), that is, the estimator is inadmissible, it is generally very difficult to find an estimator dominating the inadmissible estimator.

The function m for the James-Stein estimator is $m(w) = w^{2-p}$, which satisfies the assumptions of Theorem 2.1.

COROLLARY 2.1. *The James-Stein estimator is quasi-admissible.*

In this case, it is not difficult to find an estimator dominating δ_{JS} by taking positive part. But it is not easy to find a large class of estimators dominating the James-Stein estimator. In Section 3, we introduce an elegant sufficient condition for domination over δ_{JS} proposed by [9].

3. A class of quasi-admissible estimators improving upon the James-Stein estimator. In this section, we introduce [9]'s sufficient condition for improving upon the James-Stein estimator and propose a class of smooth quasi-admissible estimators satisfying it.

[9] showed that if $\lim_{w\to\infty} \phi(w) = p-2$ then the difference of risk functions between δ_{JS} and δ_ϕ can be written as

$$\begin{aligned} &R(\theta, \delta_{JS}) - R(\theta, \delta_\phi) \\ &= 2\int_0^\infty \phi'(w)\left(\phi(w) - (p-2) + \frac{2f(w,\lambda)}{\int_0^w y^{-1} f_p(y,\lambda)dy}\right) \\ &\qquad \times \int_0^w y^{-1} f_p(y,\lambda)dydw, \end{aligned} \tag{3.1}$$

where $\lambda = \|\theta\|^2$ and $f_p(x;\lambda)$ denotes a density of a non-central chi-square distribution with p degrees of freedom and non-centrality parameter λ.

Moreover by the inequality

$$f_p(w;\lambda)/\int_0^w y^{-1} f_p(y;\lambda)dy \geq f_p(w)/\int_0^w y^{-1} f_p(y)dy,$$

where $f_p(y) = f_p(y;0)$, which can be shown by the correlation inequality, we have

$$\begin{aligned} &R(\theta,\delta_{JS}) - R(\theta,\delta_\phi) \\ &\geq 2\int_0^\infty \phi'(w)\left(\phi(w)-\phi_0(w)\right)\left(\int_0^w y^{-1} f_p(y,\lambda)dy\right)dw, \end{aligned}$$

where

$$\phi_0(w) = p - 2 + 2f_p(w)/\int_0^w y^{-1} f_p(y)dy.$$

Since $\phi_0 = \phi_K$ given in (1.8) by an integration by parts and $\lim_{w\to\infty}\phi_K(w) = p-2$, we have the following result.

THEOREM 3.1 ([9]). *If $\phi(w)$ is nondecreasing and within $[\phi_K(w), p-2]$ for any $w \geq 0$, then $\delta_\phi(X)$ of form (1.1) dominates the James-Stein estimator.*

The assumption of the theorem above is satisfied by ϕ_K and ϕ_{JS}^+. By (3.1), we see that the risk difference at $\|\theta\| = 0$ between δ_{JS} and δ_ϕ, the limit of which is $p-2$, is given by

$$2\int_0^\infty \phi'(w)\left(\phi(w)-\phi_K(w)\right)\left(\int_0^w y^{-1} f_p(y)dy\right)dw. \tag{3.2}$$

Hence δ_K does not improve upon $\delta_{JS}(X)$ at $\|\theta\| = 0$, although $\delta_K(X)$ is an admissible generalized Bayes estimator and thus smooth. On the other hand, $\delta_{JS}^+(X)$ improves on $\delta_{JS}(X)$ significantly at the origin, but it is not analytic and is thus inadmissible by [3]'s complete class theorem. Therefore a more challenging problem is to find admissible estimators dominating the James-Stein estimator especially at $\|\theta\| = 0$.

In this paper, we propose a class of estimators

$$\delta_\alpha(X) = \left(1 - \phi_\alpha(\|X\|^2)/\|X\|^2\right)X \tag{3.3}$$

where

$$\begin{aligned}\phi_\alpha(w) &= w\frac{\int_0^1 \lambda^{\alpha(p/2-1)}\exp(-w\alpha\lambda/2)d\lambda}{\int_0^1 \lambda^{\alpha(p/2-1)-1}\exp(-w\alpha\lambda/2)d\lambda}\\ &= p-2-\frac{2\exp(-w\alpha/2)}{\alpha\int_0^1 \lambda^{\alpha(p/2-1)-1}\exp(-w\alpha\lambda/2)d\lambda}\\ &= p-2-\frac{2}{\alpha\int_0^1 (1-\lambda)^{\alpha(p/2-1)-1}\exp(w\alpha\lambda/2)d\lambda}. \qquad (3.4)\end{aligned}$$

The main theorem of this paper is as follows.

THEOREM 3.2. *1. $\delta_\alpha(X)$ dominates $\delta_{JS}(X)$ for $\alpha \geq 1$.*

2. The risk of $\delta_\alpha(X)$ for $\alpha > 1$ at $\|\theta\| = 0$ is strictly less than the risk of the James-Stein estimator at $\|\theta\| = 0$.

3. $\delta_\alpha(X)$ approaches the positive-part James-Stein estimator when α tends to infinity, that is,

$$\lim_{\alpha\to\infty}\delta_\alpha(X) = \delta^+_{JS}(X).$$

Clearly $\delta_1(X) = \delta_K(X)$. The class of δ_α with $\alpha \geq 1$ is a bridge between δ_K which is admissible and δ^+_{JS} which is inadmissible.

PROOF. [part 1] We shall verify that $\phi_\alpha(w)$ for $\alpha \geq 1$ satisfies assumptions in Theorem 3.1. Applying the Taylor expansion to a part of (3.4), we have

$$\begin{aligned}&\frac{\alpha}{2}\int_0^1 (1-\lambda)^{\alpha(p/2-1)-1}\exp(w\alpha\lambda/2)d\lambda\\ &= \sum_{i=0}^{\infty} w^i \prod_{j=0}^{i}(p-2+2j/\alpha)^{-1} = \psi(\alpha, w) \quad (say.)\end{aligned}$$

As $\psi(\alpha, w)$ is increasing in w, $\phi_\alpha(w)$ is increasing in w. As $\lim_{w\to\infty}\psi(\alpha, w) = \infty$, for any $\alpha \geq 1$, it is clear that $\lim_{w\to\infty}\phi_\alpha(w) = p-2$. In order to show that $\phi_\alpha(w) \geq \phi_K(w) = \phi_1(w)$ for $\alpha \geq 1$, we have only to check that $\psi(\alpha, w)$ is increasing in α. It is easily verified because the coefficient of each term of $\psi(\alpha, w)$ is increasing in α. We have thus proved the theorem.

[part 2] Since ϕ_α for $\alpha > 1$ is strictly greater than ϕ_K and strictly increasing in w, the risk difference of δ_α for $\alpha > 1$ and δ_{JS} at $\|\theta\| = 0$, which is given in (3.2), is strictly positive.

[part 3] Since $\psi(\alpha, w)$ is increasing in α, it converges to

$$(p-2)^{-1}\sum_{i=0}^{\infty}\left(\frac{w}{p-2}\right)^i$$

by the monotone convergence theorem when α goes to infinity. Considering two cases: $w < (\geq) p-2$, we obtain $\lim_{\alpha\to\infty}\phi_\alpha(w) = w$ if $w < p-2$; $= p-2$ otherwise. This completes the proof. □

The estimator δ_α is expressed as $X + \nabla \log m_\alpha(\|X\|^2)$ where

$$m_\alpha(w) = \left\{\int_0^1 \lambda^{\alpha(p/2-1)-1}\exp\left(-\frac{\alpha w}{2}\lambda\right)d\lambda\right\}^{1/\alpha}.$$

Since $m_\alpha(w) \sim w^{2-p}$ for sufficiently large w by Tauberian's theorem and $0 < m_\alpha(0) < \infty$, we have the following result by Theorem 2.1.

COROLLARY 3.1. *$\delta_\alpha(X)$ is quasi-admissible for $p \geq 3$.*

Needless to say, we are extremely interested in determining whether or not $\delta_\alpha(X)$ for $\alpha > 1$ is admissible. Since $\delta_\alpha(X)$ with $\alpha > 1$ is quasi-admissible, it is admissible if it is generalized Bayes, that is, there exists a measure ν which satisfies

$$\int_{R^p}\exp(-\|\theta - x\|^2/2)\nu(d\theta) = \left(\int_0^1 \lambda^{\alpha(p/2-1)-1}\exp(-\alpha\|X\|^2\lambda/2)d\lambda\right)^{1/\alpha}.$$

I have no idea on the way to construct such a measure ν so far. Even if we find that there is no ν, which implies δ_α is inadmissible, it is very difficult to find an estimator dominating δ_α for $\alpha > 1$.

References.

[1] BLYTH, C. R. (1951). On minimax statistical decision procedures and their admissibility. *Ann. Math. Statist. 22*, 22–42. MRMR0039966 (12,622f)

[2] BOCK, M. E. (1988). Shrinkage estimators: pseudo-Bayes rules for normal mean vectors. In *Statistical decision theory and related topics, IV, Vol. 1 (West Lafayette, Ind., 1986)*. Springer, New York, 281–297. MRMR927108 (89b:62106)

[3] BROWN, L. D. (1971). Admissible estimators, recurrent diffusions, and insoluble boundary value problems. *Ann. Math. Statist. 42*, 855–903. MRMR0286209 (44 #3423)

[4] BROWN, L. D. (1988). The differential inequality of a statistical estimation problem. In *Statistical decision theory and related topics, IV, Vol. 1 (West Lafayette, Ind., 1986)*. Springer, New York, 299–324. MRMR927109 (89d:62010)

[5] BROWN, L. D. AND ZHANG, L. H. (2006). Estimators for gaussian models having a block-wise structure. Mimeo,
(available at: `http://www-stat.wharton.upenn.edu/ lbrown/teaching/Shrinkage/`).

[6] Guo, Y. Y. and Pal, N. (1992). A sequence of improvements over the James-Stein estimator. *J. Multivariate Anal.* **42**, 2, 302–317. MRMR1183849 (93h:62009)

[7] James, W. and Stein, C. (1961). Estimation with quadratic loss. In *Proc. 4th Berkeley Sympos. Math. Statist. and Prob., Vol. I.* Univ. California Press, Berkeley, Calif., 361–379. MRMR0133191 (24 #A3025)

[8] Kubokawa, T. (1991). An approach to improving the James-Stein estimator. *J. Multivariate Anal.* **36**, 1, 121–126. MRMR1094273 (92b:62070)

[9] Kubokawa, T. (1994). A unified approach to improving equivariant estimators. *Ann. Statist.* **22**, 1, 290–299. MRMR1272084 (95h:62011)

[10] Kuriki, S. and Takemura, A. (2000). Shrinkage estimation towards a closed convex set with a smooth boundary. *J. Multivariate Anal.* **75**, 1, 79–111. MRMR1787403 (2001h:62034)

[11] Li, T. F. and Kuo, W. H. (1982). Generalized James-Stein estimators. *Comm. Statist. A—Theory Methods* **11**, 20, 2249–2257. MRMR678683 (84b:62078)

[12] Rukhin, A. L. (1995). Admissibility: Survey of a concept in progress. *International Statistical Review 63*, 95–115.

[13] Stein, C. (1956). Inadmissibility of the usual estimator for the mean of a multivariate normal distribution. In *Proceedings of the Third Berkeley Symposium on Mathematical Statistics and Probability, 1954–1955, vol. I.* University of California Press, Berkeley and Los Angeles, 197–206. MRMR0084922 (18,948c)

[14] Stein, C. (1974). Estimation of the mean of a multivariate normal distribution. In *Proceedings of the Prague Symposium on Asymptotic Statistics (Charles Univ., Prague, 1973), Vol. II.* Charles Univ., Prague, 345–381. MRMR0381062 (52 #1959)

Center for Spatial Information Science,
The University of Tokyo
5-1-5 Kashiwanoha, Kashiwa-shi, Chiba 277-8568, Japan
E-mail: maruyama@csis.u-tokyo.ac.jp