

Quantile regression when the covariates are functions

Herve Cardot¹, Christophe Crambes² and Pascal Sarda³

E-mail addresses: ¹cardot@toulouse.inra.fr, ²crambes@cict.fr, ³sarda@cict.fr

Abstract This paper deals with a linear model of regression on quantiles when the explanatory variable takes values in some functional space and the response is scalar. We propose a spline estimator of the functional coefficient that minimizes a penalized L^1 type criterion. Then, we study the asymptotic behavior of this estimator. The penalization is of primary importance to get existence and convergence.

Key words Functional data analysis, conditional quantiles, B-spline functions, roughness penalty.

1 Introduction

Because of the increasing performances of measurement apparatus and computers, many data are collected and saved on thinner and thinner time scales or spatial grids (temperature curves, spectrometric curves, satellite images, ...). So, we are led to process data comparable to curves or more generally to functions of continuous variables (time, space). These data are called functional data in the literature (see Ramsay and Silverman, 2002). Thus, there is a need to develop statistical procedures as well as theory for this kind

¹INRA Toulouse, Biométrie et Intelligence Artificielle, 31326 Castanet-Tolosan Cedex, France

²Université Paul Sabatier, Laboratoire de Statistique et Probabilités, UMR C 5583, 118, route de Narbonne, 31062 Toulouse Cedex, France

³Université Toulouse-Le Mirail, GRIMM, EA 3686, 5, allées Antonio Machado, 31058 Toulouse Cedex 9, France

of data and actually many recent works study models taking into account the functional nature of the data.

Mainly in a formal way, the oldest works in that direction intended to give a mathematical framework based on the theory of linear operators in Hilbert spaces (see Deville, 1974, Dauxois and Pousse, 1976). After that and in another direction, practical aspects of extensions of descriptive statistical methods like for example Principal Component Analysis have been considered (see Besse and Ramsay, 1986). The monographs by Ramsay and Silverman (1997, 2002) are important contributions in this area.

As pointed out by Ramsay and Silverman (1997), "the goals of functional data analysis are essentially the same as those of other branches of Statistics": one of this goal is the explanation of variations of a dependent variable Y (response) by using information from an independent functional variable X (explanatory variable). In many applications, the response is a scalar: see Frank and Friedman (1993), Ramsay and Silverman (1997), ... Traditionally, one deals, for such a problem, with estimating the regression on the mean i.e. the minimizer among some class of functionals of

$$E (Y - r(X))^2 :$$

As when X is a vector of real numbers, the two main approaches are linear (see Ramsay and Dalzell, 1991, for the functional linear model) or purely nonparametric (see Ferraty and Vieu, 2002, which adapt kernel estimation to the functional setting). It is also known that estimating the regression on the median or more generally on quantiles has some interest. The problem is then to estimate the minimizer among \mathcal{G} of

$$E [l(Y - g(X))]; \quad (1)$$

where $l(u) = |u| + (2 - 1)u$. The value $\alpha = 1/2$ corresponds to the conditional median whereas values $\alpha \in]0; 1[$ correspond to conditional quantiles of order α . The advantage of estimating conditional quantiles may be found in

many applications such as in agronomy (estimation of yield thresholds), in medicine or in reliability. Besides robust aspects of the median, it may also help to derive some kind of confidence prediction intervals based on quantiles.

In our work, we assume that the conditional quantile of order α can be written as

$$g_\alpha(X) = h_\alpha; X; \quad (2)$$

where $\langle \cdot; \cdot \rangle$ is a functional inner product and the parameter of the model is a function to be estimated. This is the equivalent of the linear model for regression quantiles studied by Koenker and Bassett (1978) where the inner product is the Euclidean one and the parameter is a vector of scalars. We choose to estimate the function g_α by a "direct" method: writing our estimator as a linear combination of B-splines, it minimizes the empirical version of expectation (1) with the addition of a penalty term proportional to the square norm of a given order derivative of the spline. The penalization term allows on one side to control the regularity of the estimator and on the other side to get consistency.

Unlike for the square function, minimization of function 1 does not lead to an explicit expression of the estimator. While computation of the estimator can be resolved by using traditional algorithms (for instance based on Iteratively Weighted Least Squares), the convexity of 1 allows theoretical developments.

In section 2, we define more precisely the framework of our study and the spline estimator of the functional parameter g_α . Section 3 is devoted to the asymptotic behaviour of our estimator: we study L^2 convergence and derive an upper bound for the rate of convergence. Comments on the model and on the optimality of the rate of convergence are given in section 4. Finally, the proofs are gathered in section 5.

2 Construction of the estimator

In this work, the data consist of an i.i.d. sample of pairs $(X_i; Y_i)_{i=1, \dots, m}$ drawn from a population distribution $(X; Y)$. We consider explanatory variables X_i which are square integrable (random) functions defined on $[0; 1]$, i.e. are elements of the space $L^2([0; 1])$ so that $X_i = (X_i(t); t \in [0; 1])$. The response Y_i is a scalar belonging to \mathbb{R} . Assume that H , the range of X , is a closed subspace of $L^2([0; 1])$. For Y having a finite expectation, $E(Y) < +\infty$, and for $\alpha \in]0; 1[$, the conditional α -quantile functional g_α of Y given X is a functional defined on H minimizing (1).

Our aim is to generalize the linear model introduced by Koenker and Bassett (1978). In our setting, it consists in assuming that g_α is a linear and continuous functional defined on H and then it follows that $g_\alpha(X)$ can be written as in (2). Taking the usual inner product in $L^2([0; 1])$, we can write

$$g_\alpha(X) = h_\alpha; X_i = \int_0^1 \psi_\alpha(t) X_i(t) dt;$$

where ψ_α is the functional coefficient in H to be estimated, the order α being fixed. From now on we consider, for simplicity, that the random variables X_i are centered, that is to say $E(X_i(t)) = 0$, for t a. e.

When X is multivariate, Bassett and Koenker (1978) study the least absolute error (LAE) estimator for the conditional median, which can be extended to any quantile replacing the absolute value by the convex function l_α in the criterion to be minimized (see Koenker and Bassett, 1978). In our case where we have to estimate a function belonging to an infinite dimensional space, we are looking at an estimator in the form of an expansion in some basis of B-splines functions and then minimizing a similar criterion with however the addition of a penalty term.

Before describing in details the estimation procedure, let us note that estimation of conditional quantiles has received a special attention in the multivariate case. As said before, linear modelling has been mainly investigated by Bassett and Koenker (1978). For nonparametric models, we may distinguish

two different approaches: "indirect" estimators which are based on a preliminary estimation of the conditional cumulative distribution function (cdf) and "direct" estimators which are based on minimizing the empirical version of criterion (1). In the class of "indirect" estimators, Bhattacharya and Gangopadhyay (1990) study a kernel estimator of the conditional cdf, and estimation of the quantile is achieved by inverting this estimated cdf. In the class of "direct" estimators, kernel estimators based on local t have been proposed (see Tsybakov, 1986, Lejeune and Sarda, 1988 or Fan, Hu and Truong, 1994); in a similar approach, He and Shi (1994) and Koenker et. al. (1994) propose a spline estimator. Although our setting is quite different, we adapt in our proofs below some arguments of the work by He and Shi (1994).

In nonparametric estimation, it is usual to assume that the function to be estimated is sufficiently smooth so that it can be expanded in some basis: the degree of smoothness is quantified by the number of derivatives and a Lipschitz condition for the derivative of greatest order (see condition (H.2) below). It is also quite usual to approximate such kind of functions by means of regression splines (see de Boor, 1978, for a guide for splines). For this, we have to select a degree q in \mathbb{N} and a subdivision of $[0;1]$ defining the position of the knots. Although it is not necessary, we take equispaced knots so that only the number of the knots has to be selected: for k in \mathbb{N}^2 , we consider $k-1$ knots that define a subdivision of the interval $[0;1]$ into k sub-intervals. For asymptotic theory, the degree q is fixed but the number of sub-intervals k depends on the sample size n ; $k = k_n$: It is well-known that a spline function is a piecewise polynomial: we consider here piecewise polynomials of degree q on each sub-interval, and $(q-1)$ times differentiable on $[0;1]$. This space of spline functions is a vectorial space of dimension $k+q$. A basis of this vectorial space is the set of the so-called normalized B-spline functions, that we note by $B_{k,q} = (B_1; \dots; B_{k+q})$.

Then, we estimate β by a linear combination of functions B_1 . This leads us to find a vector $b = (b_1; \dots; b_{k+q})$ in \mathbb{R}^{k+q} such that

$$b = \sum_{l=1}^{k+q} b_l B_l = B_{k+q} b. \quad (3)$$

It is then natural to look for b as the minimizer of the empirical version of (1) among functional g of the form (2) with functions B_l belonging to the space of spline functions defined above. We will however consider a penalized criterion as we will see now. In our setting, the pseudo-design matrix A is the matrix of dimension $n \times (k+q)$ and elements $\{X_i; B_j\}$ for $i = 1; \dots; n$ and $j = 1; \dots; k+q$. Even if we do not have an explicit expression for a solution to the minimization problem, it is known that the solution would depend on the properties of the inverse of the matrix $\frac{1}{n} A^T A$ which is the $(k+q) \times (k+q)$ matrix with general term $\frac{1}{n} \sum_{i=1}^n (B_j; B_l)$, where $\frac{1}{n} \sum_{i=1}^n$ is the empirical version of the covariance operator γ_X of X defined for all u in $L^2([0;1])$ by

$$\gamma_X u = E (X; u | X) : \quad (4)$$

We know that γ_X is a nuclear operator (see Dauxois et al, 1982), consequently no bounded inverse exists for this operator. Moreover, as a consequence of the first monotonicity principle (see theorem 7.1, p.58, in Weinberger, 1974), the restriction of this operator to the space of spline functions has smaller eigenvalues than γ_X . Finally, it appears to be impossible to control the speed of convergence to zero of the smallest eigenvalue of $\frac{1}{n} A^T A$ (when n tends to infinity): in that sense, we are faced with an inversion problem that can be qualified as ill-conditioned. A way to circumvent this problem is to introduce a penalization term in the minimization criterion (see Ramsay and Silverman, 1997, or Cardot et al, 2003, for a similar approach in the functional linear model). Thus, the main role of the penalization is to control the inversion of the matrix linked to the solution of the problem and it consists in restricting the space of solutions. The penalization introduced below will have another effect since we also want to control the smoothness of our estimator. For this reason, and following several authors (see references above), we choose a penalization which allows to control the norm of the derivative of order $m > 0$

of any linear combination of B-spline functions, so that it can be expressed matrixially. Denoting by $(B_{k,q})^{(m)}$ the m -th derivative of the spline function $B_{k,q}$; we have

$$(B_{k,q})^{(m)}{}^2 = G_k; \quad 8 \leq 2R^{k+q};$$

where G_k is the $(k+q) \times (k+q)$ matrix with general term $[G_k]_{j1} = hB_j^{(m)}; B_1^{(m)}$;

Then, the vector b in (3) is chosen as the solution of the following minimization problem

$$\min_{2R^{k+q}} \frac{1}{n} \sum_{i=1}^n (Y_i - hB_{k,q}; X_{i1}) + k (B_{k,q})^{(m)} k^2; \quad (5)$$

where k is the penalization parameter. In the next section, we present a convergence result of the solution of (5). Note that the role of the penalization also clearly appears in this result.

3 Convergence result

We present in this section the main result on the convergence of our estimator. The behaviour of our estimator is linked to a penalized version of the matrix $\mathcal{C} = \frac{1}{n} A^T A$. More precisely, adopting the same notations as in Cardot et al. (2003), the existence and convergence of our estimator depend on the inverse of the matrix $\mathcal{C} = \mathcal{C} + G_k$. Under the hypotheses of theorem 1 below, the smallest eigenvalue of \mathcal{C} , noted $\min(\mathcal{C})$, tends to zero as the sample size n tends to infinity. As the rate of convergence of b depends on the speed of convergence of $\min(\mathcal{C})$ to zero, we introduce a sequence $(\epsilon_n)_{n \geq 2N}$ such that the set Ω_n defined by

$$\Omega_n = \{ \min(\mathcal{C}) > c_n \epsilon_n \}; \quad (6)$$

has probability which goes to 1 when n goes to infinity. Cardot et al. (2003) have shown that such a sequence exists in the sense that under hypotheses of theorem 1, there exists a strictly positive sequence $(\epsilon_n)_{n \geq 2N}$ tending to zero as n tends to infinity and such that

$$\sup_{n \in \mathbb{N}} \|b_n\|_{\mathcal{C}^1} \leq C_0 + o_p((k_n^2 n^{-1})^{-1/2}); \quad (7)$$

with $\alpha \in]0;1[$.

To prove the convergence result of the estimator b_n , we assume that the following hypotheses are satisfied.

(H.1) $\|X\|_{\mathcal{C}^0} \leq C_0 < +\infty$; as:

(H.2) The function φ is supposed to have a p^0 -th derivative $\varphi^{(p^0)}$ such that

$$\varphi^{(p^0)}(t) = \varphi^{(p^0)}(s) + C_1 |t - s|; \quad s, t \in [0;1];$$

where $C_1 > 0$ and $\alpha \in [0;1]$. In what follows, we set $p = p^0 + \alpha$ and we suppose that $q = p - m$.

(H.3) The eigenvalues of \mathcal{K}_X (defined in (4)) are strictly positive.

(H.4) For $x \in H$, the random variable defined by $\varepsilon = Y - h; X$ has conditional density function f_x given $X = x$, continuous and bounded below by a strictly positive constant at 0, uniformly for $x \in H$.

We derive in theorem 1 below an upper bound for the rate of convergence with respect to some kind of L^2 -norm. Indeed, the operator \mathcal{K}_X is strictly non-negative, so we can associate it a semi-norm noted $\|\cdot\|_2$ and defined by $\|u\|_2^2 = \langle \mathcal{K}_X u, u \rangle$. Then, we have the following result.

Theorem 1 Under hypotheses (H.1) – (H.4), if we also suppose that there exists $\alpha \in]0;1[$ such that $k_n \rightarrow \infty$, $n^\alpha \rightarrow 0$ and $n^\alpha (k_n^2 n^{-1})^{-1/2} \rightarrow 0$ (where α is defined in relation 7)), then

(i) b_n exists and is unique except on a set whose probability goes to zero as n goes to infinity,

$$(ii) \|b_n - b\|_2^2 = O_p \left(\frac{1}{k_n^{2p}} + \frac{1}{n} + \frac{1}{k_n} + k_n^{2(m-p)} \right);$$

4 Some comments

(i) Hypotheses (H.1) and (H.3) are quite usual in the functional setting: see for instance Bosq (2000) or Cardot et al. (2003). Hypothesis (H.4) implies

uniqueness of the conditional quantile of order α .

(ii) Some arguments in the proof of theorem 1 are inspired from the demonstration of He and Shi (1994) within the framework of real covariates. Moreover, some results from Cardot et. al. (2003) are also useful, mainly to deal with the penalization term as pointed out above.

Note that it is assumed in the model of He and Shi (1994) that the error term is independent of X : condition (H.4) allows us to deal with a more general setting, as in Koenker and Bassett (1978).

(iii) It is possible to choose particular values for α and β to optimize the upper bound for the rate of convergence in theorem 1. In particular, we remark the importance to control the speed of convergence to 0 of the smallest eigenvalue of \mathcal{Q}_n by λ_n . For example, Cardot et al. (2003) have shown that, under hypotheses of theorem 1, relation (7) is true with $\lambda_n = \lambda_{k_n}$. This gives us

$$k_n^b \quad k_2^2 = O_p \left(\frac{1}{k_n^{2p}} + \frac{k_n}{n} + \lambda_n + k_n^{2(m-p)} \right) :$$

A corollary is obtained if we take $k_n = n^{1/(4p+1)}$ and $\lambda_n = n^{-2p/(4p+1)}$; then we get

$$k_n^b \quad k_2^2 = O_p \left(n^{-2p/(4p+1)} \right) :$$

We can imagine that, with stronger hypotheses on the random function X , we can find a sequence λ_n greater than λ_{k_n} , that will improve the convergence speed of the estimator. As a matter of fact, the rate derived in theorem 1 does not imply the rate obtained by Stone (1982), that is to say a rate of order $n^{-2p/(2p+1)}$. Indeed, suppose that $\lambda_1 = k_n^{2p}$, $\lambda_2 = (n - k_n)$ and $\lambda_3 = (k_n - n)$ are all of order $n^{-2p/(2p+1)}$. This would imply that $k_n = n^{1/(2p+1)}$ and $\lambda_n = n^{-1/(2p+1)}$, which contradicts the condition $\lambda_n = n^{-(1-\alpha)/2}$. Nevertheless, it is possible to obtain a speed of order $n^{-2p/(2p+1)+}$. This leads to $k_n = n^{1/(2p+1) - \epsilon(2p)}$ and $\lambda_n = n^{-1/(2p+1) - \epsilon(2p+1)}$. Then, the condition $\lambda_n = n^{-(1-\alpha)/2}$ implies $\alpha = p(1 - \epsilon) = (2p+1)\epsilon$. So finally, we get $k_n = n^{(1-\epsilon)/(2(2p+1))}$, $\lambda_n = n^{-(4p-1+\epsilon)/(4(2p+1))}$ and $\lambda_n = n^{-(p-1+p\epsilon)/(2p+1)}$. The convergence result would be then

$$k^b \quad k_2^2 = O_p(n^{p(1+\epsilon)/(2p+1)}) :$$

A final remark is that the last term $k_n^{2(m-p)}$ of the speed in theorem 1 is not always negligible compared to the other terms. However, it will be the case if we suppose that $m-p = (1+\epsilon) + (1-\epsilon) = 4(1+\epsilon)$.

(iv) This quantile estimator is quite useful in practice, specially for forecasting purpose (by conditional median or inter-quantiles intervals). From a computational point of view, several algorithms may be used: we have implemented in the R language an algorithm based on the Iterated Reweighted Least Square (IRLS). Note that even for real data cases, the curves are always observed in some discretization points, the regression splines is easy to implement by approximating inner products with quadrature rules. The IRLS algorithm (see Ruppert and Carroll, 1988, Lejeune and Sarda, 1988) allows to build conditional quantiles spline estimators and gives satisfactory forecast results. This algorithm has been used in particular on the "ORAMIP" ("Observatoire Regional de l'Air en Midi-Pyrenees") data to forecast pollution in the city of Toulouse (France): the results of this practical study are described in Cardot et al. (2004). We are interested in predicting the ozone concentration one day ahead, knowing the ozone curve (concentration along time) the day before. In that special case, conditional quantiles were also useful to predict an ozone threshold such that the probability to exceed this threshold is a given risk $1-\alpha$. In other words, it comes back to give an estimation of the α -quantile maximum ozone knowing the ozone curve the day before.

5 Proof of theorem 1

The proof of the result is based on the same kind of decomposition of b as the one used by He and Shi (1994). The main difference comes from the fact that our design matrix is ill-conditioned, which led us to add the penalization term treated using some arguments from Cardot et al. (2003).

Hypothesis (H 2) implies (see de Boor, 1978) that there exists a spline function $\tilde{f} = B_{k,q} f$, called spline approximation of f , such that

$$\sup_{t \in [0,1]} |f(t) - \tilde{f}(t)| \leq \frac{C_2}{k_n^p} \quad (8)$$

In what follows, we set $R_i = h^{-\frac{1}{p}} \tilde{f}(X_i)$; so we deduce from (8) and from hypothesis (H 1) that there exists a positive constant C_3 such that

$$\max_{i=1, \dots, n} |R_i| \leq \frac{C_3}{k_n^p}; \quad \text{as:} \quad (9)$$

The operator π_n allows to define the empirical version of the L^2 norm by $\|u\|_n^2 = h \sum_{i=1}^n u_i^2$. At first, we show the result (ii) of theorem 1 for the penalized empirical L^2 norm. Writing $b = (b^{(1)} + b^{(2)})$, we get

$$\begin{aligned} \|b\|_n^2 &= \|b^{(1)}\|_n^2 + \|b^{(2)}\|_n^2 \\ &= \frac{2}{n} \sum_{i=1}^n h b^{(1)}(X_i)^2 + \frac{2}{n} \sum_{i=1}^n h b^{(2)}(X_i)^2 \\ &\quad + 2 \|b^{(1)}\|_n^2 + 2 \|b^{(2)}\|_n^2. \end{aligned}$$

Now, using again hypothesis (H 1), we get almost surely and for all $i = 1, \dots, n$, the inequality $h b^{(2)}(X_i)^2 \leq C_0^2 C_2^2 = k_n^{2p}$. Moreover, lemma 8 of Stone (1985) gives us the existence of a positive constant C_4 that satisfies $\|b^{(2)}\|_n^2 \leq C_4 k_n^{2(m-p)}$. So we deduce

$$\begin{aligned} \|b\|_n^2 &= \|b^{(1)}\|_n^2 + \|b^{(2)}\|_n^2 \\ &= \frac{2}{n} \sum_{i=1}^n h b^{(1)}(X_i)^2 + 2 \|b^{(1)}\|_n^2 \\ &\quad + \frac{2C_0^2 C_2^2}{k_n^{2p}} + 2C_4 k_n^{2(m-p)}; \quad \text{as:} \end{aligned} \quad (10)$$

Our goal is now to compare our estimator \hat{b} with the spline approximation \tilde{f} . For that, we adopt the following transformation $\tilde{b} = \tilde{f}^{1/2} + \tilde{f}$. Then, we define on the set \mathcal{B}_n

$$f_i(\beta) = \frac{1}{n} \sum_{i=1}^n Y_i - \frac{1}{n} \sum_{i=1}^n \beta^T B_{k_{\mathcal{Q}_i}} \mathcal{C}^{1=2} + \frac{1}{2} \beta^T X_i X_i^T \beta + \frac{1}{2} \beta^T B_{k_{\mathcal{Q}_i}} \mathcal{C}^{1=2} + \frac{1}{2} \beta^T \mathcal{I}_{\mathcal{Q}_i} \beta :$$

We notice that minimizing $\sum_{i=1}^n f_i(\beta)$ comes back to the minimization of the criterion (5). We are interested by the behaviour of the function f_i around zero: $f_i(0)$ is the value of our loss criterion when $\beta = 0$. Let us also notice that the inverse of the matrix \mathcal{C} appears in the definition of f_i . This inverse exists on the set \mathcal{S}_n defined by (6), and which probability goes to 1 as n goes to infinity. Lemma 1 below, whose proof is given in section 5.1, allows us to get the results (i) and (ii) of theorem 1 for the penalized empirical L^2 norm.

Lemma 1 Under the hypotheses of theorem 1, for all $\epsilon > 0$, there exists $L = L(\epsilon)$ (sufficiently large) and $(n)_{n \geq N}$ with $n = \frac{p}{1 + (n - n_0) + \epsilon^2} = (k_n - n_0)$ such that, for n large enough

$$\inf_{\beta \in \mathcal{S}_n} \sum_{i=1}^n f_i(\beta) > \sum_{i=1}^n f_i(0) > 1 - \epsilon :$$

Using convexity arguments, this inequality means that the solution $\hat{\beta}$ exists and is unique on the ball centered in 0 and of radius L_n . As we use the one-to-one transformation $\beta = \mathcal{C}^{1=2} + \gamma$ on the set \mathcal{S}_n , we deduce the existence and the uniqueness of the solution of (5) on the set \mathcal{S}_n , which proves point (i) of theorem 1.

Now, let ϵ be strictly positive; using the convexity of function f_i , there exists $L = L(\epsilon)$ such that, for n large enough

$$\inf_{\beta \in \mathcal{S}_n} \sum_{i=1}^n f_i(\beta) > \sum_{i=1}^n f_i(0) > 1 - \epsilon : \quad (11)$$

On the other hand, using the definition of f_i and the minimization criterion (5), we have

$$\begin{aligned}
& \frac{1}{n} \sum_{i=1}^n f_i(\mathcal{C}^{1=2} b - \mathcal{C}^{1=2} ?) \\
& = \inf_{2\mathbb{R}^{k+q}} \frac{1}{n} \sum_{i=1}^n \|Y_i - h_{B_{k,q}}; X_i\|^2 + B_{k,q}^{(m)^2};
\end{aligned}$$

so we finally get

$$\frac{1}{n} \sum_{i=1}^n f_i(\mathcal{C}^{1=2} b - \mathcal{C}^{1=2} ?) - \frac{1}{n} \sum_{i=1}^n f_i(0):$$

Then, combining this with equation (11), we obtain

$$\mathbb{P} \left(\inf_{j \in L_n} \frac{1}{n} \sum_{i=1}^n f_i(j) > \frac{1}{n} \sum_{i=1}^n f_i(\mathcal{C}^{1=2} b - \mathcal{C}^{1=2} ?) > 1 \right) : \quad (12)$$

Now, using the definition of \mathcal{C} , we have

$$\begin{aligned}
& \mathbb{P} \left(\frac{1}{n} \sum_{i=1}^n \|h_{B_{k,q}}; X_i\|^2 + (b - ?)^{(m)^2} \geq L_n^2 \right) \\
& = 1 - \mathbb{P} \left(\mathcal{C}^{1=2} (b - ?) > L_n \right) \\
& \mathbb{P} \left(\inf_{j \in L_n} \frac{1}{n} \sum_{i=1}^n f_i(j) > \frac{1}{n} \sum_{i=1}^n f_i(\mathcal{C}^{1=2} b - \mathcal{C}^{1=2} ?) \right) :
\end{aligned}$$

With relation (12), this last probability is greater than 1, so we obtain

$$\frac{1}{n} \sum_{i=1}^n \|h_{B_{k,q}}; X_i\|^2 + (b - ?)^{(m)^2} = O_P \left(\frac{1}{n} \right) = O_P \left(\frac{1}{n} + \frac{1}{k_n} \right) :$$

This last result, combined with inequality (10) finally gives us the equivalent of result (ii) for the penalized empirical L^2 norm. Point (ii) (with the norm $k \cdot k_2$) then follows from lemma 2 below, which is proved in section 5.5, and achieves the proof of theorem 1 (ii).

Lemma 2 Let f and g be two functions supposed to be m times differentiable and such that

$$\|f - g\|_n^2 + \|k(f - g)^{(m)}\|^2 = O_P(u_n);$$

with u_n going to zero when n goes to infinity. Under hypotheses (H 1) and (H 3) and if moreover $\|g\|$ and $\|g^{(m)}\|$ are supposed to be bounded, we have

$$\|f - g\|_n^2 = O_P(u_n):$$

5.1 Proof of lemma 1

This proof is based on three preliminary lemmas, proved respectively in sections 5.2, 5.3 and 5.4. We denote by T_n the set of the random variables $(X_1; \dots; X_n)$. Under hypotheses of theorem 1, we have the following results.

Lemma 3 There exists a constant C_5 such that, on the set Ω_n defined by (6), we have

$$\max_{i=1, \dots, n} \|B_{k_n}^{(1=2)}(X_i)\| \leq \frac{C_5 j}{k_n n}; \text{ as:}$$

Lemma 4 For all $\epsilon > 0$, there exists $L = L_\epsilon$ such that

$$\lim_{n \rightarrow \infty} P \left(\inf_{j \neq 1} \sup_{i=1}^{X^n} (f_i(L_n) - f_i(0) - E[f_i(L_n) - f_i(0) | T_n]) > \frac{\epsilon}{n} \right) = 0:$$

Lemma 5 For all $\epsilon > 0$, there exists $L = L_\epsilon$ such that

$$P \left(\inf_{j \neq 1} \sup_{i=1}^{X^n} E[f_i(L_n) - f_i(0) | T_n] > \frac{\epsilon}{n} \right) > 1 - \epsilon:$$

These three lemmas allow us to prove lemma 1. Indeed, let L be a strictly positive real number; we make the following decomposition

$$\inf_{j \neq 1} \sup_{i=1}^{X^n} f_i(L_n) - \inf_{i=1}^{X^n} f_i(0) = A_n + B_n;$$

with

$$A_n = \inf_{j \neq 1} \sup_{i=1}^{X^n} (f_i(L_n) - f_i(0) - E[f_i(L_n) - f_i(0)]T_n)$$

and

$$B_n = \inf_{j \neq 1} \sup_{i=1}^{X^n} E[f_i(L_n) - f_i(0)]T_n:$$

Using lemma 4 and 5, we can find L sufficiently large such that, for n large enough

$$P(A_n > \frac{2}{n}) < \frac{1}{2};$$

and

$$P(B_n > \frac{2}{n}) > 1 - \frac{1}{2};$$

thus we get

$$P\left(\inf_{j \neq 1} \sup_{i=1}^{X^n} f_i(L_n) - \sup_{i=1}^{X^n} f_i(0) > 0\right) > 1 - \frac{1}{2};$$

which achieves the proof of lemma 1.

5.2 Proof of lemma 3

Using lemma 6.2 of Cardot et al. (2003), we have

$$\min_{\mathcal{C}} \|\mathcal{C}\|_5 \leq C_5^0 n + o_p((k_n^2 n^1)^{1/2}):$$

Noticing that $\|B_{k,q}\|_5^{1/2} \|X_{i1}\|^2 = \|B_{k,q}\|_5^{1/2} \|X_{i1}\| \|B_{k,q}\|_5^{1/2} \|X_{i1}\|$, we deduce that

$$\|B_{k,q}\|_5^{1/2} \|X_{i1}\|^2 = \|B_{k,q}\|_5^{1/2} \|X_{i1}\| \|B_{k,q}\|_5^{1/2} \|X_{i1}\| \leq \frac{1}{C_5^0 n} + o_p((k_n^2 n^1)^{1/2});$$

which gives us $\mathbb{E} h_{k,q}^{(1=2)}(X_i) = C_5^{(0)} j^2 = (k_n - n) + o_p(n^{(1-1)=2})$ almost surely, and achieves the proof of lemma 3.

5.3 Proof of lemma 4

Considering the definition of functions f_i and l , we have

$$\begin{aligned} & \sup_{j=1}^{X^n} f_i(L_n) - f_i(0) = \mathbb{E} [f_i(L_n) - f_i(0) | \mathcal{F}_n] \\ & = \sup_{j=1}^{X^n} \mathbb{E} [h_{k,q}^{(1=2)}(X_i) - R_i - j_i - R_i | \mathcal{F}_n] \\ & = \mathbb{E} [h_{k,q}^{(1=2)}(X_i) - R_i - j_i - R_i | \mathcal{F}_n] ; \end{aligned}$$

where $1, \dots, n$ are n real random variables independent and identically distributed defined by $Y_i = h_{k,q}^{(1=2)}(X_i)$ for all $i = 1, \dots, n$. Let us also denote $l_i(\cdot) = h_{k,q}^{(1=2)}(X_i) - R_i - j_i - R_i$. To prove lemma 4, it suffices to show that, for all $\epsilon > 0$, there exists $L = L(\epsilon)$ such that

$$\lim_{n \rightarrow \infty} P \left(\sup_{j=1}^{X^n} [l_i(\cdot) - \mathbb{E}(l_i(\cdot) | \mathcal{F}_n)] > \frac{\epsilon}{n} \right) = 0 ;$$

Let ϵ be a real number strictly positive and C the subset of \mathbb{R}^{k+q} defined by $C = \{x \in \mathbb{R}^{k+q} : |x| \leq 1\}$. As C is a compact set, we can cover it with open balls, that is to say $C = \bigcup_{j=1}^{K_n} C_j$ with K_n chosen, for all j from 1 to K_n , such that

$$\text{diam}(C_j) \leq \frac{\epsilon}{8C_5 L} ; \quad (13)$$

Hence

$$K_n \leq \frac{8C_5 L}{\epsilon} n^{k+q} ; \quad (14)$$

Now, for $1 \leq j \leq K_n$, let x_j be in C_j ; using the definition of $l_i(\cdot)$ and the triangular inequality, we have

$$\begin{aligned} & \min_{j=1, \dots, K_n} \sum_{i=1}^{X^n} [\mathbb{I}_{i(j)} - \mathbb{E}(\mathbb{I}_{i(j)})] - \mathbb{I}_{i(j)} - \mathbb{E}(\mathbb{I}_{i(j)}) \\ & 2L_n \min_{j=1, \dots, K_n} \sum_{i=1}^{X^n} \mathbb{B}_{k_{j1}}^{\mathbb{C}^{1=2}}(\mathbb{I}_{i(j)}; X_{i1}) : \end{aligned}$$

Then, using lemma 3, we get

$$\begin{aligned} & \min_{j=1, \dots, K_n} \sum_{i=1}^{X^n} [\mathbb{I}_{i(j)} - \mathbb{E}(\mathbb{I}_{i(j)})] - \mathbb{I}_{i(j)} - \mathbb{E}(\mathbb{I}_{i(j)}) \\ & 2L_n \mathbb{P}^{\frac{C_5 n}{k_{n,n}}} \min_{j=1, \dots, K_n} j - j \end{aligned}$$

this last inequality being true only on the set Ω_n defined by (6). Moreover, there exists a unique $j_0 \in \{1, \dots, K_n\}$ such that $\Omega_n \subset C_{j_0}$, which gives us with relation (13)

$$\begin{aligned} & \min_{j=1, \dots, K_n} \sum_{i=1}^{X^n} [\mathbb{I}_{i(j)} - \mathbb{E}(\mathbb{I}_{i(j)})] - \mathbb{I}_{i(j)} - \mathbb{E}(\mathbb{I}_{i(j)}) \leq -\frac{1}{4} n : \\ & (15) \end{aligned}$$

On the other hand, we have

$$\sup_{2C} j - \mathbb{I}_{i(j)} \leq L_n \sup_{2C} j \mathbb{B}_{k_{j1}}^{\mathbb{C}^{1=2}}(\mathbb{I}_{i(j)}; X_{i1})$$

and using lemma 3 again, we get, on Ω_n ,

$$\sup_{2C} j - \mathbb{I}_{i(j)} \leq \mathbb{P}^{\frac{C_5 L_n}{k_{n,n}}} : \quad (16)$$

Besides, for Ω_n fixed in C , with the same arguments as before, if we denote by $\mathcal{T}^?$ the set of the random variables (X_1, \dots, X_n, \dots) , we have

$$\begin{aligned} & \sum_{i=1}^{X^n} \text{Var}(\mathbb{I}_{i(j)} | \mathcal{T}^?) \leq \sum_{i=1}^{X^n} L_n^2 \text{Var}(j \mathbb{B}_{k_{j1}}^{\mathbb{C}^{1=2}}(\mathbb{I}_{i(j)}; X_{i1}) | \mathcal{T}^?) : \end{aligned}$$

Then, using the definition of \mathcal{B} , we remark that

$$\sum_{i=1}^n \mathbb{E}_{k,q} \mathcal{B}^{1=2} ; X_i^2 = n j^2 \quad \mathcal{B}^{1=2} G_k \mathcal{B}^{1=2} ; \quad (17)$$

which gives us

$$\sum_{i=1}^n \text{Var} (\mathbb{I}_n) \leq n L^2 : \quad (18)$$

We are now able to prove lemma 4. Using first relation (15), we have

$$\begin{aligned} & \mathbb{P} \left(\sum_{j=1}^n [\mathbb{I}_n - \mathbb{E} (\mathbb{I}_n)] > \sqrt{n} \right) \\ & \leq \mathbb{P} \left(\max_{j=1, \dots, K_n} \sum_{i=1}^n (\mathbb{I}_n - \mathbb{E} (\mathbb{I}_n)) > \sqrt{n} \right) \end{aligned}$$

and then

$$\begin{aligned} & \mathbb{P} \left(\sum_{j=1}^n [\mathbb{I}_n - \mathbb{E} (\mathbb{I}_n)] > \sqrt{n} \right) \\ & \leq \mathbb{P} \left(\sum_{i=1}^n (\mathbb{I}_n - \mathbb{E} (\mathbb{I}_n)) > \sqrt{n} \right) \end{aligned}$$

By inequalities (16) and (18), we apply Bernstein inequality (see Uspensky, 1937) and inequality (14) to obtain

$$\begin{aligned} & \mathbb{P} \left(\sum_{j=1}^n [\mathbb{I}_n - \mathbb{E} (\mathbb{I}_n)] > \sqrt{n} \right) \\ & \leq 2 \exp \left(- \frac{8 C_5 L n}{n k_n} \right) \frac{2 \sqrt{n} n^2 = 4}{2 n L^2 + 2 C_5 L n} \mathbb{P} \left(\frac{1}{k_n} \right) : \end{aligned}$$

This bound does not depend on the sample $T = (X_1, \dots, X_n, \dots)$, hence, if we take the expectation on both sides of this inequality above, we deduce

$$P \sup_{i=1}^{X^n} [i(\cdot) - E(i(\cdot) | \mathcal{F}_n)] > \frac{2}{n} \setminus \frac{\#}{n} \\ 2 \exp \frac{\frac{2}{n} \frac{2^p \overline{k_n n}}{8L^2 \frac{2^p \overline{k_n n}}{k_n n} + 4C_5 L_n}}{1 - \frac{(k_n + q)(8L^2 \frac{2^p \overline{k_n n}}{k_n n} + 4C_5 L_n)}{\frac{2}{n} \frac{2^p \overline{k_n n}}{k_n n}}} \ln \frac{8C_5 L_n}{\frac{2}{n} \frac{2^p \overline{k_n n}}{k_n n}} :$$

If we $xL = L_n = \frac{2^p \overline{k_n n}}{nk_n n^2}$, we have

$$\frac{\frac{2}{n} \frac{2^p \overline{k_n n}}{L^2 \frac{2^p \overline{k_n n}}{k_n n}}}{\frac{2}{n} \frac{2^p \overline{k_n n}}{k_n n}} = \frac{1}{k_n n^{n!+1}} + 1 ,$$

$$\frac{\frac{2}{n} \frac{2^p \overline{k_n n}}{L_n}}{\frac{2}{n} \frac{2^p \overline{k_n n}}{k_n n}} = \frac{2^p \overline{k_n n}}{n^{n!+1}} + 1 ,$$

$$\frac{\frac{k_n L^2 \frac{2^p \overline{k_n n}}{k_n n}}{\frac{2}{n} \frac{2^p \overline{k_n n}}{k_n n}}}{\frac{2}{n} \frac{2^p \overline{k_n n}}{k_n n}} = k_n^2 n^{n!+1} ! 0 ,$$

$$\frac{\frac{k_n L_n}{\frac{2}{n} \frac{2^p \overline{k_n n}}{k_n n}}}{\frac{2}{n} \frac{2^p \overline{k_n n}}{k_n n}} = \frac{k_n}{n^{n!+1}} ! 0 .$$

This leads to

$$\lim_{n!+1} P \sup_{j \geq 1} \sup_{i=1}^{X^n} [i(\cdot) - E(i(\cdot) | \mathcal{F}_n)] > \frac{2}{n} \setminus \frac{\#}{n} = 0 ;$$

and with the fact that $\frac{\#}{n}$ has probability tending to 1 when n goes to infinity, we finally obtain

$$\lim_{n!+1} P \sup_{j \geq 1} \sup_{i=1}^{X^n} [i(\cdot) - E(i(\cdot) | \mathcal{F}_n)] > \frac{2}{n} = 0 ;$$

which achieves the proof of lemma 4.

5.4 Proof of lemma 5

Let a and b be two real numbers. We denote by F_i the random repartition function of i given T_n and by f_i the random density function of i given T_n . As $E(l(i+b)J_n) = \int_{\mathbb{R}} l(s+b) dF_i(s)$, we obtain, using a Taylor linearization at first order, the existence of a quantity r_{iab} such that

$$E(l(i+a+b) - l(i+b)J_n) = f_i(0)a^2 + 2f_i(0)ab + \left(\frac{a^2}{2} + ab\right)r_{iab};$$

with $r_{iab} \rightarrow 0$ when $a, b \rightarrow 0$. If we set $L^0 = \frac{P}{2}L$ and $R_i^0 = \frac{P}{2}R_i$, this relation gives us

$$\begin{aligned} & \sum_{i=1}^n \int_{\mathbb{R}} l(i) L_n \mathcal{H}_{k_{\mathcal{P}}} \mathcal{C}^{1=2}; X_i i R_i - l(i - R_i) J_n \\ &= \frac{1}{2} \sum_{i=1}^n f_i(0) L_n^2 \mathcal{H}_{k_{\mathcal{P}}} \mathcal{C}^{1=2}; X_i i^2 + L_n^0 \mathcal{H}_{k_{\mathcal{P}}} \mathcal{C}^{1=2}; X_i i R_i^0 \\ &+ \sum_{i=1}^n \int_{\mathbb{R}} L_n^2 \mathcal{H}_{k_{\mathcal{P}}} \mathcal{C}^{1=2}; X_i i^2 + L_n^0 \mathcal{H}_{k_{\mathcal{P}}} \mathcal{C}^{1=2}; X_i i R_i^0 r_i; \quad (19) \end{aligned}$$

with $r_i \rightarrow 0$. Considering such that $j \neq 1$, we have, using relation (9)

$$\begin{aligned} & L_n^2 \mathcal{H}_{k_{\mathcal{P}}} \mathcal{C}^{1=2}; X_i i^2 + L_n^0 \mathcal{H}_{k_{\mathcal{P}}} \mathcal{C}^{1=2}; X_i i R_i^0 \\ & \frac{1}{2} L_n^2 \mathcal{H}_{k_{\mathcal{P}}} \mathcal{C}^{1=2}; X_i i^2 - \frac{C_3^2}{k_n^{2p}}; \quad \text{as:} \quad (20) \end{aligned}$$

Moreover, if we set $V_n = \sup_{j \neq 1} \max_{i=1 \dots n} |r_i|$, then with condition (H.4) $\mathbb{1}_{F_{V_n} < m} \int_{\mathbb{R}} f_i(0) = 4g = \mathbb{1}_{\mathbb{R}}$ for n large enough, and

$$\begin{aligned} & \int_{\mathbb{R}} L_n^2 \mathcal{H}_{k_{\mathcal{P}}} \mathcal{C}^{1=2}; X_i i^2 + L_n^0 \mathcal{H}_{k_{\mathcal{P}}} \mathcal{C}^{1=2}; X_i i R_i^0 r_i \\ & \frac{m \int_{\mathbb{R}} f_i(0)}{4} L_n^2 \mathcal{H}_{k_{\mathcal{P}}} \mathcal{C}^{1=2}; X_i i^2 + L_n^0 \mathcal{H}_{k_{\mathcal{P}}} \mathcal{C}^{1=2}; X_i i R_i^0 \\ & 2m \int_{\mathbb{R}} f_i(0) \frac{3}{16} L_n^2 \mathcal{H}_{k_{\mathcal{P}}} \mathcal{C}^{1=2}; X_i i^2 + \frac{C_3^2}{8k_n^{2p}} : \quad (21) \end{aligned}$$

Using inequalities (20) and (21), relation (19) becomes then

$$\begin{aligned} & \frac{1}{2n} \inf_{j=1}^{X^n} E [f_i(L_n) - f_i(0)] \mathbb{I}_n] \\ & \leq \frac{1}{2n} \inf_{j=1}^{X^n} E [f_i(L_n) - f_i(0)] \mathbb{I}_n] \\ & \leq \frac{1}{2n} \inf_{j=1}^{X^n} E [f_i(L_n) - f_i(0)] \mathbb{I}_n] \\ & \leq \frac{1}{2n} \inf_{j=1}^{X^n} E [f_i(L_n) - f_i(0)] \mathbb{I}_n] \end{aligned}$$

Now, we come back to the definition of function f_i to obtain

$$\begin{aligned} & \frac{1}{2n} \inf_{j=1}^{X^n} E [f_i(L_n) - f_i(0)] \mathbb{I}_n] \\ & \leq \frac{1}{2n} \inf_{j=1}^{X^n} E [f_i(L_n) - f_i(0)] \mathbb{I}_n] \\ & \leq \frac{1}{2n} \inf_{j=1}^{X^n} E [f_i(L_n) - f_i(0)] \mathbb{I}_n] \\ & \leq \frac{1}{2n} \inf_{j=1}^{X^n} E [f_i(L_n) - f_i(0)] \mathbb{I}_n] \end{aligned}$$

Reminding that $L^2 = 2L^2$ and taking $\epsilon = \min(\frac{5}{4} \min f_i(0); 1)$, we have $\epsilon > 0$ by hypothesis (H 4) and then

$$\begin{aligned} & \frac{1}{2n} \inf_{j=1}^{X^n} E [f_i(L_n) - f_i(0)] \mathbb{I}_n] \\ & \leq \frac{1}{2n} \inf_{j=1}^{X^n} E [f_i(L_n) - f_i(0)] \mathbb{I}_n] \\ & \leq \frac{1}{2n} \inf_{j=1}^{X^n} E [f_i(L_n) - f_i(0)] \mathbb{I}_n] \\ & \leq \frac{1}{2n} \inf_{j=1}^{X^n} E [f_i(L_n) - f_i(0)] \mathbb{I}_n] \end{aligned}$$

Using relation (17), we get

$$\begin{aligned} & \frac{1}{2n} \inf_{j=1}^{X^n} E [f_i(L_n) - f_i(0)] \mathbb{I}_n] \\ & \leq \frac{1}{2n} \inf_{j=1}^{X^n} E [f_i(L_n) - f_i(0)] \mathbb{I}_n] \\ & \leq \frac{1}{2n} \inf_{j=1}^{X^n} E [f_i(L_n) - f_i(0)] \mathbb{I}_n] \\ & \leq \frac{1}{2n} \inf_{j=1}^{X^n} E [f_i(L_n) - f_i(0)] \mathbb{I}_n] \end{aligned}$$

Moreover, for $j = 1$, the minimum of $B_{k,q}^{(1)=2}(\eta)$; $B_{k,q}^{(1)=2}(\eta)$ is obtained for $\eta = \eta^{(1)=2} = j$. Using the fact that the spline approximation has a bounded m -th derivative, we deduce the existence of a constant $C_9 > 0$ such that

$$\inf_{j=1}^n h_{k,q}^{(m)} = 2^{(m)}; \quad B_{k,q}^{(m)} \leq \frac{C_9}{n};$$

hence we obtain

$$\frac{1}{2^n} \inf_{j=1}^n \mathbb{E} [f_i(L_n) - f_i(0) | \mathcal{F}_n]$$

$$L^2 \quad \frac{9}{4} m \inf_i f_i(0) \frac{C_3^2}{k_n^{2p} 2^n} \quad 2C_9 \frac{L}{p_n};$$

that is to say

$$\frac{1}{2n} \inf_{j=1}^{X^n} E [f_i(L_n) - f_i(0) | \mathcal{F}_n] \leq \frac{9m \inf_i f_i(0) C_3^2}{4 L^2 k_n^{2p} 2_n^2} - \frac{2C_9}{L_n^p} : \quad$$

Reminding that we have fixed $L = L_n = \frac{p}{nk_n n^2}$, we get

$$\text{for } \frac{1}{n_n}; \text{ we have } \frac{1}{L^2 k_n^{2p}} \frac{k_n}{n^4 k_n^{2p}} \frac{1}{n! + 1} \rightarrow 0,$$

$$\text{for } \frac{2}{n} \leq \frac{p}{k_n}; \text{ we have } \frac{p}{L_n} \leq \frac{p}{k_n} \leq \frac{p}{n!+1} \rightarrow 0.$$

This leads to

$$\lim_{n \rightarrow \infty} P \left(\frac{1}{2n} \inf_{j \neq 1} \sup_{i=1}^{X^n} E [f_i(L_n) - f_i(0)] T_n > 1 \right) = 0;$$

which achieves the proof of lemma 5.

5.5 Proof of lemma 2

Writing $x = (x_n) + n$, we make the following decomposition

$$kf - gk_n^2 = 2k_{x_n}k kfk^2 + kgk^2 + kf - gk_n^2: \quad (22)$$

Now, let us decompose f as follows $f = P + R$ with $P(t) = \sum_{l=0}^{m-1} \frac{t^l}{l!} f^{(l)}(0)$ and $R(t) = \int_0^t \frac{(t-u)^{m-1}}{(m-1)!} f^{(m)}(u) du$. P belongs to the space P_{m-1} of polynomials of degree at most $m-1$, whose dimension is finite and equal to m . Using hypothesis (H3), there exists a constant $C_6 > 0$ such that we have $kPk^2 \leq C_6 kP k_n^2$. Then, we can deduce

$$\begin{aligned} kfk^2 &= 2kPk^2 + 2kRk^2 \\ &\leq 2C_6 kPk_n^2 + 2kRk^2 \\ &= 4C_6 kfk_n^2 + 4C_6 k_{x_n}k kRk^2 + 2kRk^2: \end{aligned} \quad (23)$$

As n is a bounded operator (by hypothesis (H1)), there exists a constant $C_7 > 0$ such that we have $k_{x_n}k \leq C_7$. Moreover, under Cauchy-Schwarz inequality, there exists a constant $C_8 > 0$ such that $kRk^2 \leq C_8 kf^{(m)}k^2$. Relation (23) gives $kfk^2 \leq 4C_6 kfk_n^2 + (4C_6 C_7 + 2) C_8 kf^{(m)}k^2$. Then, if we write $f = (f - g) + g$, we finally deduce

$$\begin{aligned} kfk^2 &\leq 8C_6 kf - gk_n^2 + (8C_6 C_7 + 4) C_8 k(f - g)^{(m)}k^2 \\ &\quad + 8C_6 k_{x_n}k kgk^2 + (8C_6 C_7 + 4) C_8 kg^{(m)}k^2: \end{aligned} \quad (24)$$

We have supposed that kgk and $kg^{(m)}k$ are bounded, so

$$8C_6 k_{x_n}k kgk^2 + (8C_6 C_7 + 4) C_8 kg^{(m)}k^2 = O(1);$$

and the hypothesis $kf - gk_n^2 + k(f - g)^{(m)}k^2 = O_P(u_n)$ gives us the bounds $kf - gk_n^2 = O_P(u_n)$ and $k(f - g)^{(m)}k^2 = O_P(u_n)$. Then, relation (24) becomes

$$\|f\|_2^2 = O_p\left(1 + \frac{u_n}{n}\right) : \quad (25)$$

Finally, we have $\|f - g\|_2 = o_p(n^{-(1/2)}) = o_p(n^{-1/2})$ from lemma 5.3 of Cardot et al. (1999). This equality, combined with equations (22) and (25) gives us $\|f - g\|_2^2 = O_p(u_n)$, which is the announced result.

Acknowledgements : We would like to thank the referees for a careful reading of a previous version of the manuscript that permits to improve some results as well as the participants to the working group "staph" in Toulouse for stimulating discussions.

Bibliography

- Bassett, G.W. and Koenker, R.W. (1978). Asymptotic Theory of Least Absolute Error Regression. *J. Amer. Statist. Assoc.*, 73, 618-622.
- Besse, P.C. and Ramsay, J.O. (1986). Principal Component Analysis of Sampled Curves. *Psychometrika*, 51, 285-311.
- Bhattacharya, P.K. and Gangopadhyay, A.K. (1990). Kernel and Nearest-Neighbor Estimation of a Conditional Quantile. *Ann. Statist.*, 18, 1400-1415.
- Bosq, D. (2000). Linear Processes in Function Spaces. *Lecture Notes in Statistics*, 149, Springer-Verlag.
- Cardot, H., Ferraty, F. and Sarda, P. (1999). The Functional Linear Model. *Stat. and Prob. Letters*, 45, 11-22.
- Cardot, H., Ferraty, F. and Sarda, P. (2003). Spline Estimators for the Functional Linear Model. *Statistica Sinica*, 13, 571-591.
- Cardot, H., Crambes, C. and Sarda, P. (2004). Conditional Quantiles with Functional Covariates: an Application to Ozone Pollution Forecasting. In *Comstat 2004 Proceedings*, ed. J. Antoch, Physica-Verlag, 769-776.

- Dauxois, J. and Pousse, A. (1976). Les Analyses Factorielles en Calcul des Probabilités et en Statistique: Essai d'étude synthétique (in French). These, Université Paul Sabatier, Toulouse, France.
- Dauxois, J., Pousse, A. and Roman, Y. (1982). A Symptotic Theory for the Principal Component Analysis of a Random Vector Function: some Applications to Statistical Inference. *J. of Mult. Analysis*, 12, 136-154.
- Déville, J.C. (1974). Méthodes Statistiques et Numériques de l'Analyse Harmonique (in French). *Ann. Insee*, 15.
- de Boor, C. (1978). A Practical Guide to Splines. Springer, New York.
- Fan, J., Hu, T.C. and Truong, Y.K. (1994). Robust Nonparametric Function Estimation. *Scand. J. Statist*, 21, 433-446.
- Ferraty, F. and Vieu, P. (2002). The Functional Nonparametric Model and Application to Spectrometric Data. *Computational Statistics*, 17, 545-564.
- He, X. and Shi, P. (1994). Convergence Rate of B-Spline Estimators of Nonparametric Conditional Quantile Functions. *Nonparametric Statistics*, 3, 299-308.
- Koenker, R.W. and Bassett G.W. (1978). Regression Quantiles. *Econometrica*, 46, 33-50.
- Koenker, R.W., Ng, P. and Portnoy, S. (1994). Quantile Smoothing Splines. *Biometrika*, 81, 673-680.
- Lejeune, M. and Sarda, P. (1988). Quantile Regression: A Nonparametric Approach. *Computational Statistics and Data Analysis*, 6, 229-239.
- Ramsey, J.O. and Silverman, B.W. (1997). Functional Data Analysis. Springer-Verlag.

- Ram say, J.O . and Silverm an, B W . (2002). Applied Functional Data Analysis. Springer-Verlag.
- Ruppert, D . and Carroll, R J. (1988). Transformation and Weighting in Regression. Chapman and Hall.
- Stone, C . (1982). Optimal Rates of Convergence for Nonparametric Models. Ann. Statist., 10, 1040-1053.
- Stone, C . (1985). Additive Regression and other Nonparametric Models. Ann. Statist., 13, 689-705.
- Tsybakov, A .B . (1986). Robust Reconstruction of Functions by the Local-Approximation Method. Problems of Information Transmission, 22, 133-146.
- Uspensky, J.V . (1937). Introduction to Mathematical Probability. New York and London.
- Weinberger, H F . (1974). Variational Methods for Eigenvalue Approximation. SIAM , Philadelphia.