

Integral Operators on Spaces of Continuous Vector-valued functions

by

Paulette Saab

Abstract Let X be a compact Hausdorff space, let E be a Banach space, and let $C(X; E)$ stand for the Banach space of E -valued continuous functions on X under the uniform norm. In this paper we characterize Integral operators (in the sense of Grothendieck) on $C(X; E)$ spaces in terms of their representing vector measures. This is then used to give some applications to Nuclear operators on $C(X; E)$ spaces.

AM S (M O S) subject Classification (1980). Primary 46E 40, 46G 10; Secondary 28B 05, 28B 20.

Supported in part by an NSF Grant DM S-87500750

Introduction Let X be a compact Hausdorff space, let E and F be Banach spaces. Denote by $C(X; E)$ the space of all continuous E -valued functions defined on X under the uniform norm. In [9] C. Swartz showed that a bounded linear operator $T: C(X; E) \rightarrow F$ with representing measure G is absolutely summing if and only if each of the values of G is an absolutely summing operator from E to F and G is of bounded variation as a measure taking its values in the space of absolutely summing operators from E to F equipped with the absolutely summing norm. In this paper we shall extend Swartz's result to the class of (Grothendieck) integral operators on $C(X; E)$ spaces. More precisely we shall show that a bounded linear operator $T: C(X; E) \rightarrow F$ with representing measure G is an integral operator if and only if each of the values of G is an integral operator from E to F and G is of bounded variation as a vector measure taking its values in the space of integral operators from E to F equipped with the integral norm. This result is then used to give some applications to Nuclear operators on $C(X; E)$ spaces.

I. Preliminaries If X is a compact Hausdorff space and E is a Banach space, then $C(X; E)$ will denote the Banach space of all continuous E -valued functions equipped with the uniform norm. It is well known [4, page 182] that the dual of $C(X; E)$ is isometrically isomorphic to the space $M(X; E)$ of all regular E -valued measures on X that are of bounded variation. When E is the scalar field, we will simply write $C(X)$ and $M(X)$ for $C(X; E)$ and $M(X; E)$. If $\mu \in M(X; E)$ and $e \in E$, we will denote by $\|\mu\|$ the variation of μ and by μe the element of $M(X)$ defined on each Borel subset B of X by

$$\mu e(B) = \mu(B)(e):$$

The duality between $M(X; E)$ and $C(X; E)$ is then defined as follows: for each $f \in C(X)$ and $e \in E$

$$(f, \mu e) = \int_X f d\mu e$$

where $f \mu e$ is the element of $C(X; E)$ defined by

$$f \mu e(x) = f(x)e \text{ for all } x \in X:$$

If B is a Borel subset of X , then 1_B will denote the characteristic function of B , and if $e \in E$ we let $1_B e$ denote the element of $C(X; E)$ defined by

$$1_B e(x) = \begin{cases} e & \text{if } x \in B \\ 0 & \text{if } x \notin B \end{cases}$$

for each $e \in E$.

If X is a compact Hausdorff space, E and F are Banach spaces, every bounded linear operator $T: C(X; E) \rightarrow F$ has a representing measure G . The measure G is defined on the σ -field of Borel subsets of X and takes its values in $L(E; F)$, the space of all bounded linear operators from E to F . The measure G is such that for each Borel subset B of X and for each $e \in E$

$$G(B)e = T(1_B e)$$

For $y \in F$, if we denote by G_y the E -valued measure on X such that for each Borel subset B of X and each $e \in E$

$$\langle y, G_y(B)e \rangle = \langle y, T(1_B e) \rangle$$

then G_y is the unique element of $M(X; E)$ that represents T_y in the sense that for each $f \in C(X; E)$

$$\langle y, T f \rangle = \int_X \langle y, f(x) \rangle dG_y(x)$$

If E and F are Banach spaces, we denote by $E \otimes F$ the algebraic tensor product of E and F endowed with the norm $\| \cdot \|$

$$\| \sum_{i=1}^n x_i \otimes y_i \| = \sup \left\{ \left\| \sum_{i=1}^n x_i \otimes y_i \right\| : \sum_{i=1}^n \| y_i \|^2 \leq 1 \right\}$$

The completion $E \hat{\otimes} F$ of $E \otimes F$ is called the injective tensor product of E and F .

A bounded linear operator $T: E \rightarrow F$ from a Banach space E into a Banach space F is said to be an integral operator if the bilinear form on $E \otimes F$ defined by $\langle e \otimes y, T e \rangle = \langle y, T e \rangle$ for $e \in E$ and $y \in F$ determines an element of $(E \hat{\otimes} F)^*$; the dual of the injective tensor product $E \hat{\otimes} F$ of the Banach spaces E and F . The integral norm of T , which we will denote by $\| T \|_{\text{int}}$, is just the norm of the bounded linear functional

induced by μ as an element of the dual space $(E' \hat{\otimes} F')$. Hence if $T : E \rightarrow F$ is an integral operator, then

$$\|T\|_{\text{int}} = \sup \left\{ \left\| \sum_{i=1}^n y_i (T e_i) \right\| : \sum_{i=1}^n \|e_i\|_{E'}^2 \leq 1 \right\}.$$

Finally, $\mathcal{I}(E; F)$ will stand for the Banach space of all integral operators from E to F equipped with the integral norm. For all undefined notions and notations we refer the reader to [3], [4] or [5].

One of the most useful tools in the study of an integral operators T between two Banach spaces E and F is its representation by a finite regular Borel scalar measure on the compact $B(E') \times B(F')$, the product of the closed unit balls of E' and F' equipped with their weak* topologies. This, of course, goes back to Grothendieck [6] and can be used to characterize integral operators by the following characterization that we will state and prove before proving the main result.

Proposition 1 A bounded linear operator $T : E \rightarrow F$ between two Banach spaces is an integral operator if and only if there exists a regular F' -valued vector measure m of bounded variation defined on the σ -field of Borel subsets of the closed unit ball $B(E')$, such that for each $e \in E$

$$Te = \int_{B(E')} e(e) dm(e).$$

In this case m can be chosen so that $\|T\|_{\text{int}} = \|m\|(B(E'))$:

Proof: Assume that there exists a regular F' -valued vector measure m defined on the σ -field of Borel subsets of $B(E')$ with $\|m\|(B(E')) < 1$ and such that for each $e \in E$

$$Te = \int_{B(E')} e(e) dm(e).$$

This in particular shows that the operator $J \circ T : E \rightarrow F'$, where J denotes the natural embedding of F into F' , has an extension \hat{T} to an integral operator from $C(B(E'))$ to F' . Hence T is an integral operator [4, page 233] and

$$\|T\|_{\text{int}} = \|\hat{T}\|_{\text{int}} = \|m\|(B(E')):$$

Conversely, suppose that $T: E \rightarrow F$ is an integral operator, then it follows from [4, page 231] that there exists a regular Borel measure on $B(E) \times B(F)$ such that for each $e \in E$ and $y \in F$

$$(Ty; Te) = \int_{B(E) \times B(F)} e(e)y(y) d(e; y)$$

and

$$\|T\|_{\text{int}} = \int \int (B(E) \times B(F)):$$

Following [4, page 234], define

$$S: E \rightarrow L_1(\cdot) \text{ by } Se(e; y) = e(e)$$

and

$$R: F \rightarrow L_1(\cdot) \text{ by } Ry(e; y) = y(y):$$

Then S and R are bounded linear operators with $\|S\|, \|R\| \leq 1$. Let Q be the restriction of R to $L_1(\cdot)$, then Q is a bounded linear operator from $L_1(\cdot)$ into F . It is immediate that $J \circ T = Q \circ I \circ S$ where $I: L_1(\cdot) \rightarrow L_1(\cdot)$ is the natural inclusion and $J: F \rightarrow F$ is the natural embedding. This in particular shows that the operator $Q \circ I$ restricted to $C(B(E) \times B(F))$ is an integral operator whose representing F -valued measure $\hat{\cdot}$ is such that

$$\hat{\cdot}(C) = Q(l_C)$$

for each Borel subset C of $B(E) \times B(F)$. Finally note that since the natural projection of $B(E) \times B(F)$ onto $B(E)$ is a continuous mapping it induces a bounded linear operator from $C(B(E))$ into $C(B(E) \times B(F))$ as follows; for $\phi \in C(B(E))$, let $\hat{\phi} \in C(B(E) \times B(F))$ be such that

$$\hat{\phi}(e; y) = \phi(e):$$

This of course shows that the operator $J \circ T$ extends to an integral operator

$$\hat{T}: C(B(E)) \rightarrow F$$

such that for each $\mu \in C(B(E))$

$$\hat{T} = Q \circ I(\cdot)$$

It is immediate that the F -valued measure m representing \hat{T} is such that

$$m(B) = Q(1_B \circ B(F))$$

and

$$\int \int(B(E)) \int \int(B(E) \circ B(F)) = k(T)_{int} :$$

Since \hat{T} extends T it follows that

$$k(T)_{int} = \int \int(B(E)) :$$

Main Result Throughout this section X is a compact Hausdorff space, E and F are Banach spaces and $T : C(X; E) \rightarrow F$ is a bounded linear operator with representing measure G . The main result of this paper gives a characterization of integral operators T in terms of some properties of G . The first step to achieve such a characterization is to show that if T is an integral operator on $C(X; E)$, then one can do a little better than Proposition 1 by representing the operator T by a regular F -valued measure of bounded variation defined on the σ -field of Borel subsets of $X \circ B(E)$ rather than on the whole unit ball of $(C(X; E))'$. The proof we present here is different from our earlier proof which relied on our result [7]. We would like to thank the referee for suggesting the following approach which relies more on basic knowledge and classical results of vector measures that can be found in [4] or [5].

Lemma 2. If $T : C(X; E) \rightarrow F$ is an integral operator, then there exists a regular F -valued measure of bounded variation on the σ -field of Borel subsets of $X \circ B(E)$ such that for each $f \in C(X; E)$ and $y \in F$

$$\langle y, T f \rangle = \int_{X \circ B(E)} \langle y, e(f(x)) \rangle d\langle x, e \rangle$$

and

$$k(T)_{int} = \int \int(X \circ B(E)) :$$

Proof: Suppose that $T: C(X; E) \rightarrow F$ is an integral operator, then the bilinear map $(f; y) \mapsto y(T(f))$ for $f \in C(X; E)$ and $y \in F$ defines an element of $(C(X; E) \rightarrow F)$. It is easy to check that $C(X; E) \rightarrow F$ embeds isometrically in $C(X \rightarrow B(E) \rightarrow B(F))$ such that if $f \in C(X; E)$ and $y \in F$

$$f(y)(x; e; y) = e(f(x))y(y)$$

for each $(x; e; y) \in X \times B(E) \times B(F)$. Hence by the Hahn-Banach theorem there exists a regular Borel measure μ on $X \times B(E) \times B(F)$ so that for each $f \in C(X; E)$ and $y \in F$

$$\langle y, T f \rangle = \int e(f(x))y(y) d\mu(x; e; y):$$

and

$$\|T\|_{int} = \|\mu\|:$$

The proof now follows the steps of Proposition 1. Indeed, since the natural projection of $X \times B(E) \times B(F)$ onto $X \times B(E)$ is continuous it induces a bounded linear operator

$$S: C(X \times B(E)) \rightarrow L_1(\mu)$$

such that if $f \in C(X \times B(E))$ then

$$S(f)(x; e; y) = f(x; e; y)$$

for all $(x; e; y) \in X \times B(E) \times B(F)$:

Let $R: F \rightarrow L_1(\mu)$ be defined by $R(y)(x; e; y) = y(y)$ and let Q be the restriction of R to $L_1(\mu)$. It is straightforward to check that the operator $Q \circ I \circ S$ is an integral operator whose representing measure is such that

$$(C) = Q(I_C \circ S)$$

for each Borel subset C of $X \times B(E)$, and

$$\|\mu\|_{X \times B(E)} = \|Q\| = \|T\|_{int}:$$

Moreover note that $C(X; E)$ embeds isometrically in $C(X \times B(E))$ by

$$f(x; e) = e(f(x))$$

for each $f \in C(X; E)$ and $(x; e) \in X \times B(E)$. It is easy to see that for each $f \in C(X; E)$

$$J(T(f)) = Q(I(S(f)))$$

hence

$$T(f) = \int_{X \times B(E)} e(f(x)) d(x; e)$$

and

$$\|T\|_{\text{int}} \leq \|Q\| \|I\| \|S\|_{\text{int}} \|\cdot\|_1(X \times B(E)).$$

This completes the proof of Lemma 2.

We are now ready to state and prove the main result of this paper.

Theorem 3 Let X be a compact Hausdorff space, let E and F be Banach spaces and let $T : C(X; E) \rightarrow F$ be a bounded linear operator with representing measure G . Then T is an integral operator if and only if for each Borel subset B of X , the operator $G(B) : E \rightarrow F$ is integral and the measure G is of finite variation as a vector measure taking its values in $I(E; F)$ equipped with the integral norm.

Proof: Assume that $T : C(X; E) \rightarrow F$ is an integral operator. Let G denote the vector measure representing the operator T , hence for each Borel subset B of X and each $e \in E$

$$G(B)e = T(1_B \cdot e)$$

It is clear at this stage, that since T is also integral [4, p 236], then for each Borel subset B of X the operator $G(B) : E \rightarrow F$ is an integral operator as the composition of T and the bounded linear operator $E \rightarrow C(X; E)$ which to each e in E associates the element $1_B \cdot e$. In what follows we shall concentrate on estimating the value of $\|G(B)\|_{\text{int}}$. For this note that by Lemma 2 there exists a regular F -valued measure defined on the σ -field of Borel subsets of $X \times B(E)$ such that, for each $f \in C(X; E)$

$$T(f) = \int_{X \times B(E)} e(f(x)) d(x; e)$$

and

$$k T k_{\text{int}} = \int \int (X \in B(E)) :$$

It follows that for each $y \in F$ and $f \in C(X; E)$

$$\int_Y y; f i = \int_{X \in B(E)} e(f(x)) d_y(x; e) :$$

where μ_y is the scalar measure on $X \in B(E)$ defined by

$$\mu_y(C) = \int (C)(y)$$

for each Borel subset C of $X \in B(E)$.

We claim that for each Borel subset B of X and for each $e \in E$

$$(*) \quad G(B)e = \int_{B \in B(E)} e(e) d(x; e) :$$

For this suppose that $y \in F$ and K is a compact G subset of X . Let $(u_n)_{n=1}^\infty$ be a sequence of continuous real valued functions so that $0 \leq u_n \leq 1$ for all $n \geq 1$ and u_n converges to 1_K pointwise. If $e \in E$, then

$$\int G(K)e; y i = \int \int G_y i(K)$$

$$= \lim_{n \rightarrow \infty} \int y; T(u_n e) i$$

$$= \lim_{n \rightarrow \infty} \int_{X \in B(E)} u_n(x) e(e) d_y(x; e) :$$

$$= \int_{X \in B(E)} 1_K(x) e(e) d_y(x; e)$$

$$= \int_{K \in B(E)} e(e) d_y(x; e)$$

Therefore

$$G(K)e = \int_{K \in B(E)} e(e) d(x; e) :$$

Moreover, since for each $y \in F$ the E -valued vector measure G_y is regular, it follows that

$$G(B)e = \int_{B \in \mathcal{B}(E)} e(e) d(x; e)$$

for all Borel subsets B of X . This proves our claim. If we denote by $p: X \times B(E) \rightarrow E$ the projection mapping which to each $(x; e)$ in $X \times B(E)$ associates e in $B(E)$, then for each Borel subset B of X , let μ_B denote the regular F -valued measure defined on the σ -field of Borel subsets of $B(E)$ as follows: for each Borel subset V of $B(E)$

$$\mu_B(V) = \int_{(B \times V)} :$$

In other words $\mu_B = \int_{\mathcal{B}(B(E))} p^{-1}$ which is the image measure of the restriction of μ to $B \times B(E)$ by p . This implies that for each $e \in E$

$$(**) \quad \int_{B(E)} e(e) d\mu_B(e) = \int_{B \times B(E)} e(e) d(x; e):$$

Equations () and () show that the measure μ_B is a regular F -valued measure that represents the operator $G(B): E \rightarrow F$. Hence by Proposition 1

$$\|G(B)\|_{int} = \|\mu_B\|(\mathcal{B}(B(E))) = \|\mu_B\|(B \times B(E)):$$

and therefore

$$(y) \quad \|G(B)\|_{int} = \|\mu_B\|(B \times B(E))$$

Hence the vector measure G representing the operator T takes its values in $I(E; F)$ and it follows from (y) that

$$\|G\|_{int}(X) = \|\mu\|(X \times B(E)):$$

Here of course $\|G\|_{int}(X) = \sup_{B_i \in \mathcal{P}} \sum_{i=1}^P \|G(B_i)\|_{int}$ where the sup is taken over all the finite partitions of X into Borel subsets of X .

Conversely, suppose $T: C(X; E) \rightarrow F$ is such that

$$G: \mathcal{B}(X) \rightarrow I(E; F) \quad \text{and} \quad \|G\|_{int}(X) < 1;$$

we need to show that T is an integral operator. For this, note that if J denotes the natural embedding of F into F^* , then by [4, p. 233] it is enough to show that $J \circ T : C(X; E) \rightarrow F^*$ is an integral operator. The mapping J induces a mapping $\hat{J} : I(E; F) \rightarrow I(E; F^*)$ defined as follows, for each $U \in I(E; F)$

$$\hat{J}(U) = J \circ U$$

this in turn induces a vector measure $\hat{G} : \Sigma \rightarrow I(E; F^*)$ such that

$$\hat{G}(B) = \hat{J}(G(B))$$

for all $B \in \Sigma$. It is immediate that \hat{G} is the measure representing the operator $J \circ T$. Since $\mathcal{J}_{\text{int}}(X) < 1$, it follows easily that $\hat{\mathcal{J}}_{\text{int}}(X) < 1$. Moreover, since $I(E; F^*)$ is isometric to the dual space of the injective tensor product $E \hat{\otimes} F$ [4, p. 237], it follows that \hat{G} is a vector measure of bounded variation taking its values in the dual space $(E \hat{\otimes} F)^*$. Since $\hat{\mathcal{J}}_{\text{int}}(X) < 1$, it follows from [4, page 7] that \hat{G} is strongly additive. Recall that for each $e \in E$ and $y \in F$, the scalar measure $\mu_{e; G_y}$ is in $M(X)$. Hence for $u = \sum_{i=1}^n e_i \otimes y_i \in E \hat{\otimes} F$, let $\mu_{u; \hat{G}}$ be the scalar measure defined on X as follows: for each Borel subset B of X

$$\mu_{u; \hat{G}}(B) = \hat{G}(B)(u)$$

$$= \sum_{i=1}^n \mu_{G(B)e_i; y_i}.$$

$$= \sum_{i=1}^n \mu_{e_i; G_{y_i}}(B)$$

This implies that for each $u \in E \hat{\otimes} F$ the measure $\mu_{u; \hat{G}} \in M(X)$. This in particular shows that $\hat{G} : \Sigma \rightarrow (E \hat{\otimes} F)^*$ is countably additive with respect to the weak topology on $(E \hat{\otimes} F)^*$. Since \hat{G} is strongly additive, it follows easily that G is countably additive. Finally, since \hat{G} is weak regular, a glance at [4, page 157] reveals that it is also regular.

It follows that \hat{G} defines an element of $C(X; E \wedge F)$, hence by [5, page 269] there exists a weak \mathcal{G}_{int} -integrable function

$$h : X \rightarrow I(E; F)' (E \wedge F)$$

such that $\|h(x)\|_{int} = 1$ \mathcal{G}_{int} -a.e., $\hat{G} = h \mathcal{G}_{int}$, and for each $f \in C(X; E \wedge F)$

$$\langle hf; \hat{G} \rangle = \int_X \langle hf(x); h(x) \rangle d\mathcal{G}_{int}(x)$$

In particular, if $\varphi \in C(X; E)$ and $y \in F$, let $f = \varphi \otimes y$ be the element of $C(X; E \wedge F)$ defined by

$$\varphi \otimes y(x) = \varphi(x) \otimes y \text{ for all } x \in X :$$

then

$$\langle \varphi \otimes y; \hat{G} \rangle = \int_X \langle \varphi(x); y \rangle d\mathcal{G}_{int}(x)$$

But we also have

$$(***)\quad \langle \varphi \otimes y; \hat{G} \rangle = \langle \varphi; \hat{G}_y \rangle$$

where \hat{G}_y is the element of $M(X; E)$ such that for each Borel subset B of X

$$\hat{G}_y(B) = \int_B \langle \varphi(x); y \rangle d\mathcal{G}_{int}(x) :$$

Hence since $\hat{G} = h \mathcal{G}_{int}$, it follows that for each Borel subset B of X

$$\hat{G}_y(B) = \int_B \langle h(x); y \rangle d\mathcal{G}_{int}(x) :$$

This implies that for each $\varphi \in C(X; E)$

$$(****)\quad \langle \varphi; \hat{G}_y \rangle = \int_X \langle \varphi(x); y \rangle d\mathcal{G}_{int}(x) :$$

It follows that if $\sum_{i=1}^n \varphi_i \otimes y_i$ is in $C(X; E \wedge F)$ with $\sum_{i=1}^n \|\varphi_i\|_{int} \leq 1$, then for each x in X

$$\|\sum_{i=1}^n \varphi_i(x) \otimes y_i\|_{int} \leq \sum_{i=1}^n \|\varphi_i\|_{int} \leq 1$$

since

$$\sum_{i=1}^{X^n} \|T'(x) - y_i\| \leq \sum_{i=1}^{X^n} \|T'(x) - y_i\| \leq 1$$

It follows then from () and () that

$$\sum_{i=1}^{X^n} \|T'(x) - y_i\| \leq \hat{G}_{int}(X) < 1$$

Hence $J \circ T$ is an element of $I(C(X; E); F)$ which implies that $T \in I(C(X; E); F)$. This completes the proof.

Applications The study of integral operators on $C(X; E)$ spaces was motivated by some problems that arose in [8] concerning nuclear operators on $C(X; E)$ spaces. Recall that an operator T between two Banach spaces Y and Z is said to be a nuclear operator if there exist sequences (y_n) and (z_n) in Y and Z respectively, such that $\sum_n \|y_n\| \|z_n\| < \infty$ and for each $y \in Y$

$$Ty = \sum_n y_n(y) z_n$$

the nuclear norm is defined by

$$\|T\|_{nuc} = \inf \left(\sum_n \|y_n\| \|z_n\| \right)$$

where the infimum is taken over all sequences (y_n) and (z_n) such that $Ty = \sum_n y_n(y) z_n$ for all $y \in Y$. We shall denote by $N(Y; Z)$ the space of nuclear operators from Y to Z under the nuclear norm. The study of nuclear operators on $C(X; E)$ spaces was initiated in [1] where some of the known results in the scalar case were extended. Nuclear operators on $C(X; E)$ spaces were also considered in [2] where it was shown that if $T : C(X; E) \rightarrow F$ is a nuclear operator then for each Borel subset B of X , the operator $G(B) : E \rightarrow F$ is nuclear. In [8] it was shown that the representing measure G of a nuclear operator is in fact of bounded variation as a vector measure taking its values in $N(E; F)$. Hence all the above known results can be summarized as follows:

Proposition 4 Let X be a compact Hausdorff space and let E and F be two Banach spaces. If T is a nuclear operator from $C(X; E)$ into F with representing measure G , then

(i) For any Borel subset B of X the operator $G(B) : E \rightarrow F$ is nuclear, and
(ii) G is of bounded variation as a vector measure taking its values in $N(E; F)$ under the nuclear norm.

Easy examples show that conditions (i) and (ii) above do not characterize nuclear operators on $C(X; E)$ spaces. As a matter of fact, it was shown in [8] that counterexamples can be given as soon as the range space F fails to have the so-called Radon-Nikodym property (RNP) see [4]. The interesting question that arises is then:

Question 5 Assume that F has the (RNP) and that $T : C(X; E) \rightarrow F$ is an operator whose representing measure G satisfies conditions (i) and (ii) of Proposition 4, is T a nuclear operator?

In what follows, we shall show how Theorem 3 can be used to give a positive answer to Question 5 in case F is assumed to be complemented in its bidual.

Theorem 6 Let F be a Banach space with the Radon-Nikodym property and assume that F is complemented in its bidual. A bounded linear operator $T : C(X; E) \rightarrow F$ is nuclear whenever its representing vector measure G satisfies the following conditions

- (i) For each Borel subset B of X , the operator $G(B) : E \rightarrow F$ is nuclear, and
- (ii) G is of bounded variation as a vector measure taking its values in $N(E; F)$.

Proof: First note that for an arbitrary Banach space F any element U in $N(E; F)$ is in $I(E; F)$ with $\|U\|_{\text{int}} = \|U\|_{\text{nuc}}$. Hence by Theorem 3 an operator $T : C(X; E) \rightarrow F$ that satisfies conditions (i) and (ii) above is indeed integral. Therefore when in addition we assume that F is complemented in its bidual F'' , the operator T becomes Pietsch integral (also called strictly integral) see [4, p.235]. Under these conditions and if F is assumed to have the Radon-Nikodym property, it is well known [4, p.175] that T becomes a nuclear operator. This completes the proof.

Remark As we have just seen, Theorem 6 is a direct application of the main result of this paper, it should be noted that Theorem 6 gives a positive answer to Question 5 when F is a dual space and when F is a Banach lattice.

Bibliography

- [1] G. Alexander, Linear Operators on the Space of Vector-Valued Continuous Functions, Ph.D. thesis, New Mexico State University, Las Cruces, New Mexico, 1972.
- [2] R. Bilyeu and P. Lewis, Some mapping properties of representing measures, Ann. Math. Pure Appl. C IX (1976), 273-287.
- [3] G. Choquet, Lecture on Analysis, Vol. II, Benjamin (1969).
- [4] J. Diestel and J.J. Uhl, Jr., Vector measures Math. Surveys 15 American Mathematical Society, (1977).
- [5] N. Dinulescu, Vector measures, Pergamon Press, (1967).
- [6] A. Grothendieck, Produit Tensoriels topologiques et Espaces Nucleaires. Memoirs of the American Mathematical Society, Vol. 16, (1955).
- [7] P. Saab, Integral representation by boundary vector measures. Canadian Math. Bull. 25 (2) (1982), 164-168.
- [8] B. Smith, Some Bounded Linear Operators on the Spaces $C(X; E)$ and $A(K; E)$, Ph.D. thesis, University of Missouri, Columbia, Missouri, 1989.
- [9] C. Swartz, Absolutely summing and dominated operators on spaces of vector valued continuous functions, Trans. Amer. Math. Soc., 179, (1973), 123-132.

University of Missouri
 Department of Mathematics
 Columbia, MO 65211