

Hyperbolic buildings, affine buildings and automatic groups

Donald I. Cartwright¹ and Michael Shapiro²

Abstract. We see that a building whose Coxeter group is hyperbolic is itself hyperbolic. Thus any finitely generated group acting co-compactly on such a building is hyperbolic, hence automatic. We turn our attention to affine buildings and consider a group which acts simply transitively and in a "type-rotating" way on the vertices of a locally finite thick building of type \tilde{A}_n . We show that this is biautomatic, using a presentation of \tilde{A}_n and unique normal form for each element of \tilde{A}_n , as described in [5].

1. Introduction.

Two standard references for the theory of automatic groups are the book [11] and the paper [2], which both contain numerous examples of automatic groups. Perhaps the "canonical" class of automatic groups is the class of word hyperbolic groups of Gromov et al. [14]. As we shall see in Section 3, any building whose underlying Coxeter group is word hyperbolic is itself hyperbolic in the word metric. In particular any finitely generated group acting co-compactly with finite stabilizers on such a building is word hyperbolic and thus automatic. Thus, from the viewpoint of automatic groups, it is natural to look next at actions on affine buildings.

The first result in this direction is provided by Gersten and Short [12] who prove that a finitely generated torsion free group that acts co-compactly discretely by isometries on a Euclidean building of dimension 2 is automatic. It has often seemed likely that the restriction on dimension could be lifted. In this paper, we prove this for finitely generated groups which act simply transitively (and in a type-rotating way) on the vertices of a thick building of type \tilde{A}_n . We call such groups \tilde{A}_n -groups. In fact, we will show more. We shall see that these groups are biautomatic. The structure in question is a symmetric automatic structure [11]. (The term "fully automatic" is used in [7] and early versions of [11].) In particular this implies that \tilde{A}_n -groups have solvable conjugacy problem [13]. Further, the structure consists of geodesics, and thus these groups have rational growth functions. Examples of finitely generated \tilde{A}_n -groups are known in dimensions $n = 2$ [6], $n = 3; 4$ [5] and $n = 5$ [Cartwright, unpublished], and for $n = 2; 3; 4$ and for any prime power q there are examples of \tilde{A}_n -groups which are arithmetic lattices in $\mathrm{PGL}(n+1; \mathbb{F}_q(X))$. It remains an open problem whether \tilde{A}_n -groups exist for every n .

The paper is organized as follows. Section 2 defines hyperbolic groups and reviews some basic background information, most of which can be found in [1]. In Section 3 we show

¹ School of Mathematics and Statistics, University of Sydney, N.S.W. 2006, Australia.

² Department of Mathematics, City College of New York, New York, NY 10031, USA

1991 Mathematics Subject Classification. Primary 20F10, 51E24. Secondary 20F05.

Key words and phrases. Automatic groups, affine buildings, hyperbolic groups.

that a building is hyperbolic (in an appropriate sense) if and only if its underlying Coxeter group is hyperbolic. Section 4 defines \tilde{A}_n -groups and reviews the necessary background from [5]. Section 5 gives the proof that finitely generated \tilde{A}_n -groups are bi-automatic. Sections 4 and 5 require no knowledge of buildings, Section 3 uses elementary results which can be found in [3, IV.3].

2. Hyperbolic groups

We say that a metric space $(X; d)$ is a geodesic metric space if for every $x, y \in X$, there is a path from x to y which realizes their distance. Such a path is called a geodesic. Following [14] we say that a geodesic metric space is δ -hyperbolic if whenever P is a point on side of a geodesic triangle with sides a , b , and c , there is a point Q on $[a, b]$ so that $d(P; Q) \leq \delta$. We say that $(X; d)$ is hyperbolic if it is δ -hyperbolic for some δ .

Now given any connected graph Γ there is a natural metric on Γ . Take each edge of Γ to be isometric to the unit interval and take the path metric which this induces on Γ . Further, given any finitely generated group G , the choice of a finite generating set $G = \langle a_1, \dots, a_k \rangle$ turns G into a directed labelled connected graph $\Gamma_G = \Gamma_G^G$. The vertices of Γ_G are the elements of G and the edges of Γ_G are $f(g; ga) : g \in G; a \in G$. We direct the edge $(g; ga)$ from g to ga and label it with a . We assume that G is closed under inverses and identify $(g; ga)$ with the inverse of $(ga; g)$. Γ_G is called the Cayley graph of G with respect to G .

We say that G is hyperbolic if $\Gamma_G = \Gamma_G^G$ is hyperbolic. While this appears to depend on G , in fact only the particular value of δ depends on G .

We will want a standard fact about hyperbolic metric spaces. Given a path γ in X and $0 < \epsilon < 1$ and $0 < \delta$, we will say that γ is a $(\epsilon; \delta)$ -quasigeodesic if for every decomposition $\gamma = \gamma_1 \cup \dots \cup \gamma_n$, the endpoints of γ_i are separated by at least $\epsilon \cdot \ell(\gamma_i)$. (Here $\ell(\gamma)$ denotes the length of γ .) If X is a δ -hyperbolic metric space then there is $N = N(\epsilon; \delta)$ so that if γ is a $(\epsilon; \delta)$ -quasigeodesic and γ' is a geodesic with the same endpoints, then γ and γ' each lie in the N -neighborhood of each other [1, 3.3]. Geodesics are simply $(1; 0)$ -quasigeodesics, and thus there is $N = N(\epsilon; 1; 0)$ so that all geodesics joining common endpoints live in a N -neighborhood of each other. We call a pair of geodesics with common endpoints a bigon.

From this one can construct a proof that any geodesic metric space quasi-isometric to a hyperbolic space is itself hyperbolic, and that in particular, hyperbolicity of a group is independent of generating set. Now it is a standard result that if a finitely generated group G acts co-compactly by isometries and with finite stabilizers on a geodesic metric space $(X; g)$, then every Cayley graph of G is quasi-isometric to X . (See, for example, [4].) In particular, when X is hyperbolic, so is G .

We have seen that the definition of a hyperbolic metric space requires checking that geodesic triangles are "thin", that is, that no side of a triangle is ever far from the union of the other two sides. Papasoglu [16], [17], has shown that in graphs it is only necessary to check bigons. Note that the endpoints of geodesics (and hence the endpoints of bigons) need not be vertices of Γ .

Theorem . (Papasoglu) Suppose that Γ is a graph and there is a constant K so that if γ, γ' is a bigon, then γ and γ' each lie in a K -neighborhood of each other. Then Γ is

hyperbolic.

Notice that the hypothesis is equivalent to the a priori stronger hypothesis that there exists K^0 so that if γ and γ^0 form a bigon then for all t , $d(\gamma(t); \gamma^0(t)) \leq K^0$. For suppose that $\gamma(t)$ is within K of $\gamma^0(t^0)$. Then the fact that these are geodesics emanating from a common point allows us to use the triangle inequality to see that $\|\gamma - \gamma^0\| \leq K$. Hence, taking $K^0 = 2K$, we have $d(\gamma(t); \gamma^0(t)) \leq K^0$.

3. Hyperbolic buildings

In his doctoral thesis [15], Moussong constructs actions of Coxeter groups on non-positively curved geodesic metric spaces. An account of this metric can be found in [9]. The metric spaces in question are locally Euclidean or locally hyperbolic complexes, and can be made negatively curved if and only if the Coxeter group in question is word hyperbolic. The actions are co-compact, by isometries and with finite stabilizers. As a scholium of his construction, one knows exactly which Coxeter groups are word hyperbolic.

Theorem (Moussong) Let $(W; S)$ be a Coxeter system. Then the following are equivalent:

1. W is word hyperbolic.
2. W has no $\mathbb{Z} \times \mathbb{Z}$ subgroup.
3. $(W; S)$ does not contain a finite sub-Coxeter system of rank ≥ 3 , and does not contain a pair of disjoint commuting sub-Coxeter systems whose groups are both infinite.

Chamey and Davis [8] have pointed out that using Moussong's metric of non-positive curvature, one can give a building a metric of non-positive curvature, and that the metric on the building is negatively curved if and only if the Coxeter group is word hyperbolic. (Construction of the metric on the building can be done along the lines of [3, VI.3].)

We give a similar characterization in terms of graphs. Given a building \mathcal{B} , there is a metric on the set of chambers of \mathcal{B} , and we will want a path metric space which reflects this metric. To do this we let \mathcal{B}^0 be the graph dual to \mathcal{B} . That is to say, the vertices of \mathcal{B}^0 are the barycenters of the chambers of \mathcal{B} . Two such vertices are connected by an edge when they lie in chambers with a common face. As usual, \mathcal{B}^0 is metrized considering each edge as isometric to the unit interval. Non-stuttering galleries of \mathcal{B} correspond to edgepaths in \mathcal{B}^0 . The decomposition of \mathcal{B} into apartments induces a decomposition of \mathcal{B}^0 into apartments which are isometric as labelled graphs to the Cayley graph of $(W; S)$, the Coxeter system of W .

Theorem 1. Suppose \mathcal{B} is a building whose apartments are the Coxeter complex of a word hyperbolic Coxeter group. Then \mathcal{B}^0 is hyperbolic.

Remark. The converse is also true. That is, if \mathcal{B}^0 is hyperbolic, the associated Coxeter group is word hyperbolic. This follows immediately from the fact that the embedding of the Cayley graph into \mathcal{B}^0 is an isometry.

Corollary. Suppose \mathcal{B} is a building whose apartments are the Coxeter complex of a word hyperbolic Coxeter group and that G is a finitely generated group which acts simplicially, co-compactly with finite stabilizers on \mathcal{B} . Then G is word hyperbolic.


Proof. G acts on \mathcal{B} , and the natural embedding of \mathcal{B}^0 into \mathcal{B} is equivariant with respect to this action. This induces an action of G on \mathcal{B}^0 . Since there are finitely many G -orbits

of chambers of \mathcal{C} and finitely many G -orbits of codimension 1 faces, the induced action on \mathcal{C}^0 is co-compact. Now G carries edges of \mathcal{C}^0 to edges of \mathcal{C}^0 and thus acts by isometries of the graph metric. Since the action of G has finite stabilizers, the (setwise) stabilizer of each chamber and codimension 1 face of \mathcal{C} is finite. These are the stabilizers of the vertices and edges of \mathcal{C}^0 . Hence, as we have outlined in Section 2, G is quasi-isometric to \mathcal{C}^0 and thus word hyperbolic.

Proof of Theorem 1. To prove the Theorem, we consider a bigon γ, γ^0 . It suffices to show that there is K so that if C is any point of \mathcal{C} , then C is within K of \mathcal{C}^0 . We distinguish three cases depending on whether both, one, or neither of the endpoints of this bigon are vertices.

Case 1: Both ends of the bigon are vertices. Now given any two chambers of \mathcal{C} , there is an apartment containing them both. Any geodesic gallery in \mathcal{C} connecting these two chambers lies in \mathcal{A} . (See [3, p.88].) It now follows that γ and γ^0 lie in a common apartment of \mathcal{C}^0 , and by the hyperbolicity of the underlying Coxeter group, we are done.

Case 2: The geodesics γ and γ^0 begin at a vertex, but do not end at a vertex. We let x and x^0 be the beginning and endpoints of our bigon. We let y and z be the last vertices of γ and γ^0 respectively, and let γ' and γ'^0 be the initial segments of γ and γ^0 ending at y and z respectively. It now follows that $\gamma' = \gamma'^0$ and that x^0 is the midpoint of an edge e . If $y = z$, γ' and γ'^0 lie in a common apartment, and we are done by Case 1. Thus we may assume $y \neq z$. Now γ' and γ'^0 cannot lie in a common apartment. For if this were so, e^{-1} would label a relator of odd length in the underlying Coxeter group, and this is impossible.



Case 2


Let \mathcal{A} be an apartment containing γ' . We consider $C = \gamma(t)$ and may suppose this is a vertex lying on γ' . We let $\pi_C = \pi_{\mathcal{A}, C}$ be the retraction onto \mathcal{A} centered at C . (See, for example [3, IV.3].) Now $d(y; z) \geq 1$ and since π_C does not increase distance, we have

$$n - 1 \leq d(x; z) \leq n;$$

where $n = \ell(\gamma')$. Now consider the path γ'^0 . This may not be an edge path, as some edge of \mathcal{C}^0 may be folded by π_C . However its length as a path is still n . We thus have a path of length n whose endpoints lie at distance at least $n - 1$. It follows that γ'^0 is a $(1; 1)$ -quasigeodesic. Since γ'^0 lies in \mathcal{A} , it now follows from the hyperbolicity of \mathcal{A} that γ'^0 lies close to γ' . We can thus find $C^0 = \gamma'^0(t^0)$ so that (C^0) lies close to C . Since $\pi_C = \pi_{\mathcal{A}, C}$ preserves distance from C , C^0 lies close to C and we are done.

Case 3: Neither end of the bigon is a vertex. In this case we let a and b , (respectively a^0 and b^0) be the first and last vertices of γ (respectively γ^0), and let γ'' (respectively γ''^0)

be the segment of γ (respectively γ^0) connecting these first and last vertices. If $a = a^0$ or $b = b^0$ or both we are reduced to previous cases, so we can assume $a \notin a^0$ and $b \notin b^0$. We let e be the edge from a to a^0 and e^0 be the edge from b to b^0 . It is easy to check that $\gamma'(\gamma) = \gamma'(\gamma^0)$. We take $n = \gamma'(\gamma)$. Thus $d(a; b^0)$ is either $n - 1$, n or $n + 1$. We let γ be a geodesic from a to b^0 .



Case 3

If $d(a; b^0) = n - 1$ then e^0 and γ form a bigon whose ends are the vertices a and b . Likewise e and γ^0 form a bigon whose ends are the vertices a^0 and b^0 . Applying Case 1 twice takes care of this situation.

If $d(a; b^0) = n$, we let y and y^0 be the midpoints of e and e^0 . We then have a bigon whose ends are a and y^0 whose sides consist of γ and γ^0 , each with a half of e^0 appended. Similarly, we have a bigon with ends b^0 and y and sides γ and γ^0 with the halves of e appended. Now we can apply Case 2 twice.

Finally, if $d(a; b^0) = n + 1$ then γ and e^0 form a bigon as do γ^0 and e , and once again we can apply Case 1 twice.

x4. Review of \mathbb{A}_n -groups.

\mathbb{A}_n -groups were introduced for general $n \geq 2$ in [5], after earlier work on the case $n = 2$ in [6]. (One dimensional buildings are trees, and thus hyperbolic.) Recall that a building is a labelable complex, that is, that each vertex v of a building of type \mathbb{A}_n has a type $\gamma(v) \in \{0, 1, \dots, n\}$, with each chamber having one vertex of each type. If g is an automorphism of Δ , and if there is an integer c such that $\gamma(gv) = \gamma(v) + c \pmod{n+1}$ for each vertex v , then g is called type-rotating. Such automorphisms form a subgroup of index at most 2 in the group of all automorphisms of Δ .

If K is a field with discrete valuation, then there is a thick building Δ_K of type \mathbb{A}_n associated with K [18, Section 9.2], [3, Section V.8], and the group $\text{PGL}(n+1; K)$ acts transitively and in a type-rotating way on Δ_K .

A group is said to be an \mathbb{A}_n -group if it acts simply transitively on the vertices of a thick building of type \mathbb{A}_n in a type-rotating way.

We now describe \mathbb{A}_n -groups. Let Σ be a projective geometry of dimension $n \geq 2$ (see [10, p. 24] or [19, p. 105], for example). For $i = 1, \dots, n$, let $\Sigma_i = \{x \in \Sigma : \dim(x) = i\}$. To avoid unnecessary abstraction, the reader may assume that Σ is the set (V) (partially ordered by inclusion) of nontrivial proper subspaces of an $n+1$ -dimensional vector space V over a field k , and that $\dim(x)$ refers to the dimension of the subspace x of V . For when $n \geq 3$, or when $n = 2$ and Σ is desarguesian, Σ must be isomorphic to (V) for some V ([10, pp. 27-28] or [19, p. 203]). Let $\sigma : \Sigma \rightarrow \Sigma$ be an involution such that $\dim(\sigma(x)) = n+1 - \dim(x)$.

for $i = 1, \dots, n$, and let T is an \tilde{A}_n -triangle presentation compatible with Γ . This means that T is a set of triples $(u; v; w)$, where $u, v, w \in V$, such that

- (A) given $u, v \in V$, then $(u; v; w) \in T$ for some $w \in V$ if and only if (u) and v are distinct and incident;
- (B) if $(u; v; w) \in T$, then $(v; w; u) \in T$;
- (C) if $(u; v; w_1) \in T$ and $(u; v; w_2) \in T$, then $w_1 = w_2$;
- (D) if $(u; v; w) \in T$, then $((w); (v); (u)) \in T$;
- (E) if $(u; v; w) \in T$, then $\dim(u) + \dim(v) + \dim(w) = n + 1$ or $2(n + 1)$;
- (F) if $(x; y; u) \in T^0$ and $(x^0; y^0; (u)) \in T^0$, then for some $w \in V$ we have $(y^0; x; w) \in T^0$ and $(y; x^0; (w)) \in T^0$.

Here T^0 denotes the "half" of T consisting of the triples $(u; v; w) \in T$ for which $\dim(u) + \dim(v) + \dim(w) = n + 1$. Given $u, v \in V$, then $(u; v; w) \in T^0$ for some $w \in V$ if and only if $(u) \neq v$. We also write T^0 for $T \cap T^0$.

We form the associated group Γ_T with a generating set indexed by V :

$$\Gamma_T = \langle a_v, g_v \mid \begin{aligned} (1) & a_{(v)} = a_v^{-1} \text{ for all } v \in V; \\ (2) & a_u a_v a_w = 1 \text{ for all } (u; v; w) \in T \end{aligned} \rangle$$

It was shown in [5] that the Cayley graph of Γ_T with respect to the generators $a_v, v \in V$, is the 1-skeleton of a thick building \mathcal{B}_T of type \tilde{A}_n . Clearly, Γ_T acts, by left multiplication, simply transitively on the set of vertices of \mathcal{B}_T . Conversely, if Γ is a group of type-rotating automorphisms of a thick building \mathcal{B} of type \tilde{A}_n , and acts simply transitively on the vertices of \mathcal{B} , then $\Gamma = \Gamma_T$ and $\mathcal{B} = \mathcal{B}_T$ for some \tilde{A}_n -triangle presentation T . This generalized earlier work on the case $n = 2$ [6].

In this paper, Γ is assumed finite. The number of $x \in V$ incident with any given $y \in V$ is denoted $q + 1$, and is independent of y . Here q is called the order of Γ , and when $\Gamma = \text{Aut}(V)$, q is the number of elements in the field k . It remains an open problem whether \tilde{A}_n -triangle presentations exist for every n . They have been found when $\Gamma = \text{Aut}(V)$ for $n = 2$ [6] and for $n = 3, 4$ [5] for any prime power q , and for $n = 5$ and $q = 2$ [Cartwright, unpublished].

Let L denote the set of all strings $u_1 u_2 \dots u_\ell$ over V such that $(u_i) + u_{i+1} = V$ for $i = 1, \dots, \ell - 1$. (The notation assumes that $\Gamma = \text{Aut}(V)$, but in general, " $(u_i) + u_{i+1} = V$ " is interpreted as "there is no $x \in V$ such that $(u_i) \neq x$ and $u_{i+1} \neq x$ ", where we write $y \neq x$ if x and y are incident, and $\dim(y) \neq \dim(x)$.) Theorem 2.2 in [5] states that $u \mapsto u_1 \dots u_\ell$, $\Gamma \ni u \mapsto a_{u_1} \dots a_{u_\ell}$, is a bijection $L \rightarrow \Gamma$. Moreover, in the notation of [11], each string $u_1 \dots u_\ell \in L$ is geodesic, and so the number ℓ is the word length $l(g)$ of $g = u$. If $g \in \Gamma$, $u \in L$ and $g = u$, then u is called the normal form of g . We shall also refer to strings in L as being in normal form. When g is the identity element 1, its normal form is the empty word, and its word length is 0, by definition. We write $d(g; g^0)$ for $l(g^{-1} g^0)$, the distance from g to g^0 in the word metric.

5. Finitely generated \tilde{A}_n -groups are automatic.

Throughout this section, let $\Gamma = \Gamma_T$ be a finitely generated \tilde{A}_n -group.

We start by showing that L is a regular language. For we can define a finite state automaton M accepting L as follows: the set S of states of M is together with an initial state s_0 (not in Σ) and a single failure state s_1 (not in Σ , and distinct from s_0); the alphabet of M is Σ ; the transition function of M is given by $(s_0; x) = x$ and $(s_1; x) = s_1$ for $x \in \Sigma$, while for $x; y \in \Sigma$, we set $(x; y) = y$ if $(x) + y = V$ and $(x; y) = s_1$ otherwise; the set Y of accept states of M is $\{s_0\}$. Clearly $L = L(M)$.

Note that Σ is a set of semigroup generators for Γ which is closed under inversion (as $a_{(x)} = a_x^{-1}$). Moreover, L has the uniqueness property (i.e., $u \mapsto u$ is a bijection $L \rightarrow \Gamma$), is obviously pre x -closed, and is symmetric (i.e., $u_1 \in L$ implies that $(u_1)^{-1} \in L$). This last property of L and our Theorem 2 below imply that Γ is biautomatic, and in fact symmetric automatic or fully automatic.


If $u = u_1 \dots u_t \in L$, let $u(0)$ be the identity element 1 in Γ , and for $1 \leq t \leq |u|$, let $u(t)$ denote the element $a_{u_1} \dots a_{u_t}$ of Γ . For $t > |u|$, let $u(t) = a_{u_1} \dots a_{u_t} \cdot u$. Let $x \in \Sigma$, and let $v_1 \dots v_k$ be the normal form of $\overline{ux} = a_{u_1} \dots a_{u_t} \cdot a_x$. By [11, Theorem 2.3.5], to show that Γ is automatic, it is enough to show that for some k (independent of u and x), $d(u(t); v(t)) \leq k$ holds for each integer $t \geq 0$. This property is called the k -fellow traveller property [2]. In fact, we show this property holds for $k = 1$.

Now let $u = u_1 \dots u_t \in L$, and let $x \in \Sigma$. The main work below is to describe the normal form $v = v_1 \dots v_k$ of $a_{u_1} \dots a_{u_t} \cdot a_x$. The description of this normal form is complicated by the fact that, given $u; x \in \Sigma$, there are the following 5 mutually exclusive cases to consider: $(u) = x$, $(u) + x = V$, $(u) \leq x$, $(u) \leq x$, and, finally, (u) and x are distinct and nonincident, with $(u) + x \notin V$.

We start with a lemma which concerns the last of these cases. In this lemma and below, we consistently identify a string u over Σ with its image u in Γ , sometimes writing $\backslash u$ in Γ for emphasis, when thinking of u as u .

Lemma 1. Suppose that $u; x \in \Sigma$, and that (u) and x are distinct and nonincident, and that $(u) + x = (s) (\notin V)$. Thus we can write $(s; (u); x^0) \in T^0$ and $(s; x; (v)) \in T^0$ for some $x^0; v \in \Sigma$, and $ux = x^0v$ in Γ . Then x^0v is in normal form. That is, $(x^0) + v = V$. Conversely, if two triples $(s; (u); x^0); (s; x; (v)) \in T^0$ are given, with $(x^0) + v = V$, then (u) and x must be distinct and nonincident, with $(u) + x = (s)$.

We illustrate this lemma with a diagram in the Cayley graph of Γ :



Proof. Firstly, $(x^0) \leq v$. Otherwise, $x^0 = (v)$, and so Axioms (B) and (C) imply that $(u) = x$, contrary to hypothesis. We shall henceforth use Axioms (A)–(E) in the definition of an \mathcal{A}_n -triangle presentation without comment, but refer to Axiom (F) when it is used. Next, suppose that $(x^0) \leq v$, but that (x^0) and v are incident. Thus $(x^0; v; w) \in T$

for some $w \in \mathcal{L}$. If $(x^0; v; w) \in T^0$, then $(w; x^0; v) \in T^0$, $(s; x; (v)) \in T^0$ and Axiom (F) imply that $(x; w; y) \in T^0$ and $(x^0; s; (y)) \in T^0$ for some $y \in \mathcal{L}$. Thus $y = u$. But then $(u; x; w) \in T$, so that (u) and x are incident, contrary to hypothesis. If $(x^0; v; w) \in T^0$, then $(s; (u); x^0) \in T^0$, $((w); (v); (x^0)) \in T^0$ and Axiom (F) imply that $((v); s; y) \in T^0$ and $((u); (w); (y)) \in T^0$ for some $y \in \mathcal{L}$. Then $y = x$, and so $(x; w; u) \in T^0$, so that again (u) and x are incident, contrary to hypothesis.

Suppose that $(x^0) \notin v$, that (x^0) and v are not incident, and that $(x^0) + v = (s^0) \notin V$. So $(s^0; (x^0); x^0) \in T^0$ and $(s^0; v; (v^0)) \in T^0$ for some $x^0, v^0 \in \mathcal{L}$. Now $(s; (u); x^0) \in T^0$, $(x^0; s^0; (x^0)) \in T^0$ and Axiom (F) show that $(s^0; s; z) \in T^0$ and $((u); x^0; (z)) \in T^0$ for some $z \in \mathcal{L}$. Similarly, $((v^0); s^0; v) \in T^0$ and $(s; x; (v)) \in T^0$ imply that $(x; (v^0); z^0) \in T^0$ and $(s^0; s; (z^0)) \in T^0$ for some $z^0 \in \mathcal{L}$. Thus $(z^0) = z$. Hence $((u); x^0; (z)) \in T^0$ and $(x; (v^0); (z)) \in T^0$, so that $z = (u); x$. Hence $z = (u) + x = (s)$. But $(s^0; s; z) \in T^0$ implies that $z \notin (s)$. This contradiction completes the proof of the first part of the lemma.

Consider the converse part. Firstly, (u) and x must be distinct, for otherwise $x^0 = (v)$ must hold, which is impossible, because $(x^0) + v = V$. Next, $(u) \not\% x$ cannot hold. For otherwise $(u; x; y) \in T^0$ for some $y \in \mathcal{L}$. This, $((u); x^0; s) \in T^0$ and Axiom F then imply that $(s; x; z) \in T^0$ and $(y; x^0; (z)) \in T^0$ hold for some $z \in \mathcal{L}$. Thus $z = (v)$, so that $(y; x^0; v) \in T^0$, which implies that $(x^0) \in v$, again a contradiction. Similarly, $(u) \$ x$ leads to a contradiction. Finally, the hypotheses imply that $(u) + x = (s)$. If $(u) + x \notin (s)$, write $(u) + x = (s^0)$, and then we can find $x^0, v^0 \in \mathcal{L}$ such that $(s^0; (u); x^0); (s^0; x; (v^0)) \in T^0$. But then $x^0 v = ux = x^0 v^0$, and both $x^0 v$ and $x^0 v^0$ are in normal form (by the hypotheses, and by the first part of the lemma, respectively). Uniqueness of normal forms now shows that $x^0 = x^0$, so that $s^0 = s$, a contradiction. This completes the proof.


Lemma 2. Let $u_1; u_2; x \in \mathcal{L}$, with $u_1 u_2$ in normal form. Then to get the normal form of $u_1 u_2 x$, there are the following 5 possibilities:

- (1) If $(u_2) + x = V$, then $u_1 u_2 x$ is in normal form.
- (2) If $(u_2) = x$, then the normal form of $u_1 u_2 x$ is u_1 .
- (3) If $(u_2) \not\% x$, then $(u_2; x; (w)) \in T^0$ for some $w \in \mathcal{L}$, and the normal form of $u_1 u_2 x$ is $u_1 w$.
- (4) If $(u_2) \$ x$, then $(u_2; x; (w)) \in T^0$ for some $w \in \mathcal{L}$. Thus $u_1 u_2 x = u_1 w$ in \mathcal{L} . There are now the following possibilities:
 - (a) Either $(u_1) + w = V$, in which case the normal form of $u_1 u_2 x$ is $u_1 w$, or
 - (b) $(u_1) \not\% w$, in which case $(u_1; w; (w^0)) \in T^0$ for some $w^0 \in \mathcal{L}$, and the normal form of $u_1 u_2 x$ is w^0 , or
 - (c) (u_1) and w are distinct and nonincident, with $(u_1) + w \notin V$. Then writing $(u_1) + w = (s^0)$, there are unique $w^0; v_1 \in \mathcal{L}$ such that $(s^0; (u_1); w^0) \in T^0$ and $(s^0; w; (v_1)) \in T^0$, and the normal form of $u_1 u_2 x$ is $w^0 v_1$.
- (5) If (u_2) and x are distinct and nonincident, with $(u_2) + x \notin V$, then writing $(u_2) + x = (s)$, there are unique $x^0; v_2 \in \mathcal{L}$ such that $(s; (u_2); x^0) \in T^0$ and $(s; x; (v_2)) \in T^0$. Thus $u_1 u_2 x = u_1 x^0 v_2$ in \mathcal{L} . There are now the following possibilities:
 - (a) Either $(u_1) + x^0 = V$, in which case the normal form of $u_1 u_2 x$ is $u_1 x^0 v_2$, or


(b) $(u_1) \neq x^0$, in which case $(u_1; x^0; (w^0)) \in T^0$ for some $w^0 \in V$, and the normal form of $u_1 u_2 x$ is $w^0 v_2$, or

(c) (u_1) and x^0 are distinct and nonincident, with $(u_1) + x^0 \notin V$. Then writing $(u_1) + x^0 = (s^0)$, there are unique $x^0; v_1 \in V$ such that $(s^0; (u_1); x^0) \in T^0$ and $(s^0; x^0; (v_1)) \in T^0$, and the normal form of $u_1 u_2 x$ is $x^0 v_1 v_2$.


We illustrate the more complicated cases:



Case 4 (c)



Case 5 (b)



Case 5 (c)

P roof. The assertions in (1) and (2) are obvious.

Let us consider the situation in (3). Then $(u_2; x; (w)) \in T^0$ implies that $w = u_2$. Thus $(u_1) + w = (u_1) + u_2 = V$, and so $u_1 w$ is in normal form.

Let us consider the situation in (4). In this case, again $w \notin (u_1)$. For otherwise, $(u_2; x; u_1) = (u_2; x; (w)) \in T^0$, contradicting the hypothesis that $u_1 u_2$ is in normal form. Also, $(u_1) \neq w$ cannot hold. For otherwise, $((w); (u_1); w^0) \in T^0$ for some $w^0 \in V$. Now $((x); (u_2); w) \in T^0$, $((u_1); w^0; (w)) \in T^0$ and Axiom (F) imply that $(w^0; (x); y)$ and $((u_2); (u_1); (y))$ are in T^0 for some $y \in V$. This again contradicts the hypothesis that $u_1 u_2$ is in normal form. We are left with the three possibilities 4(a), 4(b) and 4(c). The assertions in 4(a) and 4(b) are obvious, while in 4(c), the fact that the word $w^0 v_1$ is in normal form is immediate from Lemma 1.

Let us consider the situation in (5). In this case, (u_1) cannot equal x^0 . For otherwise $(s; (u_2); (u_1)) = (s; (u_2); x^0) \in T^0$, which contradicts the hypothesis that $u_1 u_2$ is in normal form. Also, $(u_1) \neq x^0$ cannot happen. For otherwise, $((x^0); (u_1); w^0) \in T^0$ for some $w^0 \in V$. Now $(s; (u_2); x^0) \in T^0$, $((u_1); w^0; (x^0)) \in T^0$ and Axiom (F) imply that $(w^0; s; y) \in T^0$ and $((u_2); (u_1); (y)) \in T^0$ for some $y \in V$. This last fact again contradicts the hypothesis that $u_1 u_2$ is in normal form. We are left with the three possibilities 5(a), 5(b) and 5(c). The assertion in 5(a) is obvious, by Lemma 1.

Consider the situation in 5(b). First notice that $((x^0); (w^0); u_1); ((x^0); s; (u_2)) \in T^0$ and the converse part of Lemma 1 imply that $(w^0) + s = (x^0)$. As $(s; x; (v_2)) \in T^0$, we have

$v_2 \leq s$, and so $(w^0) + v_2 = (w^0) + s + v_2 = (x^0) + v_2 = V$, the last equation holding by Lemma 1. So $w^0 v_2$ is in normal form.

Finally, consider the situation in 5(c). First observe that $(v_1) + s = (x^0)$. This follows from the converse part of Lemma 1, because $(x^0; (v_1); s^0); (x^0; s; (u_2)) \in T^0$, and because, using $(s^0; (u_1); x^0) \in T^0$, we have $(s^0) + u_2 = (u_1) + u_2 = V$. Now $(s; x; (v_2)) \in T^0$, so that $v_2 \leq s$. Hence $(v_1) + v_2 = (v_1) + s + v_2 = (x^0) + v_2 = V$, the last equation holding by Lemma 1. Lemma 1 also shows that $(x^0) + v_1 = V$, and so $x^0 v_1 v_2$ is in normal form.

Remark. In later work, we shall need a converse to Lemma 2. Let us write $x +^0 y = z$ for $x; y; z \in L$ if $x; y$ are distinct and nonincident, with $x + y = z$. Then by the converse part of Lemma 1, we have $(w^0) +^0 (x) = (w)$ in part 4(b), $(v_1) +^0 (x) = (w)$ in part 4(c), $(w^0) +^0 s = (x^0)$ in part 5(b) and $(v_1) +^0 s = (x^0)$ in part 5(c). Provided these conditions are added, the converses of parts (4) and (5) hold. For example, for 5(b), if $u_1; u_2; x; s; x^0; v_2; w^0 \in L$ and triples $(s; (u_2); x^0); (s; x; (v_2)) \in T^0$ are given with $(w^0) +^0 s = (x^0)$ and $(w^0) + v_2 = V$, then $u_1 u_2$ is in normal form and $(u_2) +^0 x = (s)$. The first of these is immediate from Lemma 1, since $(x^0; (w^0); u_1); (x^0; s; (u_2)) \in T^0$. To see that $(u_2) +^0 x = (s)$, notice that $(x^0) + v_2 = (w^0) + v_2 = V$, and so the converse part of Lemma 1 is applicable.

The normal form $v_1 \dots v_k$ of $a_{u_1} \dots u, a_x$.

We can now describe how to obtain the normal form $v_1 \dots v_k$ of $\overline{ux} = a_{u_1} \dots u, a_x$, given $u_1 \in L$ and $x \in L$. When $(u_1) + x = V$, this is obviously $u_1 \dots u, x$. If $(u_1) = x$, then the normal form of \overline{ux} is clearly $u_1 \dots u$. When $(u_1) \not\leq x$, then $(u_1; x; (w)) \in T^0$ for some $w \in L$, and the normal form of \overline{ux} is $u_1 \dots u, w$, by part (3) of Lemma 2.

When (u_1) and x are distinct and nonincident, with $(u_1) + x \notin V$, then writing $(u_1) + x = (s_1)$, we have $(s_1; (u_1); x_1) \in T^0$ and $(s_1; x; (v_1)) \in T^0$ for some $x_1, v_1 \in L$. Then $u_1 \dots u, x = u_1 \dots u, x_1 v_1$ in L . This situation may be repeated several times, with (u_{i+1}) and x_i distinct and nonincident, with $(u_{i+1}) + x_i \notin V$, and so on. Suppose this situation is repeated exactly i times. Then

we find $s_i; x_{i+1}; v_i \in L$ for $i = 0; \dots; i-1$, such that $(u_i) + x = (s_i)$, $(s_i; (u_i); x_{i+1}) \in T^0$, and $(s_i; x; (v_i)) \in T^0$ for $i = 0; \dots; i-1$ (writing $x_0 = x$). Then in

$$u_1 \dots u, x = u_1 \dots u, x_1 v_1 x_2 v_2 \dots x_i v_i \quad (4.1)$$

Then by part (5) of Lemma 2, either the word on the right in (4.1) is in normal form, or $(u_{i+1}) \leq x_i$, in which case $(u_{i+1}; x_i; (w)) \in T^0$ for some $w \in L$, and the normal form of \overline{ux} is

$$u_1 \dots u, x_1 w v_1 x_2 v_2 \dots x_i v_i \quad (4.2)$$

Finally, if $(u_i) \leq x$, then $(u_i; x; (w)) \in T^0$ for some $w \in L$. Thus $u_1 \dots u, x = u_1 \dots u, w$ in L . A sequence of exactly $i-1$ steps such as led to (4.1) may now occur, so that we find elements $s_i; w_i; v_i \in L$ for $i = 1; \dots; i-1$, such that $(u_{i-1}) + w_{i-1} = (s_{i-1})$, $(s_{i-1}; (u_{i-1}); w_i) \in T^0$ and $(s_{i-1}; w_{i-1}; (v_{i-1})) \in T^0$ for $i = 1; \dots; i-1$ (writing $w_0 = w$). Then in

$$u_1 \dots u, x = u_1 \dots u, x_1 w_1 v_1 x_2 w_2 v_2 \dots x_{i-1} w_{i-1} v_{i-1} w \quad (4.3)$$

By parts (4) and (5) of Lemma 2, either the word on the right in (4.3) is in normal form, or $(u \cdot i \cdot 1) \cdot w_i$, in which case $(u \cdot i \cdot 1) \cdot w_i \cdot (w^0)^{-1} \in T^0$ for some $w^0 \in L$, and the normal form of \overline{ux} is

$$u_1 \cdot i \cdot u_2 \cdot w^0 \cdot v \cdot i \cdot i \cdot v \quad (4.4)$$

We are now ready to prove the fellow traveller property.

Theorem 2. Let T be a finitely generated \tilde{A}_n -group. Let $u = u_1 \cdot i \cdot L$, let $x \in L$, and let $v_1 \cdot k \cdot L$ be the normal form of $\overline{ux} = a_{u_1} \cdot u \cdot a_x$. Then for each integer $t \geq 0$, we have $d(u(t); v(t)) \leq 1$. That is, either $a_{u_1} \cdot u_t \cdot a_x = a_{v_1} \cdot v_t \cdot a_x$ or $a_{u_1} \cdot u_t \cdot a_x = a_{v_1} \cdot v_t \cdot a_{x_t}$ for some $x_t \in L$. Thus T is an automatic group.

Proof. Let us discuss the most complicated case in detail, leaving the other cases to the reader. Suppose that $(u \cdot i) \cdot x$, and that the normal form v of \overline{ux} is (4.4). For $0 \leq t \leq i-2$, we have $a_{u_1} \cdot u_t \cdot a_x = a_{v_1} \cdot v_t \cdot a_x$ and

$$\begin{aligned} v(i-1) &= a_{v_1} \cdot v \cdot a_{i-1} = a_{u_1} \cdot u \cdot a_{i-2} \cdot a_{w^0} \\ &= a_{u_1} \cdot u \cdot a_{i-2} \cdot a_{u \cdot i \cdot 1} \cdot a_{w_i} \\ &= u(i-1) \cdot a_{w_i} \end{aligned}$$

and $w_i \in L$. For $i-1$, we have

$$\begin{aligned} v(i) &= a_{u_1} \cdot u \cdot a_{i-2} \cdot a_{w^0} \cdot a_{v \cdot i} \cdot v \cdot a \\ &= a_{u_1} \cdot u \cdot a_{i-2} \cdot a_{u \cdot i \cdot 1} \cdot a_{w_i} \cdot a_{v \cdot i} \cdot v \cdot a \\ &= a_{u_1} \cdot u \cdot a_{i-2} \cdot a_{u \cdot i \cdot 1} \cdot a_{u \cdot i} \cdot a_{u \cdot i \cdot 1} \cdot u \cdot a \cdot a_{w_{i-1}} \\ &= u(i) \cdot a_{w_{i-1}} \end{aligned}$$

and $w_{i-1} \in L$. For $t \geq i$, $v(t) = \overline{ux} = u(t) \cdot a_x$. This completes the proof.

References.

- [1] Alonso, J., Brady, T., Cooper, D., Ferlini, V., Lustig, M., Mihalik, M., Shapiro, M., and Short, H., Short, H., ed., Notes on word hyperbolic groups in "Group Theory From a Geometric Viewpoint", Ghyss, E., Haefliger, A., and Verjovsky A., eds., World Scientific (1990).
- [2] Baumslag, G., Gersten, S.M., Shapiro, M. and Short, H., Automatic groups and amalgams, J. Pure and Appl. Alg. 76 (1991), 229{316.
- [3] Brown, K., "Buildings", Springer Verlag (1989).
- [4] Cannon, J.W., The combinatorial structure of cocompact discrete hyperbolic groups, Geom. Dedic. 16 (1984), 123{148.
- [5] Cartwright, D.I., Groups acting simply transitively on the vertices of a building of type \tilde{A}_n , to appear, Proceedings of the 1993 Conference "Groups of Lie type and their geometries", W.M. Kantor, Editor.
- [6] Cartwright, D.I., Mantero, A.M., Steger, T. and Zappa, A., Groups acting simply transitively on the vertices of a building of type \tilde{A}_2 I, II, Geom. Dedic. 47 (1993), 143{166, 167{223.

- [7] Chamey, R., Geodesic automaton and growth functions for Artin groups of finite type, Ohio State University preprint, (1993).
- [8] Chamey, R., and Davis, M., personal communication.
- [9] Cohen, A., "Recent results on Coxeter groups", preprint, (1993).
- [10] Dembowski, P., "Finite Geometries", Ergebnisse der Mathematik und ihrer Grenzgebiete, Band 44, Springer-Verlag, Berlin, Heidelberg, New York, 1968.
- [11] Epstein, D.B.A., Cannon, J.W., Holt, D.F., Levy, S.V.F., Paterson, M.S. and Thurston, W.P., "Word processing in groups", Jones and Bartlett Publishers, Boston, 1992.
- [12] Gersten, S.M., and Short, H., Small cancellation theory and automatic groups, II, Invent. Math., 105 (1991), 641{662.
- [13] Gersten, S.M., and Short, H., Rational subgroups of biautomatic groups, Annals of Math. 134 125 { 158, 1991.
- [14] Gromov, M., Hyperbolic Groups in "Essays in Group Theory", 75 { 263, Gersten, S.M., ed., Springer Verlag, M.S.R.I. Series vol. 8, 1987.
- [15] Moussong, G., "Hyperbolic Coxeter Groups", Doctoral Thesis, Ohio State University, 1988.
- [16] Papasoglu, P., "Geometric Methods in Group Theory", Doctoral Dissertation, Columbia University, 1993.
- [17] Papasoglu, P., Strongly Geodesically Automatic Groups Are Hyperbolic, University of Warwick preprint, 1994.
- [18] Ronan, M., "Lectures on buildings", Academic Press, New York, 1989.
- [19] Tamashke, O., "Projektive Geometrie, I", Bibliographisches Institut, Mannheim, Wein, Zurich, 1969.