

# A BANACH SUBSPACE OF $L_{1=2}$ WHICH DOES NOT EMBED IN $L_1$ (ISOMETRIC VERSION)

Alexander Koldobsky

**Abstract.** For every  $n \geq 3$ ; we construct an  $n$ -dimensional Banach space which is isometric to a subspace of  $L_{1=2}$  but is not isometric to a subspace of  $L_1$ : The isomorphic version of this problem (posed by S. Kwapien in 1969) is still open. Another example gives a Banach subspace of  $L_{1=4}$  which does not embed isometrically in  $L_{1=2}$ : Note that, from the isomorphic point of view, all the spaces  $L_q$  with  $q < 1$  have the same Banach subspaces.

## 1. Introduction

A well known fact is that the space  $L_1$  is isometric to a subspace of  $L_q$  for every  $q < 1$ : It is natural to ask whether the spaces  $L_q$  with  $q < 1$  contain any Banach space structure not generated by  $L_1$ : This question was first formulated in 1969 by Kwapien [6] in the following form: need every Banach subspace of  $L_0$  be also a subspace of  $L_1$ ? Later the question was mentioned by Maurey [8, Question 124].

In 1970, Nikishin [9] proved that every Banach subspace of  $L_0$  is isomorphic to a subspace of  $L_q$  for every  $q < 1$ : Therefore, if we replace the space  $L_0$  in Kwapien's question by any of the spaces  $L_q$  with  $q < 1$  we get an equivalent question.

Since all the spaces  $L_q$  with  $q < 1$  embed in  $L_0$ ; Nikishin's result also shows that these spaces are all the same from the isomorphic Banach space point of view. Namely, every Banach space which is isomorphic to a subspace of  $L_q$  with  $q < 1$  is also isomorphic to a subspace of  $L_p$  for every other  $p < 1$ :

In this paper we show that the answer to the isometric version of Kwapien's question is negative. For every  $n \geq N$ ;  $n \geq 3$  there exists an  $n$ -dimensional Banach space which is isometric to a subspace of  $L_{1=2}$  but is not isometric to a subspace of  $L_1$ : Using this example it is easy to see that the spaces  $L_q$  with  $q < 1$  may be different from the isometric Banach space point of view. We give, however, a direct example illustrating the difference by constructing a Banach subspace of  $L_{1=4}$  which does not embed isometrically in  $L_{1=2}$ :

The isomorphic version of Kwapien's question is still open. The most recent related result seems to be a theorem of Kalton [2] who proved that a Banach space  $X$  embeds in  $L_1$  if and only if  $\ell_1(X)$  embeds in  $L_0$ :

The isometric version of Kwapien's question can be reformulated in the language of positive definite functions and stable random vectors. In fact, a Banach space  $(X; \|\cdot\|)$  is isometric to a subspace of  $\ell_p$  with  $0 < p \leq 2$  if and only if the function  $\exp(\|x\|^p)$  is positive definite [1]. The main example of this paper gives a norm such that the function  $\exp(\|x\|^{1/2})$  is positive definite, but the function  $\exp(\|x\|)$  is not positive definite. This result is close to problems of Schoenberg's type (see [4]).

## 2. The idea of the construction

Let  $f$  be an infinitely differentiable even function on the unit sphere  $S$  in  $\mathbb{R}^n$ : We spoil the Euclidean norm  $\|x\|_2$  in  $\mathbb{R}^n$  by means of the function  $f$ : Namely, for  $\varepsilon > 0$  consider the function

$$(1) \quad N_\varepsilon(x) = \|x\|_2^{1+\varepsilon} + f\left(\frac{x}{\|x\|_2}\right); \quad x \in \mathbb{R}^n:$$

One can choose  $\varepsilon$  small enough so that  $N_\varepsilon$  is a norm in  $\mathbb{R}^n$ : This follows from a simple one-dimensional consideration: if  $a, b \in \mathbb{R}$ ;  $g$  is a convex function on  $[a, b]$  with  $g'' > 0$  on  $[a, b]$  and  $h \in C^2[a, b]$  then the functions  $g + h$  have positive second derivatives on  $[a, b]$  for sufficiently small  $\varepsilon$ 's, and, hence, are convex on  $[a, b]$ :

Let  $\varepsilon_f = \sup\{\varepsilon > 0 : N_\varepsilon \text{ is a norm in } \mathbb{R}^n \text{ for every } t \in [0, \varepsilon_f]\}$ ; For each  $\varepsilon_f$ ; we denote by  $X_\varepsilon$  the Banach space with the norm  $\|x\|_\varepsilon = N_\varepsilon(x)$ :

Theorem 2 from the paper [5] shows that, for every  $q > 0$  which is not an even integer, there exists a small enough number  $\varepsilon_q$  such that the space  $X_\varepsilon$  is isometric to a subspace of  $L_q$  for every  $t \in [0, \varepsilon_q]$ : This fact was used in [5] to prove that, for every compact subset of  $(0; 1) \setminus \mathbb{N}$  there exists a Banach space different from Hilbert spaces which is isometric to a subspace of  $L_q$  for every  $q \in Q$ :

For  $q \in (0; 1]$ ; let  $\varepsilon_q = \sup\{\varepsilon > 0 : X_\varepsilon \text{ is isometric to a subspace of } L_q \text{ for every } t \in [0, \varepsilon]\}$ : If  $0 < q < p \leq 2$  then the space  $L_p$  is isometric to a subspace of  $L_q$ ; therefore,  $\varepsilon_p \leq \varepsilon_q$ : In particular,  $\varepsilon_1 \leq \varepsilon_q$  for every  $q < 1$ : Clearly,  $\varepsilon_1 = \varepsilon_f$ :

Now we can explain the idea of getting a Banach subspace of  $L_q$  with  $q < 1$  which is not isometric to a subspace of  $L_1$ : Suppose we can find a function  $f$  so that  $\varepsilon_1$  is strictly less than  $\varepsilon_f$ , and also  $\varepsilon_1$  is strictly less than  $\varepsilon_q$ : Then, for every  $2 \in (\varepsilon_1; \min(\varepsilon_f, \varepsilon_q)]$ ; the space  $X_\varepsilon$  is a Banach space with the desired property.

Similarly, for  $q < p < 1$ ; if we manage to find a function  $f$  so that  $\varepsilon_p < \varepsilon_q$  and  $\varepsilon_p < \varepsilon_f$  we get an example of a Banach space which embeds isometrically in  $L_q$  but does not embed in  $L_p$ :

The construction in [5] is based on the use of spherical harmonics and, in general, does not give a chance to calculate the numbers  $\varepsilon_q$  exactly. We are, however, able to choose a function  $f$  for which it is possible to calculate the numbers  $\varepsilon_q$  for certain values of  $q$ : Our calculations do not depend on the results from [5] mentioned above, so the paper [5] only shows a direction for constructing examples.

We shall use one simple characterization of finite dimensional subspaces of  $L_q$ :

**Proposition 1.** Let  $q$  be a positive number which is not an even integer,  $(X; \|\cdot\|_X)$  be an  $n$ -dimensional Banach space, and suppose there exists a continuous function  $b$  on the sphere  $S_n$  in  $R^n$  such that, for every  $x \in R^n$ ,

$$(2) \quad \|x\|_X^q = \int_{S_n} j(x; \cdot)^q b(\cdot) dZ$$

where  $(x; \cdot)$  stands for the scalar product in  $R^n$ :

Then  $X$  is isometric to a subspace of  $L_q$  if and only if  $b$  is a non-negative (not identically zero) function.

**Proof.** If  $b$  is a non-negative function we can assume without loss of generality that  $\int_{S_n} b(\cdot) dZ = 1$ : Choose any measurable (with respect to Lebesgue measure) functions  $f_1, \dots, f_n$  on  $[0; 1]$  having the joint distribution  $b(\cdot) dZ$ : Then, by (2), the operator  $x \mapsto (x_1 f_1, \dots, x_n f_n)$ ;  $x \in R^n$  is an isometry from  $X$  to  $L_q([0; 1])$ :

Conversely, if  $X$  is a subspace of  $L_q([0; 1])$  choose any functions  $f_1, \dots, f_n \in L_q$  which form a basis in  $X$ ; and let  $b$  be the joint distribution of the functions  $f_1, \dots, f_n$  with respect to Lebesgue measure. Then, for every  $x \in R^n$ ,

$$\|x\|_X^q = \left\| \sum_{k=1}^n x_k f_k \right\|_X^q = \int_0^1 \left\| \sum_{k=1}^n x_k f_k(t) \right\|_X^q dt = \int_{R^n} j(x; \cdot)^q d\mu(\cdot) = \int_{S_n} j(x; \cdot)^q d\mu(\cdot)$$

where  $\mu$  is the projection of  $b$  to the sphere. (For every Borel subset  $A$  of  $S_n$ ,

$$\mu(A) = \int_{\{t \in [0; 1] : \sum_{k=1}^n x_k f_k(t) \in A\}} b(t) dt$$

$$\int_{S_n} j(x; \cdot)^q b(\cdot) dZ = \int_{S_n} j(x; \cdot)^q d\mu(\cdot)$$

for every  $x \in R^n$ : Since  $q$  is not an even integer, we can apply the uniqueness theorem for measures on the sphere from [3] to show that  $d\mu(\cdot) = b(\cdot) dZ$  which means that  $b(\cdot) dZ$  is a measure, and the function  $b$  is non-negative.

The representation (2) exists for every smooth enough function on the sphere (see, for example, Theorem 1 from [5]). We are going to choose special smooth norms for which it is possible to calculate the function  $b$  exactly and then check if  $b$  is non-negative. In this way we calculate the numbers  $\alpha_q$  for these norms.

We need the representation (2) for some simple functions on the sphere.

**Lemma 1.** For every  $x = (x_1, \dots, x_n)$  from the unit sphere  $S^n$  in  $R^n$  and every  $q > 0$  we have

$$(3) \quad x_n^2 = \frac{((n+q)=2)}{2^{(n-1)=2} ((q+1)=2)} \int_{S_n} j(x; \cdot)^q \frac{n+q}{q} x_n^2 \frac{1}{q} dZ;$$

Therefore,

$$(4) \quad x_n^2 = \frac{((2n+1)=4)}{2^{(n-1)=2} ((q+1)=2)} \int_{S_n} j(x; \cdot)^{1=2} (2n+1) x_n^2 dZ;$$

and

$$(5) \quad x_n^2 = \frac{((n+1)=2)}{2^{(n-1)=2}} \int_{S_n} j(x; \cdot) j(n+1)_n^2 \, d :$$

Besides,

$$(6) \quad x_n^4 = \frac{((n+1)=2)}{2^{(n-1)=2}} \int_{S_n} j(x; \cdot) j(n+3)(n+1)_n^4 + 6(n+1)_n^2 \, d :$$

Proof. It is a well-known simple fact (see, for example, [7]) that for every  $x \in \mathbb{R}^n$  and every  $k > 0$ ;

$$(7) \quad (x_1^2 + \frac{2}{n} x_1^2)^k = \frac{((n+2k)=2)}{2^{(n-1)=2} ((2k+1)=2)} \int_{S_n} j(x; \cdot)^{2k} \, d :$$

Differentiate both sides of (7) by  $x_n$  twice, and then use the fact that  $x \in S_n$  to get

$$1 + (2k-2)x_n^2 = (2k-1) \frac{((n+2k)=2)}{2^{(n-1)=2} ((2k+1)=2)} \int_{S_n} j(x; \cdot)^{2k-2} \, d :$$

Use (7) with the exponent  $2k-2$  instead of  $2k$  to get

$$(2k-2)x_n^2 = \frac{1}{2^{(n-1)=2}} \int_{S_n} j(x; \cdot)^{2k-2} (2k-1) \frac{((n+2k)=2)}{((2k+1)=2)} \frac{((n+2k-2)=2)}{((2k-1)=2)} \, d :$$

Now use the fact that  $(x+1) = x(x)$  and put  $2k-2 = q$  to get (3).

To prove (6), differentiate both sides of (7) four times by  $x_n$  (remember that  $x_1^2 + \frac{2}{n} x_1^2 \neq 1$ ; do not factor the second and the third derivatives!), and then put  $k = 5/2$  :

$$x_n^4 + 6x_n^2 + 3 = 4 \frac{((n+5)=2)}{S_n} \int_{S_n} j(x; \cdot)^4 \, d :$$

Now use (5), (7) with  $k = 1/2$ ; and the fact that  $(x+1) = x(x)$  to get (6).

### 3. Examples

For every  $\gamma > 0$  define a function  $N$  on  $\mathbb{R}^n$  by

$$N(x) = (x_1^2 + \frac{2}{n} x_1^2)^{1/2} + \frac{x_1^2 + \frac{2}{n} x_1^2 - 2x_n^2}{x_1^2 + \frac{2}{n} x_1^2} ; x \in \mathbb{R}^n :$$

Lemma 2.  $N$  is a convex function if and only if  $\frac{1}{11}$  :

Proof. The function  $N$  is convex if and only if the following function of two variables is convex:

$$g(x; y) = (x^2 + y^2)^{1/2} + \frac{x^2 - 2y^2}{x^2 + y^2} :$$

Calculating the second derivatives of the function  $g$  we get

$$a^2 \frac{\partial^2 g}{\partial x^2} + 2ab \frac{\partial^2 g}{\partial x \partial y} + b^2 \frac{\partial^2 g}{\partial y^2} =$$

$$(x^2 + y^2)^{-7/2} (ay - bx)^2 x^4 (1 - 10 - 11^2) + x^2 y^2 (2 - 2 + 104^2) + y^4 (1 + 8 - 20^2) :$$

The function  $g$  is convex if and only if the latter expression is non-negative for every choice of  $a, b, x, y \in \mathbb{C}$  clearly, it happens if and only if  $\frac{1}{11}$  :

For  $\frac{1}{11}$  denote by  $X$  the Banach space with the norm  $N$  :

Theorem 1. Let  $n \geq 3$ : If

$$n \leq \frac{(18n^2 - 18n)^{1=2} - 3n + 1}{9n^2 - 12n - 1} - \frac{1}{6n - 4}$$

then the Banach space  $X$  is isometric to a subspace of  $L_{1=2}$ ; and, at the same time,  $X$  is not isometric to a subspace of  $L_1$ :

Proof. Let us first prove that the space  $X$  is isometric to a subspace of  $L_{1=2}$  if and only if  $\frac{1}{6n-4}$ : For every  $x \in S_n$ ; use (4) and (7) with  $k = 1=4$  to get the representation (2) (with  $q = 1=2$ ) for the norm  $N$  :

$$N^{1=2}(x) = 1 + (1 - 3x_n^2) = \sum_{j=1}^Z j(x; j)^{1=2} + 7 \sum_{j=1}^Z (6n+3) \binom{2}{n} d_j$$

Clearly, the function  $b(j) = 1 + 7 \sum_{j=1}^Z (6n+3) \binom{2}{n} d_j$  is non-negative on  $S_n$  if and only if  $\frac{1}{6n-4}$ ; and the fact we need follows from Proposition 1.

Let us show that  $X$  is isometric to a subspace of  $L_1$  if and only if  $\frac{1}{11}$ : If  $x \in S_n$  then  $N(x) = 1 + 2(1 - 3x_n^2) + \sum_{j=1}^Z (1 - 6x_n^2 + 9x_n^4)$ : We use (5), (6) and (7) with  $k = 1=2$  to get the representation (2) for  $N(x)$  with  $q = 1$  :

$$N(x) = \sum_{j=1}^Z ((n+1)=2) j(x; j) b\left(\frac{2}{n}\right) d_j$$

where

$$b\left(\frac{2}{n}\right) = (1 + 8 \sum_{j=1}^Z 20^2) + \sum_{j=1}^Z (n+1) (48 \sum_{j=1}^Z 6) - 9 \sum_{j=1}^Z (n+3) (n+1)^2 :$$

The function  $b$  is a quadratic function of  $\frac{2}{n}$  with negative first coefficient. By Proposition 1, the space  $X$  embeds in  $L_1$  if and only if  $b$  is non-negative for every  $\frac{2}{n} \in [0; 1]$ : Clearly, it happens if and only if both numbers  $b(0) = 1 + 8 \sum_{j=1}^Z 20^2$  and  $b(1) = 1 + (2 - 6n) + \sum_{j=1}^Z (1 + 12n - 9n^2)$  are non-negative. Since  $1 + 12n - 9n^2$  is a negative number for every  $n \geq 2$  and we consider only positive numbers; it is clear that the condition is that  $\frac{1}{11}$ : To prove the theorem, it suffices to note that, for every  $n \geq 3$ ;  $\frac{1}{6n-4}$  is less than  $\frac{1}{11}$ ; and  $\frac{1}{n} < \frac{1}{6n-2} < \frac{1}{6n-4}$ :

Remarks. 1. Since every two-dimensional Banach space is isometric to a subspace of  $L_1$  it is impossible to construct a two-dimensional space which the property of Theorem 1. In our example,  $\frac{2}{n} = \frac{1}{11}$  which coincides with the bound from Lemma 2.

2. The author is unable to apply the scheme from Section 2 to every  $q < 1$ ; although it is very likely the same idea works. For instance, getting an example for  $q = 3=4$  is a matter of calculating the eighth derivative of the function  $(x_1^2 + \dots + x_k^2)^{1/4}$ .

Theorem 1 shows also that the spaces  $L_q$  with  $q < 1$  may have different Banach subspaces. The Banach subspaces of  $L_{1=2}$  constructed in Theorem 1 can not be isometric to subspaces of  $L_q$  for all  $q < 1$ : In fact, if a Banach space  $(X; \|\cdot\|)$  is isometric to a subspace of  $L_q$  for every  $q < 1$  then, by a theorem from [1], the function  $\exp(-\|x\|^q)$  is positive definite for every  $q < 1$ : The function  $\exp(-\|x\|)$  is then positive definite as a pointwise limit of positive definite functions, and the space  $X$  (by the same result from [1]) is isometric to a subspace of  $L_1$ :

On the other hand, we can show the difference directly. For every  $\epsilon > 0$  define a function  $N_\epsilon$  on  $\mathbb{R}^3$  by

$$N_\epsilon(x) = (x_1^2 + x_2^2 + x_3^2)^{1=2} \left( 1 + \frac{x_1^2 + x_2^2 - 2x_3^2}{x_1^2 + x_2^2 + x_3^2} \right)^{\frac{1}{4}}; \quad x \in \mathbb{R}^3:$$

The proof of the following theorem is similar to the proofs of Lemma 2 and Theorem 1.

Theorem 2. The function  $N_\epsilon$  is a norm if and only if  $\epsilon > \frac{1}{23}$ : The corresponding Banach space  $X_\epsilon$  is isometric to a subspace of  $L_{1=4}$  if and only if  $\epsilon > \frac{1}{26}$ ; and  $X_\epsilon$  is not isometric to a subspace of  $L_{1=2}$  if  $\epsilon > \frac{1}{28}$ : Thus, for  $\frac{1}{28} < \epsilon < \frac{1}{26}$ ; the space  $X_\epsilon$  is a Banach subspace of  $L_{1=4}$  which does not embed isometrically in  $L_{1=2}$ :

## References

1. J. Bretagnolle, D. Dacunha-Castelle and J. L. Krivine, Lois stables et espaces  $L_p$ , Ann. Inst. H. Poincaré Probab. Statist. 2 (1966), 231{259.
2. N. Kalton, Banach spaces embedding into  $L_0$ , Israel J. Math. 52 (1985), 305{319.
3. M. Kanter, The  $L_p$ -norms of sums of translates of a function, Trans. Amer. Math. Soc. 179 (1973), 35{47.
4. A. Koldobsky, Schoenberg's problem on positive definite functions, Algebra and Analysis 3 (1991), 78{85 (English translation in St. Petersburg Math. J. 3 (1992), 563{570).
5. A. Koldobsky, Common subspaces of  $L_p$ -spaces, Proc. Amer. Math. Soc., to appear.
6. S. Kwapień, Problem 3, Studia Math. 38 (1970), 469.
7. P. Levy, Theory de l'addition de variable aleatoires, Gauthier-Villars, Paris, 1937.
8. B. M. aurey, Theoremes de factorisation pour les operateurs lineaires a valeurs dans les espaces  $L_p$ , Asterisque 11 (1974).
9. E. M. Nikishin, Resonance theorems and superlinear operators, Russian Math. Surveys 25 (1970), 124{187.

Division of Mathematics, Computer Science, and Statistics, University of Texas at San Antonio, San Antonio, TX 78249, U.S.A.  
E-mail address: koldobsk@ringer.cs.utsa.edu